ᓄᓇᕗᑦ ᐅᒪᔨᓕᕆᔨᖅᑎᑦ ᑲᑎᒪᔨᖏᑦ
ᑲᑎᒪᓴᐅᑎᖅ: ᓇᐅᓕᐅᓯᒪᔨᓕᐳᖅ ᑲᑎᒪᓂᖅ RM001-2020
ᐃᖅᓯ 11, 2020 (9:00 ᐅᑦᓯᓴᐅᔪᑦ–6:00 ᐅᵒᓄᓴᐅᔪᑦ)
ᐃᖃᓗᐃᑦ, ᓄᓇᕗᑦ

	ᓇᓴᐅᒃ:	ᑲᒪᒋᔨᖅᖃᖅ:	ᓇᓂᔨᐱᑎᓐᓄᑦ:	ᐅᓂᒃᑲᖅᑐᖅ	ᐱᕕᖅᖃᖅ
9:00 AM - 9:02 AM	1	ᒪᑐᐃᖅᑕᖅ ᑲᑎᒪᓂᖅ		ᐃᖅᔨᕿᐅᑕᖅ	2 ᒥᓇᖅ
9:02 AM - 9:03 AM	2	ᓇᓗᓇᐃᖅᔪᒍᑦ ᐊᑐᑐᐊᓗᐊᑐᖅᑲᖅᑕᑦ		ᐃᖅᔨᕿᐅᑕᖅ	1 ᒥᓇᖅ
9:03 AM - 9:05 AM	3	ᑲᑎᒪᓴᐅᑎᖅ: ᖁᒥᓱᑭᖅᑕ ᐊᓚᒐᓗ ᐊᕐᕈᖅᑕᑦ	1	ᐃᖅᔨᕿᐅᑕᖅ	2 ᒥᓇᖅ
		ᐱᓕᓴᕐᓂᖅᑕ ᐊᕿᑎᓕᓄᓵᑦ–ᑲᕿᒪᓐᔪᑦ ᓄᓇᕗᑦ (GN-DOE: ᐱᓴᐊᑕᐅᔪᑦ/ᐃᓱᒪᓕᐅᑭᒍᑦ			
9:05 AM - 10:00 AM	3	ᖁᑭᖅᑕᒐᓴᒥ ᑐᓴᑐᓄ ᐊᐅᓪᓚᕐᓂᕐᒧᑦ ᕿᖏᓗᒍᑦ [ᐃᓱᒪᓕᐅᑭᑦ]	2	ᑲᕿᒪᓐᔪᑦ ᓄᓇᕗᑦ	55 ᒥᓇᖅ
10:00 AM - 10:15 AM		ᖁᑭᖅᑐᑦ			15 ᒥᓇᖅ
10:15 AM - 11:00 AM	4	ᖁᓄᓇᑕᖅᑐᖅᑎᑦ ᐊᕿᑦᓯᓕᓵᓐᑦ ᐊᔪᓂᐅᔨᒥᓄ ᐊᕿᕈᑐᓄ ᖁᓄᐅᓴᒪᒐᓱᐊᖅᑐᖅᓄᓇᑦ ᐱᑕᖅᖁᕐᑕᖅ [ᑐᓴᕋᕐᕋᑦ]	3	ᑲᕿᒪᓐᔪᑦ ᓄᓇᕗᑦ	45 ᒥᓇᖅ
		ᐃᒪᖃᐅᑕᓕᓵᓐᑦ ᑲᓇᑕᒥ (ᐃᒪᖃᐅᑕᓕᓵᓐᑦ): ᐱᓴᐊᑕᐅᔪᑦ/ᐃᓱᒪᓕᐅᑭᒍᑦ			
11:00 AM - 11:45 AM	5	ᑲᒪᒋᔨᓄᑦ ᕿᖏᓗᒍᑦ ᐃᕐᓂᓐᑲ, ᐅᐊᓐᓇᖅᓯᓂ ᐊᑦᓭᓐᑎᑲ ᐱᑕᖅᖁᔭᑐᓐᑲ, ᑲᓇᑕᒥ [ᐃᓱᒪᓕᐅᑭᑦ]	4	ᑲᕿᒪᓐᔪᑦ ᓄᓇᕗᑦ	45 ᒥᓇᖅ
11:45 AM - 1:00 PM		ᐅᑦᓗᑭᒥᓴᓇᖅ			1 ᐃᑲᓴᖅ & 15 ᒥᓇᖅ
1:00 PM - 1:30 PM	6	ᑲᑎᑦᑐᑦ ᐊᐳᖅᑎᑦ ᐱᔾᐅᓂᒃᑯᑦ ᖁᓄᐃᑦᓗᐊᓂᖅᑎᑦ ᒥᑭᔨᑕᐅᓂᖅᑎᑦ ᐊᓚᒐᓗ ᐊᕐᕈᔨᑕᐅᓂᖅᑎᑦ ᑭᖑᑦᑕᐃᑦ ᐃᓄ 2020-2021 ᐊᓴᔭᒥ ᐱᖓᓱᓇᖓᓄ ᐊᓚᒐᓗ ᐅᓇᓱᓇᖓᓄ ᖁᓄᐅᓴᓴᕆᐊᐅᓪᓚᑦ ᓇᓕᖅᑭᓱᓄ [ᐃᓱᒪᓕᐅᑭᑦ]	5	ᐃᒪᖃᐅᑕᓕᓵᓐᑦ ᑲᓇᑕᒥ	30 ᒥᓇᖅ

1:30 PM - 2:00 PM	7	ᐃᓱᒪᓕᐅᕐᓯᑎᑦ ᐊᓚᓗᓐ ᐊᑐᓕᔪᖅᑎᑦ ᐅᑭᐅᖅᑕᕐᑐᒥ ᐅᒃᓴᓂᖅ ᐱᖃᕐᓇᐅᔪᖅᓴᐅᔪᓐᓇᖅᑐᑦ ᑭᓕᓚᕐᓂᖅ ᒥᓂᔅᓴᓂᖅ ᒥᓚᓗᓐ ᐊᖏᔪᑦᑕᐅᓂᕐᓂᖅ ᑭᖓᔪᖅᓯᐊᕐᓂᖅ ᐃᓕᐊᓂ ᐊᓚᓗᓐ ᖃᐅᓂᑯᖓᓂ ᓄᓇᑦᓯᒪᔪᓂ ᓇᓕᓐᓂᑕ [ᐃᓱᒪᓕᐅᑉᑐᑦ]	6	ᐃᒪᖏᐅᑕᓕᓚᔨᓂ ᐅᓇᑎᑕ	30 ᒥᓇᕐ
2:00 PM - 2:30 PM	8	ᐱᖏᓂᖅ ᐱᖃᑎᖅᑭᑲᖃᐅᓂᖅ ᓇᓗᓇᐃᖏᑕᓂᖅ ᒥᑭᔪᑕᐅᓂᖏᓂᖅ (*Pandalus borealis)* ᓄᓇᕗᑦ ᐊᓚᓗᓐ ᓄᓇᕕᖅ ᐅᓇᓱᓐᓂᖅ [ᐃᓱᒪᓕᐅᑉᑐᑦ]	7	ᐃᒪᖏᐅᑕᓕᓚᔨᓂ ᐅᓇᑎᑕ	30 ᒥᓇᕐ
2:30 PM - 3:00 PM	9	ᐅᑎᕐᑐᑦ ᐊᔪᖓᑕᑦ ᐱᖃᐅᓂᖏᖅ ᖃᐅᓄᑦᓗᐊᓂᖓᓕᑦ ᒥᑭᔪᑕᐅᓂᖏᓂᑦ ᑭᖓᔪᓂᐊᓂᑦ ᒪᓂ 2020 ᐊᓱᕐᔪᒥ ᑭᖓᔪᓂᓕᓂᐊᕐᓇᖅᑎᒥ 0 [ᐃᓱᒪᓕᐅᑉᑐᑦ]	8	ᐃᒪᖏᐅᑕᓕᓚᔨᓂ ᐅᓇᑎᑕ	30 ᒥᓇᕐ
3:00 PM - 3:15 PM		**ᖃᐅᕐᑐᑦ**			15 ᒥᓇᕐ
3:15 PM - 3:45 PM	10	ᐃᓕᓴᐅᑐᕐᐊᕐᓇᖅᑎᑦ ᐃᓕᓴᐅᐱᓐᓂᐊᕐᓚᐃᑦ [ᑐᓴᕋᕐᓴᖅ]	9	ᐃᒪᖏᐅᑕᓕᓚᔨᓂ ᐅᓇᑎᑕ	30 ᒥᓇᕐ
3:45 PM - 4:15 PM	11	ᐃᒪᖏᐅᑕᓕᓚᔨᓂ ᐅᓇᑎᑕ – ᖃᓄᐃᑕᓂᓂᖓᓂ [ᑐᓴᕋᕐᓴᖅ]	10	ᐃᒪᖏᐅᑕᓕᓚᔨᓂ ᐅᓇᑎᑕ	30 ᒥᓇᕐ
		ᐊᐊᑎᑕᓕᓚᔨᓂ ᐊᓚᓗᓐ ᓱᑕᐅᑦ ᐊᓱᓴᓂᓂᖅᓯᓂᓂᖅ ᑲᒪᔨᓂ ᐅᓇᑎᑕ: **ᐱᔪᑎᑕᐅᔪᑦ/ᐃᓱᒪᓕᐅᑉᑐᑦ**			
4:15 PM - 5:00 PM	13	ᐃᓂᓂᐅᓂᖏᓕ ᓇᓕᖓᑭᓂ ᐊᐅᓕᕐᓱᖃᑎᑎᑦ ᑲᑎᒪᓴᕆᓪᓕᑦ: ᐊᐅᓕᕐᓱᓂᕐᒧᑦ ᕿᖏᓇᒧᑦ ᖃᖅᖄᓱᐅᔨᑉᑐᖅᑎ ᐊᓚᓗᓐ ᐃᓂᖃᑎᐊᕐᓇᖅᑎ ᑎᓕᐅᐊᕐᓇᖅᐅᓕᑎ [ᐃᓱᒪᓕᐅᑉᑐᑦ]	11	ᐊᐊᑎᑕᓕᓚᔨᓂ ᐅᓇᑎᑕ	45 ᒥᓇᕐ
5:00 PM - 5:30 PM	14	ᖃᓄᐃᑕᓂᓂᖓᓂ ᑐᓴᕋᕐᓴᖅᖃᓄᐃᓕᐊᕐᓂᓂᑦ ᓄᓇᕗᑦᑎ ᐱᓪᓗᑎᑦ ᐃᓱᒪᑎᑉᓴᑦ ᑎᑎᕋᕐᑕᐅᓂᖓᓂᑦ ᓇᓪᓴᑦᑐᕐᓂᖓᓂᑦ ᑐᓗᓚᐊᓇᕐᑐᑎᑦᑐᑦ ᐊᒋᔪᑦ ᓚᐊᒪᓴᖅᑲᓂᕐᒧᑎᓪᓗ ᐅᒪᓕᕐᓂᓕ ᐊᑦᑕᓇᕐᑐᑎᑦᑐᓗᓂ ᒪᓕᑦᑎᓴᐊᕐᓇᖅ (ᐊᑦᑕᓇᕐᑐᑎᑦᑐᓗᓕᖅᑲᓪᓚᕆᐊᑐᑦ) [ᑐᓴᕋᕐᓴᖅ]	12	ᐊᐊᑎᑕᓕᓚᔨᓂ ᐅᓇᑎᑕ	30 ᒥᓇᕐ
		ᖃᑎᖅᑎᐅᓂ ᐅᒪᓕᓕᓚᔨᓂ ᑲᑎᒪᔨᓂᖓᓂ: ᐱᔪᑎᑕᐅᔪᑦ/ᐃᓱᒪᓕᐅᑉᑐᑦ			
5:30 PM - 6:00 PM	15	ᑐᑭᕋᐅᑦ ᐊᒥᓱᑖᓚᐊᕐᑐᓂᑦ ᐊᐳᓪᓂᑎ ᑕᓀᐊᖅᓴᓂᑎ ᐊᖏᓇᓱᓕᑐᓂᓂ ᓂᐲᖏᓐᓂᑕ ᖃᑎᖅᑎᐅᓂ ᐅᐊᖏᓐᓂ 15 ᐅᐃᖏᓗ 20 ᓂᐲᖏᓐᓂᑦ [ᐃᓱᒪᓕᐅᑉᑐᑦ]	13	ᖃᑎᖅᑎᐅᓂ ᑲᑎᒪᔨᓂ	30 ᒥᓇᕐ
	16	ᓄᖅᑲᖅᑐᖅ RM001-2020			

ᐅᔨᓇᖏᑦ ᑐᓂᓯᐅᔾᔪᖅ

ᓄᓇᕗᒻᒥ ᐅᒪᔨᓕᕆᔨᖅᑎᑦᑕᐃᓐᓇᓕᒫᑦ ᑲᑎᒪᔨᖏᑦᓱᓄᑦ

ᐅᔨᓇᖏᑦ

ᑐᓴᐅᒪᔭᕐᓂᖅᓴᓂᑦ: **ᐃᓕᓴᕐᕕᖏᓐᓇᖅᑐᑦ: X**

ᐱᔭᕆᑦ: ᖃᑭᖅᑕᓗᓪᓕᒥᑦ ᑐᑦᑐᑦ ᐊᐅᓪᓚᑕᐅᓂᖏᓱᓄᑦ ᐸᖅᓇᑲᑕᐅᓕᒫᖅ

ᐱᓕᕆᕕᖏᓐᓇᓕᔪᑦ:

- ᑎᓕᐸᐱᓇ 2014–ᒥᑦ ᖃᑭᖅᑕᓗᓪᓕᒥᑦ ᑐᑦᑐᑕᐅᔨᓇᓴᖅᑐᑦ ᐸᖅᓇᑲᑕᐅᓪᓚᑕᐅᓴᕐᓕᐅᖅ ᑲᑐᑦᓴᐅᓪᓗᓂᓗ ᐊᖏᓇᓕᑦᑐᑕᓪᓚᖅᑎᓱᓄ ᐊᑦᑐᖅᑕᐅᓕᒫᔪᓂ, ᖃᑭᖅᑕᓗᓪᓕᒥᑦ ᐅᒪᔨᓕᕆᔨᖅᑎᑦᑕᐃᓐᓇᓕᒫᑦ ᑲᑎᒪᔨᖏᑦᓱᓄᑦ, ᓄᓇᕗᑦ ᑐᖓᓚᐃᔭᑦᓂᓗ. (ᓄᓇᕗᑦ ᑐᖓᓚᐃᔭᑦ), ᐊᓪᒪ ᑲᓇᑕᒥᑦ ᒥᖅᖢᐃᐸᖅᓯᐊᕐᓕᓚᐅᔭᐃᑦ.

- ᖃᑭᖅᑕᓗᓪᓕᒥ ᑐᑦᑐᕕᓚᐅᔭᐃᑦ ᑲᑎᒪᔨᖏᑦ ᑐᓂᓴᐅᓕᐅᖅᑐᑦ ᓄᓇᕗᑦᑐᑦ ᐅᒪᔨᓕᕆᔨᖅᑎᑦᑕᐃᓐᓇᓕᒫᑦ ᑲᑎᒪᔨᖏᑦᓱᓄᑦ ᐊᖅᑭᖅᑕᐅᔪᓂᑦ ᐃᓚᒌ 2016-ᒥᑦ. ᑲᑎᒪᔨᖏᑦ ᑐᑦᓯᕋᓕᐅᖅᓯᒪᔪᑦ ᐊᐅᑎᓕᕆᓚᐅᔭᐃᓐᓇᓱᓄᑦ ᖃᐅᔨᓴᓕᓯᐅᖃᖅᑲᓐᓂᕐᓗᑎᑦ ᑎᑎᕋᖅᑕᐅᔾᔨᓐᓇᐅᑎᓪᓗᒍ ᐸᖅᓇᑲᑕᒥᓐᓄᑦ ᓯᓚᐅᑉᐳᓐᓂᒃᑯᑦ ᖃᐅᔨᓴᓕᓯᐅᖃᖅᑎᓪᓗᑎᑦ.

- ᐊᖅᑭᑦᐊᐅᖅᑕᐅᓯᒪᔪᑦ ᐊᐅᓚᑦᑎᓂᖓᓐᓇᒧᑦ ᐸᖓᓇᐅᑎᑦ ᓇᑦᓯᐅᑦᔨᓚᐅᔭᖅᑐᑦ ᓯᑎᐱᓇ 2019–ᒥᑦ ᐃᓚᖃᓚᐅᔭᖅᓴᓂᓗ ᖃᐅᔨᓴᕿᑎᐱᓕᓂᖓᓐᓂᑦ ᑲᑎᒪᔨᖏᑦᓯᓯᒪᔪᑦ ᓇᐃᔭᕐᑎᑎᓂᖓᓐᓂᑦ ᐱᐊᓂᑦᑕᐅᓕᒫᑦ 2015-2018. ᑲᑎᒪᔨᖏᑦᓯᓯᒪᔪᑦ ᓇᐃᔭᐃᑦᔨᓚᑎᑦ ᐊᑐᑦᑕᐅᓚᐅᔭᖅᑐᑦ ᖃᐅᔨᓴᕿᑕᐅᓪᓚᑎᑲ ᓄᑦᓱᓄᑦ ᖃᓄᐃᓐᑐᓂᖓᓪᓚᓗ ᐊᖏᑎᓪᓚᓗ ᐊᑦᓇᓪᓚᓗ. ᐊᖅᑭᑦᐊᐅᖅᓯᒪᔭᐃᑦ ᓇᓗᓇᐃᔨᓯᓚᐅᕆᓂᖅ ᑐᓗᖅᑕᕿᑕᐅᔨᓴᓂᑦ ᓇᓗᓇᓗᓇᐃᔪᐃᓂᖓᓐᓂᑦ ᐊᒥᓯᓂᓪᓚᐊᖅᓂᖓᓐᓂᑦ ᒪᑕᒃᓴᑎᑲ ᑲᑎᒪᔨᖏᑦᓯᓯᒪᔪᑦ ᓇᐃᔭᐃᑦᔨᓚᑎᑦᑎᓂᖓᓐᓂᑦ.

- ᓯᓂᓪᓕ ᐊᖅᑭᑦᐊᐅᖅᓯᒪᔪᑦ ᑎᑎᕋᖅᑕᐅᑎᐅᕐᖓᖅᑐᑦ ᐊᐅᓚᑦᑎᓂᖓᓐᓇᒧᑦ ᐸᖓᓇᐅᑎ ᐃᓚᓕᐅᔾᔨᓚᐅᖅᑐᑦ ᑕᓪᓕᒪᓂᑦ ᐊᖏᓇᓕᖓᓐᓇᒧᑦ ᐊᐅᓚᑦᑎᓂᖓᓐᓇᒧᑦ ᐊᖏᓇᓕᖓᓐᓇᓱᓕ ᐊᐅᓚᑦᑎᓂᖓᓐᓇᒧᑦ ᖃᐅᔨᓴᐅᑎᓕᐅᖅᑐᑦᑐᖅᑐᑦ ᖃᐅᔨᓴᕿᑕᐅᓪᓚᑎᑲᓂᖅ ᖃᑭᖅᑕᓗᓪᓕᒥᑦ ᒪᓕᒪᓂᑲᓂᖅ ᐊᒥᓱᓕᑕᐅᓪᓚᑎᑲᓂᖅ ᐊᑐᖅᑐᑦ ᐊᒥᓱᓕᑕᐅᓪᓚᑎᑲᓂᖅ ᐊᑐᖅᑐᑦ. ᑕᒪᓐᓇᐊ ᐃᓗᓕᓂᖓᓐᓂᑦ ᐱᓴᖅᑕᐅᓕᐅᖅᑐᑦ ᐸᖓᓇᐅᑎᑦ ᒪᓕᒪᓂᑲᓂᖅ ᑐᓴᕐᓴᐅᓂᖅ ᐊᒥᓱᓕᑕᐅᓪᓚᑎᑲᓂᖅ.

ᐃᓐᓇ ᖃᐅᔨᓴᕿᑎᓂᖓᓐᓂ:

- ᑲᑎᒪᔨᐅᖅᑎᓪᓚᓂᑦ ᔪᑎᐱᓇ 2019–ᒥᑦ ᑲᑎᒪᓂᑐᐃᓐᓇᒥᑦ, ᓄᓇᕗᒻᒥ ᐅᒪᔨᓕᕆᔨᖅᑎᑦᑕᐃᓐᓇᓕᒫᑦ ᑲᑎᒪᔨᖏᑦᓱᓄᑦ ᑐᓴᕐᓴᐅᓂᖅᓴᓂᑦ ᖃᑭᖅᑕᓗᓪᓕᒥᑦ ᐅᒪᔨᓕᕆᔨᖅᑎᑦᑕᐃᓐᓇᓕᒫᑦ ᑲᑎᒪᔨᖏᑦᓱᓄᑦ ᑐᓴᕐᓴᐅᓂᖅᓴᓂᑦ ᑐᖓᓚᐃᔭᑦᓂᓗ ᐊᖏᓇᓕᑦᑐᑕᓪᓚᖅᑎᓱᓄᑦ, ᐃᓚᓕᐅᔾᔨᓚᐅᖅᑐᑦ ᐊᐅᓚᑦᑎᓂᖓᓐᓇᒧᑦ ᐸᖅᓇᑲᑕᐅᓪᓚᑎᑲᓂᖅ ᐊᐅᓚᑦᑎᓂᖓᓐᓇᒧᑦ, ᐊᖏᓇᓕᑦᑐᑕᓪᓚᖅᑎᓱᓄᑦ ᐊᐅᓚᑦᑎᓂᖓᓐᓇᒧᑦ ᐸᖅᓇᑲᑕᐅᓪᓚᑎᑲᓂᖅ, ᐊᖏᓇᓕᑦᑐᑕᓪᓚᖅᑎᓱᓄᑦ ᑎᑎᕋᖅᑕᐅᓪᓚᑎᑲᓂᖅ ᐸᖅᓇᑲᑕᐅᓪᓚᑎᑲᓂᖅ, ᐊᖏᓇᓕᑦᑐᑕᓪᓚᖅᑎᓱᓄᑦ ᑎᑎᕋᖅᑕᐅᓪᓚᑎᑲᓂᖅ ᐸᖅᓇᑲᑕᐅᓪᓚᑎᑲᓂᖅ ᐊᑐᖅᑐᑦ ᐊᐅᓚᑦᑎᓂᖓᓐᓇᒧᑦ ᐊᖏᓇᓕᑦᑐᑕᓪᓚᖅᑎᓱᓄᑦ ᐊᐅᓚᑦᑎᓂᖓᓐᓇᒧᑦ, ᐊᐅᓚᑦᑎᓂᖓᓐᓇᒧᑦ ᐊᖏᓇᓕᑦᑐᑕᓪᓚᖅᑎᓱᓄᑦ

- ᐱᑕᖅᓗᐊᖏᓐᓂᖅᓱᓚᓄᑦ ᐊᐅᓪᓚᑦᑎᓂᖅᓴᒻᒪᑦ ᖃᓄᐃᓕᓚᐅᕕᑎᐅᓴᕐᓇᐃᑦ ᐊᑦᑕᖅᓴᐃᐅᑎᓄᑦ ᖁᑭᖅᑕᓗᓂᑦ ᑐᑐᖅᑕᓂᑦ.
- ᐊᐅᓪᓚᑦᑎᓂᖅᓴᒻᒪᑦ ᐸᖅᓇᐅᑎ ᓄᑕᖅᐳᓚᐊᖅᓭᒪᖏᑦᑐᑦ ᐃᓪᓚᖃᖅᓂᐅᐊᕐᓗᓂ ᐊᓯᔨᖅᑕᐅᕕᔪᓂᑦ ᖁᑭᖅᑕᓗᓂᑦ ᑐᑐᑕᐅᕕᔪᓇᖅᑐᑦ ᑭᓯᓕᕐᓂᑦ.

- ᒥᓂᔾᑕ ᐊᕿᑎᓪᓚᕆᔨᖅᓄᓄᑦ ᑎᑎᕋᓚᐅᖅᑐᑦ ᓄᓇᕗᒻᒥ ᐅᓚᔪᓪᓚᕆᔨᕋᐊᖅᓄᑦ ᑲᑎᒪᔨᖅᑎᑦᑎᓄᑦ ᐅᖃᓕᒫᖅ 6, 2020 ᖃᓚᒋᔭᖅᖢᓂᐅᕐᓗᑎᒃ ᓄᓇᕗᒻᒥ ᐅᓚᔪᓪᓚᕆᔨᕋᐊᖅᓄᑦ ᖃᓄᓚᔨᖅᑎᑦᑎᓂᑦ ᐃᓕᓴᕆᑎᖅᑎᓂᑦ ᐱᓕᕆᔨᑎᖅᑎᓂᑦ ᐊᐅᓚᑦᑎᓂᖅᓴᒻᒪᑦ ᖃᓄᐃᓕᓚᐅᕕᑐᐊᕐᓂᑦ ᐃᓚᒌᑦ ᐳᑐᐃᐅᕕᓂᑦ:

  - “ᐱᓕᕆᓪᓚᖅᓂᑦ ᐊᐅᓪᓚᑦᑎᓂᖅᓴᒻᒪᑦ ᖃᓄᐃᓕᓚᐅᕕᔨᐅᓕᐊᕐᓂᑦ ᓕᓐᓂᑦ ᖃᓄᓪᓚᒥᑦ ᐊᖏᔪᖅᑕᐅᕕᔪᓇᖅᑐᑦ ᕿᓐᓂᐊᕐᓂᖅᓯᖅᑎᑦ, ᒪᓕᒃᓴᓂᖅ ᖃᐅᔨᓴᓕᕆᓂᖅ, ᐃᓄᐃᑦ ᖃᐅᔨᒪᔭᕐᔪᖅᑐᖅ, ᑐᓴᕆᓕᓕᓐᓂᖅᓱᓚᐅᕕᑎᐅᓴᕐᓂᑦ, ᐃᓕᓴᕐᐅᑎᓕᒪᓕᐅᕕᖏᓐᓂᑦ ᐊᐅᓪᓚᑦᑎᓂᖅᓴᒻᒪᑦ ᐸᖅᓇᐅᑎᑦ. ᑕᐃᒪᐃᑦᑐᑐ ᐅᓚᔪᓪᓚᖅᓴᒻᒪᑦ ᐊᐅᓪᓚᑦᑎᓂᖅᓴᒻᒪᑦ ᓄᓇᕗᒻᒥᑦ ᐱᕐᕆᐊᖃᖅᖃᐸᖅᓯᒪᒋᑦ ᓄᓇᕗᒻᒥ ᐅᓚᔪᓪᓚᕆᔨᕋᐊᖅᓄᑦ ᖃᓄᐃᓕᓚᐅᕕᑐᐊᕐᓂᑦ ᐊᐅᓚᑦᑎᓂᖅᓴᒻᒪᑦ ᐃᓗᖃᖅᑐᓂᑦ ᐊᐅᓚᑦᑎᓂᖅᓴᒻᒪᑦ ᐊᐅᓚᑦᑎᓂᖅᓴᒻᒪᑦ. ᐃᓂᓪᑎᓐᓂᐳᒍᑦ ᐱᓕᕆᓪᓚᖅᓴᒻᒪᑦ ᐊᐅᓪᓚᑦᑎᓂᖅᓴᒻᒪᑦ ᖃᓄᐃᓕᓚᐅᕕᑎᐅᓴᕐᓂᑦ ᖃᓄᐃᓕᓚᐅᕕᑐᐊᕐᓂᑦ ᐸᖅᓇᐅᑎᓯᒪᓕᐅᕕᖏᓐᓂᑦ ᐊᐅᓚᑦᑎᓂᖅᓴᒻᒪᑦ ᖃᓄᐃᓕᓚᐅᕕᑐᐊᕐᓂᑦ ᖃᓄᐃᓕᓚᐅᕕᑎᐅᓴᕐᓂᑦ ᐊᐅᓚᑦᑎᓂᖅᓴᒻᒪᑦ ᐸᖅᓇᐅᑎᑎᒋᐊᖅᐸᑦ, ᐅᖃᓕᓂᖅᓴᐃᓚᐅᕕᑐᐊᕐᓂᑦ ᐊᐅᓚᑦᑎᓂᖅᓴᒻᒪᑦ ᐊᐅᓚᑦᑎᓂᖅᓴᒻᒪᑦ ᐃᓚᒌᑦ ᐱᓕᕆᓪᓚᖅᓴᒻᒪᑦ ᖃᓄᐃᓕᓚᐅᕕᑐᐊᕐᓂᑦ ᐊᐅᓚᑦᑎᓂᖅᓴᒻᒪ

ᑲᓐᓄᐊᕆᒥᑦ ᖃᐅᔨᓴᓂᖅᓂᕐᓂᑦ. ᐃᖅᑲᓇᐅᑎᒃᓯᑦ ᐃᖅᑲᐃᑎᑦᑎᓪᓗᐅᖅᓴᑦ ᓄᓇᕐᔪᐊᓇ 26, 2018–ᒥᑦ ᑐᓴᕋᐃᓗᓂᑦ ᑲᑎᒪᓪᓗᐅᖅᓯᓂᖅᓂᓂᖅ ᐅᖃᐅᔨᓴᕐᓂᖅ.

- ᖃᐅᖅᑕᓗᓂᒃᑯᑦ ᖃᐅᔨᓴᖅᑎᑦ ᖃᐅᖅᑕᑦᔪᓂᑦ ᓄᓇᓕᓂᒃᓄᐊᓪᓗᐅᖅᓴᑦ, ᑐᓴᑐᕐᔪᓂᑦ ᖃᐅᖅᑕᑦᔪᓂᑦ ᑐᓴᑐᓂᒃᓂᑦ, ᐅᓄᔪᒪᓕᕐᓂᑦ 7–18, 2019–ᒥᑦ, ᐅᖃᐅᔨᓴᖃᓕᕆᓂᖅᑐᓂᒃ ᑲᑎᒪᔨᐅᓴᑐᓂᑦ ᓇᐃᔭᖅᓯᒪᔨᓂᑦ 2015-2018, ᖃᐅᖅᑕᑦᔪᓂᑦ ᑐᓴᑐᓂᒃᓂᑦ ᐊᐅᓪᓕᑎᓂᒃᓴᑦ ᓯᓚᐅᑉᐳᓂᖅ, ᖃᓄᐃᓕᑦᑐᓂᒃᓯᑦ ᐊᖏᔪᓕᐊᑦᑐᓂᑦ ᔪᓚᐅᑦᑎᓂᒃᓴᓂᓪᓗ ᖃᐅᔨᓴᕐᓗᑎᒃᓴᑦ.

- ᖃᐅᔨᓴᓇᔪᓂᖃᓕᐅᖅᓴᑦ ᐊᖏᓇᔪᑐᑕᓕᕆᔨᒃᑯᑦ ᑭᒡᒋᐊᒡᒋᑦ, ᖃᐅᖅᑕᒡᔾᔨᐊᖏᑦ, ᓯᓚᐅᖅᑐᑦ, ᐃᖃᓗᓐᓂᑦ, ᑭᖑᓂᓂᑦ, ᓴᓂᕆᔨᖏᑦ, ᐃᒡᓗᓕᖏᑦ, ᐅᑦᐱᐊᖅᖏᑦ ᐊᒡᒪᒪ ᒥᑦᑎᒪᒡᕕᖏᑦ. ᑭᔪᐊᓂᑦᑕᐅᖅ, ᖃᐅᔨᓴᓇᔪᓂᖃᖅᑎᓪᓗᐅᖅᓴᑦ, ᐊᑕᖅᔪᒪᔪᖅ ᒥᑦᑐᓵᓇᐅᖅᑎᒡᓴᑦ ᔪᓕ ᐱᓪᓗᒍ ᑲᖏᖅᓱᓕᕆᓂᖅᓂᓂᖅ. ᖃᐅᔨᓴᓇᔪᓂᖃᖅᑕᐃᓵᓇᐅᓕᖅᓴᑦ ᑲᖏᖅᓱᓕᕆᓂᖅᓂᓂᖅ ᐊᒡᒪᒪᓕᐊᑦᑐᓂᑦ ᓴᓇᔨᓴᒥᒃ ᐊᒡᒪᒪᓕᐊᑦᑐᓂᑦ ᐃᓚᒋᔭᐅᓂᖅ ᓴᓇᔨᓴᒥᒃ 27, 2019–ᒥᑦ.

- ᐅᖃᐅᔨᓴᖏᓐᓄᑦᑎᓂᒃᑯᑦ ᓂᕆᐊᖅᑕᐅᔪᓕᒥᑦ ᐊᐅᓚᓵᕐᔪᖏᓐᓄᑦᑐᓂᒃᓴᑦ ᐸᖓᓂᕐᒥᓂ ᐅᓴᓀᓐᓂᑦ ᖃᐅᖅᑕᑦᔪᓂᑦ ᐱᓪᓗᐅᖅᓴᑦ, ᐊᓂᒍᑎᓕᓪᓗᐅᖅᓯᒪᔪᓂᑦ ᐊᐱᓚᕆᐅᓚᐅᖅᓴᖅᓯᒪᓂᒃᑯᑦ ᐅᖃᐅᔨᓴᖏᓐᓄᑦᑐᓂᒃᓴᑦ ᐃᓚᒋᔭᐅᓂᖅ 15, 2019–ᒥᑦ.

- ᖃᐅᖅᑕᓂ ᐅᒪᔨᓕᑎᓪᓗᒃᑯᑦ ᑲᑎᒪᔨᖏᑐᓂ ᐊᒍᑎᓕᑎᓪᓗᒃᑯᑦ ᑭᖓᔪᓕᒥᑦ ᑎᑎᕋᖅᑕᓕᐅᖅᓴᑦ ᑐᓴᐅᕋᖅᓴᓂᑦ ᐅᖃᐅᔨᓴᖏᓐᓄᑦᑐᓂᒃᓴᑦ ᐱᔨᓴᓕᓂᑦ ᐅᖃᖅᓯᓕᕆᓂᖅᑕᐅᖅᑐᓴᖅᓯᓂᓕᓂᑦ. ᐊᒍᑎᓕᑎᓪᓗᒃᑯᑦ ᐊᖏᔪᓕᐊᑦᑐᓂᑦ ᐅᖃᐅᔨᓴᖏᓐᓄᑦᑐᓂᒃᓴᑦ ᐱᔨᓴᓕᓂᑦ ᐅᖃᖅᓯᓕᕆᓂᖅᑕᐅᖅᑐᓴᖅᓯᓂᓕᓂᑦ ᐅᖃᐅᔨᓴᖏᓐᓄᑦᑐᓂᒃᓴᑦ ᐱᔨᓴᓕᓂᑦ ᐅᖃᐅᔨᓴᖏᓐᓄᑦᑐᓂᒃᓴᑦ ᐅᖃᐅᔨᓴᖏᓐᓄᑦᑐᓂᒃᓴᑦ ᐱᔨᓴᓕᓂ

ᓄᓇᕗᑦ ᑲᓇᒪᓐᓚ: ᐊᓇᑎᓕᓵᔨᖏᑦ

ᖃᑭᖅᑕᓗᒍ ᑐᑦᑐᓕᓵᓂᖅᑎᑦ ᐊᐅᓪᓕᔾᔪᔭᑎᑦᕼᓂᑦ ᐸᖅᓇᐅᑎᑦᕼᖅ

ᐱᓕᓵᖅᑎᒋᔭᓂᖅ ᓇᓗᓇᐃᖅᓱᒪᖅᓯᓗᒍᑦ ᖃᑭᖅᑕᓗᒍ ᑐᑦᑐᓂᑦ

ᐊᖏᔪᓇᔭᓂᖅ ᑲᔪᔾᔪᑕᒪᔭᓇᖅᓯᓗᒍ

GN DOE

ᑎᒃᐱᐃᓚ 2019

ᓄᐃᓇᓕᔪᑎᑦ

ᖃᑭᖅᑕᓗᒍ ᑐᑦᑐᓕᓚᓂᖅᑎᑦ ᐊᐅᓪᓕᓯᔪᓂᕋᓂᑦ ᕿᖏᐅᑎᓕᐅᕈᒪᓕᓪᓕᐅᖅᑐᑦ ᐱᑕᖅᖃᔨᓪᓚᑎᑦ ᐊᑐᐊᒐᕋᓂᑦ ᐊᓯᒪ ᐊᑐᓕᖃᔨᔪᓂᕋᓂᑦ ᐃᔨᒪᕋᔨᔾᑎᓄᓪᓗᑦ ᐊᑐᖅᑕᐅᖅᖃᓂᒃ ᐱᔨᔪᑎᑦᓚᓪᓕᑦ ᐊᐅᓪᓕᓯᔪᓂᑦ ᐃᓕᓴᕐᓂᖅᑐᓂᑦ ᐊᑐᖅᑕᐅᖅᖃᓂᒃ ᐱᖓᓱᖏᑦᑐᓂᑦ ᐊᐅᓪᓕᓯᔪᓂᑦ ᐊᓯᒪ ᖃᐅᔨᓴᖅᑕᐅᖅᖃᖅᑕᕐᓂᖏᓐᓄᓪᓗᑦ ᑐᑦᑐᓄᐃᑦ ᓴᓇᓕᖅᑐᓂᒃ ᐊᑐᐃᓐᓇᐅᔪᓂᑦ ᐃᓄᐃᑦ ᖃᐅᔨᔪᑎᑎᒍᑦ ᐊᑐᐃᓐᓇᐅᔪᓂᑦ ᖃᐅᔨᔪᐅᑎᓂᑦ. ᑎᒥᓴᒪᓂᐊᖅᑕᖏᓐᓄ ᐊᐅᓪᓕᓯᔪᑎᓄᑦ ᕿᖏᐅᑎᓕᐅᕈᑦᔨ ᐊᑐᐊᖃᖅᑐᑦᓚᐅᖅᑎᓂᒃ ᑎᑎᕋᖅᑕᐅᖅᑐᓂᒃ ᐊᑐᐃᓐᓇᐅᔪᓂᑦ ᑎᑎᕋᖅᑕᐅᖅᑐᓂᒃ ᐃᓚᖓᑦ ᐊᑐᐃᓐᓇᐅᔪᓂᑦ ᐃᓚᖓᑦ ᖃᐅᔨᔪᐅᑎᓂᑦ ᖃᐅᔨᔪᖓᑎᓂᒃ ᐊᑐᐃᓐᓇᐅᔪᓂᑦ ᐃᓚᖓᑦ ᖃᐅᔨᔪᐅᑎᓂᑦ ᐊᑐᐃᓐᓇᐅᔪᓂᑦ ᐃᓚᖓ

ᖃᓂᖅᓴᐅ

ᓄᐃᑲᓇᔪᑎᑦ 1

ᖃᓂᖅᓴᐅ 2

1.0 ᑐᖏᑦᓴᐃᓪᑦ 3

2.0 ᐃᓕᐅᔪᑦ 4

3.0 ᖃᑎᖅᑕᐊᖅᓂᖅ/ᐅᓪᓗᖅᓯᐅᑎᑦ 4

4.0 ᖃᑭᖅᑕᓗᓴᒥ ᑐᑦᑐᐃᑦ 5

4.1 ᐊᒥᓱᓂᖏᑦ 5

4.2 ᑐᓂᓂᖅᓴᖅᑕᐅᓂᖏᓐᓄ 6

4.3 ᐄᓀᓇᐅᔪᖅ ᖃᓄᐃᓕᓐᓄᑦᓂᖏᓐᓄ 8

5.0 ᐊᖏᔪᓱᓀᓂᖅ 10

5.1 ᐊᖏᔪᓱᓀᓂᖅ ᐱᓪᓗᒍ ᐊᐅᓚᑦᓯᐸᑎᓕᕆᓂᖅ 10

5.2 ᐄᓀᓇᐅᔪᖅ ᖃᓄᐃᓕᓐᓄᑦᓂᖏᓐᓄ 11

6.0 ᖃᐅᔨᓴᖅᑭᓀᓂᖅ 11

6.1 ᓄᓇᓕᑦᓂᒥᐅᑕᓂᓂᑦ ᐃᓕᑎᕐᐅᑎᑦᓯᓂᖅ ᖃᐅᔨᓴᖅᑭᓀᓂᑲᓪᓚᑦ 12

6.2 ᐃᓄᐃᑦ ᖃᐅᔨᓴᒪᕐᒪᖅᑎᑦ 12

6.3 ᐊᖏᔪᑕᐅᔪᑦ ᐅᖃᕆᐊᖅᑐᑕᐅᖃᖅᑕᑦᓂᖏᑦ 12

6.4 ᑲᑎᖏᓐᓄᐅᖅᑐᑦ ᖃᓄᐃᓕᓐᓄᓂᖏᑦ 12

6.4.1 ᖃᓄᐃᑎᒥ ᐊᒥᓱᓂᖏᑦ, ᐅᓄᕐᓯᓂᓕᐊᓂᖏᑦ/ᐊᒥᓱᕈᖅᐸᑦᓕᐊᓂᖏᑦ, ᐊᓪᓚ ᖃᓄᐃᑎᒥ ᐊᓯᐅᓴᑦᓕᐊᑎᒥᓂᖏᑦ 12

6.4.2 ᑲᑎᕆᓂᑦ ᐱᓪᓗᒥᑦ ᓇᓗᓇᐃᕐᐃᑦᓯᐊᖅᓂᖅ 12

6.4.3 ᑎᒥᖏᑕ ᐃᓗᖏᑦ ᖃᓄᐃᓕᓂᓂᖏᑦ 13

6.5 ᐅᖓᓯᓂᓂᖏᑕ ᐱᕕᑐᓂᖏᑦ 13

6.6 ᑭᖑᑐᐃᓂᓇᐃᑦ ᐸᕕᓴᓂᖏᑦ 13

6.7 ᓂᕈᑐᓯᑦ 13

6.8 ᐄᓀᓇᐅᔪᖅ ᖃᓄᐃᓕᓐᓄᓂᖏᓐᓄ 14

7.0 ᓇᓂᑎᑕᐃᑦ 16

ᐃᓕᓐᓂᐅᓴᐅᓯᒪᔪᖅ 1: 2019/2024 ᖃᑭᖅᑕᓗᓴᒥ ᑐᑦᑐᓂᑲ ᖃᓄᐃᓕᓐᓂᐅᑐᑎᓄᑦ ᐸᖅᓇᐅᑎ 20

2014/2015 ᐊᐅᓚᑦᓯᐊᑎᑦᓴᓂᑦ ᖃᓄᐃᓕᓐᓂᐅᖅᓂᖅ 20

ᐊᑐᓕᖅᑯᐅᔪᑦ 20

ᐊᐅᓚᑦᓯᓂᐊᑐᑎᑦᓴᓂᑦ ᐸᖅᓇᐅᑎ 20

ᖃᐅᔨᓴᐅᑎᐅᑉᓴᖅᑐᑐᓂᑦ ᐸᖅᓇᐅᑎ 20

ᐃᓕᓐᓂᐅᓴᐅᓯᒪᔪᖅ 2: ᒪᓕᓴᓕᓇᔪᑦ ᐊᓪᓚ ᐃᓗᓕᖅᑎᑦ 22

ᖃᑭᖅᑕᓗᒍ ᑐᑦᑐᓕᓴᓂᖅᑎᑦ ᐊᐅᓪᓚᕆᔨᕐᓂᕐᓄᓂᑦ ᐸᖁᓕᐅᑎᑦᓂᖅ

1.0 ᑐᖏᑲᓂᐅᑦ

ᑐᖏᓪᓗᐊᑕᐅᔪᖅ ᖃᑭᖅᑕᓗᒍ ᑐᑦᑐᓕᓴᓂᖅᑎᑦ ᐊᐅᓪᓚᕆᔨᕐᓂᕐᓄᓂᑦ ᐸᖁᓕᐅᑎᓪᓕᐅᑦᓴᓂᖅᑐᓂᑦ (BICMP) ᐊᑐᖅᑕᐅᔪᓴᓗᖅᑐᒥᑦ ᐊᑐᐊᓚᓪᐅᕈᒪᓪᓗᑎᑲᑎᒃ ᐊᖏᓕᓇᑦᑕᐅᓯᓂᐊᖑᓴᓇᑦ ᖃᓇᖅ ᐊᒻᒪ ᖃᐅᔨᓴᖅᑕᐅᓗᑎᒃ ᖃᑭᖅᑕᓗᒍ ᑐᑦᑐᓕᓴᓂᖅᑎᑦ ᑐᑦᑐᐃᑦ (ᑕᐸᓯᓂ ᖃᑭᖅᑕᓗᒍ ᑐᑦᑐᓕᓴᓂᖅᑎᑦ ᑐᑦᑐᐃᑦ). ᐊᐅᓪᓚᕆᔨᕐᓂᕐᓄᓂᑦ ᐊᑐᓕᖏᔨᔪᑎᐅᓪᓚᕆᓪᓕᐊᓱᑦ ᐃᓗᐊᓂ ᐸᖁᓕᐅᑎᓪᓕᐊᑦ ᐊᑐᖅᑕᐅᔪᖏᐅᓪᓚᕆᓪᓕᐊᓱᑦ ᑐᑦᑐᐃᑦ ᐃᓕᓯᓪᓚᕆᖏᐅᓪᓗᑎᒃ ᐃᓂᓯᓴᑦᑐᓂᑦ ᐱᐅᔪᖏᓪᓗᑎᒃ ᐃᓂᓯᓴᑦᑐᓂᑦ, ᐃᓐᓇᐅᑦᑐᑦᑐᐃᑦ ᐱᐅᔪᓯᓯᓂᖅᑐᓂᑦ ᐊᑐᖅᑕᐅᓯᓂᐊᖑᓴᓇᑦ ᐃᓐᓇᐅᑦᑐᑦᑐᐃᑦ, ᐃᓂᓯᓴᑦᑐᓂᑦ ᐊᑐᐃᓐᓇᐅᓪᓚᕆᓪᓕᐊᓱᑦ ᐃᓐᓇᐅᑦᑐᑦᑐᐃᑦ, ᐃᓂᓯᓴᑦᑐᓂᑦ ᐅᐸᐸᑦᑐᓂᑦ ᐊᑐᐃᓐᓇᐅᓪᓚᕆᓪᓕᐊᓱᑦ ᐃᓐᓇᐅᑦᑐᑦᑐᐃ


2.0 ᐃᓕᐅᔪᑦ

ᖃᑭᖅᑕᓗᒍ ᑐᑦᑐᓕᓴᓂᖏᑦ ᐊᐅᓪᓚᕆᔨᐸᑎᓂᖅᓯᓂ ᐸᖅᓯᓐᔪᐊᑎᓕᓂ ᓄᓇᒃᓵᒋᐅᓕᓚᒥᓂ; ᑭᓯᐊᓂᓕ, ᐃᓄᐃᑦ ᐊᖏᓪᓗᓯᑎᑦ ᖃᓄᖅ (10) ᓄᓇᓕᵒᓂᑦ ᑐᑦᑐᑎᓯᐸᑦᑐᓂᵒᒃ ᖃᑭᖅᑕᓗᒍ (ᐃᖏᓗᐃᑦ, ᑭᒋᒥᑦ, ᑭᵒᖏᓚᐃᑦ, ᐸᵒᓂᵒᖅᑐᖅ, ᖃᑭᖅᑕᑦᔪᐊᖅ, ᑲᑭᖅᓴᐃᓚᒃ, ᒥᑦᑎᒪᒋᓕᓂᵒ, ᐃᐢᐱᐊᐊᖅᓴᒃ, ᐃᒡᓗᓂᵒ ᐊᖏᓚ ᓴᓂᕋᖅᓴ) ᐊᑦᑐᖅᑕᐅᓂᖅᐸᖅᓯᓂᐊᖅᑐᑦ. ᑲᑐᔨᓂᵒᓄᑦ ᐊᐅᓪᓚᕆᔨᓂ ᑲᑐᔨᖅᑎᑦᑕ ᐃᑲᔪᓂᓂᐊᖅᑐᑦ ᐱᓂᓕᵒᓄᑦ ᐊᐅᓪᓚᕆᔨᓂ ᐱᖓᐊᖅᑎᑦᑕᑐᑦ, ᖃᑭᖅᑕᓗᒍ ᐅᓪᓚᕆᔨᓂ ᑐᑦᑐᑎᓯᐸᑦᑐᓂᵒᒃ ᖃᑭᖅᑕᓗᒍ ᑐᑦᑐᑎᓯᐸᑦᑐᓂᵒᒃ ᐊᐅᓪᓚᕆᔨᓂ ᑐᑦᑐᑎᓯᐸᑦᑐᓂᵒᒃ (HTOs), ᖃᑭᖅᑕᓗᒍ ᐅᒪᔨᓪᓚᕆᔨᓄᑦ ᑲᑎᒪᔨᖅᑐᑦ (QWB), ᓄᓇᓕᵒᓂ ᑐᵒᖏᓚᐃᓄᑦ (NTI), ᓄᓇᓕᓗᒍ ᐅᒪᔨᓪᓚᕆᔨᓄᑦ ᑲᑎᒪᔨᖅᑐᑦ (NWMB), ᐊᖏᓗ ᒥᐅᔭᓕᓪᓚᕆᐃᓚᓄᑦ ᑲᓇᒥᑦ (PC). ᐊᐅᓪᓚᕆᐅᓂᖅᑐᑦ ᖃᑭᖅᑕᓗᒍ ᑐᑦᑐᑎᓯᐸᑦᑐᓂᵒᒃ ᐊᐅᓪᓚᕆᔨᓄᑦ ᑲᑎᒪᔨᖅᑐᑦ ᑐᑦᑐᑎᓯᐸᑦᑐᓂᵒᒃ ᑲᑐᔨᓂᵒᓄᑦ ᐊᐅᓪᓚᕆᔨᓂᖅ ᑲᑐᔨᖅᑎᑦᑕ ᐱᓕᕆᖅᑎᑦᑕᑐᑦ ᑲᑐᔨᖅᑎᑦᑕᑐᑦ ᐊᐅᓪᓚᕆᔨᓂᖅ

4.0 ᖃᑭᖅᑕᓗᒍ ᑐᑦᑐᐃᓕᑦ


4.1 ᐊᒥᓱᓕᕆᑦ

ᐃᓄᐃᑦ ᖃᐅᔨᓴᓕᑐᖅᓯᕆᑦ ᐊᒻᒪ ᐃᑦᑕᖏᓕᓪᓕᔨᓄᐃᓪᓄᑦ >ᐊᖏᓴᖅᑕᐅᓂᒪᔪᑦ ᓂᕆᔨᓕᐊᔪᓚᐅᕆᖅᑕᑦᑕ ᐃᓂᒥᓕᕆᓕᕆᓂᕆᑦ ᐃᓂᒪᓕᐅᑕᓂᒪᔪᑦ ᖃᑭᖅᑕᓗᒍ ᑐᑦᑐᐃᓕᑦ ᐊᖏᔪᖅᓴᓂᒪᓕᕆᓂᕆᑦ ᐊᒥᓱᓕᕆᔨᑦ ᐊᖏᔪᖅᓴᓂᒪᓕᕆᓂᕆᑦ ᐊᒥᓱᓕᕆᔨᑦ ᐅᓄᖅᓴᖅᑐᑎᒃ, ᐊᓂᔨᓴᕐᑎᑎᒃ ᐊᖏᕐᕋᐃᑦ ᐊᓂᔪᖅᑐᑎᒃᓴᖅᑐᑎᒃ 50-ᓂᑦ 90-ᓄᑦ. ᐱᓗᐊᖅᓱᐊᑎᓕᒋᑦ, ᑐᑦᑐᐃᑦ ᐊᒥᓱᓕᕆᓕᓕᒫᖅᑐᑦ ᖃᓄᖅᑐᑎᒃ ᐱᑕᖅᓗᐊᖅᑎᓪᓗᐊ ᐊᒻᒪ ᖃᑦᓯᓴᐅᑎᓕᒋᑦ (ᑲᑎᐅᐅᓕ ᐊᓂᕐᑕᓗ 2015; ᐅᖏᓪᓕᓗ 1998; ᓂᑕᑕᓂ 1991). ᑕᒪᑲᓂᐊ ᐊᖏᓂᐊᓚᖏᓂᕆᑦ ᐊᓂᔨᐅᑎᐅᓂᒪᔪᑦ ᖃᐅᓂᓂ ᐊᓂᕐᑕᓂ ᐅᑭᐅᓂᕆᑦ ᐊᒥᓱᓕᕆᔨᑦ ᐱᐅᓂᕐᓴᐅᓂᒪᔪᑦ ᐊᒻᒪ ᖃᑦᓯᓴᐅᑎᓕᒋᑦ (ᑲᑎᐅᐅᓕ ᐊᓂᕐᑕᓗ 2015; ᐅᓪᓚᕆᓕᓗ 1998; ᓂᑕᑕᓂ 1991). ᑕᒪᑲᓂᐊ ᐊᖏᓂᐊᓚᖏᓂᕆᑦ ᐊᓂᔨᐅᑎᐅᓂᒪᔪᑦ ᖃᐅᓂᓂ ᐊᓂᕐᑕᓂ ᐅᑭᐅᓂᕆᑦ ᓇᒥᐅᓕᓪᓕᖅ (ᐅᖏᓪᓕᓗ 1997; ᓴᓂ 2001; ᒪᐊᖏᓄ ᐊᒻᒪ <ᐃᑲᑦ 2000). ᑕᒪᑲᓂᐊ ᐊᑐᖅᑕᐅᓕᖅᑐᑦᑐᑦ ᐃᓂᒪᒪᔨᐅᔪᑦ ᐱᐅᓂᐅᔨᓂᒪᔪᑦ ᐊᒻᒪ ᐱᔪᓂᖅᓴᐅᑦᑐᓂᑦ ᑐᓂᓯᓂᐅᓂᑦ ᐊᒻᒪ ᐊᓂᕐᑕᓂ ᐅᑭᐅᓂᕆᑦ ᐃᓂᒪᔪᑦ ᐅᓄᖅᓴᖅᑐᑦᑐᑦ ᐊᒻᒪ ᐃᓂᒪᔪᑦ ᐊᓂᕐᑕᓂ ᐅᑭᐅᓂᕆᑦ ᐃᓂᒪᔪᑦ ᐊᒻᒪ




ᓇᓗᓇᐃᖅᑎᑦ 2. ᖃᓄᐃᖅᑎᓂᓗᓴᕐᓂᖅᓯᑦ ᓇᐸᖅᑐᖅᓯᓂᖅᑐᑦ ᑐᓂᑐᐃᓐ ᑲᑎᓇᓂᑦ ᓇᓗᓇᐃᔨᐃᔪᑦ ᐊᑦᓯᑦᑐᓂᑦ/ᐊᐅᐸᖅᑐᖅᓯᑦ, ᐃᓕᓴᕐᓂᐊᔪᑦ/ᓴᖅᓯᖅᑐᖅᓯᑦ, ᖃᓄᓯᑦᑐᑦ/ᑐᖓᔪᐊᖏᓪᓚᔪᖅ ᐊᖏᒪ ᓄᖓᕐᐸᕐᓂᐊᔪᑦ/ᐊᐅᐳᐸᕐᓇᖅᑐᖅ ᐊᒥᓱᓂᓴᓕᕐᓂᖅᑐᓄᓂ ᐊᓯᔨᐸᕐᓂᐊᔾᔨᒃᑯᑦ. (ACCWM 2014).

ᓄᐊᖃᓱᓇᐅᖅᑐᒍᑦ ᐊᒥᓱᓯᖅᐸᕐᓂᐊᓂᖅᑐᓂᓂ ᑐᑭᓯᐊᕈᑎᓂᓂ ᐊᐱᖅᓄᑎᒍᑦ ᐊᖅᓄᖅᑐᖅᑐᑐᒃ ᐁᖅᑭᖃᓱᒪᔪᓂᖅ ᐊᐱᖅᓄᑎᒍᑦ 2015-ᒥᑦ 2019-ᒧᑦ, ᐊᑐᖅᑕᐅᔪᓐᓇᕆᔨᖅᑐᓂᖅ ᓇᓗᓇᐃᔨᐃᑎᐅᓪᓗᑎᒃ ᑐᑦᓯᐃᓐ ᐃᓕᓴᕐᓂᐊᓕᖓᒪᓪᓗᒥ, ᑕᐃᒪᐃᑐᐃᓐᓇᖏᓪᓗᖓᒪᓪᓗᒥ ᐅᐸᓴᓂᓂᖅ ᐃᓕᓴᕐᓂᐊᓕᖓᒪᓪᓗᒥ. ᑭᓱᐊᓂᓕ, ᐊᑐᖅᑕᐅᔪᓐᓇᖅᑐᓂᑦ ᓇᓗᓇᐃᖅᑕᐅᓱᓕᔪᓂ ᐊᑐᖅᑐᖅᑎᑦ ᐁᖅᑭᖃᓱᒪᔪᓂᖅᑐᑦ ᐊᖏᓪᓗᒥ ᐱᓕᕆᔨᓴᖅᓯᓐᓂᖅᑐᑦ ᓇᓗᓇᐃᖅᑕᐅᓱᓕᔪᓂ ᐊᐅᓪᓚᕆᔨᓴᖅᓯᓐᓂᖅᑐᑦ ᐃᓕᓴᕐᓂᐊᓕᖓᒪᓪᓗᒥ ᐊᐅᓚᑦᓯᕐᓗᒍ ᐃᓕᓴᕐᓂᐊᓕᖓᒪᓪᓗᒥ ᐱᓕᕆᔨᓴᖅᓯᓐᓂᖅᑐᑦ ᖃᓄᐃᓕᓴᕐᓂᐊᓕᖓᒪᓪᓗᒥ ᐊᐅᓚᑦᓯᕐᓗᒍ



ᓇᓗᓇᐃᑦᑯᖅ 3. ᐃᓄᐃᑦ ᖃᐅᔨᒪᔭᑐᖃᖅᑭᒃᑯᑦ ᖃᐅᔨᓴᐅᑎᐅᑭᒡᓗᑦ ᐃᓕᑭᔾᔪᓕᑎᒃ ᐃᓄᐃᑦ ᖁᑭᖅᑕᓗᒍᑎᐅᑕᐃᑦ ᐊᖏᕐᕋᐃᑦ 1990–ᖃᓗᓕ ᓱᓕᔨᓂᖅᓴᓂᓕ ᓄᖏᔪᓐᓇᒧᑦ (1993) ᐊᑐᐃᓇᓚᑕᐃᓕᐅᖅᑐᑦ ᐱᖓᓱᓂᑦ ᖁᑭᖅᑕᓗᒍᑎᒥ ᑐᑦᑐᑐᒥᑦ ᑎᑎᖅᖏᓚᕐᑐᓂᒃ ᐅᕝᕙᓚᓇᓂᑦ ᓄᓇᐃᑦ ᐊᕙᑎᖏᓇᓂᕐᑐᓂᑦ ᓇᒥ.



ᓇᓗᓇᐃᓐᓄᑦ 4. ᑐᑦᑐᐃᑦ ᑲᑎᖏᓐᓄᐸᐃᑦ/ᐃᓕᖓᑦ ᐊᒥᓱᐃᑦ ᐊᖏᕿᙵᒫᒃ ᐅᖏᓱᓂᖅᓯᓂᑦ ᓇᓗᓇᐃᖏᑦᓱᐊᕐᓂᓂᖅ ᒪᓕᑦᑐᓕᑦ ᓂᐱᓕᐅᖅᑕᐅᓱᒪᔪᓂᖅ ᖃᐅᔨᓴᐅᑎᐃᓇᓂᖅ 1987-ᒥᑦ 1994-ᒧᑦ (ᐱᓗᐊᖅᑐᒥ ᓂᒋᖏᓂ ᖃᑭᖅᑕᓗᐸᑦ), ᐊᒻᒪ 2008-ᒥᑦ 2011-ᒧᑦ (ᐅᐊᖏᓇᖏᓂ ᖃᑭᖅᑕᓗᐸᑦ) (ᑲᑎᐱᐅᓪ ᐊᓱᖏᓪᓗᓇ 2015). ᓇᓗᓇᐃᖃᑦᓱᒪᔪᖅᑐᓂᖅ ᓇᓗᓇᐃᖃᐃᔪᓂᑦ ᐱᓪᓚᔪᑎᑎᓂᔪᖅ ᐊᑐᖅᑕᐅᓪᓚᐅᖅᑐᓂᖅ ᓴᖅᑭᔪᔪᑎᐅᓗᑎᑐᑦ ᐅᔪᓂᖏᓐᓄᑦ ᓯᓚᑦᓴᐅᓂᖅᓱᒋᑦ ᐊᐃᑦᑕᐃᓗᓂᖅᓱᒋᑦ ᑐᑭᖅᑕᓗᐸᒥ ᑐᑦᑐᐃᑦ ᑕᑯᔪᔪᓂ (ᑲᑎᐱᐅᓪ ᓂᒋᖏᓂ 2015).


4.3 ᐃᓕᓴᐅᔪᖅ ᖃᓄᐃᓕᓚᒋᓕᕐᓂᖏᓐᓄᑦ

ᑭᑭᖅᑕᓗᓪᓕ ᑐᑦᑐᐃᑦ ᐃᓕᓴᓐ ᐊᐅᐸᑦᑐᒥᑦ ᐊᑐᖅᑐᑦ (2019). 2000–ᓂᑦ ᐊᖏᕿᙵᐃᑦ ᐱᑎᐊᓕᕆᔨᖅᑎᓕᓚᓂᑦ, ᐃᓄᐃᑦ ᖃᐅᔨᓴᓕᒃᑐᖏᓱᓂᑦ ᓄᐊᑕᐅᓕᐅᖅᑐᑦ ᐊᖏᔪᓂᕐᓂᓄᑦ ᓇᑭᐅᐃᓴᓐᓇᖅ ᑭᑭᖅᑕᓗᓪᓕᒥᑦ ᐃᓂᓕᐅᑕᓇᓂᕐᒥᑦ ᑐᑦᑐᐃᑦ ᓄᖅᑭᑦᑐᓕᕐᓂᖅᓱᒋᑦ ᐊᖏᔪᓂᕐᓂᓄᑦ ᑭᑭᖅᑕᓗᓪᓕᒥᑦ 1990-ᖏᑦᓱᓄᑦ. ᐱᔨᔪᑎᐅᓚᓂ ᓇᓚᓇᓕᐅᕐᓱᓂᖅᑐᑦᑕᑦ ᓄᖅᑭᐸᑦᓯᓕᐊᓂᖏᓱᓄᑦ ᐅᓂᓱᓴᖅᑭᓕᐅᖅᑐᑦᑕᑦ ᐱᓪᓚᑦᑕᑦ ᐊᖏᔪᓂᕐᓂᓄᑦ ᑭᑭᖅᑕᓗᓪᓕᒥᑦ ᓇᓕᐅᑐᓂᖅᓱ, ᖃᐅᔨᓴᓇᐊᓕᓕᐅᖅᑐᑦᑕᑦ ᐊᑐᖅᑕᐅᓚᓂᑎᓄᑦ ᐃᓄᐃᑦ ᖃᐅᔨᓴᓕᒃᑐᖏᓱᓂᑦ ᐊᒥᓱᓂᖏᓱᓄᑦ ᓇᓕᐅᑕᒋᐅᑕᓚᓂᑎᓄᑦ ᐃᓚᓂᕐᓇᓂᓂᕐᓂᑦ ᑭᑭᖅᑕᓗᓪᓕᒥᑦ ᑐᑦᑐᐃᑦ ᑕᑯᔪᔪᓂ ᐅᐸᑦᑐᒥᑦ ᐱᓪᓚᑦᑕᑦ ᐃᓕᓴᐅᔪᖅᑐᖏᓱᓂᑦ ᐊᖏᔪᓂᕐᓂᓄᑦ ᐃᓚᓂᕐᓇᓂᓂᕐᓂᑦ ᐊᒻᒪ ᓂᓂᓕᐊᒥ 2014. ᐃᓕᓂ 2014 ᑎᑭᑎᓕᓚᒍ 4,652-ᖓᓕᐅᖅᑐᑦ (3,462-ᓂᑦ 6,250-ᓄᑦ; 95 ᐳᓴᕐᑎᒍᑦ ᖁᓂᖖᕐᓂᔪᑎᖅᑐᖅᑎᑎᓄᑦ) ᑐᑦᑐᐃᑦ ᑭᑭᖅᑕᓗᓪᓕ, ᐱᖅᓂᐅᑎᓕᓚᓂᑦ ᖃᓄᑦᑐᒋᑦᑐᑦ ᑭᓚᓕᔭᓪᓕ ᑭᓚᓕᔭᓪᓕ ᓄᓇᒥᒃᑐᓂᖅ ᐃᒪᖅᓗᓂ. ᐊᒥᓱᓂᖏᓱᓄᑦ ᑐᑦᑐᐃᑦ ᓂᒋᖏᓂᓂᓱᓄᑦ ᑭᑭᖅᑕᓗᓪᓕ ᐊᒻᒪ ᑭᓚᓕᔭᓪᓕ (ᑎᑎᕆᔨᓴᓂ 1; ᓇᓗᓇᐃᖃᑦᓱᒪᔪᖅ 5; ᑲᑎᐱᐅᓪ ᐊᓱᖏᓪᓗᓇ 2015). ᑕᒪᓐᓇ ᐊᑕᐅᓴᖃᑦᓱᑎᓕᓚᓂᑦ ᑐᑦᑐᐃᑦ ᓄᖅᑭᐸᑦᓯᓕᐊᓂᖏᓱᓄᑦ 95 ᐳᓴᓂᑦ ᐅᖏᓕᒋᓂᑐᑦᑕᑦ ᒪᓕᑦᑐᓕᑦ 1991–ᒥᑦ ᖃᓄᖅ ᐱᐅᓂᖏᓱᓄᑦ ᓇᓕᐅᑕᕐᑕᐅᓴᖃᑦᓱᒪᔪᓂᖅ ᐱᓱᐅᓴᖃᑦᓱᒪᔪᓂᖅ ᐃᓄᐃᑦ ᖃᐅᔨᓴᓕᒃᑐᖏᓱᒧᑦ (ᓂᔪᓴᓗᓂ 1998). ᐃᓕᓂ 2014 ᑎᑭᑎᓕᓚᒍ 159-ᓂᑦ 622-ᓅᖖᓕᐅᖅᑐᑦ (95 ᐳᓴᕐᑎᒍᑦ ᖁᓂᖖᕐᓂᔪᑎᖅᑐᖅᑎᑎᓄᑦ) ᑐᑦᑐᐃᑦ ᐅᐊᖏᓪᓚᑦ ᑭᑭᖅᑕᓗᓪᓕ (ᑎᑎᕆᔨᓴᓂ 1; ᓇᓗᓇᐃᖃᑦᓱᒪᔪᖅ 4). ᑐᑦᑐᐃᑦ ᑕᒫᓂ ᐊᑦᑕᓇᖅᑐᒍᑦᑐᑦ ᓄᖅᑕᐅᑐᑐᑦᓄᐊᓕᖏᓱᖅᑎᑎᓄᑦ.

ᑎᑎᕃᓯᒪᔪᓂ 1. ᓇᓕᐅᑦᑕᕃᓯᒪᔪᑦ ᑐᑦᑐᐃᑦ ᐱᒋᖏᕃᓂᐅᓂᑦ ᖃᐅᔨᓴᕃᑦᓴᐅᔪᓂ (ᓇᓗᓇᐃᒃᓯᑎᖅ 5) ᐃᓐᓂᐊᕆᒥᑦ ᐊᒻᒪ ᒪᕐᓯᒥᑦ 2014-ᒥ ᕿᑭᖅᑕᓗᓴᒥ ᖃᐅᔨᓴᕃᑦᓂᒥᑦ.

ᖃᐃᕃᓯᖅ (ᖃᐅᔨᓴᕃᓴᐅᔪᓯᖅ)	ᐊᑦᑐᐃᑦ ᓇᓕᐅᑦᑕᖅᑕᐅᓯᒪᓂᒥᑦ	95 ᐳᓴᖏᔪᑦ ᕿᓯᒪᖏᕐᔾᔪᑎᓕᒃ ᑭᒃᓕᖏᓐ (ᐊᑦᑐᐃᑦ ᐅᖏᓯᓂᖏᑦ ᓇᓕᐅᑦᑕᕃᓯᒪᔪᑦ)
<u>ᕿᑭᖅᑕᓗᖏᒥ ᐅᐊᖏᓇᖏᓕ</u>		
ᐃᒃᐱᑭᑦᑐᕃᐊᖅ, ᓯᓚᓯᐅᑦᕃᖏᒥ	6	1-30
ᓄᓗᔾᔪᒃ	224	96-521
ᐅᐊᓴᓇᐅᑉ ᕿᑎᐊ ᕿᑭᖅᑕᓗᓴᒥ	85	31-230
ᖃᑎᒌᓗᑦᑕ	**315**	**159-622**
<u>ᓂᑦᓯᒥ ᕿᑭᖅᑕᓗᒃ</u>		
ᕿᑎᐊ ᕿᑭᖅᑕᓗᓪᓕ	1,091	662-1,798
ᓯᑯᓯᓪᓕᖅ	216	48-972
ᓴᐅᒥᒃ	887	467-1,686
ᑭᓴᖏᐃᑦ	539	256-1,138
ᑭᖏᖏᓀᐃᑦ	1,603	1,158-2,220
ᖃᑎᑦᑐᑦᑕᑦ (+ᑭᖏᖏᓀᐃᑦ)	4,337	3,169-5,935
ᖃᑎᑦᑐᑦᑕᑦ (- ᑭᖏᖏᓀᐃᑦ)	**2,734**	**1,777-4,207**
ᕿᑭᖅᑕᓗᓴᒥ ᖃᑎᑦᑐᑦᑕᑦ	**4,652**	**3,462-6,250**

ᐊᒥᓯᐃᑦ ᐊᖏᕐᖃᑎᑎᕋᖅᑎᓪᓗᓂᑦ ᐃᓄᐃᑦ ᐊᖏᔪᓯᑦᑎᑦ ᐊᒻᒪ ᖃᐅᔨᓴᕃᑎᑦ ᓯᓇᑐᐃᓐᓇᕐᓂᖅ ᖃᑦᓯᓴᓐᓇᐅᓂᕐᓴᐅᑦᑐᓂᖏᓂᑦ ᕿᑭᖅᑕᓗᓴᒥ ᑐᑦᑕᐃᑦ ᐃᓕᓴᕐᕕᖏᓂᖅ. ᐊᒥᓯᐊᓗᐃᑦ ᐃᓄᐃᑦ ᑐᓴᐅᒪᑕᐅᓯᐊᓪᓚᕆᑦᑐᑦ 2013-ᒥ ᐊᓂᒋᑎᓪᓗᓴᐃᓯᑦ ᕿᑭᖅᑕᓗᓴᒥ ᑐᑦᑕᐃᑦ ᐃᓄᐃᑦ ᖃᐅᔨᓴᕃᔪᖏᑦ ᖃᐅᔨᓴᕃᓂᖅᓯᒪᓂᑦ (ᐊᖏᔪᓯᑦᑎᑦ-ᐃᓚᓇᑐᖃᐃᑦ ᐊᐱᕃᓯᑕᐅᓗᑎᒃ) ᐊᒻᒪ ᓄᓇᓕᓂᒥᐅᑕᐃᑦ ᐃᓯᒪᒋᔭᖏᓂᑦ ᑐᓴᕆᓯᓂᖅ ᐊᐱᖅᓯᑎᑦᓯᓯᒪᓂᑦ, ᐊᒻᒪ 2012-2015 ᓄᓇᓕᓂᑦ ᐊᒻᒪ ᐊᖏᔪᓯᑦᑎᑦ ᒥᑭᒋᐊᒥᓂᖅᑎᓪᓗᒍ ᑲᑎᕕᖅᑎᑦᓂᓪᓗᒃ ᑐᓴᐅᒪᑎᑦᓯᔪᑎᐅᓕᐅᖅᑐᑦ ᐃᓯᒪᓕᐅᖅᑐᑦ ᑐᑦᑕᐃᑦ ᐊᒥᓱᓂᑦ ᓄᖏᓪᓕᐊᓂᖅᓯᒪᓂᑦ, ᐅᖃᓕᑦᑐᐊᓕᒫᓂᖅ ᐊᒥᓱᓂᑦ ᓄᖏᓪᓕᐊᓂᖅᓯᒪᓂᑦ ᐃᓯᒪᒋᔭᖏᓂᖅᓯᒪᓂᑦ. ᑭᓱᐊᓂᓕ, ᐃᓯᒪᓕᐅᖅᑐᑦ ᑐᑦᑕᐃᑦ ᐊᓱᐊᓄᑦ ᓄᓯᒪᓂᖅᓂᓕ, ᐅᖃᓕᑦᑐᐊᓕᒫᓕᒫᓂᖅᓯᒪᓕᑕ ᑕᐃᒪᓐᓇᖅ ᓯᓚᑐᐃᓐᓇᕐᓂᖅ ᖃᐅᔨᓴᕃᑎᑦ ᐃᓯᒪᓕᐅᖅᑎᓪᓗᓂᑦ. ᑭᓱᐊᓂᓕ, ᐃᓯᒪᓕᐅᖅᑐᑦ ᑐᑦᑕᐃᑦ ᐊᓱᐊᓄᑦ ᓄᓯᒪᓂᖅᓂᓕ, ᐅᖃᓕᑦᑐᐊᓕᒫᓕᒫᓂᖅᓯᒪᓕᑕ ᖃᐅᔨᓴᕃᑎᑦ ᐃᓯᒪᓕᐅᖅᑐᑦ ᐊᖏᔪᓯᑦᑎᑦ ᐊᐱᕃᓯᑕᐅᓗᑎᒃ ᓯᓚᑐᐃᓐᓇᕐᓂᖅ ᐊᒻᒪ ᐊᖏᔪᓯᑦᑎᑦ ᒥᑭᒋᐊᒥᓂᖅᑎᓪᓗᒍ ᑕᐃᒪᓐᓇᖅ



ᓇᓗᓇᐃᒋᑦ 5. ᖃᐅᔨᓴᕕᐅᓂᐊᖅᑐᑦ ᖁᑭᖅᑕᓗᒍ ᑐᑦᑐᓂᑦ ᖃᐅᔨᓴᓂᖅ – ᐃᓕᓴ 2014 ᐊᐃᓱᓗᓯᓂᓚᕆᐅᖅᑐᑦ ᖃᓄᕆᐅᓂᕐᓂᓇᓕᒃ ᐊᒻᒪ ᑲᒪᑦᐊᓂᐅᒋᓴ ᐊᖏᓇᓂᑕᐅᕐᓴᖅᑐᓂᑦ ᓇᓚᐅᑦᑎᐊᕐᓂᖅᑐᓂᑦ.

5.0 ᐊᖏᔪᓂᖅ

5.1 ᐊᖏᔪᓂᖅ ᐱᒃᓗᒍ ᐊᐅᓚᑦᓯᐊᕐᓇᓂᖅ

ᑐᑦᑕᐃᑦ ᐊᑐᖅᑐᒪᓇᐊᔭᒪᒡᒪᑦ ᐱᖃᓯᑐᖃᐅᖅᓯᒪᑦ ᓄᓇᕗᒻᒥ ᐊᒻᒪ ᓂᖅᑭᐅᕐᓂᑐᑎᑦ ᓄᓇᕗᒻᒥᑦ ᐃᓄᓐᓄᑦ ᐸᖏᓐᓄᑦ ᐃᒪᓕᒃ 1000-ᓄᑦ (ᓯᑕᒃᑕᓐ 1991). ᖃᓄᑦ (10) ᖁᑭᖅᑕᓗᒍ ᓄᓇᓕᒃ ᐊᒻᒪ ᐊᖏᕐᓂᔭᖅᑎᑦ ᓄᓇᕗᒻᒥᐅᑕᐃᑦ ᑕᑎᑎᔨᖅᖃᓂᒪᒡᒪᑦ ᖁᑭᖅᑕᓗᒍ ᑐᑦᑐᓂᑦ ᓂᕆᔨᒋᑦ, ᐱᓕᕆᔨᒃᑯᑦ ᐊᒻᒪ ᐃᓕᓴᕐᓂᐊᕐᓂᖅᑎᑦ. ᐅᖃᓄᐃᓕᓴᕐᓂᑦ ᐊᖏᓇᓂᑕᐅᕐᓴᖅᑐᓂᑦ ᐊᑐᐃᓄᖅᑲᓇᕐᓂᖅᑐᓂᑦ ᖁᑭᖅᑕᓗᒍ ᑐᑦᑐᓂᑦ, ᐱᓗᐊᖅᑐᒥ ᐅᕙᑦᑎᐊᕐᓂᖅᐳᖅᑕᖅᑐᑦᓯᒪᓂᑦ ᐊᒻᒪ ᐃᓕᓴᕐᓂᐊᕐᓂᖅᑐᓂᑦ. ᐊᖏᓇᓂᖅ, ᐃᓕᒪᖅᐳᓗᓂ ᐊᓯᖏᓐᓄᑦ ᐱᑕᖅᑭᓂᕐᑐᐃᓐᓇᕐᓂᓗ ᓯᐢᓗ ᓂᖅᑭᑐᓯᓄᑦ ᐱᖅᐳᓂᖅᓂᕐᓂᑦ ᐊᒻᒪ ᐃᓕᒍᑦ ᐱᐅᓯᖅᑎᑦ, ᐊᒥᓯᓂᕆᓂᑦ ᓄᓇᕆᑦᐸᑦᑐᓂᖅ, ᑐᑦᑐᓂᑦ ᓄᓇᕗᒻᒥᑦ ᐃᓕᓴᕐᓂᐊᕐᓂᖅᑐᓂᑦ. ᐊᖏᓇᓯᑎᓂᖅ ᐊᐳᖅᑐᓂᑦ ᐃᓕᓴᕐᓂᐊᕐᓂᖅᑐᓂᑦ ᐊᒻᒪ ᐃᓕᒍᑦ ᐱᐅᓯᖅᑎᑦ, ᐃᓕᓴᕐᓂᐊᕐᓂᖅᑐᓂᑦ ᐊᒻᒪ ᐃᓕᓴᕐᓂᐊᕐᓂᖅᑐᓂᑦ ᐊᖏᓇᓂᑕᐅᕐᓴ

ᐊᖏᕕᖓᑕᐅᔪᓐᓇᒃ ᐊᒥᓱᑦᑐᓪᓗᒍᓇᔭᖅᑐᓂᒃ ᐃᓐᓇᐅᔪᓐ ᐊᑐᖅᑕᐅᔪᒥᒃ ᐊᑦᓱᓂᐅᐅᔪᒥᒃ. ᐊᖏᓇᓂᓕᒫᑦ ᐊᐅᓪᓚᓱᓂᖅ ᐱᖅᓱᐅᔨᔪᓂᓇᖅᑐᑦ ᐊᐅᓪᓚᓱᔨᐸᓂᒃ ᐱᓪᓚᓂᒃ: ᑲᑎᒪᓂᒃ ᐱᔨᐅᔭᓇᖅᑎᑕᐅᔪᒃ ᐊᖏᓇᓂᐅᔪᓂᓇᖅᑎᑕᐅᔪᑎᒃ (TAH), ᓄᖅᑲᖏᑎᑦᓱᓂᑦ, ᐊᖏᓇᓂᓕᒫᑦ ᐊᐅᓪᓚᓱᔨᐸᓂᒃ(MUs), ᑯᑕᖅᖃᑦᓴᓂᒃ (NQLs), ᐋᒪ ᐊᑎᒍᒻᒥᐅ ᐃᓗᐊᓂ ᖃᖏᑕᐅᑎᓪᓗᒍ ᐱᑐᐃᓇᐅᔭᖏᔨᔪᑎᖃᖅᓂᖅ. ᐱᒋᐊᖅᑎᑦᓂᓂᖅ ᑲᑎᓪᓚᓱᔨᐸᓂᒃ ᐱᖅᓱᐅᔨᔪᓇᖅᑎᑕᐅᔪᒃ ᐊᖏᓇᓂᐅᔪᓂᓇᖅᑎᑕᐅᔪᓂᒃ ᐅᑭᖅᑐᑦᓱᓂᒃ ᐊᖅᑭᑦᑕᐅᔪᓂᒃ ᖃᖅᑎᑦᓱᓂᒃ ᑐᑦᓱᓂᒃ ᑐᑦᓯᐊᓂᒃ ᐊᖏᓇᓂᐅᔪᓇᖅᓯᒪᓕᒫᑦ, ᑯᑕᖅᖃᑦᓴᓂᒃ ᐊᖅᑭᑦᑕᐅᔪᓂᒃᑐᑦᓱᓂᒃᑐᑦᓯᐊᓂᒃ ᐱᒋᐊᖅᑎᑦᓂᓂᖅ ᓇᒥ ᐊᒥᓱᑦᑐᓪᓗᒍᓇᔭᖏᔨᔪᑎᖃᖅᓂᖅ ᐊᖅᑭᑦᑕᐅᔪᓂᒃ ᐊᖅᑭᑦᑕᐅᔪᓂᒃ ᐊᖏᓇᓂᐅᔪᓂᓇᖅᑎᑕᐅᔪᓂᒃ ᐊᖅᑭᑦᑕᐅᔪᓂᒃ ᐊᖏᓇᓂᐅᔪᓂᓇᖅᑎ

6.1 ᓄᓇᕗᑦᓂᒥᐅᑕᓂᕐᓂ ᐃᓕᑎᕕᐅᑎᑦᓯᓂᖅ ᖃᐅᔨᓴᖅᑭᓐᓂᖏᑦ

ᖃᐅᔨᓴᖅᑭᓕᕐᕕᖅᑐᑦ ᐊᕙᑎᓕᓚᕆᖏᑦ ᐱᖃᑎᖃᖅᑭᓴᑐᑎᑉ ᓄᓇᕗᑦᓂᒥᐅᑕᓂᕐᓂ ᐊᒻᒪ ᓄᓇᕗᖅᖃᕐᓂᕐᓂ ᐊᖏᔪᓯᑎᑦ ᒥᑭᑦᓂᐊᒻᒪᖅᑎᓪᓗ ᑲᑎᓴᖃᑎᒋᔭᐅᓂᖏᑦ (ᑕᓯᖏᓂᑦ ᐅᑦᑐᑎᑦᓴᑦ ᑲᑦᓴᐃᐅᓪ 2015; ᔪᐊᑦᑖᑦ 2014; ᔨᐊᓐᑭᓕᓪ ᐊᓯᖏᑦᓗ 2012; ᔨᐊᓐᑭᓕᓪ ᐊᒻᒪ ᔪᐊᑦᑖᑦ 2012; ᓄᓇᕗᒥᑎ ᐊᕙᑎᓕᓚᕆᖏᑦ 2013). ᐱᖃᓯᐅᑦᔨᓂᑦ Δ) ᐱᓕᓚᐊᑦᓴᐃᑦ ᐃᓗᓕᖅᑭᓯᓕᓂᖅ ᐊᒻᒪ ᖃᓄᐃᓕᓚᒪᓂᕐᓂᑦᓂᖅ ᖃᐅᔨᓴᐅᑎᑦ, ᐱ) ᑐᓴᐅᒪᑎᑦᓯᓂᐅᑕᑎᖃᓂᖅ ᐱᖃᓯᒋᑦᓂᖅ ᐊᒻᒪ ᐃᓚᖃᑎᒌᑦᓴᓂᖅᑐᑦ ᐊᒻᒪ ᑎᑭᓕᓚᕆᖏᑦ ᐃᓕᓴᕐᕕᖏᓂᑦᓴᓂᑦ ᐊᒻᒪ ᑎ) ᐃᓕᓕᒃᕕᐅᓪᓗᓂ ᖃᐅᔨᓴᖅᑐᓄᑦ ᐊᑐᑕᐅᓗᓂᑦ ᐃᓄᓇᑦ ᐱᓕᓚᐊᑦᓴᓂᖅᑐᑦ ᐊᒻᒪ ᐃᓕᓴᕐᕕᖏᓂᑦᓴᓂᑦ ᐊᒻᒪ ᐃᓄᓇᑦ ᐱᓕᓚᐊᑦᓴᓂᖅᑐᑦ ᐊᒻᒪ ᐃᓚᖃᑎᒌᑦᓴᓂᖅᑐᑦ ᓴᓕᑦᓂᖅᑐᑦ ᐊᒻᒪ ᐃᓕᓴᕐᕕᖏᓂᑦᓴᓂᑦ ᐊᒻᒪ ᑎ) ᐃᓕᓕᒃᕕᐅᓪᓗᓂ ᖃᐅᔨᓴᖅᑐᓄᑦ ᐃᓄᓇᑦ ᐱᓕᓚᐊᑦᓴᓂᖅᑐᑦ ᐊᒻᒪ ᐃᓕᓴᕐᕕᖏᓂᑦᓴᓂᑦ ᐊᒻᒪ ᑎ) ᑐᓴᐅᒪᑎᑦᓯᓂᐅᑕᑎᖃᓂᖅ ᐊᒻᒪ ᐃᓕᓴᕐᕕᖏᓂᑦᓴᓂᑦ ᐊᒻᒪ ᑎ) ᐃᓕᓕᒃᕕᐅᓪᓗᓂ ᖃᐅᔨᓴᖅᑐᓄᑦ ᐃᓄᓇᑦ ᐱᓕᓚᐊᑦᓴᓂᖅᑐ

ᖃᑉᑭᖅᑕᓗᒍ ᐊᑐᖅᑕᐅᔾᔨᐊᖏᖅᑐᒃᓯᒪᓕᒃ ᐊᖏᖅᓯᐅᒃᑐᓂᒃᓴᓪᓗᒍ ᐊᐱᓇᓱᓂᖅᓴᒻᒥᒃ ᐊᐅᓪᓚᕆᓯᐅᑎᓂᒃ ᐱᓪᓚᕆᐊᑎᐅᐅᔪᖅ (ᓄᓇᕗᑦᓴᒻᒥᒃ ᐊᓂᖓᓪᓚᐅᔪᓂᖅ ᐱᖓᒪ ᓄᓇᓪᓚᐅᔪᓂᖅ 2011). ᐊᐅᓪᓚᑕᐅᑎᓪᓗᓂᒃ ᑭᑉᑭᑦᑕᒍ ᐊᑕᐅᓂᕆᑎᓪᓗᓂᒃ ᑲᑎᕐᓂᐊᓕᖅᑎᑦ, ᐊᐃᓪᑐᖅᓯᒪᐅᖅᑎᓪᓗᓂᒃ, ᐅᓪᓚᓇᖅᑐᖅᑭᓇᖏᖅᑐᑦᑎᒃ ᓄᖓᒋᓂᕐᓂᐊᕐᓇᖏᖅᑐᑦᑎᒃ ᐊᓕᖅᑐᓂᒃᓴᓂᒃ ᑲᑎᕐᓂᐊᓕᖅᑎᑦ ᑲᑎᕐᓂᐊᓕᖅᑎᑦ.

6.4.3 ᑎᒥᖅᑕ ᐃᓗᖅᑦ ᖃᓄᐃᓕᖅᖢᑦ

ᑎᒥᖅᑕ ᐃᓗᖅᑦ ᖃᓄᐃᓕᖅᖢᖅᓂᒃ ᐅᖃᐅᓯᐅᑎᓪᓗᓂᒃ ᐱᓵᔪᑎᐅᔪᓂ ᖃᓄᐃᓕᓕᒫᓂᖅᑦ ᐊᑕᐅᓯᓂ ᐅᒪᔪᐃᓂ ᐃᓗᐊᓂᒃᑐᖅᓂ, ᓇᓗᓇᐃᔨᐃᑎᐅᐅᓯᓂᓚᒡᒪᑦ ᐅᕐᓴᕐᓂᖅᑐᑦᑎᒃ ᖃᓄᐃᓕᓕᒫᓂᖅᑐᓂᒃ, ᐋᓐᓂᐊᖅᑐᖅᑭᖅᑐᓂᒃ ᐊᒻᒪ ᖃᓄᑎᒥᖅᑐᓂᒃ ᑲᑎᕐᓂᐊᓕᖅᑎᑦᑐᓂᒃ ᐋᖅᑭᑦᓯᒪᓂᖅᑦ. ᑎᒥᖅᑕ ᐃᓗᖅᑦ ᖃᓄᐃᓂᖅᑦ ᐊᒻᒪ ᖃᓄᐃᓕᓕᒫᓂᖅᑦ ᑐᑎᑕᖅ. ᖃᐅᔨᒪᓂᖃᖅᑎᓪᓗᓂᒃ ᑐᑎᑕᖅ. ᖃᐅᔨᓴᓂᖅᑐᓂᒃ ᑐᑎᑕᖅ ᖃᐅᔨᓴᓂᖅᑐᓂᒃ ᐊᑐᖅᑐᓂᒃ ᐊᑐᖅᑐᓂᒃ ᐊᑐ

ᑎᑎᕆᔨᒪᔪᓂ 2. ᓄᓇᑐᐃᓇᖅᓯᓂᑦ ᖃᐅᔨᓴᐅᑎᓂᑦ ᐱᓕᕆᔨᐅᑎᖃᕐᓗᑐᑦ ᓇᓗᓇᐃᔾᐃᑎᓕᑐᑦ ᖃᓄᐃᓇᐅᑎᖃᖅᑐᓂᑦ ᑐᑦᑐᒃᓯᓄᓂᐅᕙᖅᑕᑦᑐᑦ, ᑐᑦᑐᐃᑦ ᐃᓂᒋᔭᐅᓂᑦ ᐊᒻᒪ ᑲᑎᓐᓂᑦ ᐊᒻᒪ ᐱᓕᕆᔨᐅᑎᓕᕐᓂᐊᕐᓂᑦᑐᑦ ᐊᑐᖅᑐᓂᑦ ᐱᓕᕆᔨᐅᑎᐅᓄᓕᐅᑦ ᐊᑐᖃᖅᑕᐅᓗᑎᑦ ᐱᓕᕆᔨᐅᑎᐅᓄᕙᑦ.

ᖃᐅᔨᓴᖃᑦᑕᑦᓂᖅ	ᐊᒥᓱᓕᕆᔨᓂᖅᓂᑦ ᖃᓄᐃᑎᑎᓂᑦ ᓇᓗᓇᐃᔾᐃᑎᓕᑐᑦ	ᐊᒥᓱᐸᖅᐸᓕᐅᓂᑦ	ᑲᒪᓕᓱᐅᓂᖅ ᐊᖏᓄᓇᓕᑕᐅᔨᓴᖃᖅᑐᑦ ᖃᓄᐃᓕᐅᓂᑦ	ᐃᓕᑦᑎᓂᑦ ᓄᓇᒥ	ᐊᒥᓱᓕᕆᔨᐅᑎᓂᑦ ᖃᓄᐃᑎᑎᓂᑦ ᓇᓗᓇᐃᔾᐃᑎᓕᓱᐅᓂᖅ	ᐃᓂᒋᔭᐅᑎᓂᑦ ᓂᕈᐊᖏᓂᑦ	ᐅᒪᐃᐃᑦ ᖃᓄᐃᓇᓕᑎᑦ ᑎᒥᖁᑎᑦ ᐃᓗᓐᓇᒧᑦ	ᐊᖏᓄᓇᓕᑕᐅᓂᖅᑐᑦ ᖃᑦᓱᐅᓂᖅᑐᑦ	ᖃᓄᐃᑎᑎ ᐅᖏᓱᓂᖅᓯᑕᖅᑐᑦ ᖃᓄᐃᓕᓚᓂᖅᑐᑦ
ᖃᖏᓪᑕᖃᑎᑎᑕᐅᓱᒪᔪᓄᑦ ᖃᓄᐃᔾᐃᖏᓂᖅ ᐅᓴᒥᓱᒪᔪᓂᖅ	X*	X*	X*	X	X	X*	X	X	
ᖃᑦᓱᐅᓂᖅᑐᑦᓂᑦ ᖃᓄᐃᔾᐃᖏᓂᖅ ᐊᐱᑦᓱᓂᖅᓄᑦ	X*	X	X						
ᑐᑦᑐᐃᑦ ᖃᓄᓂ ᐊᒥᓱᓂᖅᑐᓂᑦ ᖃᓄᐃᔾᐃᖏᓂᖅᓄᑦ ᓇᓗᓇᐃᔾᐃᓂᖅ	X	X*	X						
ᓄᓇᕗᒻᒥᖅ ᖃᓄᐃᑕᑐᓂᖅᑐᑦ ᖃᓄᐃᔾᐃᖏᓂᖅ					X*				
ᖃᓄᓕᕆᓂᖅᖁᑦ ᖃᓄᐃᔾᐃᖏᓂᖅ							X		
ᖃᓄᐅᑎᑦ ᐅᖃᓱᓂᖃᖏᓂᖅᑐᓂᑦ ᓇᓗᓇᐃᔾᐃᓂᖅᓄᑦ						X*			X
ᐊᖏᓄᓇᓕᑕᐅᑐᓂᖁᑦ ᖃᓄᐃᔾᐃᖏᓂᖅ		X*	X*					X	

* ᐃᓕᐃᓇᓚᖏᓂᓂ ᑭᓱᐊᓂ ᐅᕐᖢᓗᓂᓂᑦ ᐃᓕᓕᐅᔪᐅᓂᐊᖏᑎᓕᑐᑦ ᐊᓱᖏᓂᓇᑦ ᐱᓕᕆᐊᖏᓱᒪᔪᓇᑦ ᐅᕐᖢᓗᓂᓂᑦ ᐊᕐᕋᒍᒎᓂᑦ

6.8 ᐃᓚᓇᐅᔪᖅ ᖃᓄᐃᑕᖏᓕᕐᓂᖏᓐᓄᑦ

ᖃᑎᖒᐊᖏᑕᐅᑦᓱᐊᑭᓇᖅᑐᑦ ᐊᒻᒪ ᖃᐅᔨᓴᖏᑕᐅᑲᓂᕐᓗᑎᑦ ᐊᑐᖏᑕᐅᓱᒪᔪᓂᑦ ᖃᐅᔨᓴᐅᑎᐃᓂᕐᓂᑦ ᖃᑭᖏᑦᑐᓗᒍ ᑐᑦᑐᓂᑦ ᑕᒪᔨᐅᔪᓇᖅᑐᑦ ᐅᕙᓂ ᑲᓚᐅᔾᔨᒪᔪᓂ (2015). ᐅᕙᓂᓱᐊᖏᓚᐅᑕᐅᓕᖏᖏᖅᖃᖏᑦ ᖃᐅᔨᓴᖏᑎᓂᑦ ᓱᓇᑐᐃᓇᖅᓂᑦ ᖃᐅᔨᓴᐅᑎᐃᓐᓇᓂᑦ (1940-1970) ᑕᑎᑎᔪᓂᖏᖅᑐᓂᑦ ᖃᑦᓱᐅᔪᐃᓐᓇᐅᑎᓂᕐᓂᓱᓂᑦ ᓇᓕᐅᑦᑕᐸᑎᑎᓂᑦ ᐱᑕᖃᓕᐅᖏᑦᓱᒪᒪᑕ ᐊᒻᒪ ᐅᕙᓂᓱᐊᖏᑐᓴᐅᓕᐅᖏᖅᑐᓂᑦ. 2012-ᖑᒪᓕᓗᒍ, ᐊᕙᑎᓕᓗᒃᑯᑦ ᖃᐅᔨᓴᕆᐅᓱᒪᔪᓂ ᓂᕐᖏᓐᓄ ᖃᑎᖒᐊᖏᓕᒃ ᐊᒻᒪ ᓇᓕᐅᑦᑕᐸᓱᒪᒪᕐᖏᖅᑐᑎᑎᑦ ᖃᐅᔨᓴᐅᑎᑎᓂᕐᓂᑦ. 1,555-ᓂᑦ 3,093-ᓄᑦ (95 ᐳᓴᖏᑎᒍᑦ ᖃᑯᕙᖓᓱᐊᖏᖅᖃᖏᑦᑐᑎᑎᑦ; ᔮᐊᐟᑭᓴ ᐊᓱᖏᓕᓚᓇ 2012; ᐊᓱᐅᔨᕆᓱᒪᔪᓂ ᓇᓕᐅᑦᑕᐸᖏᑕᐅᓱᒪᔪᓂ ᐅᕙᓂ ᑲᓚᐅᔾᔨᒪᔪᓂ (2015). ᐃᖏᕐᓂᐊᓇᖅ ᐊᒻᒪ ᐃᓕᖏᓂ 2014, ᓱᓕᓕᒫᕆᓱᐊᒥᑦ ᖃᑎᖒᐊᖏᑕᐅᓕᒥᑦ ᑐᑦᑐᓂᑦ ᖃᓄᐃᑕᖏᓕᕐᓂᖏᓐᓄᑦ ᖃᐅᔨᓴᐅᑎᖃᖅᑕᐅᓗᑎᑦ ᐃᓚᖓᓕᒫᑦ ᖃᐅᔨᓴᐅᑎᐃᓐᓇᓂᑦ ᐊᒥᓱᓂ ᖃᓄᐃᑕᖏᓕᕐᓂᖏᓐᓄᑦ ᐊᑐᖃᖅᑕᐅᓗᑎᑦ (ᐳᓚᐃᓕᓴᐅᓱᒪᔪᓂ 2014). ᓄᓇᕗᒻᒥᖅ ᐊᖏᓄᓇᓕᑎᑦ ᐊᒻᒪ ᒥᑭᐅᓴᓂᐊᖏᑎᑦ ᑲᑐᔾᔨᖃᖅᑎᑦᑐᑦ ᐃᓚᓕᐅᑎᐅᓄᕐᓗᒍ ᖃᓄᐃᑕᖏᓕᕐᓂᖏᓐᓄᑦ ᐊᑐᖃᖅᑕᐅᓗᑎᑦ ᐱᓕᕆᔨᐅᑎᐃᓐᓇᓂᑦ ᐊᒻᒪ ᐱᓕᕆᔨᐅᑎᐃᓐᓇᓂᑦ ᐃᓚᖓᓕᒫᑦ ᐊᑐᖃᖅᑕᐅᓗᑎ

ᐱᓕᕆᐊᒃᑎᓂᒃ ᓇᓗᓇᐃᔨᐃᑎᐅᔨᔪᓇᖅᑐᑦ ᑲᑎᕆᓂᒃ ᓇᓗᓇᐃᔨᓯᐊᕐᓗᒪᓂᐅᕐᖏᓐᓂᑦ ᐊᖏᓪᓕᓕᒧᐊᖅᑕᐅᔨᓚᐅᓪᓕᖅᓯᓂᒃ, ᐱᓕᕆᐊᒃᑎᓂᒃᓯ 2008-2011-ᓂ ᐊᐃᒪ 1987-1994-ᓂ ᐅᔨᒥᓯᖏᓐᓄᑦᓯᓂᒃᓯᑦ ᐃᓕᖏᔪᓂᒃ ᐊᖏᕐᕋᒍᑦ ᐃᓗᐊᓂ ᓄᖅᑕᒥᓂᔪᓄᑦ ᐃᓗᐊᓂ ᐊᖏᕐᕋᒍᑦ ᐊᐅᓚᐅᖅᓯᒪᓂᒃ ᐊᐃᒪ ᓂᒥᔪᓂ ᖁᑭᖅᑕᓗᓴᐅᑦ. ᐊᖏᕐᕋᒍᑦᑐᓂᒃ ᓄᓕ ᐅᔨᒥᓯᖏᖅᑕᓚᐊᖏᔪᓂᖅᓂᒃ ᐊᖏᕐᕋᒍᐃᓂᒃ ᖃᓄᖅ (10) ᐃᓚᐃᐸᐸᑕ ᑕᒪᔭᐲᔪᓂᒃ ᐊᒥᓄᓂᒃ ᐊᐃᒪ ᐊᒥᓄᓯᓂᒃᓯᑦ ᐅᔨᓄᒪᔪᓂᒃ ᖁᑭᖅᑕᓗᓴᐅᑦ ᓇᓗᓇᐃᔨᖅᓯᒪᓂᐊᖏᔪᓂᒃ ᑕᑯᔾᔨᓂᒃ ᐃᓕᓴᐲᐸᑕᓐᓇ, ᓇᓗᓇᐃᔨᖅᑕᐅᓗᑎᒃ ᐃᓂᓕᓴᓂᐅᔪᓂᒃ ᖃᓄᖅ ᐊᖏᕐᕋᒍᑦ ᐊᑐᖅᐸᓂᖅᓂᒃ.

ᑐᑦᑐᐃᑦ ᖃᓄᐃᓕᓴᓂᖏᓐᓂᒃ ᖃᐅᔨᓴᖅᐸᓪᓕᐅᖅᑐᑦ ᖁᑭᖅᑕᓗᓴᐅᑦ ᐊᑯᓂᖏᓐᓂᒃ ᐊᖏᕐᕋᒍᐃᑦ 2012 ᐊᐃᒪ 2014 (ᔫᐊᖏᓪᓗ 2014). ᑕᕐᓇ ᐱᓕᕆᐊᒃᑎᓪᓕᐅᖅᑐᑦ, ᐱᓕᕆᐊᒃᑎᖅᑐᑎᒃ ᓄᓇᖏᐊᖏᑎᐅᑕᐃᑦ ᐅᑭᐅᖅᑕᖅᑐᑐᓂ ᑭᓇᐅᔪᖅᓂᖏᑦ ᐊᐃᒪ ᖃᐅᔨᓴᖅᑎᓕᓚᓂᖏᑦ (CARMA) ᐱᓕᕆᓵᑦ ᐱᓕᕆᐊᒃᑎᖅᑎᑦ, ᐊᖏᔪᓇᓂᕆᑎᒃ ᐱᑎᑦᓂᓕᕆᓂᑐᒃ ᓄᐊᑦᓂᔭᕐᓂᒥᓂ, ᐅᐸᓗᖓᐃᓕᖅᑕᐅᓕᕆᓂᑐᒃ ᐊᐅᓪᓚᖅᑎᑕᐅᓕᓚᐅᖅᑐᑦ ᖃᐅᔨᓴᖅᓯᐃᓚᓄᓂᒃ. ᐱᓕᕆᐊᑦᕿᒥᑦ ᐊᖏᔪᓂᒃᑐᓂᒃ ᐊᐅᓚᐅᖅᑐᑦ 2014–ᒥ ᑐᑦᑐᐃᑦ ᐊᒥᓇᓂᖏᓐᓂᓂ ᓇᓚᐅᑦᑕᐅᓇᓂᖏᓐᓂᑦ ᑐᑦᑐᐃᑦ ᖁᑭᖅᑕᓗᓴᐅᑦ ᐃᓚᐃᐸᐸᑕᑦᑐᑦ. ᒪᓴᐊᐃ ᐱᖃᓇᐅᓴᐅᐅᔨᑦ: 1) ᑲᑎᑦᑐᐃᑦ ᐊᖏᔪᑦᑕᐅᓇᓚᐊᖏᖃᓚᓕᑦᑐᔪᑦ ᓇᓗᓇᐃᔨᓯᒪᓕᕆᔨᓴᖅᓯᐊᖏᓪᓕᖅᑐᑦ ᐅᑎᑦᓇᐊᕐᓇᓂᖏᓐᓂᑦ ᑐᑦᑐᐃᑦ ᖁᑭᖅᑕᓗᓴᐅᑦ ᐊᒥᓇᓂᖏᓐᓂᓂ ᓇᓚᐅᑦᑕᐅᓇᓂᖏᓐᓂᑦ ᐊᐃᒪ 2018–ᒥ ᐅᐱᓴᖏᓕᖏᑐᒍ ᐊᐃᒪ ᖃᐅᔨᓴᖅᑎᑕᐅᓇᓚᐅᖅᑐᑦ (ᒪᓕᑕᑎᒍᑦ ᐊᐃᒪ ᒪᓕᑕᑎᐅᑦ ᓇᓚᐅᑦᑕᐅᓇᓚᐅᖅᑐᑦ), 2) ᐊᒥᓇᐃᑦ ᐊᑕᐅᓇᓴᓇᓕᑦᑐᑦ ᐊᐃᓴᐲᐸᑕᑦᑐᑦ ᓇᓚᐅᑦᑕᐅᓇᓚᐅᖅᑐᑦ ᐅᑭᐅᒥᑦ ᐊᓂᒍᓂᓚᖅᑕᐅᓇᓚᐅᖅᑐᑦ, 3) ᓇᓕᒧᐲᐸᓕᑦᑐᓂᒃ ᐊᐃᒪᓕᖅᑐᑦ ᐊᐅᓚᐅᖅᑐᑦ ᐅᑦᑕᐃᓘᖅᑐᑦ ᐱᓕᕆᐊᒃᑎᖅᑐᑦ, 4) ᖃᓄᐃᓪᓂᖏᓐᓂᒃ ᐊᓪᓂᐊᖏᖅᓯᒪᓕᕆᓂᖏᓐᓂᒍ ᑐᑦᑐᐃᑦ. ᖁᑭᖅᑕᓗᓴᐅᑦ ᑐᑦᑐᐃᑦ ᐊᐅᓪᓚᑕᐅᔨᓴᐅᑎᓪᓗᒍᓐᓇᖅᑐᑦ ᑐᑭᓯᐊᓂᖅᑐᑦ ᐊᐅᓪᓚᑕᐅᔨᓴᐅᑎᓪᓗᒍᓐᓇᖅᑐᑦ. ᑐᑭᓯᐊᓂᖅᑐᑦ ᐊᐃᒪ ᐃᓕᓴᐲᐸᑕᑦᑐᑦ ᖃᓄᐃᓕᐅᕈᑎᑦᑎᓂᒃ ᐊᐃᒪ ᐃᓕᓴᐲᐸᑕᑦᑐᑦ

7.0 ᓄᓇᕗᑦ

Advisory Committee for Cooperation on Wildlife Management (ACCWM). 2014. The Cape Bathurst, Bluenose-west, and Bluenose-east Barren-ground caribou herds management plan. Yellowknife, NT.
ᐅᖃᐅᓯᕕᒃᑐᑦ ᑲᑎᒪᔨᖏᑦᑑᓂ ᑲᑐᔪᖃᑎᒌᓂᓗᒍᑦ ᐃᒪᓕᕆᓂᖅᓯᒪᑦ ᐊᐅᓚᕋᓂᖅ (ACCWM). 2014. ᖃᓄᐃᓕᓐᓂ ᓴᓇᓀᐃᓐᓂ ᒥᑦᓴᓂᒥᐅᑕᐃᑦ, ᑐᑦᑐᔨᖅᑐᑦ ᐃᓄᒃᐸᑦᓂ ᐱᖓᓱᓀᖅᐸᓯᐊᓂ ᒥᑦᓴᓂᒥᐅᑕᐃᑦ, ᑐᑦᑐᔨᖅᑐᑦ ᐃᓄᒃᐸᑦᓂ ᑲᑎᒪᔨᖏᑦᑑᓂ ᒥᑦᓴᓂᒥᐅᑕᐃᑦ ᓀᐊᖅᑐᖅᓯᖏᑦᑐᒥ ᑐᑦᑐᐃᑦ ᑲᑎᓗᓂᑦ ᐱᓕᓐᔪᑦ ᐊᐅᓚᕋᓯᐅᑎᓂᑦ ᐸᑦᓀᐅᑎ. ᔨᐊᓗᓀᐃ, ᓄᓀᕋᐸᖅ.

Agreement, Nunavut Final (NLCA) 1993. Agreement between the Inuit of the Nunavut Settlement Area and her Majesty the Queen in right of Canada. Ottawa: Indian Affairs and Northern Development Canada and the Tunngavik Federation of Nunavut. 282 pp.
ᐊᖏᕈᑎᑎ, ᓄᓇᕗᓕᒧᑦ ᖃᖓᓕᓕᖅᐸᖅ (ᓄᓇᕗᓕᒥ ᓄᓇᑦᓴᓂᒃᑯᑦ ᐊᖏᕈᑎᑎ) 1993. ᐊᖏᕈᑎᑕᐊᖓᓕᓄᑦᑐᖅ ᐱᓕᓐᓂᑦ ᐃᓄᐃᑦ ᓄᓇᕗᒥ ᓄᓇᕐᓂᖅᓴᖏᓐᓂ ᐊᐸᑎᐊᓂᔪᓂᑦ ᐊᒻᒪ ᐃᒃᓱᓇᐃᑦᑐᖅ ᑯᐃᒪ. ᐊᑐᐢ: ᐱᓕᕆᔨᐅᑎᓕᕆᓂᑦ ᐊᒻᒪ ᐅᑭᐅᖅᑕᖅᑐᑐᒥ ᐊᒻᒪ ᑐᖅᖏᓐᓇᖅ ᑲᑐᔪᖃᑎᒌᓂᑦ ᓄᓇᕗᓕᒧᑦ. 282L<ᐱᒪᖅ.

Alaska Department of Fish and Game. 2001. Caribou management report of survey-inventory activities 1 July 1998-30 June 2000. C. Healy, editor. Project 3.0. Juneau, Alaska.
ᐊᓯᔪᓐᓇ ᐃᖅᑲᓕᑕᓴᖏᑦ ᐊᒻᒪ ᐅᒪᔨᓕᑕᓴᖏᑦ. 2001. ᑐᑦᑐᓂᑦ ᐊᐅᓚᕋᓯᐅᑎᓂᑦ ᐅᓂᒃᑲᓕᐊᑦ ᖃᐅᔨᓴᖅᑕᐅᓚᑎᒃ ᓇᓴᖅᑕᐅᓚᑎᒃ ᐊᑐᖅᑕᐅᔨᕐᓯᓂᑦ ᔨᓚᐃ 1, 1998-ᔨᓂ 30, 2000. ᓄ. ᕼᐃᓕᓕ, ᑎᑎᕆᖅᑎ. ᐱᓕᕆᐊᑦᓴᖅ 3.0. ᔨᓇᓂ, ᐊᓯᔪᓐᓇ.

Beverly and Qamanirjuaq Caribou Management Board (BQCMB). 2005. Beverly and Qamanirjuaq Caribou Management Plan, 2005-2012. BQCMB. Stonewall, MB. 17pp.
ᑎᓴᕆᐅᑦ ᒥᑦᓴᓂᒥᐅᑕᐃᑦ ᑐᑦᑐᐃᑦ ᐊᒻᒪ ᖃᒪᓂᔨᐊᖅ ᑐᑦᑐᓂᑦ ᐊᐅᓚᑦᓄᔨᑦ ᑲᑎᒪᔨᖏᑦᑑᓂ.(BQCMB) 2005. ᑎᓴᕆᐅᑦ ᒥᑦᓴᓂᒥᐅᑕᐃᑦ ᑐᑦᑐᐃᑦ ᐊᒻᒪ ᖃᒪᓂᔨᐊᖅ ᑐᑦᑐᓂᑦ ᐊᐅᓚᑦᓄᔨᓂᓗᒍᑦ ᐸᑦᓀᐅᑎ, 2005-2012. ᑎᓴᕆᐅᑦ ᒥᑦᓴᓂᒥᐅᑕᐃᑦ ᑐᑦᑐᐃᑦ ᐊᒻᒪ ᖃᒪᓂᔨᐊᖅ ᑐᑦᑐᓂᑦ ᐊᐅᓚᑦᓄᔨᓂ ᑲᑎᒪᔨᖏᑦᑑᓂ. ᓄᑖᒃᐅᐊᓗᑎ(BQCMB), ᓯᓚᑐᐸ. 17L<ᐱᒪᖅ.

Boulanger, J., M. Campbell, D. Lee, M. Dumond, and J. Nishi. 2014. A double observer method to model variation in sightability of caribou in calving ground surveys. Rangifer In prep.
ᐅᓪᓗᐃᖅᓯᒪᑦ, ᔨᑦ. ᐃᐊᓐᑦ. ᑲᑦᐱᐅᐅᓪᓗ, ᑎᑎ. ᓪᓚ, ᐃᐊᓐᑦ. ᑐᑎᓪᓗᑦ, ᐊᓐᑦ. ᓂᓯ. 2014. ᒪᖁᑉᓂᒃ ᖃᐅᔨᓴᖅᑎᖃᖅᑐᒃ ᐱᓕᓀᓗᐯ ᐅᑦᑐᑎᐅᓚᓂ ᐊᓯᔪᑦᓂᖅᑐᓂᑦ ᑕᒪᓕᕆᐅᓂᖅᓂᒃᓂᑦ ᑐᑦᑐᓂᑦ ᓄᖁᖃᖅᓴᖅᓂᒃᓂᑦ ᖃᐅᔨᓴᖅᑎᓕᓓᑦ. ᑭᐊᐃᓐᓂᖅᓴᖁ ᐅᐸᓚᓀᒪᔨᐅᑎᐅᓴᖅ.

Campbell, M. ,Goorts, J., Lee, D.S., Boulanger, J. and Pretzlaw, T. 2015. Aerial Abundance Estimates, Seasonal Range Use, and Demographic affiliations of the Barren-Ground Caribou (Rangifer tarandus groenlandicus) on Baffin Island – March 2014. Government of Nunavut Department of Environment Technical Report Series – No: 01-2015. Government of Nunavut, Department of Environment, Iqaluit, NU. 179pp.
ᑲᒪᐱᐅᓕ, ᐃᐊᒪ., ᔪᐊᖅᑕᕆ, ᔨ., ᓪᓚ, ᑎ.ᐃᐊᖅ, ᓯᓚᑐᐃᖅᓯᒪᑦ, ᔨ. ᐊᒻᒪ ᓯᖅᓱᓪᓚ, ᑎ. 2015. ᖃᓄᒪᑕᓴᑦ ᐊᒥᓱᓂᖓᓂᓂᓂ ᓇᓪᐅᑦᑕᑐᑎᑦ, ᐊᖅᑦᔪᒍᑦ ᐃᓪᓕᓄᓂ ᖃᓄᐱᑎ ᐅᓄᓱᑎᑎᔪᒥᑦ ᐊᑐᖅᐸᓂᖓᓂᓂᓂ, ᐊᒻᒪ ᖃᓄᐃᐸᑦᓴᐅᔪᑎᑎᐅᓂᓂᓂᓗ ᑲᑎᐃᒪᒪᓕᑕ ᓇᐅᑦᑕᐅᑐᑎᑦ ᐊᐅᑐᖅᐸᓂᖓᓂᓂᓂ ᑐᑦᑐᓂᑦ ᖃᓄᐃᐸᑦᓴᐅᔪᑎᑎᐅᓂᓂᓂᓗᒍᑦ

Department of Environment (DOE) 2015. Community and Hunter and Trapper Organization Consultations on Baffin Island Caribou: December 2013- January 2014. Government of Nunavut, Department of Environment, Iqaluit, NU. 42pp.
ᐊᕐᕌᒍᓕᕆᔨᒃᑯᑦ (DOE) 2015. ᓄᓇᓕᖅᓂ ᐊᒃᑐᐊᓯᔪᓐᓇᑦ ᐊᒻᒪ ᒥᑭᐊᖅᓂᐊᖅᑎᑦ ᑲᑎᒪᔨᖅᑕᐅᓂᒃ ᑐᑭᓯᒪᓕᕆᓂᖅᓂᓂ ᑐᓴᐅᒪᓕᕆᔨᒋᐊᕐᓂᖅ ᓄᓇᖃᕐᓂᐊᓕᒻᒥ ᐃᓕᓴᕐᓂᒃ: ᑎᓯᐱᐊᓗ 2013-ᑲᓱᓄᐊᓗ 2014. ᓄᓇᓕᖓᒥ ᓴᖅᓯᒃᑯᑦ, ᐊᕐᕌᒍᓕᕆᔨᒃᑯᑦ, ᐃᖃᓄᐃᑦ, ᓄᓇᓕᕐᓂ. 42ᓯᐸᐱᖃᖅ.

Department of Environment . 2013. Management Plan for Peary Caribou in Nunavut, 2014-2020. Government of Nunavut, Department of Environment, Iqaluit, NU. 37pp.
ᐊᕐᕌᒍᓕᕆᔨᒃᑯᑦ. 2013. ᐊᐅᓪᓚᕆᐊᑎᓂᑦ ᐸᕐᓇᐅᑎ ᐅᐊᖏᓇᒥ ᑐᑦᑐᓂᑦ ᓄᓇᓵᒥ 2014-2020. ᓄᓇᓕᖓᒥ ᓴᖅᓯᒃᑯᑦ, ᐊᕐᕌᒍᓕᕆᔨᒃᑯᑦ, ᐃᖃᓄᐃᑦ, ᓄᓇᓕᕐᓂ. 37ᓯᐸᐱᖃᖅ.

Department of Environment. 2013. Working Together for Baffin Island Caribou. Workshop Report (August 2013). Government of Nunavut, Department of Environment, Iqaluit, NU. 17pp.
ᐊᕐᕌᒍᓕᕆᔨᒃᑯᑦ. 2013. ᐱᓕᕆᖃᑎᒌᓂᖅ ᑭᑭᖃᑕᒻᒥᒥ ᑐᑦᑐᐃᑦ ᐱᓕᕆᔨᒃᑯᑦ. ᑐᑭᓯᐊᕈᑎᓕᕆᓂᖅᑎᑦ ᐅᓂᒃᑲᓕᐊᑦ (ᐋᒪᓗ 2013). ᓄᓇᓕᖓᒥ ᓴᖅᓯᒃᑯᑦ, ᐊᕐᕌᒍᓕᕆᔨᒃᑯᑦ, ᐃᖃᓄᐃᑦ, ᓄᓇᓕᕐᓂ. 17ᓯᐸᐱᖃᖅ.

Department of Environment. 2014. Working Together for Baffin Island Caribou. Workshop Report (November 2014). Government of Nunavut, Department of Environment, Iqaluit, NU. 34pp.
ᐊᕐᕌᒍᓕᕆᔨᒃᑯᑦ. 2014. ᐱᓕᕆᖃᑎᒌᓂᖅ ᑭᑭᖃᑕᒻᒥᒥ ᑐᑦᑐᐃᑦ ᐱᓕᕆᔨᒃᑯᑦ. ᑐᑭᓯᐊᕈᑎᓕᕆᓂᖅᑎᑦ ᐅᓂᒃᑲᓕᐊᑦ (ᔪᓐᐃᐱᐊᓗ 2014) ᓄᓇᓕᖓᒥ ᓴᖅᓯᒃᑯᑦ, ᐊᕐᕌᒍᓕᕆᔨᒃᑯᑦ, ᐃᖃᓄᐃᑦ, ᓄᓇᓕᕐᓂ. 34ᓯᐸᐱᖃᖅ.

Department of Environment. 2015a. Community and Hunter and Trapper Organization Consultations on Baffin Island Caribou (December 2013 - January 2014). Government of Nunavut, Department of Environment, Iqaluit, NU. 39pp.
ᐊᕐᕌᒍᓕᕆᔨᒃᑯᑦ. 2015a. ᓄᓇᓕᖓᒥ ᐊᒻᒪᓗᓯᔪᓐᓇᑦ ᐊᒻᒪ ᒥᑭᐊᖅᓂᐊᖅᑎᑦ ᑲᑎᒪᔨᖅᑕᐅᓂᒃ ᑐᑭᓯᒪᓕᕆᓂᖅᓂᓂ ᑐᓴᐅᒪᓕᕆᔨᒋᐊᕐᓂᖅ ᓄᓇᖃᕐᓂᐊᓕᒻᒥ ᑐᑦᑐᐃᑦ (ᑎᓯᐱᐊᓗ 2013-ᑲᓱᓄᐊᓗ 2014). ᓄᓇᓕᖓᒥ ᓴᖅᓯᒃᑯᑦ, ᐊᕐᕌᒍᓕᕆᔨᒃᑯᑦ, ᐃᖃᓄᐃᑦ, ᓄᓇᓕᕐᓂ. 39ᓯᐸᐱᖃᖅ.

Department of Environment. 2015b. HTO Consultations on Baffin Island Caribou Management Plan. March and April 2015. Government of Nunavut, Department of Environment, Iqaluit, NU. 24pp.
ᐊᕐᕌᒍᓕᕆᔨᒃᑯᑦ. 2015b. ᐊᒻᒪᓗᓯᔪᓐᓇᑦ ᐊᒻᒪ ᒥᑭᐊᖅᓂᐊᖅᑎᑦ ᑲᑎᒪᔨᖅᑕᐅᓂᒃ ᐱᓵᓕᕆᔨᒃᑯᑦ ᓄᓇᓕᖓᒥ ᑐᑭᓯᒪᓕᕆᓂᖅᓂᓂ ᑐᑦᑐᐃᑦ ᐊᐅᓪᓚᕆᐊᑎᓂᑦ ᐸᕐᓇᐅᑎ. ᐃᓕᓴ ᐊᒻᒪ ᐃᐳᑭ 2015. ᓄᓇᓕᖓᒥ ᓴᖅᓯᒃᑯᑦ, ᐊᕐᕌᒍᓕᕆᔨᒃᑯᑦ, ᐃᖃᓄᐃᑦ, ᓄᓇᓕᕐᓂ. 24ᓯᐸᐱᖃᖅ.

Ferguson, M. A. D. 1993. Working With Inuit To Study The Population Ecology Of Baffin Island Caribou. Information North. 8 pp.
ᐅᖃᐅᓯᖅ, ᐃᐊᓱ.ᐊᐃ.ᑎ. 1993. ᐱᓕᕆᖃᑎᒌᖃᑦᑐᖅ ᐃᓄᓐᓄᑲ ᖃᐅᔨᓴᖃᑕᐅᓗᑎᑲ ᐅᒪᔨᐃᑦ ᓇᔨᓴᕐᑕ ᐊᒻᒪ ᔪᓕᑎᕐᑕ ᐊᕐᕌᒍᕐᓂ ᑭᑭᖃᑕᒻᒥᒥ ᑐᑦᑐᐃᑦ. ᑐᑭᓯᐊᕈᑎᓕᕆᓂᖅ ᐅᑭᐅᖃᖅᑕᖃᑎᒌᑦ. 8ᓯᐸᐱᖃᖅ.

Ferguson, M.A.D. 1997. Arctic tundra caribou and climatic change: questions of temporal and spatial scales. Geoscience Canada, 23(4): 245-252.
ᐅᖃᐅᓯᖅ, ᐃᐊᓱ.ᐊᐃ.ᑎ. 1997. ᐅᑭᐅᖃᖅᑕᒥᐃᑦ ᓇᓵᖅᑐᖃᓂᐃᑦ ᑐᑦᑐᐃᑦ ᐊᒻᒪ ᓯᓚᐅᐳᑦ ᐊᓯᔨᐳᓪᓕᐊᓂᖅᓴᓗ: ᐊᐱᖅᓄᑎᑦ ᐊᕐᕌᒍᓕᕆᔨᓂᑦ ᐊᒻᒪ ᐃᓂᒋᔭᐅᓂᒃ ᐅᖅᓴᓯᓂᖅᓂᓂ. ᓄᓇᖃᐊᒥᑦ ᖃᐅᔨᓴᖅᑎᓪᓗᒍᑦ ᑲᓇᑕᒥ, 23(4): 245-252.

Ferguson, M.A.D. 2005. *Draft* South Baffin Caribou Management Plan 2006-2020. Government of Nunavut, Department of Environment, Iqaluit, NU. 19pp.
ᐅᖃᐅᓯᖅ, ᐃᐊᓱ.ᐊᐃ.ᑎ. 2005. ᑎᑎᕋᑕᐅᑭᐊᖅᓴᓚᐅᑎᑎᓂ ᓂᑭᖅᓴᓂ ᑭᑭᖃᑕᒻᒥᒥ ᐊᐅᓚᕆᐊᑎᓂᑦᓴᓂᒃ ᐸᕐᓇᐅᑎ 2006-2020. ᓄᓇᓕᖓᒥ ᓴᖅᓯᒃᑯᑦ, ᐊᕐᕌᒍᓕᕆᔨᒃᑯᑦ, ᐃᖃᓄᐃᑦ, ᓄᓇᓕᕐᓂ. 19ᓯᐸᐱᖃᖅ.

Ferguson, M.A.D., and Messier, F. 2000. Mass emigration of Arctic tundra caribou from a traditional winter range: population dynamics and physical condition. Journal of Wildlife Management, 64(1): 168-178.
ᐅᖃᐅᓯᖅ, ᐃᐊᓱ.ᐊᐃ.ᑎ., ᐊᒻᒪ ᒪᓂᐊᐃ, ᐃᐊᖅ. 2000. ᐊᐅᓪᓚᕆᐊᑎᓂᑦᓴᓂᒃ ᐅᑭᐅᖃᖅᑕᖃᑎᒌᑦ ᐃᓄᓐᓄᑲᓂᑦᓴᓂᒃ ᑐᑦᑐᐃᑦ ᐅᑭᐅᔨᖅᑯᑦ ᐃᓂᒋᔭᐅᓂᒃᑲᓐᓂᑕᓂᑦᓴᓂᒃ: ᑐᑦᑐᐃᑦ ᑲᑎᖃᓐᓇᓂᑦ ᖃᓄᖅ ᐃᓗᓕᖅᑐᓂᒃ ᖃᓄᐃᓕᐅᑕᐅᓂᖅᓴᓂᒃᑎᑦ ᐊᒻᒪ ᑎᒥᖅᑐᓂ ᖃᓄᐃᓕᓴᕐᓂᖅᓂᓂ. ᐅᖃᓪᓚᒪᓕᐃᑦ ᐱᓚᐅᑦᓂᒃ ᐊᐅᓚᕆᐊᑎᓂᑦᓴᓂᒃ, 64(1): 168-178.

Ferguson, M.A.D., Gauthier, L., and Messier, F. 2001. Range shift and winter foraging ecology of a population of Arctic tundra caribou. Canadian Journal of Zoology, 79: 746-758.
ᑫᕐᒍᓱᓕ, ᐃᐊᓚ.ᐊᐃᐃ.ᑎ., ᓚᑎᐅᕐ, ᐃᐊᓕᓵ. ᐊᓚᒪ ᒪᓯᐊᐃᐃ, ᐃᐊᐢ. 2001. ᓇᒧᑦ ᑰᑕᑦᔨᓂᖏᑦ ᐊᓚᒪ ᐅᑭᐅᐳᑯᓂ ᓂᖏᓯᐅᕐᓂᕐᓕᕐᑕᖏᑦ ᖃᓄᖅᓱᓂᕐᑕᖏᑦ ᑲᑎᖅᓯᓚᔨᓂ ᐅᑭᐅᖃᑕᖃᑐᑦ ᓄᑦᖢᓂᕐᑕᖏᑦ ᓄᑦᔪᑎᓕᔨᖅ ᓂᖏᔪᓕᕐᑕᖏᑦ ᑐᑦᑐᐃᑦ. ᖃᓇᑕᒥ ᐅᖃᓕᒫᓕᕐᓯᒪᔪᖅ ᓂᖏᔪᑎᓕᓴᓂᖏᑦ, 79: 746-758.

Ferguson, M.A.D., Williamson, R.G., and Messier, F. 1998. Inuit knowledge of long-term changes in a population of Arctic tundra caribou. Arctic, 51(3): 201-219.
ᑫᕐᒍᓱᓕ, ᐃᐊᓚ.ᐊᐃᐃ.ᑎ., ᐅᐃᓕᐊᓱᓕ, ᐊᖏᕐᓴ., ᐊᓚᒪ ᒪᓯᐊᐃᐃ, ᐃᐊᐢ. 1998. ᐃᓄᐃᑦ ᖃᐅᔨᒪᓕᕐᓂ ᐊᖏᕐᕿᒍᓱᓕᖅ ᐊᓯᔨᐅᑎᐅᔨᓯᒪᔪᓂᑦ ᐅᑭᐅᖃᑕᖃᑐᑦ ᓇᒃᑯᑐᖃᕐᓂᑐᑦ ᑐᑦᑐᐃᑦ, 51(3): 201-219.

Geist, V. 1998. Deer of the world: their evolution, behavior, and ecology. Mechanicsburg, PA: Stackpole Books.
ᑭᕐᑦ, ᓀ. 1998. ᑐᑦᑐᓂᐃᑦ ᓄᓇᕐᔪᐊᖏᑎᐅᑕᐃᑦ: ᐊᓯᔨᔪᓕᕐᓂᐊᓂᔨᓂᖏᑦ ᐊᖏᕐᕿᒍᓱᓕᖅ, ᐱᐅᓯᖏᑦ ᐊᓚᒪ ᓇᔪᖃᓂᑕ ᓯᓚᑎᖏᑕᓗ ᖃᓄᖅᓱᓂᕐᑕᖏᑦ ᓄᓇᕐᔪᐊᖏᑎᐅᑕᐃᑦ: ᐊᓯᔨᔪᓕᕐᓂᐊᓂᔨᓂᖏᑦ ᐊᖏᕐᕿᒍᓱᓕᖅ ᐊᓯᔨᐅᑎᐅᔨᓯᒪᔪᓂᑦ ᐅᑭᐅᖃᑕᖃᑐᑦ ᓇᒃᑯᑐᖃᕐᓂᑐᑦ ᑐᑦᑐᐃᑦ ᐱᐅᓯᖏᑦ ᐊᓚᒪ ᓇᔪᖃᓂᑕ ᓯᓚᑎᖏᑕᓗ ᖃᓄᖅᓱᓂᕐᑕᖏᑦ ᐊᖏᕐᕿᒍᓱᓕᖅ, ᓄᓇᕐᔪᐊᖏᑎᐅᑕᐃᑦ ᐊᓯᔨᐅᑎᐅᔨᓯᒪᔪᓂᑦ ᐅᑭᐅᖃᑕᖃᑐᑦ ᐃᓕᓴᕐᓂᐊᓂᔨᓂᖏᑦ ᐅᑭᐅᖃᑕᖃᑐᑦ ᐊᓯᔨᐅᑎᐅᔨᓯᒪᔪᓂ

MacPherson, AH and TH Manning. 1968. Mercy Mission; caribou for the hunters of Southampton island. Eskimo Churchill Manitoba Canada. Volume 78, 10pp.
ᒪᖅᒧᓴᓂ, ᐊᐃᕼ ᐊᖏᒪ ᑎᕼ ᐃᓂᓚ. 1968. ᓇᓕᓕᔨᖅᑐᑕᓗᓂᖅ; ᑐᑦᑐᐃᑦ ᐊᖅᓯᓇᓱᑎᓄᑦ ᓴᓕᓂᑦ ᖁᑭᖅᑕᖏᓚᓂᒥᐅᑕᓄᑦ. ᐃᐊᓱᑭᒧᐃ ᑯᔨᐊᐊᖏᓗᓂᖅ, ᐃᓇᑐᐸ ᑲᓇᑕ. ᐅᖅᑲᓕᒪᔪᐃᑦ 78, 10ᒪᐟᐱᒐᖅ.

Morneau, C., and Payette, S. 2000. Long-term fluctuations of a caribou population revealed by tree-ring data. Canadian Journal of Zoology, 78: 1784-1790.
ᒧᐊᖅᓂᐅ, ᓱ., ᐊᖏᒪ <ᐃᒋᐊᑦ, ᐃᐊᖓ. 2000. ᐊᖅᖃᒧᑉᓂᓗᓂᑕᖏᑦ ᐊᑭᐸᑎᑐᑎᓄᑦ ᑐᑦᐃᑦ ᐊᒥᓱᓂᖅᓄᓂᒃ ᓴᖅᑭᑕᐅᓱᒪᔪᑦ ᓇᐸᖅᑐᐃᑦ ᐃᓗᐊᓂᑐᑎᔪᓂᑦ ᐊᖏᒪᓗᑭᑕᐃᑦ ᐃᓗᐊᓂᑦᑐᓄᑦ ᐊᖏᒪᓗᓀᑦᑎᔪᓂᑦ ᖃᐅᔨᓴᐃᐅᑎᐃᓱᒪᔪᓂᑦ. ᑲᓇᑕᒥ ᐅᖃᓕᒫᖅ ᓂᖅᔨᒪᔪᓂᑦ, 78: 1784-1790.

Northwest Territories, Department of Environment and Natural Resources. 2011. Caribou forever – our heritage, our responsibility. A barren-ground caribou management strategy for the Northwest Territories 2011-2015. Northwest Territories, Environment and Natural Resources. 47pp.
ᓄᓇᕐᓂᐊᖅ, ᐊᐅᑎᓕᓚᐅᓂᑦ ᐊᖏᒪ ᓄᓇᒥᐅᑕᓕᓚᐅᓂᑦ. 2011. ᑐᑦᑐᐃᑦ ᑕᐃᒪᖕᓕᒫᓕᒐᖅ – ᐅᐸᑎᓵᓄᑦ ᖃᒪᑲᓂᐃᑦ, ᐃᐸᐱᑎᓚᐊᓕᓄᑦ. ᓇᐸᖅᑐᖃᖕᖏᑎᑐᑕᒥ ᑐᑦᑐᓂᑦ ᐊᐅᓕᓂᓱᑎᑕᓂᑦ ᐅᐸᓗᖕᒪᐃᐃᓂᖅ ᓄᓇᕐᓂᐊᖅᓐᒪᑦ 2011-2015. ᓄᓇᕐᓂᐊᖅ, ᐊᐸᑎᓕᓚᓂᖅ ᐊᖏᒪ ᓄᓇᒥᐅᑕᓕᓚᓂᖅ. 47ᒪᐟᐱᒐᖅ.

Porcupine Caribou Management Board (PCMP). 2010. Harvest management plan for the Porcupine caribou herd in Canada. PCMB. Whitehorse, YT. 45p.
ᐊᓯᔪᑲᐅᐸ ᐁᑲᓂᓗ ᒥᑦᓴᓂᒥᐅᑕᓂᑦ ᑐᑦᑐᓂᑦ ᐊᐅᓕᑦᓂᓂᒥᑦ ᑲᑎᒪᔨᖏᑦ (PCMB) 2010. ᐊᖏᒪᓇᓂᑐᓕᓚᓂᓪᒧᑦ ᐊᐅᓕᑦᓂᔨᐸᑎᑦᓴᑦ <ᖃᓇᐅᑎ ᐊᓯᔪᑲᐅᐸ ᐁᑲᓂᓗ ᒥᑦᓴᓂᒥᐅᑕᓂᑦ ᑐᑦᑐᓂᑦ ᑲᑎᖏᓚᔨᓂᑦ ᑲᓇᑕᒥ. ᐊᓯᔪᑲᐅᐸ ᐁᑲᓂᓗ ᒥᑦᓴᓂᒥᐅᑕᓂᑦ ᑐᑦᑐᓂᑦ ᐊᐅᓕᑦᓂᓂᒥᑦ ᑲᑎᒪᔨᖏᑦ (PCMB). ᐅᐊᐃᑦᕼᐅᐊᖏᓪ, ᐁᑲᓂᒥ. 45ᒪᐟᐱᒐᖅ

Rettie, W.J. 2010. Porcupine Caribou Herd: A review of the design of the harvest reporting program. Government of Canada, Parks Canada, Western and Northern Service Center, Winnipeg. 16pp.
ᕿᑎ, ᑐᐃᔨ.ᐋ. 2010. ᐊᓯᔪᑲᐅᐸ ᐁᑲᓂᓗ ᒥᑦᓴᓂᒥᐅᑕᓂᑦ ᑐᑦᑐᐃᑦ ᑲᑎᑌᓂᑦ: ᖃᒥᖏᑭᐊᖏᓂᖅ ᖃᓄᕐᓚᑦᓂᓚᓂᓂᑦ ᐊᖏᒪᓇᓂᓂᕐᓂᓂᒥᑦ ᐅᓂᓴᑲᕐᑐᑎᐅᐸᔨᖅ. ᒪᐃᒃᓚᓄᑦ ᑲᓇᑕᒥ, ᒥᖏᒍᐃᓇᖏᐃᓕᓚᓂᖅ ᑲᓇᑕ, ᐱᖓᓕᓇᒥ ᐊᖅᑎ ᐅᐊᓕᓇᒥ ᐱᓕᓯᓇᕐᐃᓕᓴ, ᐅᐃᓂᐸᐃᑉ. 16ᒪᐟᐱᒐᖅ

Stenton, D.R. 1991. Caribou population dynamics and Thule culture adaptations on southern Baffin Island, N.W.T. Arctic Anthropology, 28(2) : 15-43.
ᓱᑕᔭᑕ, ᑎ.ᐊᑦ. 1991. ᑐᑦᑐᐃᑦ ᐊᒥᓱᓂᖅᓄᓂᑦ ᖃᓄᖅ ᐃᓗᓕᖅᑎᑦ ᖃᓄᐃᖃᑦᑕᐅᑎᓂᓚᐊᖏᑦᑎᑦ ᐊᖏᒪ ᓱᐅᔭᓕᐊᔨᓴᒥᐅᑕᑐᑦ ᐱᓕᒪᓂᑦᑐᓄᑦ ᐃᓕᑦᑐᓂᑦᑎᓄᑦ ᐃᓕᓴᕐᓂᓂᒥᑦ ᐃᓕᓴᕐᓂᓂᒥᑦ ᐃᓕᓴᕐᓂᓂᒥ

ᐃᓕᓴᐅᔪᖅᑐᑦ 1: 2019/2024 ᖃᐅᖅᑕᓗᒍ ᑐᑦᑐᓂᒃ ᖁᓇᐃᓕᓴᐅᑎᓂᑦ <ᖁᓇᐅᑎ

2014/2015 ᐊᐅᓚᑦᔨᕐᑎᕐᓂᑦ ᖁᓇᐃᓕᓴᐅᖁᔭᓂᖅ

ᐃᖁᔨ 2014–ᒥ ᑐᑦᑐᓂᑦ ᖁᐅᔪᖁᔭᓂᐅᓚᐅᖅᑐᖅ ᓇᓗᓇᐃᖁᔨᐸᑎᐅᓚᐅᖅᑐᖅ ᖃᐅᖅᑕᓗᒍᒥᑦ ᑐᑦᑐᑭᑦᑐᓂᑦᑐᖅᑐᑦ ᓄᖅᑲᖅᓯᓚᐅᖅᑐᑦ ᑕᕝᕙᐊᐃᓐᓇᖅ ᐊᐅᓚᑦᔨᕐᑎᕐᓂᑦ ᖁᓇᐃᓕᓴᐅᓇᑦᓯᐊᓚᐅᖅᑐᑦ ᐃᓇᐅᑦ ᑭᐅᒍᑐᐃᓐᓇᐃᑦ ᐊᖏᔪᓕᕐᓂᐊᕐᓚᐅᖅᑐᑦ ᓄᖅᑲᖅᓯᓚᐅᖅᑐᑦ ᓄᓇᕗᒻᒥᐅ ᐃᓐᓇᐃᓕᓴᐅᓇᑦᓯᐊᓚᐅᖅᑐᑦ ᐊᐅᓚᑦᔨᕐᑎᕐᓂᑦ ᖁᓇᐃᓕᓴᐅᓇᑦᓯᐊᓚᐅᖅᑐᑦ ᐃᓐᓇᐃᓕᓴᐅᓇᑦᓯᐊᓚᐅᖅᑐᑦ ᓄᓇᕗᒻᒥᐅ ᐃᓐᓇᐃᓕᓴᐅᓇᑦᓯᐊᓚᐅᖅᑐᑦ ᐊᐅᓚᑦᔨᕐᑎᕐᓂᑦ ᖁᓇᐃᓕᓴᐅᓇᑦᓯᐊᓚᐅᖅᑐᑦ

ᑎᑎᕆᔨᒪᔪᓂ A1. ᑐᑦᑐᓂᓵ ᖃᐅᔨᓴᐅᑎᐅᔪᒪᔪᑦ ᐊᒻᒪ ᑐᑭᔪᑎᐊᑭᑎᐅᔪᑦ ᐱᓴᐅᔪᒪᓂᐊᖅᑐᑦ ᐊᑦᕿᒍᓄᑦ ᑕᓪᓕᒪᓄᑦ. ᖃᓄᐃᓕᓂᐅᑭᑎᐅᓚᕆᖅᑐᑦ ᒪᓕᓵᕆᖅᑐᑦ ᑭᓇᐅᕆᐃᐃᓗ ᐊᑐᖅᑐᑦᓴᓂᑦ ᐊᒻᒪ ᐊᖏᓇᔪᑦᑎᓂᑦ ᒥᑭᑎᐊᓂᐊᖅᑎᓂᑦ/ᓄᓇᓕᓵᓂᑦ ᖃᒥᑦᑭᓂᕆᑦ ᐊᒻᒪ ᐅᖃᐅᔪᐅᔪᓂᑦ. ᖃᐅᔨᒪᒋᐊᓕᓵ ᖃᑦᔪᐅᓕᓵᓂᖏᓵᓂᑦ ᓇᓕᐅᑦᑕᑭᑎᓕᐅᑦᓂᖅ ᐊᖏᑐᔪᑭᓗᒪᑦ ᐊᒻᒪ ᓄᓇᕕᒪᒥ ᒪᕼᒪᓵᔪᑦ ᐱᓕᑎᓂᓕᕆᓂᑦᑐᓗᒃ ᐅᖃᑭᓴᓇᓕᐅᑦᓂᖅᑐᓗᒃ ᐃᓕᓴᕐᓂᐊᕐᓂᖅᑐᑦ ᐊᑐᖅᑐᑦᓴᓂᑦ ᐊᑦᕿᒍᓄᑦ ᑕᓪᓕᒪᓄᑦ (5) (ᐊᑦᕿᒍᓄᑦ ᖃᓄᓕᓵᓚᓵᓂᑦ (10) ᓇᓗᓇᖅᖢᑎᑐᖅ) ᖃᐅᔨᓴᓪᓚᕆᓂᖅᑎᓵᓂᑦ ᐱᔪᓐᖃᖅᓯᓗᓂᑦ ᑭᓇᐅᔨᓄᑦ ᐊᑐᖅᑐᑦᓴᓂᑦ.

ᖃᓄᐃᓗᓯᖏᑐ	ᐱᒋᐊᖅᑎᑦᓂᔪᐃᖏᓄᑦ ᐅᓪᓗᑦᓂᔪᐅᑐᑎ				
	2019/2020	2020/2021	2021/2022	2022/2023	2023/2024
ᐅᓴᒥᓪᓂᔪᐃᓂᑎᒥᑦ ᐱᓪᓚᕐᔪᐊᑎ		**X**	**X**	**X**	**X**
ᖃᑦᓂᐅᓂᖅᑭᓂᓂᑦ ᖃᐅᔨᓴᓪᓂᖅ ᐊᐱᕐᓂᔪᓂᖅᓴᑦ	**X**	**X**	**X**	**X**	
ᓄᓇᓪᓚᓂᖅ ᖃᓄᐃᑦᑐᓂᖅᑭᓂᑦ	**X**	**X**			
ᖃᓂᒪᔪᓪᓚᓂᓂᒥᑦ ᖃᐅᔨᓴᓪᓂᖅ			**X**	**X**	**X**
ᖃᓄᑎᑎ ᐅᖓᓯᓄᓂᖃᖅᓂᖅᑭᓂᑦ ᓇᓗᓇᐃᔾᔨᐃᓂᖅ		**X**	**X**		
ᖃᑦᓂᐅᑦᓂᓂᖅᑭᓂᑦ ᓇᓪᑐᐅᑦᑕᓂᖅᑎᑦ ᖃᐅᔨᓴᓪᓂᖅ ᐊᐱᕐᓂᔪᓂᖅᑎᑦ					**X**
ᐊᖓᔪᓂᑦᑐᓪᓚᓂᓂᒥᑦ ᖃᐅᔨᓴᓪᓂᖅ			**X**	**X**	**X**

ᐃᓕᓴᐅᔪᓐᓇᖅ 2: ᒪᓴᒪᓴᓗᐊᑦ ᐊᒻᒪ ᐃᓚᓴᖏᑦ

ᖃᑭᖅᑕᓚᓐ ᑐᑦᑐᓴᓂᖅᑎᑦ ᐊᐅᓪᓚᑦᔨᐊᑎᕆᓂᑦ <ᖁᓘᐅᑎᑦᓇᖅ ᒪᓪᓕᑐᖅ ᐊᒻᒪ ᐃᑲᔪᑦᖃᑕᐅᓚᓂ ᒪᓴᒪᑎᒍᑦ ᐃᒪᐃᓴᖏᓐᓄᑦᔨᓂᑦ: ᑲᓇᑕᒥ ᓄᓇᓕᒻᒧᑦ ᐱᖁᔭᕐᕕᐊᓂᑦ (CNA), ᓄᓇᓕᓐᓂ ᓄᓇᑕᖁᓂᑦᓂᑦ ᐊᖏᑭᑐᑎᓂᑦ (NLCA), ᐊᒻᒪ ᓄᓇᓕᓐᓂ ᐅᒪᔨᓴᓴᓂᖅᑎᑦ ᐱᖁᓂᔭᕐᕕᐊᑎᒍᑦ.

ᑎᑎᖅᖁᓪᓕᔪᒥ 5–ᒥ ᓄᓇᓕᓐᓂ ᓄᓇᑕᖁᓂᑦ ᐊᖏᑭᑐᑎᒥ ᐃᓪᓕᓴᐅᑕᐅᓪᓕᔪᓂ, ᑐᕐᒪᓚᑲᐅᔪᓂ ᐊᒻᒪ ᐊᖓᓂᓚᑲᐅᔪᓂ ᓄᓇᓕᓐᓂ ᓇᓗᓇᐃᔭᖏᓪᓕᔪᑦ ᐱᔨᑎᒋᔭᐅᑎᖏᑦ ᐱᐅᖓᓇᐅᑎᒍᑦ ᐊᑕᐅᑦᓂᓄᑦ ᐊᒻᒪ ᐊᐅᓚᑦᔨᓂᑦᓗᑦ ᐃᓕᓴᐅᑕᐅᓪᓕᔪᓂ ᐊᐅᑦᓂᓄᑦ ᓄᓇᓕᓐᓂ ᐃᓪᓕᓴᐅᑕᐅᓪᓕᔪᑦ ᐊᖏᑭᑐᑎᒍᑦ ᐅᒪᔨᓴᓴᓂᖅᑎᑦ ᐊᒻᒪ ᐊᐅᖁᑎᓴᓂᖅᑎᑦ. ᓇᓗᓇᐃᖁᓪᓕᔪᑦᓴᓗᒍ, ᐃᓕᖏᑦ ᐱᒻᒪᓚᐅᓂᖏᑦ ᑎᑎᕐᖅᖁᓪᓕᔪᒥ 5 ᑎᑎᕐᖁᓪᓕᔪᑦ ᐊᑐᑎᖅᑐᑦ ᖃᑭᖅᑕᓚᓐ ᑐᑦᑐᓴᓂᖅᑎᑦ ᐊᐅᓪᓚᑦᔨᐊᑎᕆᓂᑦ <ᖁᓘᐅᑎᓕᕐᓯᒪᔪᑦ ᐱᖓᓱᐅᔪᓂᑦ:

ᐃᓗᓕᓂᐅᕐᓂᐊᖅᑐᑦ

- 5.1.1 a) “ᐅᒪᔨᐃᑦ ᐊᐳᐸᓂᖏᑦ” ᑐᐳᖅᑐᑦ ᓇᓕᓂᐊᑐᐃᓇᖅ ᐅᒪᔨᐃᑦ ᐅᕐᓂᒍᓐᓂᑦ ᓇᓕᓂᐊᑐᐃᓇᖅ ᐊᐳᐸᑦ ᐃᓚᒋᐅᓂᑦ ᐊᒥᓗᓂ ᐃᓗᓕᓂᑦᑐᑦ ᐊᐳᐸᑦᑐᓗᓂ ᓗᑦᓯ ᐊᒥᓗᐃᑦ ᐅᒪᔨᐃᑦ ᐅᕐᓂᒍᓐᓂᑦ ᐊᒥᓗᓀᓂᓂᖅᑐᓂᑦ;
- 5.1.2 e) ᐱᓂᓕᓂᐅᑎᑎᓀᐅᓇᖅᑐᑦᑕ ᐅᒪᔨᓂᑦ ᐊᐅᓚᑦᔨᓀᐱᓕᓂᑦ ᐃᑲᔪᑎᖅᑐᑦᑐᓗᓂᖅ ᐃᓄᐃᑦ ᐊᖏᑭᑐᑎᓀᐅᓇᖅᑐᑦᓂᓂᑦ ᐱᓂᓐᓇᐅᑎᒍᑦ ᐊᒻᒪ ᓀᐅᓕᓂᐅᔪᓐᓇᖅᑐᓂᑦ, ᐊᒻᒪ ᖃᐅᔨᒪᔨᐅᑎᓀᐊᖅᑐᓂᖅ ᐃᓄᐃᑦ ᐱᓂᓐᓇᐅᑎᒍᑦ ᐃᑲᔪᑎᖅᑐᑦ ᐊᐅᓚᑦᔨᓀᐱᓕᓂᑦ ᐃᑲᔪᑎᖅᑐᑦᑐᓗᓂᖅ ᐱᐅᓂᒃᑯᑦ ᐊᐅᓚᑦᔨᓀᐱᓕᓂᑦ ᐊᒻᒪ ᓴᓇᔨᐅᓂᖅᑐᑦ ᐃᑲᔪᑎᖅᑐᑦ ᐃᓄᐃᑦ ᐊᓂᒍᓐᓇᖅᑐᑦᑐᓗᓂᖅ
- 5.1.2 f) ᐱᓕᓐᓄᑦᔨᓂᖅᑐᓚᐅᓪᓕᑦ ᐅᒪᔨᓂᑦ ᐊᐅᓚᑦᔨᓀᐱᓕᓐᓂᑦᑐᓂᑦ ᐊᒻᒪ ᓄᓇᒥᑦ ᐊᐅᓚᑦᔨᓀᐱᓕᓐᓂᑦᑐᓂᑦ ᐊᖏᓪᓚᓐᖁᑎᒍᑦ ᓴᓇᔨᔨᓴᓀᖅᑐᓂᑦ ᓄᑕᖅᓯᒪᔪᖅᑐᓀᐅᓇᖅᑐᓂᖅ ᑭᓇᐅᔭᓕᖅᑐᑎᒍᑦ
- 5.1.2 g) ᐅᒪᔨᓂᑦ ᐊᐅᓚᑦᔨᓀᐱᓕᑦ ᐊᒻᒪ ᐱᓕᓐᓂᐊᖅᖁᓪᓕᔪᑦ ᐃᓄᐃᑦ ᐱᓂᓐᓇᐅᑎᒍᑦ ᐊᖏᓪᓚᓐᖁᑎᒍᑦ ᐊᐅᓚᑕᐅᔪᑦᑐᓂᑦ ᐊᒻᒪ ᒪᓕᓐᓂᐊᖅᑐᑦ ᐃᓗᓕᓂᐊᑐᐃᓕᓂᑦᑐᓗᓂᖅ ᐱᐅᓂᒃᑯᑦ ᐊᐅᓚᑦᔨᓀᐱᓕᓐᓂᑦᑐᓂᑦ

ᐱᐅᓂᖅᑎᑦᓂᖅ

- 5.1.5 ᐃᓗᓕᓂᐅᕐᓂᐊᕐᓂᑦ ᐱᐅᓂᒍᓐᓇᖅᑐᓂᑦᒧᑐᒃᓴᑦ ᐅᑯᐊ:
  c) ᒪᑭᒪᓂᓪᓂᖅ ᐊᑐᖅᑐᒃᒪᓚᐅᔪᓂᑦ, ᖃᓄᐃᖅᓯᓂᑦᑐᓂᓄᑦ, ᐅᒪᔨᓄᑦ ᒪᑭᒪᓂᓪᓂᔨᐃᑎᒍᑦ ᐊᐅᓚᑦᔨᓀᐱᓕᓐᓂᖅᑐᓄᑦ ᐊᖏᑭᑐᑎᓀᐅᓇᖅᑐᓂᑦ ᓚᓂᑐᑎᒍᑦ ᑐᓂᓯᓂᖅᑐᓂᑦ ᑐᓴᐅᒪᓂᖅᑐᑦ ᐃᓗᓕᓂᐊᑐᐃᓕᓂᑦᑐᓗᓂᖅ ᐅᑎᓪᓗᒥ; ᐊᒻᒪ
  d) ᐅᑎᖅᑎᑕᐅᓚᓂᓂᑦᑐᓂᑦ ᐊᒻᒪ ᐅᒪᓕᓂᔪᓚᑎᑕᐅᖅᑲᓂᓂᓂᑦᑐᓂᑦ ᓄᖓᑦᑐᑦ ᑐᑦᑐᐃᑦ ᐊᒥᓗᓂᖅᑐᓂᑦ ᐅᒪᔨᐃᑦ ᐊᒻᒪ ᐅᒪᔨᓄᑦ ᐃᓚᒋᐅᓂᑦ.

ᑲᑎᓴᒥᑦ ᐱᓱᐅᓂᓇᖅᑎᑕᐅᔪᑦ ᐊᖏᑭᑐᑎᓀᐅᓇᖅᑎᑕᐅᓚᐅᖅᑐᓄᑦ

- 5.6.16 ᒪᓕᓴᒥᑦ ᑎᑎᕐᖁᓪᓕᔪᑦᑐᑦ ᐅᔾᔨᓂ, ᓄᓇᓕᓐᓂ ᐅᒪᔨᐃᓕᓂᑦ ᑲᑎᒪᔨᖏᑦ ᐱᓂᓐᓇᐅᑎᖅᑎᑎᑕᐅᓂᑦ ᐊᖏᓪᓚᓐᖁᑎᒍᓐᓇᖅᑐᓂᑦ, ᐊᓂᔨᐸᐅᑎᓀᐅᓇᖅᑐᑦᓂᑦ ᐅᔾᔨᓴᓂᓐᓂᑦ ᐃᓘᖅᑕᐅᓂᖏᓐᓂᑦ, ᖃᖅᑐᒥᓂᐅᓂᖏᓐᓄᑦ ᑕᐱᒪᐃᑎᐅᖃᖅᑐᓂᖅ ᐊᒻᒪ ᐊᑐᖅᑐᐅᓂᖅᑐᓂᑦ ᐅᒪᔨᐃᑦ ᐊᒻᒪ ᐅᒪᔨᓄᑦ ᐱᓕᓐᓂᐊᖅᑐᑦ ᐃᓗᓕᓂᐊᑐᐃᓕᓂᑦᑐᓗᓂᖅ ᐊᖏᓪᓚᓐᖁᑎᒍᓐᓇᖅᑐᓂᑦ ᐊᖏᓪᓚᓐᖁᑎᒍᓐᓇᖅᑐᓂᑦ ᐅᑎᓪᓗᒥ ᐊᑐᖅᑐᐅᓂᖅᑐᓂᑦ ᐊᒻᒪ ᐅᒪᔨᓄᑦ ᐃᓚᒋᐅᓂᑦ ᐊᒻᒪ ᐅᒪᔨᓄᑦ ᐃᓗᓕᓂᐊᑐᐃᓕᓂᑦᑐᓗᓂᖅ
- 5.6.17 ᑲᑎᓴᒥᑦ ᐱᓱᐅᓂᓇᖅᑎᑕᐅᔪᑦ ᐊᖏᓪᓚᓐᖁᑎᒍᓐᓇᖅᑎᑕᐅᓚᐅᖅᑐᓄᑦ ᐃᓚᒋᐅᑦ ᐊᒥᓗᑦ ᐅᔾᔨᓴᓂᓐᓂᑦ ᑐᓴᐅᒪᓂᑦ ᐊᒥᓗᐃᑦᑐᑦᓂᓂᑦ ᓇᓗᓇᐃᔭᖏᓪᓕᔪᓀᐅᓇᖅᑐᓂᑦ ᓇᓕᓐᓂᐊᖅᑐᑦ... a) ....ᓄᓇᓕᓐᓂᑦ ᑲᑎᓴᒥᑦ ᐱᓱᐅᓂᓇᖅᑎᑕᐅᔪᑦ ᐊᖏᓪᓚᓐᖁᑎᒍᓐᓇᖅᑎᑕᐅᓚᐅᖅᑐᓄᑦ, ᐊᒻᒪ...b)... ᓄᓇᓕᓗᒍᓂᓐᓂᖅᑐᓂᑦ ᑲᑎᓴᒥᑦ ᐱᓱᐅᓂᓇᖅᑎᑕᐅᔪᑦ ᐊᖏᓪᓚᓐᖁᑎᒍᓐᓇᖅᑎᑕᐅᓚᐅᖅᑐᓄᑦ.
- 5.6.20 ᓇᓕᐅᑦᑕᖅᑕᐅᓪᓚᓂᑦ ᐃᓚᒋᖅᑕᐅᓪᓚᒪᑐᓴᐅᒪᖅᑐᓄᑦ ᐱᓂᓐᓇᐅᑎᖅᑎᑎᑕᐅᓂᑦ ᐊᖏᓪᓚᓐᖁᑎᒍᓐᓇᖅᑎᑕᐅᓚᐅᖅᑐᓄᑦ. ᑲᑎᓴᒥᑦ ᐱᓱᐅᓂᓇᖅᑎᑕᐅᓚᐅᖅᑐᓄᑦ ᐊᖏᓪᓚᓐᖁᑎᒍᓐᓇᖅᑎᑕᐅᓚᐅᖅᑐᓄᑦ ᐊᐳᖅᖃᖅᑐᓂᖅᑐᑦᓂᓂᑦ ᐅᔾᔨᓴᓂᓐᓂᑦ ᑐᓂᓯᓂᖅᑐᓂᑦᓂᓂᑦ, ᐱᓱᐅᓂᓇᖅᑎᑕᐅᓚᐅᖅᑐᓄᑦ ᐃᓗᓕᓂᐊᑐᐃᓕᓂᑦᑐᓗᓂᖅ ᐅᓗᓕᓂᐊᑐᐃᓕᓂᑦᑐᓗᓂᖅ ᐃᓚᒋᐅᖅᑐᓂᑦ ᐊᖏᓪᓚᓐᖁᑎᒍᓐᓇᖅᑎᑕᐅᓚᐅᖅᑐᓄᑦ.

ᓇᓗᓇᐃᓚᐊᖅᑐᓂᖅᑐᓄᑦ ᐃᓚᒋᐅᓂᓪᓗᑦ ᐱᔨᑎᒍᑦᓴᑦ ᑎᑎᕐᖁᓪᓕᔪᖅᑐᓂᑦ, ᖃᖅᑐᒥᓂᐅᓚᐅᖅᑐᑦ ᐊᖏᓪᓚᓐᖁᑎᒍᑦ <ᖁᓘᐅᑎᓀᐅᓇᖅᑐᓂᑦ ᐊᑐᖅᑐᐅᓂᖅᑐᓂᑦ ᐃᓄᐃᑦᑐᓂᑦᑐᓂᑦ ᐃᓕᓴᐅᑕᐅᓪᓕᔪᑦᑐᓂᑦ ᒪᓕᓐᓂᑦ ᐊᑐᖅᑐᐅᓂᖅᑐᓂᑦ ᐃᓕᓴᐅᑕᐅᓪᓕᔪᑦᑐᓂᑦ ᐊᒻᒪ ᐅᒪᔨᓄᑦ ᐊᖏᓪᓚᓐᖁᑎᒍᑦ ᖃᖅᑐᒥᓂᐅᓚᐅᖅᑐᑦ ᐊᒻᒪ ᐅᒪᔨᓄᑦ ᐱᓂᓐᓇᐅᑎᒍᑦ ᐊᖏᓪᓚᓐᖁᑎᒍᓐᓇᖅᑎᑕᐅᓚᐅᖅᑐᓄᑦ.

Consultations with Hunting and Trapping Organizations on the Baffin Island Caribou Composition Survey Results, Future Research Recommendations, and Draft Management Plan

January 7-18, 2019 and May 27, 2019

ᐊᖏᑎᓕᓱᐅᑦ, ᓄᓇᕗᑦ ᑲᖏᒪᑦᖁᑎᑦ, ᐃᖃᓗᐃᑦ, ᓄᓇᕗᑦ

ᐊᐅᓪᓚᑎᓂᖅᓯᓐᓄᖅᑐᑦ ᓇᐃᓐᓇᕐᓂᒪᓕᕆᑦ

ᓄᓇᕗᑦ ᑲᖏᒪᒃᑯᖓᓂ, ᐊᖏᑎᓪᓗᓵᓴᒃᑯᑦ ᑭᖑᓕᖅᑐᓄᖅᑎᖓᓂ ᑐᓴᕐᓯᓚᐅᕐᖓᓪᓚᐃᓐᓇᖅᑐᑦ ᖃᐅᔨᓴᖅᑭᑦ ᐅᓇᐊᓗ 7-18, 2019 ᐊᒻᒪᓴ ᒪᐃ 27, 2019 ᑲᖏᖅᑐᐃᒪᐃᖅᑭᑦ ᓂᓪ ᐱᕐᓂᑎᑦᓴᒍ ᐅᓇᐊᓗᑦ.

ᐱᕐᓯᑕᐅᓪᓚᕆᖅᑐᖅ ᑐᓴᕐᓂᖅᓱᑦ ᑕᒃᑯᐊ ᐊᖏᓇᓂᖅᑐᓪᓚᓵᓴᒃᑯᑦ ᑐᓴᐅᒪᕐᐊᕐᑯᓪᓕᑦ ᖃᒃᐅᓵᓴᕐᑎᐊᓂᖅᓯᓐᓄᑦ ᑲᑎᑦ ᐊᒥᓱᓂᖅᑐᓂᑦ ᑲᑎᒃᓴᖅᑎᐊᑦᑕᐅᓯᒪᔪᓪᓗ ᖃᒃᐅᓵᓴᕐᑎᐊᑦᑕᐅᓚᑦ 2014–ᒥᑦ ᐅᓪᓗᒥᓗᑦ ᖃᐅᖅᑕᒍᖏᑦ. ᐊᖏᑎᓪᓗᓵᓴᒃᑯᑦ ᓴᖅᑭᕕᑎᓪᓚᐅᖅᓴᑦ ᓇᓂᕋᕕᑎᓂᖅᓴᑦ ᓯᐅᑎᓂᓕᕆᓂᑦ ᖃᒃᐅᓵᓴᖅᓱᖏᓪᓗᑦ ᖃᐅᖅᑕᒍᖏᑦ ᐃᓕᐅᓪᓗᓂ ᖃᒃᐅᓵᓕᔪᐊᑎᒃᓴᖅᑐᓂᑦ–ᖁᓱᖓᓂᓯᓱᓄᑦ ᐱᓕᓚᐊᒃᓴᑦ. ᑐᓴᓕᐅᖅᑕᐅᓴᑦ ᑐᓴᕐᓂᒃᑎᓪᓗᓕ ᑕᒪᑐᒪᓕ ᐃᒃᔨᑎᑎᕆᐅᓂᐊᖅᓴᖅ ᓄᓇᕗᑦ ᑲᖏᖅᑐᓄᖅᑎᖓᓂᑦ ᓴᖅᑭᕕᑎᓂᖅᓱᖏᓪᓗᑦ ᐸᓴᓇᐃᓂᖅᓱᑦ ᖃᒃᐅᓵᓴᖅᑎᐊᑦᑕᐅᓯᒪᔪᓪᓗ ᖃᒃᐅᖅᑕᒍᓕᔨᑦ ᑐᓴᐅᒪᕐᐊᕐᑯᓪᓕᑦ.

ᐅᓇ ᐅᓂᒃᑲᖅ ᓇᐃᓚᓕᑎᐊᓕᕆᑦ ᐅᖃᒃᐅᓕᐅᓕᐅᖅᑐᓂᑦ ᐃᓕᐅᔨᓄᑦ ᑐᓴᕐᓂᒃᑎᓪᓗᓕ.

ᐱᑭᐊᖅᓂᖅ

ᐅᓇ ᐅᓂᒃᑲᖅ ᐊᐅᑎᑕᓱᓕᕆᔨᑦ ᐱᓇᔪᑎᐊᑭᖅᓇᓂᒋᓕᖅᑭᓂᑦ ᑕᓪᓕᖅᓭᒪᖅᑕᑐᓂᑦ ᐃᓕᑕᐅᕆᐃᓇᓱᓭᓪᓕᔪᑦ ᑐᓴᕆᐊᖅᑎᓕᔪᒋᓕᖅᑐᓂᑦ ᑐᓴᕆᓭᓕᐊᖅᑎᓕᖅᑐᑦ ᑲᑎᓕᖅᑎᖅᑲᓕᑎᒃ ᐊᖏᔪᓱᐅᑕᓱᓕᕆᔨᒪᓂᑦ ᑭᒪᑭᑭᑦ, ᖃᑭᖅᑕᖅᔪᐊᖅ, ᐸᖅᓂᒃᑐᖅ, ᐃᖅᑲᐃᐃᑦ, ᑭᖅᑲᐃᒪᒃ, ᐃᓗᓕᓕᒃ, ᐃᒃᐱᐊᖅᔪᒃ ᒥᑎᑕᒪᓕᕆᓂᖅᒥᓕᒫᓂ. ᐊᐃᑦᑕᖏᓚᑲᐊᖅᓕ, ᑕᒪᑐᒪᓂ ᑐᑭᓯᓕᕆᓂᑕᒥᑦ ᓯᓚᒧᑦ ᑲᑎᓪᓗᒍᑭᓕᕆᓂᖅᑲᐅᓕᐅᖅᑐᑦ ᑲᖏᔪᓂᑦᑐᐅᓚᐅᖅᑎᑦᑎᓂᖅ ᐊᖏᔪᓱᐅᑕᓱᓕᕆᔨᒪᓂᑦ ᒪᐃ 27, 2019.

ᐃᓯᒪᑎᖃᖅᑎᑦ ᓴᖅᑭᑕ ᐅᕐᓂ ᐊᖃᐸᑐᐊᒪᓗᒃᑖᖅᑎᓪᓗᑦ ᐊᐅᑎᑕᓱᓕᕆᔨᑦ, ᓄᓇᕗᒻᒥᓐᓄᑦ ᑲᕐᒪᓕᕆᑦ.

ᐃᓗᓕᖅᑎᑦ

ᐊᐅᓚᕋᐱᓂᖅᓯᒪᔨᖅᑐᑦ ᓇᐃᓐᓇᖅᓯᒪᔪᑦ i

ᐱᒋᐊᖅᓂᖅᓱᒃ ii

ᐃᓗᓕᖅᑎᑦ iii

1.0 ᐅᓂᒃᑲᑦ ᐱᓱᒋᑦ ᐊᑦᑭᓴᓯᒪᓂᖅᓱᓗᓐ 4

2.0 ᑐᐸᕋᓯᖅᓂᒍᑦ ᐱᓱᒋᑦ 4

2.1 ᑲᑎᒪᓕᕐᑦ ᐊᑦᑭᓴᓯᕋᓯᒪᓂᖅᑎᑦ 4

3.0 ᐅᓂᒃᑲᐅᓯᐅᒋᑦ ᓄᓇᓕᒃᑎᑐᑦ 5

3.1 ᑭᒡᒋᐅᑦ 5

3.2 ᖃᑭᖅᑕᖅᖢᐊᖅ 6

3.3 ᕿᓂᓂᖅᑐᖅ 7

3.4 ᐃᖅᑲᓗᐃᑦ 9

3.5 ᑭᖏᓚᐃᑦ 10

3.6 ᓴᓂᕆᔨᒃ 12

3.7 ᐃᓐᓗᓕᓄᒃ 14

3.8 ᐃᒃᐱᐊᖅᖢᒃ 15

3.9 ᒥᕆᑎᒪᑕᓄᒃ 17

3.10 ᑲᖏᕐᓂᖅᑐᐃᓚᐱᒃ 18

4.0 ᓇᐃᓐᓇᖅᓯᒪᓂᖅᓱᓗᓐ 19

5.0 ᐅᐃᒍᐊᑭᓯᖅ 1 21

1.0 ᐅᓂᒃᑯᓪ ᐱᓱᒋᑦ ᐊᖏᕈᒪᓂᖅᓱᓗᓐ

ᐅᓇ ᐅᓂᒃᑯᖅ ᐊᖏᕈᒪᔪᖅᓯᒪᔪᖅ ᓇᐃᐅᓐᖏᓕᒫᓂᓗᑎᓗ ᐅᖏᒃᐅᓯᐅᖏᒃᑕᑕᒪᐅᖅᑐᑦ, ᐊᐱᖅᑯᑎᑦ, ᐃᓯᒪᓗᑎᑎᕐᓂᑦ, ᖃᒃᓄᐃᑦᐅᑕᒪᐊᖏᔪᓂᖅᓗ ᐊᖅᙱᓚᓯᑎᑦᓂᑦ ᑭᐅᑦᓗᑎᒃ ᓇᐃᐅᓐᖏᓕᒫᓂ 2015-2018 ᑲᑎᕋᖅᓯᒪᔪᓄᑦ ᖃᒃᐅᔨᓴᕈᑎᐃᓂᖅᓄᑦ, ᑐᒃᑐᓂ ᖃᒃᐅᔨᓴᕈᑕᐅᖅᑯᑐᑦ, ᒎᓐᓇᓗ ᔪᑲᑦᑕᖅᓯᒋᑦ ᑎᑎᕋᖅᑕᐅᖅᓯᒪᓂᖅᑐᑦ ᑐᑭᖅᑕᓗᓐᓂᑦ ᑐᑭᑐᖅᑯᓗᒃᑐᑦ ᑐᑭᓯᐅᑎᓗᒃ ᑐᓴᕐᓂᖅᓯᒪᓂᖅᑐᑦ ᑐᑭᑐᖅᑯᓗᒃᑐᑦ ᑐᑭᓯᐅᑎᓗᒃ

ᖃᐅᔨᓴᕈᑎᐃᓕᓐᓂᓕ ᖃᑭᖅᑕᕐᓗᓐᓂᒃ. ᑲᒪᐸᓗᑦ ᓴᖅᑭᕈᑕᐅᓚᐅᕐᖓᖅ ᑐᓴᑐᓕ ᖃᐅᔨᓴᖃᓐᓂᕐᒧᑦ ᐱᓕᕆᐊᖓᓕᓂ ᑭᓚᓕᓐᓂᒃᑎᑦ, ᖃᓄᖅᑕᓱᓕᒃᑯᑦ ᓇᐃᓴᐃᓂᓕ (ᐊᒥᓱᓕᓐᓂᑲᓐᓇᑦ), ᑲᑎᓱᒪᔪᑦ ᖃᐅᔨᓴᖃᑎᑦ ᐊᒻᒪᓗ ᐅᖃᓪᓚᕆᑦ ᖃᐅᔨᓴᖃᑎᓕᓂ. ᐅᖃᐅᓪᓚᖅᒪᑦ ᓄᖅᓄᖅᓱᐅᐃᓇᒪᔨᐅᖅᑐᑦ ᐊᐱᖅᑯᑎᐅᓴᓂ. ᓇᖃᑦᓴ ᓇᐃᔅᑐᒥᑦ ᓇᓗᓇᐃᖏᐃᓪᒪᔨᐅᖅᑐᖅ ᓱᓕᑦᓭᓂᖅᓯᒪᔨᐅᑎᓄᑦᓱᖅᑐᒥᑦ ᖃᑭᕈᓴᑦ ᑐᓴᖅᑭᕐᓂᓕ ᐊᐅᓪᓚᐅᑎᓂᒃ ᐸᖃᓇᐅᑎᒥᑦ ᑐᓴᖅᑭᕐᓂᐊᓚᐅᖅᑐᓂᓗ ᐃᓱᒪᒋᔭᐅᓂ, ᐃᓱᐊᓕᑕᐅᔨᓂᒃ ᐊᐱᕐᓇᑎᑕᐅᑎᓪᓗᒥᒃ ᐸᖃᓇᐅᑎᐅᑦ ᑐᓴᖅᑭᕐᓂᐊᓚᐅᖅᑐᓂᒃ ᐅᓂᒃᑳᖅᑐᓂᒃ ᐊᐱᖅᑯᑎᐅᓴᖅᑭᕐᓂᐊᓚᐅᖅᑐᓂᒃᑐᓂᖅᓱᓂᒃ ᐊᐱᖅᑯᑎᐅᓴᖅᑭᕐᓂᐊᓚᐅᖅᑐᓂᒃᑐᓂᖅᓱᓂᒃ ᐊᐱᖅᑯᑎᐅᓴᖅᑭᕐᓂᐊᓚᐅᖅᑐᓂᒃ ᐃᓕᑦᖃᓄᖅ ᐅᓴᖅᑭᕐᓂᐊᓚᐅᖅᑐᓂᒃ ᐊᐱᖅᑯᑎᐅᓴᖅᑭᕐᓂᐊᓚᐅᖅᑐᓂᒃᑐᓂᖅᓱᓂᒃ ᐃᓱᒪᒋᔭᐅᓂ ᐊᐱᖅᑯᑎᐅᓴᖅᑭᕐᓂᐊᓚᐅᖅᑐᓂᒃ ᐅᓴᖅᑭᕐᓂᐊᓚᐅᖅᑐᓂᒃᑐᓂᖅᓱᓂᒃ ᐊᐱᖅᑯᑎᐅᓴᖅᑭᕐᓂᐊᓚᐅᖅᑐᓂᒃ ᐃᓱᒪᒋᔭᐅᓂ

ᐅᖅᑲᐅᓯᕐᓴᐅᑦ ᐊᐱᖅᑯᑎᕐᓴᐅᑦ:

ᐊᖏᓚᓂᑐᑕᓵᓴᕆᔨᑦ ᐃᓱᒪᑎᒃᑲᒪᐅᖅᑐᑦ ᐋᖅᑭᖅᓯᒪᓂᖅᓯᒋᐊᓂᑦ ᑲᑎᒪᔨᖅᓯᒪᔪᓂᑦ ᖃᐅᔨᓴᕿᑎᐃᓇᓂᖅ ᑕᐃᖅᑯᐊᓗ <ᓴᕿᑎᐅᓯᒪᔪᑦ ᑐᓴᕋᓪᓕᐅᖅᑐᑦ >ᖅᑭᑐᑉᐳᖅ ᖁᑭᖅᑕᕐᓗᓂᕐᒥᑦ. ᐊᖏᓚᓂᑐᑕᓵᓴᕆᔨᑦ ᐃᓱᒪᓕᐅᖅᑐᑦ ᖃᐅᔨᓴᖅᓯᐃᓂᖅ ᓯᐅᓂᕿᑎᖅᓯᐃᓂᖅ ᐊᑐᖃᑕᐅᓴᖅᓯᓐᓂᑦ ᐊᒻᒪᓗ ᑐᓴᕋᓯᓯᒪᓂᖅᓱᑦ ᐱᔭᐊᑎᖅᑲᓕᐅᖅᓯᒪᓂᖅᓱᑦ ᖃᑎᒌᑦ ᐊᖏᓚᖅᑕᐅᔪᓐᓇᖅᑐᑦ. ᐃᓱᒪᓗᑎᒃᑲᒪᐅᖅᓯᒪᔪᑦ ᐱᓕᕆᔨᑦᑐᖅᑐᑦ ᖃᐅᔨᓴᖅᑎᑦᓯᐊᕐᓂᖅ ᐊᐅᓚᑦᓯᐊᕐᓂᖅ ᐊᓕᖅᑐᑦ ᐊᖏᓚᖅᑕᐅᔪᓐᓇᖅᑐᑦ ᐃᓕᓴᕐᓂᖅ ᐊᐅᓚᑦᓯᐊᕐᓂᖅ ᐊᖏᓚᖅᑕᐅᔪᓐᓇᖅᑐᑦ ᐃᓱᒪᓕᐅᖅᑐᑦ ᐊᐅᓚᑦᓯᐊᕐᓂᖅ ᐃᓱᒪᓕᐅᖅᑐᑦ ᐊᐅᓚᑦᓯᐊᕐᓂᖅ ᐊᖏᓚᖅᑕᐅᔪᓐᓇᖅᑐᑦ

- ᓄᓇᕗᑦ ᑐᖓᓯᖏᓐᓄᑦ, ᐃᓕᓴᕐᕕᖏᓐᓄᑐᖃᓕᕐᓂᖅᓯᓐᓴᑦ ᐊᐅᓚᑦᑎᓂᖅᓴᑦ ᐅᓕᖅᐅᓵᕐᕕᓐᓄᑦ; ᓱᐅᑭᓪ ᕿᐃᓪᓓᑦ
- ᓄᓇᕗᓴᒥ ᐅᒪᔨᓚᓕᔪᕐᕕᖏᓐᓄᑦ ᑲᑎᒪᔨᖅᑭᑦ, ᐅᒪᔨᓂᖅ ᓇᐅᔨᓴᓇᖅᑎ ᑲᐃᐅᓪ ᓚᑭᓱ
- ᓇᑉᑉᑕᓂ ᐅᒪᔨᓚᓕᔪᐅᓂᖅᓯᓐᓴᑦ, ᐅᒪᔨᓚᓕᓂᕐᓴᑦ ᐅᓇᐅᕐᓇᕐᐃᓐᓇᖅ: ᒪᐃᓂᓪ ᕿᖓᓴᓵᖅ
- ᑐᐊᔪᓐ, ᑕᐃᑐᓱ ᐊᓴᓇᑦᓪᓚᖅ
- ᓇᑎᐅᒃ ᐊᖓᓇᓱᑐᑕᓕᔪᐅᓂᖅᓯᓐᓴᑦ ᑲᑎᒪᔨᖅᑭᑦ
    - ᐋᓪᓚᔪᓵ ᑯᓯᑦ
    - ᕐᐃᑕ ᑯᓱᐊᒃ
    - ᓗᐊᓱ ᐊᓕᓇᖅᑐᓇᖅᑐᓇᖅ
    - ᐅᓴᐃᐱ ᐊᐅᓚᕿᐊᖅ
    - ᕐᓇᓇ ᕿᓂᓱᒋᖅ
    - ᓴᐃᐱᑎ ᓄᕿᖓᓚᖅ
    - ᕐᕉᐃᔭ ᓄᕿᖓᓚᖅ

ᐅᓇᐅᓱᓴᐃᑦ ᐊᐱᖅᑯᑎᓴᐃᓪᓗ:

ᐊᖓᓇᓱᑐᑕᓕᔪᐅᓂᖅᓯᓐᓴᑦ ᐅᓇᐅᓱᓇᖅᑐᓚᐅᖅᑐᑦ ᐊᒥᓱᐅᔪᓂᒃᑯᓪᓕᐊᔪᓂᑦ ᐅᓴᓕᓱᒪᔭᖏᑐᓂᑦ ᐱᑎᐊᖏᓂᑎᓂᑦ 2000. ᐃᓱᒪᓕᑕᐅᓚᐅᖅᓴᑦ ᐊᒥᓱᓂᖏᑦ ᐊᒪᖅᑯᑦ ᑕᑯᐅᓱᒪᔨᐅᑦ ᓇᐃᔨᐃᑎᑦᓕᑎᑦ ᐊᒥᓱᓵᖏᑐᓂᖏᑦᓕᓗ ᑐᖓᑐᐃᑦ ᑕᐅᙵᓂ. ᐊᙵᑎᑦᓕᓕᔪᖅᑯᑦ ᓇᓗᓇᐃᔭᓱᓚᐅᖅᑐᑦ ᐅᓴᓕᓚᐅᖅᑕᖏᑐᓂᑦ ᖃᓴᓱᓇᐃᑦ ᐊᒪᖅᑯᑦ ᓇᐃᔨᐃᑎᑦᓕᑎᑦ 2012–ᒥᑦ ᐅᑦᓗᒥᓱᑦ ᐊᖏᑐᐃᓗᐊᔭᓱᐅᑦᓭᓇᑎᓐᓗ ᐊᒪᖅᑯᐃᑦ ᐃᓐᓇ. ᐊᖓᓇᓱᑐᑕᓕᔪᐅᓂᖅᓯᓐᓴᑦ ᑐᓴᖅᑎᑎᓚᐅᖅᑐᑦ ᑐᖓᐃᑦ ᓇᒡᓐᓂᖏᓂᓂᑦ ᑭᖓᓐᓂᓂᑦ ᐊᙵᒋᓂᑦ ᖃᑉᑉᑕᓕᕐᔪᐊᑦ ᐊᒪᒪᓕᓂᑕᐅᖅᑐᖅ ᐊᖓᓇᓱᓕᐃᐅᖅᑎᖅᑕᖏᑐᑐᖅᓯᒪᓕᖅ.

ᐊᖓᓇᓱᑐᑕᓕᔪᐅᓂᖅᓯᓐᓴᑦ ᐅᖅᑎᐅᓱᖅᑐᓚᐅᖅᑐᑦ ᓇᒃᑕᖅᓚᐃᐅᙵᖅᑐᑦ ᑐᖓᐃᓂᑦ ᖃᑉᑉᑕᒡᓗᓐᓂᒥᑦ. ᖃᑉᑉᑕᓕᑕᓂ ᐅᒪᔨᓚᓕᔪᕐᕕᖏᓐᓄᑦ ᑐᓴᖅᑎᑎᕐᑕᓚᐅᖅᑐᖅᓯᓐᓴᑦ ᑐᓴᐅᒪᔪᑐᖅᑐᓂᖅᓯᓐᓴᑦ ᐅᓴᖅᑎᐅᓱᖅᑕᖏᑐᓂᑦ ᐃᓇᐅᑦᓂᓂᑦ. ᖃᑉᑉᑕᓕᑕᓂ ᐅᒪᔨᓚᓕᔪᕐᕕᖏᓐᓄᑦ ᑐᓴᖅᑎᑎᕐᑕᓚᐅᖅᑐᖅᓯᓐᓴᑦ ᐃᓇᐅᑦᓂᓂᑦ ᐃᓇᐅᒪᔪᑐᖅᑐᓂᖅᓯᓐᓴ

- ᓄᓇᕗᑦ ᑲᖏᒪᒃᑯᐊᓂᓪᓗᐊᖏᑎᓕᕐᓂᒃᑯᑦ, ᖁᑭᖅᑕᕐᓗᓐᓂᑦ ᓂᕆᔨᑎᓂᒃ ᖃᐅᔨᓴᖅᑎᑐᖅ: ᐅᓐᓄᓐᓴᓕ
- ᓄᓇᕗᑦ ᑲᖏᒪᒃᑯᐊᓂᓪᓗᐊᖏᑎᓕᕐᓂᒃᑯᑦ, ᖁᑭᖅᑕᕐᓗᓐᓂᑦ ᓂᕆᔨᑎᓂᒃ ᖃᐅᔨᓴᖅᑎᑐᖅ: ᒥᓵ ᐅᒪᐳ
- ᓄᓇᕗᑦ ᑲᖏᒪᒃᑯᐊᓂᓪᓗᐊᖏᑎᓕᕐᓂᒃᑯᑦ, ᖁᑭᖅᑕᕐᓗᑦ ᓂᕐᐳᑎᓂ ᐊᐅᓪᓚᕆᔨᐅᑲᐃᓐᓇᖅᑐᓂᖅᓯᒪᓕᖅᑐᖅ: ᐊᑦᑕᐊ ᐅᐃᓕᐊᓴ
- ᓄᓇᕗᑦ ᑐᖓᓴᕐᓂᒃᑯᑦ, ᐃᖅᑲᐅᓇᐃᕐᖅᑐᑎᓂᓪᓗᑦ ᐊᐅᓚᕆᑎᓂᓪᓗᑦ ᐅᖅᑲᐅᕐᔪᐃᐅᓐ; ᓱᐅᑦ ᑭᐃᓕᓴ
- ᓄᓇᕗᓴᒥ ᐅᒪᔨᓪᓗᖅᓯᐊᖅᑯᑦ ᑲᑎᒪᔨᖏᑦ, ᐅᒪᔨᓂᖅ ᖃᐅᔨᓴᖅᑎᑐᖅ ᑲᐃᐅᑦ ᓇᕆᓱ
- ᖁᑭᖅᑕᓂ ᐅᒪᔨᓪᓗᖅᓯᐊᖅᑯᑦ ᑲᑎᒪᔨᖏᑦ, ᐅᒪᔨᓂᓪᓗᑦ ᐅᖅᑲᐅᕐᔪᐃᐅᓐ: ᒪᐃᔪᑦ ᑭᖑᓴᓕᓕ
- ᑐᓴᓕ, ᑕᐃᑐᓱ ᐊᖅᓇᑲᑦᓂᒃ
- ᐸᖕᓂᖅᑐᒥᑦ ᐊᖓᔪᓱᖅᑐᑕᓪᓗᐊᖏᑎᓕᕐᓂᒃᑯᑦ ᑲᑎᒪᔨᖏᑦ
    - ᕆᐃᕕᑎᑎ ᓇᐅᔨᖅ
    - ᓴᓂ ᒪᐃᒃ
    - ᐃᒃ ᖁᓕᖕᒫᖅ
    - ᔨᐊᕐᓴ ᖃᑲᖅᑲᓱᖅ
    - ᐱᓕ ᐃᑐᐊᖕᖓᑦ
    - ᑲᓕ ᖃᑲᐱᒃ
    - ᐸᑎ ᖁᓕᖕᒫᖅ

<u>ᐅᖅᑲᐅᓱᐊᕆᐃᑦ ᐊᐱᖅᑯᑎᐊᕆᐃᑦᓗᓐ:</u>

ᐊᖓᔪᓇᓱᐊᓕᓪᓗᓕᕐᑯᑦ ᐃᓱᒪᑎᓴᖅᑲᓇᐅᖅᑐᑦ ᐊᒥᓱᓂᖓᑎᑦᑕ ᓇᓕᐅᑦᑕᖅᑕᐅᓱᒪᔨᕐᑦ 2014–ᒥᑦ, ᓇᓕᐅᑦᑕᖅᑕᐅᓱᒪᔨᕐᑦ ᓄᖅᓀᐅᖏᓂᖓᓂᑦ ᒪᓕᓚᓐᓂᑎᒃ ᑲᑎᓴᓱᖅᓱᒪᔨᔪᓄᑦ ᐃᓕᑎᓴᐅᔨᒪᓂᖏᓂᕐᒥᓂᓗ ᓄᓇᕗᑦ ᑲᖏᒪᒃᑯᑎᑦ ᓇᐃᔾᐃᑎᑎᓕᖅᑐᓄᑦ. ᐊᖓᔪᓇᓱᐊᓕᓪᓗᓕᕐᑯᑦ ᐃᓱᒪᑎᓴᖅᑲᓇᐅᖅᑐᑦ ᑕᒪᐃᔨᓂᖅᓱᒪᔨᔪᓄᑦ ᐃᓚᖓᔪᖅᑐᓂᖏᓂᕐᒥᓂᓗ ᐃᓚᖓᔪᖅᑐᓂᖏᓂᕐᒥᓂᓗ ᐃᓱᒪᑎᓴᖅᑲᓇᐅᖅᑐᑦ

ʕᑯᖏᓚᓬᐅᐃᑦ ᐃᑲᓴᖅᑐᖅᑕᐅᓇᕈᑎᑲᒃ ᐊᖏᔪᓇᓴᑐᓕᓂᓱᐃᓐᓄᑦ ᐊᑐʕᑕᐅᑦᓯᓗᑎᑦ ᕿᑭᖅᑕᓗᖓᑎᑦ ᐃᓐᓇᑕᐅᓂᕆᓇᐃᓐᓄᑦ ᐊᑐʕᑕᐅᑦᓯᓗᑎᑦ ᐅᓪᓗᕈᓘᑎᑦ ᐊᑐʕᑕᐅᓯᓗᑎᑦ ᐊᑐʕᑕᐅᓇᖅᑐᓂ

ᐅᖃᐅᓯᕐᓴᐅᑎᑦ ᐊᐱᖅᑯᑎᕐᓴᐅᑦᓯ:

ᑲᑎᒪᓂᖅ ᐃᖃᓗᓐᓂᑦ ᐅᖃᐅᓯᖃᓕᐅᖅᑐᑦ ᑲᑎᓐᓂᕐᓂ ᓇᐃᓴᐃᓪᓚᑎᕐᓚᐅᓂᑦ ᐃᓄᐃᑦ ᖃᐅᔨᒪᔪᑐᑎᓂᕐᓂᓗ, <ᓐᓇᐅᑦᓗ ᐱᑕᖃᓚᐊᖃᐅᓐᓂᕐᓐᓂᑦ, ᑭᓯᓕᓚᓪᓚᑎᖃᓚᐊᖃᐅᓐᓂᕐᓐᓂᓪᓗᓂ, ᓯᐅᒃᓗ ᐅᓐᓱᓯᑐᒥᑦ ᖃᐅᔨᓴᓴᑎᓂᑦ, ᐃᑲᔪᑕᐅᓂᐊᖅᑐᑦ ᐃᓄᐃᑦ ᖃᐅᔨᒪᔪᑐᑎᓂᕐᓐᓂᑦ.

ᐊᖏᓇᓯᑐᑕᓚᐅᑉᑯᑦ ᐅᖃᐅᓯᖃᓕᐅᖅᑐᑦ ᐃᓯᐃᓗᔨᑎᒥᓂᖅ ᐱᓪᐊᑎᓕᑦ <ᓐᓇᐅᓐᓇᓐᓂᑦ ᐊᖏᓇᓯᔪᓐᓇᓐᓂᕐᒥᑦ ᐊᐳᐅᖕᓕᐅᑎᓕᕎᕆᖅᑐᑦ ᐊᒥᓯᐊᓐᓇᐅᖅᑕᑕ <ᓐᓇᐅᑦ. ᐊᐢᑎᓕᑕᓚᐅᑉᑯᓐᓇᓐᓂᑦ ᑭᓴᖃᑐᑎᐃᓚᐅᖅᑐᑦ ᐊᒥᓯᓗᐊᓂᑦ <ᓐᓇᓐᓂᑦ <ᓐᓇᐅᑦᓗᑕ ᓇᖃᐅᔨᐊᑭᓐᓇᖅᑐᑦᓯᑦ ᑭᓯᐊᓂᓕ ᐊᖏᑦᑐᑎᐃᓚᐅᖅᑐᑦᖃᖅ ᐊᒥᓯᔪᓂᑦ <ᓐᓇᐅᑦᓗᑕ ᓇᒥᐅᔨᓚᐅᖅᑐᑦ ᑭᓯᐊᓂᓕ <ᓐᓇᐅᑦᓗᑕ ᓇᑭᓕᐅᔨᓴᓴᑎᓂᑦ ᑭᓯᐊᓂᓕ ᐃᓄᐃᑦ ᖃᐅᔨᓴᓴᑎᓂᑦ ᐃᓄᐃᑦ ᖃᐅᔨᒪᔪᑐᑎᓂᕐᓐ

- ᓄᓇᕗᑦ ᑲᑕᒪᒃᑯᐊᓂᕐᓂᖅ–ᐊᕐᑎᑕᖅᓯᒪᔪᑦ, ᖃᐅᖅᑕᕐᓗᒃᑐᑦ ᓂᕆᔨᑎᓂᒃ ᖁᑕᐅᔪᖅᑎᑐᒃ: ᐅᓐᓄᑦ ᓇᒃᑲᓴᖅ
- ᓄᓇᕗᑦ ᑲᑕᒪᒃᑯᐊᓂᕐᓂᖅ–ᐊᕐᑎᑕᖅᓯᒪᔪᑦ, ᖃᐅᖅᑕᕐᓗᒃᑐᑦ ᓂᕆᔨᑎᓂᒃ ᖁᑕᐅᔪᖅᑎᑐᒃ: ᒥᔪᔭ ᐅᓪᓴᐅ
- ᓄᓇᕗᑦ ᑲᑕᒪᒃᑯᐊᓂᕐᓂ-ᐊᕐᑎᑕᖅᓯᒪᔪᑦ, ᖃᐅᖅᑕᕐᓗᑉ ᐅᐊᖏᓇᖅᓄᓂᒃ ᐊᐅᓪᓚᕐᑎᓗᒃ ᓱᓕᑦ ᐅᓐᓴᐊᖓ
- ᓄᓇᕗᑦ ᑐᙵᑲᕆᒃᑯᑦ, ᐃᖅᑲᓇᐃᔨᖅᑐᑕᕐᓂᖏᓐᓇᑦ ᐊᐅᓪᓚᕐᑎᓂᖏᓐᓇᑦ ᐅᖅᑲᐅᖓᐃᓄᐅ; ᓱᐅᑉᑦ ᕗᐃᓪᔪ
- ᓄᓇᕗᓴᒥ ᐅᒪᔨᓪᓚᕐᔪᐊᒃᑯᑦ ᑲᑎᒪᔨᖏᑦ, ᐅᒪᔨᓂᒃ ᖁᑕᐅᔪᖅᑎᑐᒃ ᑲᐃᐅᑉ ᓇᕐᔪ
- ᖃᐅᖅᑕᓂ ᐅᒪᔨᓪᓚᕐᑎᑕᖅᓯᒪᔪᑦ, ᐅᒪᔨᓂᒃ ᖁᑕᐅᔪᖅᑎᑐᒃ: ᒪᐃᑯᑉ ᕗᖓᔪᓇᖅ
- ᑐᐃᓴᓄ, ᑕᐃᑐᓱ ᐊᖏᓐᓇᒃᑉᓪ
- ᐊᐃᐃᕐᖅ ᐊᖕᓇᓱᑲᑐᓪᓚᕐᑎᓂᒃ ᑲᑎᒪᔨᖏᑦ
    - ᐋᓂ ᓱᐃᓚ
    - ᐋᑕᒥ ᓄᓇ
    - ᒋᓇ >ᑐᒍᖅ
    - ᓱᒥᑦ ᓱᐃᓚ
    - ᐃᔨᓱᐊᖅ ᐲᑕ
    - ᑕᑭᐊᑐᒃ ᓄᓇ
    - ᓂᕐᐅᓂᖅᑐᓂᓪᓚ ᑐᑕᑭᑦᑕᓪᓚᑐᒃ ᐃᓐᓇᑐᓂᒃᑯᑦ

<u>ᐅᖅᑲᐅᓱᐊᕆᐃᑦ ᐊᐱᖅᑯᑎᐊᕆᐃᓕᓪᓗ:</u>

ᐅᖅᑲᐅᓱᐅᓕᓪᓚᐊᑕᓕᐅᖅᑐᑦ ᐊᐃᐃᕐᖅ ᐊᕐᓇᓱᑲᑐᓪᓚᕐᑎᓂᒃᑯᓂᓂᑦ ᐃᓕᖅᓕᐅᖅᑐᑦ ᐃᓱᐊᒻᒪᕆᑎᓕᕐᓂ ᐸᓐᓂᑐᐊᓂᑦ ᐊᕐᓇᓱᑲᑎᕐᓂᕐᑕᐊᖅᑲᓐᓂᖅᓯᓂᓂᖅ, ᑲᑎᓱᒪᔨᑦ ᓇᐃᔨᖅᑕᐅᓱᒪᔨᓕᕐᓗ ᖁᑲᐅᔨᓴᕐᑕᐅᓯᒪᓂᖏᓐᓇᑦ ᐊᒥᓱᓕᖅᐸᓕᓂᐅᑉᐳᓂᑦ, ᓇᓗᒋᓂᑦ ᐅᓴᖏᓂᓂᐊᖅᑎᑦ ᖃᐅᖅᑕᕐᓗᒃᑐᑦ ᐅᐊᖏᓇᖅᓄᓂᑦ, ᐱᓯᓕᐊᓂᐅᔨᓂᑦ, ᐊᐅᓚᓂᑕᓴᓂᓪᓗ ᑐᓴᑦ ᖃᐅᖅᑕᒥᑦ.

ᖃᐅᖅᑕᓂᑦ ᐃᒪᖏᐅᑕᓕᕐᓂᑕᓴᐅᔪᑦ ᑲᑎᒪᔨᖏᑦ ᐊᐱᕆᓚᐅᖅᑐᑦ ᐊᑐᑎᖅᑲᓚᐊᕐᓴᖏᓐᓂᑦ ᓄᕐᕋᑦᕠᐊᖏᓚᖅᑐᑦ ᐊᒃᓚᒪ ᐸᖏᓐᓂᑦᕠᐊᖏᓚᖅᑐᑦ ᐊᒥᓱᓂᖏᓐᓂᑦ ᐃᖅᑲᐃᔨᓱᑎᑦᑎᓪᓗᑐᑎᓪᓗ ᑎᑎᖅᑲᒥᑦ 1990–ᖏᓐᓂᓂᓴᖅᑐᒥᑦ 42 ᓄᕐᕋᑦᕠᐊᖏᓄᓕ ᑭᓱᐊᓂᓚ ᑭᖑᓕᖅᖄᑐᒥᑦ ᐅᐊᑐᑕᐅᓱᒪᔨᓕᕐᖅ. ᒪᐃᐅᒃ ᐅᖏᓚᖅ ᐅᖅᑲᓕᐅᖅᑐᖅᖄᓂ ᐊᒥᓱᓂᖏᓐᓂᓚᐅᖅᓯᓂᕐᓂᑦ ᐊᑐᖅᑕ ᐊᖅᑭᑦᑕᓂᑦ ᐃᖅᑲᐅᔪᖅᑎᑐᒃᑯᓂᓂᑦ ᐊᕐᓇᓱᑲᑎᕐᓂᕐᑕᐊᖅᑲᓐᓂᖅᓯᓂᓂᖅ.

ᐊᖏᓚᐅᖅᑐᓕᕐᓂᑕᓴᐅᔪᑦ ᐊᐱᕆᓚᐅᖅᑐᑦ <ᐅᓂᖅᑲᐅᓂᖅᓴᐅᓚᖅᑎᑦᑎᓪᓗ ᐊᖅᓚᓕᑦᑕᖅᑲᖅᐅᓂᖅᓴᐅᓚᖅᑎᑦᑎᓪᓕᑦᑐᓐᓄᑦ ᐅᖏᑦᓕᓂᓂᑦ ᓇᐃᒪᐃᓂᐅᓕᐅᖅᑐᑦ ᐃᓱᒪᓕᐅᓚᓕᑦᑐᓐᓂᑐᐱᓪᓗ <ᐅᓂᑐᐊᓂᑦ ᐊᖏᓚᐅᖅᑐᑦ ᐊᕐᓇᓱᑲᑎᕐᓂᕐᑕᐊᖅᑲᓚᐊᕐᓴᖅ ᐊᒥᓱᓂᖅᓴᐅᓂᖅᓴᐅᓚᖅᑎᑦᑎᓪᓕᑦ <ᐅᓂᑦ. ᓇᓇᕗᑦ ᑲᑕᒪᒃᑯᖏᓂᑦ ᓇᓗᓇᐃᖅᓯᒪᓕᐅᖅᑐᑦ ᑲᑎᒪᔨᖏᑦ ᓇᐃᔨᐃᓱᐊᑦ ᐊᖅᑭᑦᑕᕐᓂᖅᓱᒪᒋᐊᖅ ᐃᓕᓴᕐᓯᒪᔨᓕᕐᖅ ᐃᓇᐅᑦ ᖁᑕᐅᔪᖅᑎᑐᒃᑯᓂᓂᑦ ᓇᓕᕐᓂᓂᖅᓴᐅᓂᖅᓴᐅᓚᖅᑎᑦᑎᓪᓕᑦᑐᓐᓄᑦ ᐃᓱᒪᓕᐅᓚᓕᑦᑐᓐᓂᑐᐱᓪᓗ ᐊᐱᕆᓚᐅᖅᑐᑦ ᖁᑕᐅᔪᖅᑎᑐᒃᑯᓂᓂᑦ ᐊᕐᓇᓱᑲᑎᕐᓂᕐᑕᐊᖅᑲᓐᓂᖅᓯᓂᓂᖅ ᐊᖏᓚᐅᖅᑐᑦ ᖁᑕᐅᔪᖅᑎᑐᒃᑯᓂᓂᑦ ᐊᕐᓇᓱᑲᑎᕐᓂᕐᑕᐊᖅᑲᓐᓂᖅᓯᓂᓂᖅ ᐊᐱᕆᓚᐅᖅᑐᑦ ᖁᑕᐅᔪᖅᑎᑐᒃᑯᓂᓂᑦ ᐊᕐᓇᓱᑲᑎᕐᓂᕐᑕᐊᖅᑲᓐᓂᖅᓯᓂᓂᖅ ᐊᖏ

ᐱᑎᐊᓪᓂᕐᓂᓕᒃ ᑲᑎᒪᓂᕆᓯᒪᓕᕐᕆᑕ ᐅᖃᑲᓯᕐᖃᖏᓂᐊᕐᓗᑎᒃ ᐊᖏᓐᓇᓯᑕᐅᕙᒃᑐᑦ ᑲᑎᕐᓴᑎᑦ ᐊᑲᐅᓪᓕᐅᕐᓂᕐᓗᒍᑦ. ᑭᓯᐊᓂᓕ, ᓄᓇᕗᑦ ᓚᕙᒪᓂᕐᑎᑦ ᐅᖃᓕᐅᕐᖃᑦ ᓄᕐᓗᐅᖏᓐᓂᓇᕐᔪᓕᑎᑦ, ᓂᕆᔨᖃᑎᒌᑦ ᐊᒥᓱᓕᕆᓂᖃᕐᓂᖃᑦ ᐊᖃᓪᓚᕐᔪᐊᓂᖃᑦ ᐊᖏᓐᓇᓯᑕᐅᕙᒃᑐᑦ.

ᐊᖏᓐᓇᓯᑐᑦ ᐊᐱᓕᓕᐅᕐᖃᑦ ᑐᑐᐃᑦ ᖃᑭᖃᓕᒍᓱᑦ ᐅᐊᖏᓇᖏᑦᑐᑦ, ᓄᑦᖁᕐᖃᓱᑦ, ᓇᑦᓄᐃᕐᓂᐊᕐᓱᑦ ᐃᓇᓇᐃᕐᓂᐊᕐᓱᑦᑐᓂ ᐱᒻᒥᓂᕐᓂᑦ ᐊᑲᐅᕙᓪᓕᐊᑦᑐᑦᑎᓇᓂᐊᕐᓱᑦ ᐱᓕᕆᓂᐊᕐᓱᑦ. ᐊᖏᑎᓕᓚᕐᔪᐊᓂᑦ ᓇᓇᐃᕐᓂᐊᕐᓱᑦᑐᑦ ᐊᑲᐃᓂᐅᑦᑕᓂᐊᕐᓱᑦᑐᑦ ᐊᕐᓯᔪᐊᖏᓐᓂᖏᓂᑦ ᐱᓕᕆᓂᐊᕐᓱᑦᑐᓂᑦ ᐃᑲᔪᓂ ᖃᓄᐃᓕᑐᓂᑦ ᐊᓐᑐᐃᓂᕐᓂᐊᕐᓱᑦᑐᓂᑦ ᐃᓕᓴᕐᓂᐊᕐᓱᑦᑐᓂᑦ ᐃᓚᖓᑦ ᐃᓕᓴᕐᓂᐊᕐᓱᑦᑐᓂᑦ ᐃᓕᓴᕐᓂᐊᕐᓱᑦᑐᓂᑦ ᐃᓚᖓ

- ᓯᐊᕐᓴ ᐃᓄᒃᓱᒃ
- ᓯᐃᓗ ᐊᖏᓐᓇᓯᐊᖅ

ᐅᖏᐅᓱᒃᓴᐃᑦ ᐊᐱᖅᑯᑎᒃᓴᐃᓪᓗ:

ᐅᖏᐅᓱᐅᓪᓚᐅᕐᖓᑦ ᓴᓂᖏᔭᖓᒥᐅᑦ ᐊᖅᐳᓱᒃᑐᑕᓂᓚᔨᒃᑯᖓᓐᓄᑦ ᐃᓕᖏᓚᐅᕐᖓᑦ <ᖅᓂᓐᓇᖅᓂᑦ ᐊᖅᐳᓱᓂᖅᒥᑦ, ᑲᑎᓪᓗᑎᑦ ᐊᖅᐳᓱᒃᑕᐅᔨᓐᓇᕐᖓᑦ ᐁᖏᑲᒃᓱᒪᓂᖅᑎᑦ ᓱᐊᓕᒃᑎᖏᑕᐅᔪᖅᑯᖅᓂᖅᑐᓂᒃᓗ ᓄᓇᕗᖅᓄᑦ, ᑲᑎᕐᓱᕐᓱᒪᓂᖅᑎᑦᑕ ᓇᐃᓴᐃᔨᐊᑎᑦ, ᐅᖅᓱᓱᒃᑐᑦᓚᓪᓗ ᖃᐅᔨᓴᖅᓂᖅᓗᑦ ᖁᑎᓴᖅᓂᖅᓗᑦ ᓴᒋᖅᓱᓱᕐᖓᐅᓂᐅᒃ ᒥᖅᓴᖅᓗᓂᖅ.

ᐊᖅᐳᓱᒃᑐᑕᓂᓚᔨᒃᑯᑦ ᐅᖏᑕᐅᖏᖅᑐᑦ ᐃᓱᒪᑎᑉᖏᖅᑐᓂᒃᑯᓂᖅᑐᓂᒃ ᓇᐃᓕᓂᖅᑐᓂᒃ ᓇᓕᓕᒫᓂᖅᑐᓂᒃ ᐃᓕᑕᐅᓚᔨᒃᑯᖓᓐᓄᑦ ᓇᓕᓕᒫ

ᓄᒋᖅᓯᓕᕐᕆᓄᑦ ᒥᑲᓂᐸᓕᕐᕆᓕᕐᕆᓐᓂᑦ ᐅᕙᑲᓂᐸᓕᕐᕆᕆᑎᒋᐅᓕᖅᑎᑐᓄᓗ ᕿᐊᖏᓚᐳᐃᓕᑦ ᑲᑎᒪᔨᖅᑎᒋᑦ ᐊᒋᐅᓚᐊᖅᑐᑦ ᐊᒋᐅᓚᐊᑕᐃᓇᓂᑐᓄᓗ ᑐᓴᐅᒋᐊᖅᑐᑦ 2008-2011 ᒎᓕᐊ ᕿᐊᖏᓚᐳᖏᓐᓄ ᐃᓕᓴᕆᐊᕙᒪᖕᖏᓐᓇᓂᐅᑦ ᓴᐳᑎᒋᐅᓕᐅᕗᖅᑐᖅ ᐅᒪᔨᒍᑦ. ᓄᓇᕗᑦ ᓚᖓᒪᓂᖏᓐᓂ ᑭᐅᓕᐅᕗᖅᑐᑦ ᑕᐃᒪᐃᓕᐅᖏᓂᕆᐊᑦᑕᓂᑐ ᐃᓕᓴᕆᐊᑎᒋᐅᓚᖓᓂᓂᐊᓄᑦ ᒎᓪᓗᓂ ᐊᑐᕋᐅᒪᖕᖏᓐᓂ ᐱᓕᒪᓐᓂᐊᓂᖏᓐᓂ ᕿᐊᖏᓚᐳᓕᖕᖏᐊᓂᖓᓂ. ᕿᑭᖅᒎᓗᖓᑦ ᐅᒪᔨᓕᐅᕗᖅᑐᓂ ᑲᐃᒪᓕᖅᑐᓄᓗ ᐃᓕᓴᕆᐊᑎᒋᐅᓚᖓᓂᓂᐊᓄᑦ ᐊᒋᐅᓚᐊᖅᑐᓂᓂ ᑐᓴᐅᒋᐊᖅᑐᑦ ᐊᑐᕐᓗᒍᑦ ᐱᓕᒪᓐᓂᐊᓂᖏᓐᓂ ᕿᐊᖏᓚᐳᖏᓐᓇᓂᐅᑦ ᑐᓴᐅᒋᐊᖅᑐᑦ ᐊᒋᐅᓚᐊᖅᑐᑦ ᐊᒋᐅᓚᐊᑕᐃᓇᓂᑐᓄᓗ

ᐅᖃᐅᓯᖏᓚᐅᕐᖏᑦ ᐃᓐᓇᓕᕐᒥᑦ ᐊᖏᔪᓕᓴᒃᑖᓂᖅᓯᒪᔪᓂ ᐃᓕᖏᓚᐅᕐᖏᑦ ᑐᓴᕋᕐᓯᓂᖏᓐᓄᑦ ᓄᐅᕐᑕᖅᓯᓚᑦ, ᐊᖏᔪᓕᓴᒃᑖᓂᖅᓯᒪᔪᓂ ᐃᓕᐅᕙᓪᓕᐊᑎᓪᓗᒥᑦ, ᐅᖃᓗᓕᒫᑦ ᖃᐅᖅᑕᕐᓯᓚᖅ ᖃᐅᖅᑕᖅᓯᓚᖅ ᓄᓇᕗᑦ ᐃᓐᓇᓕᕐᒥᑦ ᐃᓐᓇᓕᕐᒥᑦ ᐃᓐᓇᓕᕐᒥ

- ᓄᓇᕗᑦ ᑲᑎᒪᔨᖏᑕᓂᑦ–ᐊᕐᑎᑕᖅᓱᒪᔨᑦ, ᖃᐅᔨᒪᓂᖅᑐᑦ ᓂᕆᐊᑎᓂᒃ ᖃᐅᔨᓴᖅᑎᑐᖅ: ᒥᔪᑦ ᑲᓪᓚᕆ
- ᓄᓇᕗᑦ ᑲᑎᒪᔨᖏᑕᓂ-ᐊᕐᑎᑕᖅᓱᒪᔨᑦ, ᖃᐅᔨᒪᓂᖅᑐᑦ ᐅᐊᖏᓇᖅᑐᓂᒃ ᐊᐅᓪᓚᑎᓵᓂ ᓯᑲᑦ ᓴᓐᓴᕐ
- ᓄᓇᕗᑦ ᑲᑎᒪᔨᖏᑕᓂ- ᐊᕐᑎᑕᖅᓱᒪᔨᑦ, ᐅᒪᓯᓪᓚᓵᓂ; ᐃᒪᐅᐳ ᐊᑭᔨᑐᓴᐅ
- ᓄᓇᕗᑦ ᑐᖓᑲᕆᔨᑦ, ᐃᓕᑲᖅᐱᓴᖅᑐᑕᓪᓚᖓᓐᓇᑦ ᐊᐅᓪᓚᑎᓂᖅᓯᓐᓇᑦ ᐅᕝᕙᓴᐅᐃᓵᓂ; ᓯᐅᑕᑦ ᕐᐃᓴᓵ
- ᓄᓇᕗᓴᒥ ᐅᒪᓯᓪᓚᓵᔭᕐᐳᒪᔨᑦ ᑲᑎᒪᓵᖏᑦ, ᐅᒪᓯᓂᖅ ᖃᐅᔨᓴᖅᑎ ᑲᐃᐅᑦ ᓗᓯᓄ
- ᖃᐅᔨᖅᑕᓂ ᐅᒪᓯᓪᓚᓵᔨᑦ ᑲᑎᒪᓵᖏᑦ, ᐅᒪᓯᓂᖅᓱᒪᔨᑦ: ᒪᐃᔨᑦ ᐅᖓᔪᓴᐅ
- ᑐᐃᓴᓵ, ᑕᐃᑐᓯ ᐊᖏᓐᓇᒃᑕᒃ
- ᐃᑲᔨᖅᑎᑦ ᐊᖓᔪᓇᓯᑐᖅᑕᓪᓚᓵᔨᑦ ᑲᑎᒪᓵᖏᑦ
    - ᕿᓪᓗᓇ ᖃᑲᐅᓐᓇᖅ
    - ᔨᐃᑕ ᖃᑲᒪᓂᖅ
    - ᑯᐅᓇ ᐃᓄᔪ
    - ᐅᓪᓕᐅ ᑕᖅᑐ
    - ᕿᑦ ᐃᔪᖓᑎᐊᖅ
    - ᔨᐊᓇ ᐅᔨᔨᑯᓴᒃ
    - ᓴᓂᐅᓄ ᐸᐅᓪᓯᓯ

<u>ᐅᖃᑲᐅᓯᕐᓴᐃᑦ ᐊᐱᖅᑯᑎᕐᓴᐃᑦᓪᓗ:</u>

ᐅᖃᑲᐅᓯᖃᓕᐅᖅᑐᑦ ᐃᑲᔨᖏᑦ ᐊᖓᔪᓇᓯᑐᓪᓚᓵᔨᑯᖓᓐᓂᓂᑦ ᐃᓕᖃᓕᐅᖅᑐᑦ ᓇᐃᓴᐃᖃᑎᐅᕐᒪᖅᑐᓂᑦ, ᐊᖓᔪᓴᓪᓗᐊᓂᑦ ᐊᖓᔪᓇᓯᓂᓂᑦ, ᐊᖓᔪᓇᓯᑐᓪᓚᓵᔨᑯᑦ ᐃᓕᐅᕐᒪᓂᖓᓐᓂᒃ ᓇᐃᓴᐃᑎᓪᓗᒃᑐᑦ, ᐃᓚᐅᐅᑎᓯᓂᖓᓐᓂᓪᓗ ᐅᓴᖓᓐᓂᑦ ᖃᐅᔨᓴᖅᑎᖓᓂᖅᑐᑦ ᖃᐅᔨᓴᖅᑎᓂᖓᓂᑦ. ᐅᖃᑲᐅᓯᖃᓕᐅᖅᑐᑦ ᐊᓯᓚᓕᓕᐅᖅᑐᑦ ᑭᖓᔪᓂᓪᓗᒃ ᐊᐱᖅᑯᑎᕐᓴᐃᑦ ᖃᐅᔨᓴᖅᑎᖓᓂᖅᑐᑦ ᐊᐱᖅᑯᑎᕐᓴᐃᑦ ᓯᓚᑦ ᐊᐱᖅᑯᑎᕐᓴᐃᑦ ᖃᐅᔨᓴᖅᑎᖓᓂᖅᑐᑦ ᐊᐱᖅᑯᑎᕐᓴᐃᑦ ᑕᒪᐃᑯᐊ ᐊᐢᑐᐃᐊᖏᓪᓗᒃᑐᑦ ᖃᐅᔨᓴᖅᑎᖓᓂᖅᑐᑦ. ᐅᖃᑲᐅᓯᖃᓕᐅᖅᑐᑦ ᐊᓯᓚᓕᓕᐅᖅᑐᑦ ᑭᖑᓪᓕᐊᑦ ᐊᐢᑐᐃᐊᖏᓪᓗᒃᑐᑦ ᖃᐅᔨᓴᖅᑎᖓᓂᖅᑐᑦ ᐊᐱᖅᑯᑎᕐᓴᐃᑦ ᖃᐅᔨᓴᖅᑎᖓᓂᖅᑐᑦ ᐊᖓᔪᓇᓯᑐᓪᓚᓵᔨᑯᑦ ᖃᐅᔨᓴᖅᑎᖓᓂᖅᑐᑦ ᐊᐱᖅᑯᑎᕐᓴᐃ

ᐅᖃᐅᓯᕆᓕᔪᓐᓇ ᓄᓇᕗᑦ ᑲᑎᒪᔨᖏᑕᖅᑐᑦ. ᐊᖏᓂᔪᒃᑐᑕᓂᕆᔨᒃᓯᑦ ᐅᖃᐅᓯᖃᓕᕕᒃᓴᖅᑎᑦ ᖃᓄᖅᓱᓕᕕᒃᑐᐃᓐᓇᖅᑎᑦ ᐳᖓᓕᒫᑦ ᑲᑎᒪᓂᖅᑎᑦ ᑲᑎᒪᒫᖅᐸᑕ.

ᐅᖃᐅᓯᐅᖏᖃᓕᒫᖅᑎᑦᑐᑦ ᑎᑎᕋᖃᑕᐅᕈᐊᖅᓱᖃᓯᒪᓕᒻᒥᐸ ᐊᐅᓕᕆᑎᓂᖅᓗᑦ <ᖅᓀᐅᑎᑎᑦ ᑐᓴᕋᑎᑎᓕᑦᑕᑦ. ᐊᖏᓂᔪᒃᑐᑕᓂᕆᔨᒃᓯᑦ ᐅᖃᓕᒫᖅᑐᑦ ᐅᖃᖅᑎᑎᖃᐳᓕᒫᓕᑦᑕᐅᒃ ᐊᐅᓕᕆᑎᓂᖅᓗᑦ <ᖅᓀᐅᑎᑎᑦ ᐃᓕᐊᓂᑦ ᑲᑎᒪᓕᕕᑦ ᑲᑎᒪᒫᕐᐸᑕ.

3.9 ᒥᕆᒪᒐᓗᒃ

ᐅᓪᓗᖅᓯ: ᓴᓂᐊᓗ 18, 2019

ᑭᖑᓕᖅᑐᑎᒃ:

- ᓄᓇᕗᑦ ᑲᑎᒪᔨᖏᑕᓂᓕᒻᐊᑎᑎᓕᕆᔨᒃᓯᑦ, ᖁᑭᖅᑕᒋᓕᖅᓯᑦ ᓂᕐᔪᑎᓂᒃ ᖃᐅᔨᓴᕐᕕᒃ: ᑲᓂ ᓇᖅᑲᓴ
- ᓄᓇᕗᑦ ᑲᑎᒪᔨᖏᑕᓂᓕᒻᐊᑎᑎᓕᕆᔨᒃᓯᑦ, ᖁᑭᖅᑕᒋᓕᖅᓯᑦ ᓂᕐᔪᑎᓂᒃ ᖃᐅᔨᓴᕐᕕᒃ: ᒥᕐᓵ ᑲᒪᐸ
- ᓄᓇᕗᑦ ᑲᑎᒪᔨᖏᑕ-ᐊᐅᑎᑎᓕᕆᔨᒃᓯᑦ, ᖁᑭᖅᑕᒋᑲᐸ ᐅᐊᖅᓇᖅᓱᓄᑦ ᐊᐅᓕᕆᑎᓕᐸ ᓯᑲᑦ ᑲᓄᓴᓇ
- ᓄᓇᕗᑦ ᑐᖅᑲᕕᒃᓯᑦ, ᐃᖃᖅᓇᐃᓴᖅᑐᑕᓂᕆᓐᓂᖓᓕᑦ ᐊᐅᓕᕆᑎᓂᖅᓗᑦ ᐅᖃᐅᓴᕐᔪᐃᓐᓇᐅᒃ; ᓯᐅᕗᑦ ᕘᐃᓴᓵ
- ᓄᓇᕗᓴᒥ ᐅᓚᔪᓕᕆᔨᒃᓯᑦ ᑲᑎᒪᓂᖅᑎᑦ, ᐅᓚᔪᓂᒃ ᖃᐅᔨᓴᕐᕕᒃ ᑲᐃᐅᑦ ᓇᑦᓯ
- ᖁᑭᖅᑕᓂ ᐅᓚᔪᓕᕆᔨᒃᓯᑦ ᑲᑎᒪᓂᖅᑎᑦ, ᐅᓚᔪᓂᒃ ᖃᐅᔨᓴᕐᕕᒃ: ᒪᐃᑯᑦ ᕘᖓᓴᓵᓴ
- ᑐᐃᓴᓄ, ᐊᐃᕐᓀᓪᓕᕿᐊᑦ ᑯᑦᓗ
- ᒥᕆᒪᒐᓗᖅᑎᑦ ᐊᖏᓂᔪᒃᑐᑕᓂᕆᔨᒃᓯᑦ ᑲᑎᒪᓂᖅᑎᑦ
  - ᐊᐃᒥ ᖁᓕᖅᑎ
  - ᑕᐃᕐᐱᑎ ᖃᒪᓂᖅ
  - ᐃᓕᐃᓵᓴ ᓇᓗᓴ
  - ᐃᐅᓇᓴ ᐅᑦᑐᓀ
  - ᒎᓂᐊᓕ ᖃᑯᐊᖅ
  - <ᓄᐃᓕ ᐃᓄᐊᕐ
  - ᐃᓅᑉᐹ ᐃᓄᐊᕐᖃᖅ

ᐅᖃᐅᓯᐅᖏᖃᓕᑦ ᐊᐱᖅᑯᑎᖏᖃᓕᒻᓗ:

ᐅᖃᐅᓯᖃᓕᒫᖅᑐᑦ ᒥᕆᒪᒐᓗᒃ ᐊᖏᓂᔪᒃᑐᑕᓂᕆᔨᒃᓯᑯᓂᓐᓂᒃ ᐃᓕᖃᓕᒫᖅᑐᑦ ᓇᐃᖏᐃᓕᐅᕐᐅᖃᓕᒫᖅᑐᑦᑎᑦᑕᒫᖅᑐᑦ, ᐅᖓᓯᓇᖅᑐᓕᑦ ᖃᐅᔨᓴᖓᓂᒃᓴᑦ ᖁᑭᖅᑕᒋᓕᖅᓯᑦ ᐅᐊᖅᓇᖅᓱᓄᑦ, ᐊᖏᑐᐃᓐᓇᐅᑉᐸᓗᒃᑐᑦ ᐃᓕᓴᕐᓂᖅᓱᓕᑦ ᐊᐅᓚᑦᑎᓐᓂᒃᑐᑦ.

ᐊᖏᓂᔪᒃᑐᑕᓂᕆᔨᒃᓯᑦ ᐃᓇᐹᒐᑎᖃᓕᕆᖅᑐᑦ ᐊᔪᖖᑎᖃᒃᓗᐊᕆᒃᓴᖓᓂᒃᓴᑦ ᐃᑲᖃᖅᑎᓂᑦ 2018–ᖓᑎᓪᓗᒍ ᑯᑦᑕᓴᓂᖅᓂᒃᓗ ᐊᖏᓂᔪᓂᖅᓂᑦ ᑐᖅᑐᓂᓗᒃᓂᑦ ᓇᓗᓇᐃᔪᓂᖅᓂᑦ. ᐃᓕᓐᓂᑦ ᐅᖃᐅᓯᖃᓕᕆᖅᑐᑦ >ᖅᑐᑐᓂᒃᓂᑦ ᐊᔨᑲᖖᑎᑦᑕᓗ ᐱᓕᕆᔨᓂᖅᓂᑦ ᓇᓗᓇᐃᔪᓐᓇᕆᕿᓂᕐᕕᑯᓂᓐᓂᑦ ᐊᖏᓴᓕᓗᓂᖅᓂᑦ ᐊᔪᓐᓇᖅᑐᓂᑦ ᐊᐅᓚᑦᑎᓐᓂᒃᑐᑦ ᓇᓗᓇᐃᔪᓕᕆᔨᒃᓯᑦ.

ᐊᖏᓇᓱᖅᑐᓕᕆᔨᒃᓯᑦ ᒥᕋᒪᒋᓕᓐᖏᑦ ᐃᓱᐊᓗᑎᕋᕆᐊᖅᑐᑦ ᐊᖅᑐᐃᓂᕆᔨᓂᑦ ᓄᓗᑦᑎᓂᑦ ᐅᕋᖅᑕᐅᓗᓂᑦ ᓄᓗᑦᑎᓂᑦ ᐃᑲᔪᖅᑕᐅᔨᕆᐊᕋᕆᑲᒥᖏᓪᓗᑎᒃ ᐅᔨᕋᖏᓂᐊᖏᓂᓕᕆᔨᓄᑦ. ᓂᖏᓴᐅᒪᓂᕋᑲᒥᐅᖏᑦᑐᓂ ᐊᖏᕋᖅᑕᐅᓂᖏᓐᓂᑦ ᐱᓕᕆᐊᖏᒧᐅᓂᕆᐊᓂᖏᓐᓂᑦ ᐊᒥᓱᓕᖅᑐᓂᖏᓐᓂᑦ ᐊᖏᕋᖅᑕᐅᒧᖏᓂᓪᓗ ᒥᓂᑕᒪᑦ. ᐊᖏᓇᓱᖅᑐᓕᕆᔨᒃᓯᑦ ᐃᓱᒪᕋᕆᔨᖅᑕᑦ ᐊᐃᒃᓱᐊᐸᒪᓪᓗᑎᒃ ᕿᑭᕋᕐᒪᓂᑦ ᐅᐊᖏᓇᖅᓯᓂᑦ ᓂᒋᐊᓂᖅᓱᓂᑦ ᐊᐃᒃᓱᐊᓕᓗᑎᒃ ᐊᐅᓚᕆᐊᖅᑐᓂ.

ᐊᒧᑎᓕᕆᔨᒃᓯᑦ ᐅᕋᖅᑖᓕᐅᕋᖅᑎᑦᓗᓂᑦ ᑭᒧᑦᓕᖓᑦ ᑐᑐᓕᕆᔨᑦ ᑐᓴᐅᑦ ᕋᕆᐊᐅᑦ ᕋᖅᐅᔨᓴᕋᐅᓂᖏᓐᓂᑦ ᐅᕋᖅᐅᓂᕋᖅᑕᐅᓂᖏᓐᓂᑦ ᕋᔪᖏᓂᔪᕐᓂᐊᓂᖏᓐᓇᓕᑦ ᐱᓕᕆᔨᓂᑦ ᕋᖅᓯᓗᓐᓄᑦ ᑕᒃᔪᐊ ᕋᔪᖏᓂᔪᕐᓇᓕᑦ ᐃᓕᓴᐅᓂᖏᓐᓂᑦ. ᐃᓱᐊᓗᑎᒃᓯᑦ ᕋᔪᖏᓂᔪᕐᓇᓕᑦ ᐅᒪᓕᑦ ᓴᓇᖅᑎᓯᓕᕆᔨᓂᖏᓐᓂᑦ ᕋᔪᖏᓂᔪᕐᓇᓕᒃ. ᓄᓇᕗᑦ ᓴᖅᓪᓚᒃᑯᓂᑦ ᕿᑭᕋᕐᒪᓂᑦ ᑕᒃᔪᐊ ᕋᖅᑐᐃᓂᑦ ᕋᔪᖏᓂᔪᕐᓇᓕᕐᓂᐊᓕᓱᐃᓂᖏᓐᓂᖅᓂᑦ ᐊᑐᓵᖅᓴᓐᓂᖅᓗᓐ ᐅᒪᓂᒋᒧᖏᓂᖅᓂᑦ. ᓄᓇᕗᑦ ᓴᖅᓪᓚᒃᑯᓂᑦ ᕋᔪᖏᓂᔪᕐᓇᓕᑦ ᐅᕋᖅᑖᓕᐅᕋᖅᑎᑦᓗᓂᑦ ᓴᖅᓪᓗᓐᓄᑦ ᕋᖅᑐᐃᓂᑦ ᐋᒃᑐᓂᑦ ᕋᔪᖏᓂᔪᕐᓇᓕᒃ ᕿᑭᕋᕐᒪᓂᑦ ᐃᓕᓴᐅᓂᖏᓐᓂᑦ ᓴᓇᖅᑎᓯᓕᕆᔨᓂᖏᓐᓂᑦ. ᐃᓱᐊᓗᑎᒃᓯᑦ ᕋᖅᑐᐃᓂᑦ ᕋᔪᖏᓂᔪᕐᓇᓕᒃ, ᕋᖅᑐᓗ ᐱᖅᑕᐅᓇᖅᑕᖅᑐᖅ ᐱᓇᐅᔨᓴᕆᔨᖅᑐᑦ.

ᐅᖅᑕᐅᓂᕐᓴᖅᑕᐅᓕᖅᑐᓂᑦ ᑎᑎᕐᖅᑕᐅᑎᐅᔪᖅᑐᓂᖅᓱᓕᓚᒃ ᐊᐅᓚᕆᐊᓂᕐᓗᑦ ᕿᖏᓇᐅᑎᑦ ᑐᖓᔪᓂᖅᓴᖅᑐᑦ. ᑲᑎᒪᔨᑦ ᐅᓇᐃᕐᓂᓚᕆᐊᖅᑐᓂᑦ ᓂᕆᔨᓕᕐᓂᑦ ᐊᐅᓚᕆᐊᓂᕐᓗᑦ ᕿᖏᓇᐅᑎᑦ ᐃᓚᖏᓪᓗᑎᒃ ᐃᓚᖏᓪᓕᐊᓂᖅᓴᖅᑐᑦ ᐃᓕᓴᐅᓂᖏᓐᓂᑦ ᓄᓇᕗᑦ ᓴᖅᓪᓚᒃᑯᓂᑦ ᑲᑎᒪᔨᑦ ᐃᓚᖏᓪᓕᐊᓂᖅᓴᖅᑐᑦ ᐃᓕᓴᐅᓂᖏᓐᓂᑦ ᓄᓇᕗᑦ ᓴᖅᓪᓚᒃᑯᓂᑦ ᑲᑎᒪᔨᑦ ᐃᓚᖏᓪᓕᐊᓂᖅᓴᖅᑐᑦ ᐃᓕᓴᐅᓂᖏᓐᓂᑦ ᐃᓱᐊᓗᑎᒃᓯᑦ ᐅᕋᖅᑖᓕᐅᕋᖅᑎᑦᓗᓂᑦ ᐊᐅᓚᕆᐊᓂᕐᓗᑦ ᕿᖏᓇᐅᑎ

ᐊᖏᓇᓱᑐᑕᓚᕆᔨᒃᓴᓕ ᐃᓕᑎᕆᐅᕐᓂ ᓇᓗᓇᐃᕆᐃᓕᐅᕆᖅᑐᑦ ᐃᓱᒪᑎᕆᖏᖅᓯᑎᓗᓐᓄᑦ ᓇᐃᓴᐃᓂᐊᓘᓐᓂ ᑲᓇᒪᓂᖏᓐᓄᓐᓂᑦ, ᐸᕿᑎᓕᑦ ᑐᖏᑎᖅᓱᒪᒪᕐᓂ ᐳᕆᓭᑐᐃᓐᓄᑦ ᖁᓕᖅᑕᑭᑐᓗᕐᓂᑦ, ᐅᖏᓐᓱᓇᒥᑦ ᖃᐅᔨᓴᖅᓂᒪᑦ ᖃᐅᔨᓴᓕᕐᓂᑦ, ᐊᕆᑐᐃᓂᐅᕐᓂᓗ ᐱᙳᓕᓂᐊᖅᓂᑦ, ᐊᕆᑐᐃᓂᐅᕐᓂᓗ ᑐᓴᐅᒃᑐᓂᑦ.

ᐊᖏᓇᓱᑐᑕᓚᕆᔨᒃᓴᖅᓂ ᐊᙳᑎᓕᑕᓚᕆᔨᒃᓴᖅᓯᓗ ᐅᖃᕆᔨᖃᓕᐅᕆᖅᑐᑦ ᐊᒥᓱᓂᖏᓐᓂᒃᓂ ᑲᑎᖃᓱᖏᓐᓂᒃᓂ ᓇᐃᓴᐃᓂᐊᓘᓐᓂ ᑲᓇᒪᓂᖏᓐᓄᓐᓂᑦ ᐱᓕᒪᓐᓂᐅᕐᓂᑦ ᐊᕆᑐᐃᓂᐅᕐᓂᑦ ᐱᓕᒪᓐᓂᐅᕐᓂᑦ, ᑕᒪᓐᓇ ᐃᓐᓇ ᐱᐅᓐᓂᐅᕐᓂᑦ.

ᐊᖏᓇᓱᑐᑕᓚᕆᔨᒃᓴᓕ ᐃᓕᑎᕆᑦ ᐊᙳᑎᓕᑕᓚᕆᔨᒃᓴᖅᓯᓗ ᐅᖃᕆᔨᖃᓕᐅᕆᖅᑐᑦ ᐅᓂᒪᓱᒃᓴᓕ ᖃᐅᔨᓴᖅᓂᒪᓕᒃᑐᑦ ᖃᐅᔨᓴᓕᕐᓂᑦ ᑎᑎᖅᑲᖅᓯᒪᔪᑦ ᖃᐅᔨᓴᑎᑐᑦ ᐊᙳᑎᓕᑦ ᖃᖅᔪᓂᑦ, ᐊᖅᔪᑎᖃᕐᔪᑦ, ᐊᕆᑐᐃᓂᐅᕐᓂᓗ ᐱᙳᓕᓂᐊᑎᒋᔭᓂᑦ. ᐃᓕᑎᕆᐅᕐᓂᑦ ᐱᐅᖅᓴᓕᕐᓂᖅᓯᑦ ᐅᖃᕆᔨᐅᕐᔪᑦ ᑕᖃᓴᖅᑎᑕᐅᓯᒪᔪᓂᒃᑐᑦ ᐊᖅᔪᑎᑦ ᑐᑐᓕᑦ ᐊᐅᓚᓕᕐᓂᑦ ᐱᐅᓐᓂᐅᕐᓂᑦ ᐱᐅᓐᓂᐅᕐᓂᓗ. ᐅᖃᕆᔨᖃᖅᑲᖅᑎᑦᑎᓂᒃᑐᑦ ᐊᖅᔪᑎᑦ ᖃᐅᔨᓴᖅᓂᒪᓕᒃᑐᑦ ᐱᙳᓕᓂᐊᑎᒋᔭᓂᑦ ᐃᓕᑎᕆᐅᕐᓂᑦ ᐱᙳᓕᓂᐊᑎᒋᔭᓂᑦ ᐊᖅᔪᑎᑦ ᐊᐅᓚᓕᕐ

ᑲᑎᒪᓕᖅᓯᒋᑦ ᐃᓗᐊᓕᓂᑦ ᐅᖃᖅᑲᑎᖅᑲᓕᐅᖅᑐᑦ ᐊᖏᔪᓕᓴᕐᓗᓂᒡᓗᓂᑦ ᑕᕆᔨᐊ ᓄᓇᕗᑦ ᓴᖅᒪᓕᕆᔨᓯᒋᑦ ᑐᓴᕐᕋᖅᑲᓕᐅᖅᑎᓪᓗᓂᑦ ᐅᖃᐅᓯᖅᑲᖅᓯᐊᕐᓗᑎᒃ ᓯᓚᐅ ᐊᑐᖅᑕᒥᑦ ᑲᑎᓪᓗᓂᑦ ᐊᖏᔪᓕᑕᐅᔪᓚᐅᖅᑐᓂᑦ.

ᐅᖃᐅᓯᕆᔨᖃᖅᑐᖃᑲᓕᐅᖏᓪᓕᐅᑐᖅ ᐊᖏᔪᓕᓴᕐᑐᓗᓴᕆᔨᓂᑦ ᑐᓴᓚᐊᖅᐸᐅᓯᒪᔪᓂᑦ ᓯᓂᓪᓕᖏᓪᓗᑎᒃ ᐊᐅᓪᓕᐊᑎᓂᖓᓐᓇᓯᑦ <ᖏᓕᐅᑦᓂᑦ ᑕᕆᔨᐊᓗ ᐊᖏᔪᓕᓴᕐᑐᓗᓴᓪᓕᐊᑦ ᐅᖃᓪᓕᐅᖅᑐᑦ ᐅᖃᖅᑲᑎᖅᑲᕈᒪᓪᓗᑎᒃ ᐃᓗᐊᓂᑦ ᓄᕐᓯᐅᓴᓪᓕᐅᖅᑲᐅᑎᒃ ᓄᓇᕗᑦ ᓴᖅᒪᓕᕆᔨᓐᓄᓐᓂᑦ.

5.0 ᐅᐃᒍᐊᑭᓯᖅ 1

ᖃᑭᖅᑕᓗᓐᒥ ᑐᕆᑐᐃᑦ ᒥᖅᓴᓄᑦ
ᑐᓴᐅᒪᖅᑎᒋᔭᖅ ᐊᖏᔪᓇᓯᖅᑏᑦ
ᒥᑭᐊᖅᓂᐊᖅᑏᑦᓗ ᖃᑐᐅᖅᑎᒋᔭᖅᓯᓂᑦ (HTO)

ᐃᓂ ᓚᖓᔪ
ᖃᑭᖅᑕᓗᓐᒥ ᐊᕕᐅᑐᖅᓯᒪᔪᒥᑦ ᓂᕆᔨᑎᓂᖅ ᐅᒪᔨᓕᓪᓚᕆ
ᒥᕋᑎᒪᑕᓪᖅ
Jringrose@gov.nu.ca
(867) 899-7576

ᐅᕋᖅᐅᓯᕋᖅᑕᓗᐊᑕᕋᓗᓂ

ᓇᓗᓇᐃᔨᐃᓂᖅ

ᕋᖅᐅᓴᓴᕋᓂᖅ ᐊᐱᕋᖅᓯᕋᓂᖅᑯᑦ
ᕋᖅᓄᐃᑦᑐᓂᖅᓱᒃ

ᕋᖅᓴᓯᐅᓂᖅᑎᖅᓂᑦ ᕋᖅᐅᓴᓴᕋᓂᖅ
ᐊᐱᕋᖅᓯᕋᓂᖅᑯᑦ

2015-2018 ᕋᖅᐅᓴᔨᐅᓯᒪᔪᑦ

2019 ᐅᐱᕋᖅᓱᒋᑦ ᕋᖅᐅᓴᓴᕋᓂᖅ ᐊᐱᕋᖅᓯᕋᓂᖅᑯᑦ

ᑐᖅᑐᓂᖅ ᕋᖅᐅᓴᓴᕋᓂᖅ

ᑎᑎᕋᖅᑕᐅᑎᐊᖏᓪᓚᐅᑦ
ᐊᐅᓪᓚᕆᓂᕐᒧᑦ ᐸᕋᓇᐅᑎᓄᑦ
ᐅᕋᖅᐅᓯᐅᓯᒪᔪᑦ



ᓇᓗᓇᐃᔨᐃᓂᖅ



ᐅᓂ ᓱᖁᑦ

John Ringrose

ᐊᐃᔨᖏᓱᒪᓕᕆᓂ ᐅᒪᓕᖅᓂᐊᖅᑎ
ᖃᑭᖅᑕᓗᒃ
ᐊᐅᖅᑕᓂᕐᓂᐅᔪᓂ
ᓄᓇᕗᑦ ᓴᖅᒪᓴᒃᑯᑦ

Regional Wildlife Biologist
Baffin Region
Department of Environment
Government of Nunavut

Avikturviani Huradjanut Qauyihaiyi
Qiqiktaaluk
Avatiliqiyitkut
Nunavut Kavamanga

P.O. Box/C.P. 400
Pond Inlet, NU X0A 0S0
867-899-7576

iringrose@gov.nu.ca

www.gov.nu.ca

ᖃᐅᔨᓴᖅᑕᐅᔪᑦ ᑭᓯᓂᖅᑎᑦ ᐊᒻᒪ ᖃᐅᔨᓴᕕᓯᑦ


ᒪᖃᑦᓗᐊᑦᑕᒃ ᐊᔪᐅᑲᖏᑦᑐᑦ ᑐᐅᑐᐃᑦ ᖃᐅᔨᓴᖅᑕᐅᔪᑦ
ᑐᐅᑐᖃᐅᑎᐊᖅᑐᑦ- ᑐᐅᑐᐃᑦ ᖃᐅᔨᓯᐅᓂᖏᑦ
ᖃᐅᓄᐃᑦᓴᓂᖅ- ᖃᐅᔨᓯᐅᓂᖏᑦ <ᖅᓂᑦ, ᓄᖅᖃᓕᑦ, ᓄᖅᖃᐃᑦ ᐊᒻᒪ ᓄᖅᖃᐃᑦ ᐅᒪᔪᑦ
ᐅᑭᐊᖏᓴᓂᑦ ᖃᐅᐅᔨᖅᓯᐅᑐᑦ ᐊᐅᐅᓕᖓᑦ <ᖅᓂᖅᓄᑦ: ᓄᖅᖃᓕᑦ
ᖃᐅᔨᓯᐅᓂᖏᓐᓂᐅ ᐊᒻᒪ ᐊᐅᔪᖅᑎ ᐊᓂᔪᖅᑎᑦᑕᐃᑐᑦ ᓄᖅᖃᖅᓄᑦ ᐅᑭᐅᓴᓂᑦ ᐊᓐᓇᓴᒪᓪᓕᐊᑦ
ᐅᐱᖅᓯᐊᓴᓂᑦ ᖃᐅᐅᔨᖅᓯᑕᖅᑐᑦ ᓄᖅᖃᖅᓄᑦ ᐅᑭᐅᓴᓂᑦ ᐊᓐᓇᓴᒪᓪᓕᐊᑦ

ᖃᑭᑦᑕᓗᒃᑲᒥᒃ ᑐᐅᑐᐃᑦ ᐊᒥᓱᓂᖏᑦ 2014-ᒥᑦ ᑭᖑᓪᓕᖅᑭᒥ ᐱᐊᓂᖅᑕᐅᓪᓕᐅᖅᓱᖅ

ᖃᐅᓄᐃᑦᓴᓂᖅᑎᑦ ᖃᐅᔨᓴᓂᓪᓕᐅᖅᓱᑦ 2015 -2018.
ᑭᖑᓪᓕᖅᑎᑦ ᖃᐅᔨᓴᖅᑕᐅᔪᖅᑕ ᐱᐊᓂᖅᑕᐅᓪᓕᐊᓂᖅᑐᖅ
ᐃᓕᓴ/ᐊᐃᑦᓪᓕ 2019-ᒥ


ᖃᐅᔨᓪ ᐊᖏᑭᒃᓴᖅᑕᕐᒪᑦ ᖃᐅᔨᓴᒃ ᑐᑐᖃᕐᓂᖅᑕᐅᔪᓂᒃ?

1) ᓂᕆᐊᖅᓯᒪᔭᒃ ᓇᐅᒃᑯᑦ ᖃᐅᖓᑕᓂᐊᖅᓂᖅᑭᓂᒃ ᐊᒥᓱᓂᖓᑦ**
ᖃᑭᖅᑕᒻᒥᑯᑦ



ᖃᐅᔨᓪ ᐊᖏᑭᒃᓴᖅᑕᕐᒪᑦ ᖃᐅᔨᓴᒃ ᑐᑐᖃᕐᓂᖅᑕᐅᔪᓂᒃ? ᐊᒥᓱᓂᖓᑦ**

2) ᑐᑭᓕᐊᖅᑐᓂᒃ ᐃᓕᐅᖅᑲᐃᓐᓇᑐᒃ ᖃᑭᖅᑕᒪᔭᑦ
ᖃᓄᖅᑕᕐᕆᐊᐅᑲᑎᒪᓕᒃ



ᖃᒃᓱᐃᓂᖅ ᐊᐃᑯᓗᐃᑦᑐᑦᓯᐊᖅ ᖃᓄᐃᓕᖅᑐᑦ ᑲᑎᒪᔨᖅ ᑐᓴᕐᓯᒪᓕᕆᔨᒃᑯᑦ?


3) ᐃᑦᓯᓴᓐᓂᑦ ᐃᑦᑐᐊᖅᑲᓐᓂᖅ
ᑐᓴᑦᑐᖅᓯᒪᔪᓕᑦ


ᐊᒥᓱᓕᕐᓂᑦ**


ᖃᒃᓱᐃᓂᖅ ᐊᐃᑯᓗᐃᑦᑐᑦᓯᐊᖅ ᖃᓄᐃᓕᖅᑐᑦ ᑐᓴᑦᑐᖅᓯᒪᓕᕆᔨᒃᑯᑦ?

4) ᑐᓴᐃᑦ ᑕᑯᐅᕆᔨᑦ
ᓴᓱᖅᑕᕆᔨᒧᓐᓂᖅ

ᐊᒥᓱᓕᕐᓂᑦ**


ᖃᒥᓱᓂᖅ**
ᖁᑎᓂᒃᑯᐊᓂᖅᓯᒪᔪᑦ ᖁᑦᓴᓂᒃ ᑐᑐᖁᖅᓂᕐᖁᑎᐅᔨᓂᒃ?
5) ᖁᑎᐅᔨᓴᓐᓇᖅᑐᒃᑯᑦ ᓇᓴᐅᑎᑦ ᑕᑯᓐᓂᑕᒃᑯᑦ



2014–ᒥ
ᖁᑎᐅᔨᓴᖅᑕᐅᓪᓚ
ᐅᖅᑐᑦ ᑐᑐᐃᑦ
ᐅᓄᖅᓂᖓᑦ
Subpopulation
Individuals Observed
We are 95% certain the actual number of Caribou lay between these two values
ᖃᒥᓱᓂᖅ**
North Baffin
Borden Peninsula 1 1 -30
Mary River 49 96 - 521
North Central Baffin 13 31 - 230
Total 63 159 - 622
South Baffin
Central Baffin 197 662 – 1,798
Foxe Peninsula 20 48 - 972
Hall Peninsula 176 467 – 1,686
ᑭᖓᑕᐃᑦ 91 256 -1,138
Prince Charles Island 557 1,158 – 2,220
Total 824 3,169 – 5,935
Total (-Prince Charles Island) 267 1,777 – 4,207
Other areas
Melville Peninsula 26 88 - 551
Baffin Island + Melville P. 1,130 3,661 – 6,484
Baffin Island Total 1,104 3,462 – 6,250
Nunavut



ᑐᐃᑐᑦ ᖃᕆᔨᐅᓂᖅᑎᓂᒃ ᖃᐅᐅᓚᒪᔪᖅᑦ. (ᐊᒥᓱᓂᖅᑎᑦ)
ᖃᕆᔨᐅᔾᔪᑦ ᕿᖏᓂᑦ (ᓄᖅᓇᐃᑦᑐᑦ, ᕿᖏᓂᑦ), ᓄᖅᓇᐃᑦ, ᓄᖅᓇᒥᑦ ᐊᓱᖏᑎᒍᓐᓇ?

ᖃᐅᓇᐃᑕᖏᓐᓄᖏᓐᓄᑦ**


1) ᖃᐅᓂᖅᓯᑎᑦ ᑐᐃᑐᐃᑦ ᑲᑎᖏᓚᐅᑦ (ᐊᐋᖕᓇᓱᐊᓂᑐᑕᓚᐅᕐᓂᐅᑉᑯᑦ



ᖃᑦᓴᓂᐅᔨᒪᔪᑦ ᐊᑐᐊᓂᒃ (ᓄᓇᕗᑦᑐᓂᒃ, ᐊᑐᐊᓂᒃ), ᓄᓇᕗᑦ, ᓄᓇᕗᒻᒥᓂ ᐊᓯᖏᑦᓯᓗ?

ᖃᓄᐃᓕᖅᓯᒪᔪᖅ


2) ᑕᐅᔭᓐᓇ ᖃᖅᖢᑕᕆᒡᓗᑎᒃ ᑐᒥᖏᕐᓂᒃ ᐅᕐᕕᒃᓴᖅᓂᒃ ᑐᓴᑐᓂᒃ ᖃᓄᖅᓗᑎᒃ



ᖃᓄᐃᓕᖅᓯᒪᔪᑦ ᐊᑐᐊᓂᒃ (ᓄᓇᕗᑦᑐᓂᒃ, ᐊᑐᐊᓂᒃ), ᓄᓇᕗᑦ, ᓄᓇᕗᒻᒥᓂ ᐊᓯᖏᑦᓯᓗ?

ᖃᓄᐃᓕᖅᓯᒪᔪᖅ

3) ᖃᓄᐃᓕᖅᓂᖏᑦ ᐃᓚᓕᐅᑎᓗᑦ ᓇᓂᑐᐃᓇᖅᑐᖅ ᑕᒻᒧᐊᐅᑎᑕᐅᔨᓂᒃ ᐊᑕᖏᑎᖅᓗᑦ ᐊᑎᓕᓂᖏᓐᓇᓂᒃ




ᖃᑉᓯᐅᓪᓗᓂᓪᓗᓂ ᖃᑉᐳᔪᓴᕈᓯᖅ 2015-2018

ᑎᑭᑕᐅᓇᓯᐊᖅᑐᑦ


1) ᑐᑭᓯᐊᒪᓪᓗᑎᒃ ᖃᑉᓯᐃᖏᕐᕋᐊᖅᑐᓂᒃ ᑐᑦᑐᖓᓕᓂᒃ ᓄᖅᖢᓕᐳᖃᑉᕋᖅᑐᑦ ᐊᒻᒪ ᑭᓯᓂᖓᕐᕋᑦ ᒪᓕᒃᓗᑎᕐᕋᑦ; ᓯᖅᓗ ᓇᓕᐊᒃ ᑐᒃᑐᖃᑎᒋᑦ ᕿᖓᓂᒪᓗᐅᖏᕐᕋᓱᑦ, ᕿᖓᓂᐅᕐᓂᓪᓚᑦ, ᓄᖅᓴᕐᕋᐅᓪᓚᑦ, ᐃᓄᖅᑐᖃᑎᒋᓪᓚᑦ, ᐊᒻᒪ ᓄᖅᖃᖃᖅᕿᑦ.

2) ᑐᑭᓯᐊᒪᓪᓗᑎᒃ ᐃᖅᓂᐅᖅᑐᑦ ᐊᖅᑯᑎᕐᓂᓂᒃ ᑐᒃᑐᖃᑎᒋᑦ ᑭᓯᓂᖓᕐᕋᑦ ᒪᓕᓃᓗᑦ· ᐊᒻᒪ ᐅᐱᖅᓯᓴᑯᑯᑦ ᖃᑉᓯᐃᓕᓚᖓᓂᓪᓗᒃ ᖃᑉᐳᔪᓴᕈᐅᔪᑦ, ᑐᑭᓯᐊᓇᓚᓗᑎᒃ ᓀᓗᓀᐃᐱᖅᓯᒪᔪᑦ ᓄᖅᖃᖃᖅᕿᑦ ᐊᒻᒪ ᐅᑭᐅᓂᖅᑯᑦ ᐊᓐᓀᐅᕐᓂᖅᑐᑦ ᐅᓄᖅᓯᓂᖅᓴᓕᓕᖅᑐᑦ ᐊᒻᒪ ᐅᑭᐅᓂᖅᑯᑦ ᐅᓄᖅᓯᓂᖅᓴᓕᓕᖅᑐ


ᖃᒃᓄᐃᒋᓐᓂᓐᓂᓕᓂ ᖃᐅᔨᓴᖅᑭᓱᓂ ᕿᑭᖅᑕᕐᓗᓐᒥ ᕿᑎᖅᓯᓱᐊᓂ



ᖃᒃᓄᐃᒋᓐᓂᓐᓂᓕᓂ ᖃᐅᔨᓴᖅᑭᓱᓂ ᕿᑭᖅᑕᕐᔪᑦ ᓂᒋᐊᓂ




ᖃᑲᓇᐸᓐᓚᓂᓐᓚᓂ ᖃᑎᔨᓴᕈᓯᓂ 2015-2018



**ᓴᓗᔨᓕᒪᖏᐃᐅᔪᓂ
ᐃᓂᓐᒋᑦ

ᖃᑲᓇᐸᓐᓚᓂᓐᓚᓂ ᖃᑎᔨᓴᕈᓯᓂ ᖃᑎᔨᓴᕈᑕᐅᔪᓂ



ᖃᑎᔨᓴᕈᓯᓂ



ᖃᑎᔨᓴᕈᓯᓂ

ᐱᑕᓚᐊᖓᕆᔨᐅᔪᓂ ᓄᖅᕿᐃᑦ: ᓄᖅᕿᐃᑦ ᖃᑎᔨᓴᕈᓂᐅᓐᓂᑦ ᓄᓇᕐᔪᐊᒥ ᓂᖅᖁᓗᐊᖏᑎᑐᑦ ᐅᕐᖏᓪᓗᓂᑦ ᐅᓄᖅᓂᖅᓯᓚᐊᔪᓂ:

- 70-90 ᓄᖅᕿᐃᑦ
- 50-70 ᐅᑭᐊᒃᓴᓄᑦ
- 30-50 ᐅᑭᐅᖅᑎᑦᓗᒍ

40 <ᓐᓂᑦ:100 ᓄᖅᕿᐃᑦ ᓇᓗᓇᐃᓄᑦᑎᑦᓄᐅᔾᔨᑦ <ᓐᓂᑦ ᐃᓕᐅᑎᖏᑐᓄᑦ ᖃᑎᔨᐅᓕᖅᖢᖅᓯᓂᐅᓐᓂᒃ ᑐᑐᖏᑦ ᓄᖅᕿᓚᐅᔾᔨᑦᓗᒍ (ᑐᐱ 2001).

ᒥᓄᑦᐊ
ᓇᓗᓇᐃᖅᑕᐅᓂᒪᑐᐃᓐᓇ
ᖃᓄᑦᑐ?

ᖃᑦᓴᐅᓂᖅᓯᓂᑦ ᖃᐅᔨᓴᖅᓂᖅ ᐊᐱᖅᓱᖅᓯᖅᓂᑦ ᖃᐅᔨᓴᐅᓱᒪᔪᑦ – ᓂᒋᖅᒥ

ᐅᑭᐊᒃᓴᖅ

Table 4 Number of observed caribou by demographic group during Baffin Island composition surveys 2015-2018.

Year	2015				2016				2016				2017				2017				2018			
Season	Fall				Spring				Fall				Spring				Fall				Spring			
Location	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island
Calves Observed	55	28	133	49		23	82	49	54			81	47	1	114	92	86				21	18	31	155
Cows Observed	77	39	189	64		67	328	222	94			196	120	1	351	249	139				36	33	161	401
Calves/100 Cows	71	72	70	77		34	25	22	57			41	39	100	32	37	62				58	55	19	39
Yearlings Observed	N/A	N/A	N/A	N/A		10	76	29	N/A			42	23	0	57	75	17				5	7	37	100
Bulls Observed	76	29	126	46		25	204	151	54			126	64	6	133	181	74				38	40	73	277
Bulls/100 Cows	99	74	67	72		37	62	68	57			64	53	600	38	73	53				106	121	45	69
Bull + Cows	153	68	315	110		92	532	373	148			322	184	7	484	430	213				74	73	234	678
Adults + Yearlings Observed	153	68	315	110		102	608	402	148			364	207	7	541	505	230				79	80	271	778
Total Observed (Calves, Yearlings and Adults)	208	96	448	159		125	690	451	202			445	254	8	655	597	316				100	98	302	933

Not completed

40 ᐊᖓᔪᖅᓴᐃᑦ : 100 ᐊᖏᓐᓴᐃᑦ

40 ᐊᖓᔪᖅᓴᐃᑦ : 100 ᐊᖏᓐᓴᐃᑦ

40 ᐊᖓᔪᖅᓴᐃᑦ : 100 ᐊᖏᓐᓴᐃᑦ

ᐊᑲᐅᔪᓐᓇᖅᑐᑦ

ᑎᓪᓗᒡᓗᖕᖓ ᐊᖓᔪᖅᓴᐃᑦ ᐃᓚᒋᔭᖏᑦᑐᓂ ᐊᖏᓐᓴᐃᓂ ᓄᓪᐊᑭᐅᓐᖏᖏᑦᑐᓂ

ᖃᑦᓴᐅᓂᖅᓯᓂᑦ ᖃᐅᔨᓴᖅᓂᖅ ᐊᐱᖅᓱᖅᓯᖅᓂᑦ ᖃᐅᔨᓴᐅᓱᒪᔪᑦ – ᓂᒋᖅᒥ

ᐅᐱᖏᓐᓇᖅ

Table 4 Number of observed caribou by demographic group during Baffin Island composition surveys 2015-2018.

Year	2015				2016				2016				2017				2017				2018			
Season	Fall				Spring				Fall				Spring				Fall				Spring			
Location	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island
Calves Observed	55	28	133	49		23	82	49	54			81	47	1	114	92	86				21	18	31	155
Cows Observed	77	39	189	64		67	328	222	94			196	120	1	351	249	139				36	33	161	401
Calves/100 Cows	71	72	70	77		34	25	22	57			41	39	100	32	37	62				58	55	19	39
Yearlings Observed	N/A	N/A	N/A	N/A		10	76	29	N/A			42	23	0	57	75	17				5	7	37	100
Bulls Observed	76	29	126	46		25	204	151	54			126	64	6	133	181	74				38	40	73	277
Bulls/100 Cows	99	74	67	72		37	62	68	57			64	53	600	38	73	53				106	121	45	69
Bull + Cows	153	68	315	110		92	532	373	148			322	184	7	484	430	213				74	73	234	678
Adults + Yearlings Observed	153	68	315	110		102	608	402	148			364	207	7	541	505	230				79	80	271	778
Total Observed (Calves, Yearlings and Adults)	208	96	448	159		125	690	451	202			445	254	8	655	597	316				100	98	302	933

Not completed

? ᓄᖏᐃᑦ : 100 ᐊᖏᓐᓴᐃᑦ

30–ᒥᑦ 50–ᒧᑦ ᓄᖏᐃᑦ : 100 ᐊᖏᓐᓴᐃᑦ

30–ᒥᑦ 50–ᒧᑦ ᓄᖏᐃᑦ : 100 ᐊᖏᓐᓴᐃᑦ

ᐊᑲᐅᔪᓐᓇᖅᑐᑦ

ᑎᓪᓗᒡᓗᖕᖓ ᓄᖏᓗᓂ ᐊᖏᓗᐅᒪᓕᑎᑎᒃ ᐅᑭᐅᐸ ᖃᓄᐃᓕᐅᕐᓂᑦ

ᖃᑦᓴᐅᓂᖅᓯᓂᑦ ᖃᐅᔨᓴᖅᓂᖅ ᐊᐱᖅᓱᖅᓂᖅᓯᒪᓂᑦ ᖃᐅᔨᓴᐅᔨᒪᔪᑦ – ᖃᑭᑎᐊᓂ

ᐅᑭᐊᒃᓴᖅ

Table 4 Number of observed caribou by demographic group during Baffin Island composition surveys 2015-2018.

Year	2015				2016				2016				2017				2017				2018			
Season	Fall				Spring				Fall				Spring				Fall				Spring			
Location	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island
Calves Observed	55	28	133	49		23	82	49	54			81	47	1	114	92	86				21	18	31	155
Cows Observed	77	39	189	64		67	328	222	94			196	120	1	351	249	139				36	33	161	401
Calves/100 Cows	71	72	70	77		34	25	22	57			41	39	100	32	37	62				58	55	19	39
Yearlings Observed	N/A	N/A	N/A	N/A		10	76	29	N/A			42	23	0	57	75	17				5	7	37	100
Bulls Observed	76	29	126	46		25	204	151	54			126	64	6	133	181	74				38	40	73	277
Bulls/100 Cows	99	74	67	72		37	62	68	57			64	53	600	38	73	53				106	121	45	69
Bull + Cows	153	68	315	110		92	532	373	148			322	184	7	484	430	213				74	73	234	678
Adults + Yearlings Observed	153	68	315	110		102	608	402	148			364	207	7	541	505	230				79	80	271	778
Total Observed (Calves, Yearlings and Adults)	208	96	448	159		125	690	451	202			445	254	8	655	597	316				100	98	302	933

Not completed

40 ᐊᖏᓴᓪᓗᐃᑦ : 100 ᐊᖏᓐᓇᓪᓗᐃᑦ ? ᐊᖏᓴᓪᓗᐃᑦ : 100 ᐊᖏᓐᓇᓪᓗᐃᑦ ? ᐊᖏᓴᓪᓗᐃᑦ : 100 ᐊᖏᓐᓇᓪᓗᐃᑦ

ᐊᑲᐅᔪᖏᓪᓗᒍ

2015–ᒥ ᑎᓕᐅᕐᓂᖅᓯ ᐊᖏᓴᓪᓗᐃᑦ ᐃᓚᒪᓂᖅᓯᓂ ᐊᖏᓐᓇᓪᓗᓯ ᓄᓕᐊᑭᓐᓇᓂᖅᓯᓂ

ᖃᑦᓴᐅᓂᖅᓯᓂᑦ ᖃᐅᔨᓴᖅᓂᖅᓯᒪᓂᑦ ᖃᐅᔨᓴᐅᔨᒪᔪᑦ – ᖃᑭᑎᐊᓂ

ᐅᐱᖏᓪᓗᖅ

Table 4 Number of observed caribou by demographic group during Baffin Island composition surveys 2015-2018.

Year	2015				2016				2016				2017				2017				2018			
Season	Fall				Spring				Fall				Spring				Fall				Spring			
Location	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island
Calves Observed	55	28	133	49		23	82	49	54			81	47	1	114	92	86				21	18	31	155
Cows Observed	77	39	189	64		67	328	222	94			196	120	1	351	249	139				36	33	161	401
Calves/100 Cows	71	72	70	77		34	25	22	57			41	39	100	32	37	62				58	55	19	39
Yearlings Observed	N/A	N/A	N/A	N/A		10	76	29	N/A			42	23	0	57	75	17				5	7	37	100
Bulls Observed	76	29	126	46		25	204	151	54			126	64	6	133	181	74				38	40	73	277
Bulls/100 Cows	99	74	67	72		37	62	68	57			64	53	600	38	73	53				106	121	45	69
Bull + Cows	153	68	315	110		92	532	373	148			322	184	7	484	430	213				74	73	234	678
Adults + Yearlings Observed	153	68	315	110		102	608	402	148			364	207	7	541	505	230				79	80	271	778
Total Observed (Calves, Yearlings and Adults)	208	96	448	159		125	690	451	202			445	254	8	655	597	316				100	98	302	933

Not completed

30–ᒥᓵ 50–ᒍᑦ ᓄᖏᐃᑦ : 100 ᐊᖏᓐᓇᓪᓗᐃᑦ 30–ᒥᓵ 50–ᒍᑦ ᓄᖏᐃᑦ : 100 ᐊᖏᓐᓇᓪᓗᐃᑦ 30–ᒥᓵ 50–ᒍᑦ ᓄᖏᐃᑦ : 100 ᐊᖏᓐᓇᓪᓗᐃᑦ

ᐊᑲᐅᔪᖏᓪᓗᒍ

ᑎᓕᐅᕐᓂᖅᓯ ᓄᖏᓄᑦ ᐊᓐᓇᐅᒪᓕᕆᔨ ᐅᑭᐅᒋ ᖃᓄᐃᓕᖅᑎᑦᑐᑦ

ᖃᑦᓴᐅᓂᖅᓯᓂᑦ ᖃᐅᔨᓴᖅᓂᖅ ᐊᐱᖅᓱᖅᓂᖅᑎᑦ ᖃᐅᔨᔪᐅᓱᒻᒪᓕᒃ – ᑭᖑᓪᓗᖅᑎᒃ

ᐅᑭᐊᒃᓴᖅ

Table 4 Number of observed caribou by demographic group during Baffin Island composition surveys 2015-2018.

Year	2015				2016				2016				2017				2017				2018			
Season	Fall				Spring				Fall				Spring				Fall				Spring			
Location	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island
Calves Observed	55	28	133	49		23	82	49	54			81	47	1	114	92	86				21	18	31	155
Cows Observed	77	39	189	64		67	328	222	94			196	120	1	351	249	139				36	33	161	401
Calves/100 Cows	71	72	70	77		34	25	22	57			41	39	100	32	37	62				58	55	19	39
Yearlings Observed	N/A	N/A	N/A	N/A		10	76	29	N/A			42	23	0	57	75	17				5	7	37	100
Bulls Observed	76	29	126	46		25	204	151	54			126	64	6	133	181	74				38	40	73	277
Bulls/100 Cows	99	74	67	72		37	62	68	57			64	53	600	38	73	53				106	121	45	69
Bull + Cows	153	68	315	110		92	532	373	148			322	184	7	484	430	213				74	73	234	678
Adults + Yearlings Observed	153	68	315	110		102	608	402	148			364	207	7	541	505	230				79	80	271	778
Total Observed (Calves, Yearlings and Adults)	208	96	448	159		125	690	451	202			445	254	8	655	597	316				100	98	302	933

Not completed

40 ᐊᖏᓴᓪᓚᐃᑦ : 100 ᐊᖏᓪᓚᐃᑦ ? ᐊᖏᓴᓪᓚᐃᑦ : 100 ᐊᖏᓪᓚᐃᑦ ? ᐊᖏᓴᓪᓚᐃᑦ : 100 ᐊᖏᓪᓚᐃᑦ

ᐊᑲᐅᑎᕐᓯᒪᓂᖅᑎᑦ

2015–ᒥ ᑎᓕᐅᕆᔨᖅ ᐊᖏᓴᓪᓚᐃᑦ ᓄᓚᒪᓂᖓᓂᓐ ᐊᖏᓪᓗᓪᓘᓐ ᓄᓕᐊᑭᓐᓇᖅᓯᓂᖓᓂᓐ

ᖃᑦᓴᐅᓂᖅᓯᓂᑦ ᖃᐅᔨᓴᖅᓂᖅᑎᑦ ᐊᐱᖅᓱᖅᓂᖅᑎᑦ ᖃᐅᔨᔪᐅᓱᒻᒪᓕᒃ – ᑭᖑᓪᓗᖅᑎᒃ

ᐅᐱᖅᓯᖅ

Table 4 Number of observed caribou by demographic group during Baffin Island composition surveys 2015-2018.

Year	2015				2016				2016				2017				2017				2018			
Season	Fall				Spring				Fall				Spring				Fall				Spring			
Location	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island
Calves Observed	55	28	133	49		23	82	49	54			81	47	1	114	92	86				21	18	31	155
Cows Observed	77	39	189	64		67	328	222	94			196	120	1	351	249	139				36	33	161	401
Calves/100 Cows	71	72	70	77		34	25		57			41	39	100	32		62				58	55	19	
Yearlings Observed	N/A	N/A	N/A	N/A		10	76	29	N/A			42	23	0	57	75	17				5	7	37	100
Bulls Observed	76	29	126	46		25	204	151	54			126	64	6	133	181	74				38	40	73	277
Bulls/100 Cows	99	74	67	72		37	62	68	57			64	53	600	38	73	53				106	121	45	69
Bull + Cows	153	68	315	110		92	532	373	148			322	184	7	484	430	213				74	73	234	678
Adults + Yearlings Observed	153	68	315	110		102	608	402	148			364	207	7	541	505	230				79	80	271	778
Total Observed (Calves, Yearlings and Adults)	208	96	448	159		125	690	451	202			445	254	8	655	597	316				100	98	302	933

Not completed

30–ᒥᐢ 50–ᒍᐢ ᓄᖅᕃᐢ : 100 ᐊᖏᓪᓚᐃᑦ 30–ᒥᐢ 50–ᒍᐢ ᓄᖅᕃᐢ : 100 ᐊᖏᓪᓚᐃᑦ 30–ᒥᐢ 50–ᒍᐢ ᓄᖅᕃᐢ : 100 ᐊᖏᓪᓚᐃᑦ

ᐊᑲᐅᑎᕐᓯᒪᓂᖅᑎᑦ

ᑎᓕᐅᕆᔨᖅ ᓄᖅᕃᓄᑦ ᐊᖏᓇᐅᒪᓕᑎᑎᒃ ᐅᑭᐅᒋ ᓇᒪᓐᕆᑎᑐᖅ

ᖃᑦᓴᐅᓂᖅᓯᓂᑦ ᖃᐅᔨᓴᖅᓂᖅ ᐊᐱᖅᓱᖅᓯᖅᓂᖅ ᖃᐅᔨᓴᐅᓱᒪᔪᑦ – ᐊᑭᓐᓇᖅ

ᐅᑭᐊᒃᓴᖅ

Table 4 Number of observed caribou by demographic group during Baffin Island composition surveys 2015-2018.

Year	2015				2016				2016				2017				2017				2018			
Season	Fall				Spring				Fall				Spring				Fall				Spring			
Location	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island
Calves Observed	55	28	133	49		23	82	49	54			81	47	1	114	92	86				21	18	31	155
Cows Observed	77	39	189	64		67	328	222	94			196	120	1	351	249	139				36	33	161	401
Calves/100 Cows	71	72	70	77		34	25	22	57			41	39	100	32	37	62				58	55	19	39
Yearlings Observed	N/A	N/A	N/A	N/A		10	76	29	N/A			42	23	0	57	75	17				5	7	37	100
Bulls Observed	76	29	126	46		25	204	151	54			126	64	6	133	181	74				38	40	73	277
Bulls/100 Cows	99	74	67	72		37	62	68	57			64	53	600	38	73	53				106	121	45	69
Bull + Cows	153	68	315	110		92	532	373	148			322	184	7	484	430	213				74	73	234	678
Adults + Yearlings Observed	153	68	315	110		102	608	402	148			364	207	7	541	505	230				79	80	271	778
Total Observed (Calves, Yearlings and Adults)	208	96	448	159		125	690	451	202			445	254	8	655	597	316				100	98	302	933

Not completed

40 ᐊᖓᔪᖅᓵᐃᓪᓕᐊᓂ : 100 ᐊᖏᓐᓂᐃᓪᓕᐊᓂ

40 ᐊᖓᔪᖅᓵᐃᓪᓕᐊᓂ : 100 ᐊᖏᓐᓂᐃᓪᓕᐊᓂ

? ᐊᖓᔪᖅᓵᐃᓪᓕᐊᓂ : 100 ᐊᖏᓐᓂᐃᓪᓕᐊᓂ

ᐊᑲᐅᑎᕐᓗᓂᑦ

2015–ᒥ ᑎᓕᐅᓚᐅᖅᑐᑦ ᐊᖓᔪᖅᓵᐃᓪᓕᐊᓂ ᓇᓱᒪᓂᖅᑎᓂ ᐊᖏᓐᓂᔅᓴᖏᓂ ᓄᓕᐊᑭᓐᓇᖏᓂᖅᓂ

ᖃᑦᓴᐅᓂᖅᓯᓂᑦ ᖃᐅᔨᓴᖅᓂᖅ ᐊᐱᖅᓱᖅᓯᖅᓂᖅ ᖃᐅᔨᓴᐅᓱᒪᔪᑦ – ᐊᑭᓐᓇᖅ

ᐅᐱᖏᓪᓗᖅ

Table 4 Number of observed caribou by demographic group during Baffin Island composition surveys 2015-2018.

Year	2015				2016				2016				2017				2017				2018			
Season	Fall				Spring				Fall				Spring				Fall				Spring			
Location	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island	North Baffin Island	Central Baffin Island	Prince Charles Island	South Baffin Island
Calves Observed	55	28	133	49		23	82	49	54			81	47	1	114	92	86				21	18	31	155
Cows Observed	77	39	189	64		67	328	222	94			196	120	1	351	249	139				36	33	161	401
Calves/100 Cows	71	72	70	77		34	25	22	57			41	39	100	32	37	62				58	55	19	39
Yearlings Observed	N/A	N/A	N/A	N/A		10	76	29	N/A			42	23	0	57	75	17				5	7	37	100
Bulls Observed	76	29	126	46		25	204	151	54			126	64	6	133	181	74				38	40	73	277
Bulls/100 Cows	99	74	67	72		37	62	68	57			64	53	600	38	73	53				106	121	45	69
Bull + Cows	153	68	315	110		92	532	373	148			322	184	7	484	430	213				74	73	234	678
Adults + Yearlings Observed	153	68	315	110		102	608	402	148			364	207	7	541	505	230				79	80	271	778
Total Observed (Calves, Yearlings and Adults)	208	96	448	159		125	690	451	202			445	254	8	655	597	316				100	98	302	933

Not completed

30–ᒥᑦ 50–ᒧᑦ ᓄᖅᓴᐃᑦ : 100 ᐊᖏᓐᓂᐃᓪᓕᐊᓂ

30–ᒥᑦ 50–ᒧᑦ ᓄᖅᓴᐃᑦ : 100 ᐊᖏᓐᓂᐃᓪᓕᐊᓂ

30–ᒥᑦ 50–ᒧᑦ ᓄᖅᓴᐃᑦ : 100 ᐊᖏᓐᓂᐃᓪᓕᐊᓂ

ᐊᑲᐅᑎᕐᓗᓂᑦ

ᑎᓕᐅᓚᐅᖅᑐᑦ ᓄᖅᓴᓄᑦ ᐊᓐᓇᐅᒪᓗᑎᒃ ᐅᑭᐅᖓ ᖃᓄᐃᓇᕐᓂᑐᑦ

ᖃᓄᐃᒥᖅᓯᖅᓯᓂ ᖃᐅᔨᓴᕐᓂ
2019-Γ ᐅᐱᕐᖁᑎᑦ

ᐃᓐᓇ <ᕐᓇᐃᒥᖅᑐᑦ ᐊᑦᓴᖅᑐᓚᑦᓴᓂᒃ ᐅᐱᕐᖁᑎᑦ
ᖃᐅᔨᓴᓪᖅᑐᓄᑦ ᐃᓕᔨ/ᐊᐃᓗᒥ 2019–Γ

ᖃᐳᖅᑕᕐᔪᑉ ᓂᒋᐊᓄᑦ ᖃᑎᖅᖢᓕᐊᓄᑦᓗ <ᕐᓇᐃᔪᑦ

ᓄᔪᓴᒥᓚᔨᑎᓂᒃ ᐅᕐᖃᕐᕿᑎᖅᖃᖅᑐᑦ ᖃᐳᖅᑕᕐᔪᑉ
ᐅᐊᖏᓇᖏᓂᒃ ᐃᑲᔪᖅᑐᑐᔪᑦ ᐱᓕᕆᑎᒋᓪᓗᒋᑦ

ᓴᓇᕐᕿᑎᑦᓴᓂᒃ ᑐᓴᑐᐃᑦ ᖃᐳᖅᑕᕐᔪᖓᒥ
ᖃᐅᔨᓴᕐᕿᓗᒋᑦ

– <ᕐᓇᖏᓪᓗᓂᒃ ᐅᑎᖅᑎᖏᓇᓂᖏᓪᓗᒋᑦ

ᖁᖏᓚᒋᔭᖅᔪᑦ ᖁᐅᐊᓴᓴᖁᓂᒃᑐᑦ

- ᑐᖅᑐᓂᖅ ᑕᐅᑐᖏᓂᒥᑦ ᑐᓗᕈᑎᖁᖏᑕᖅᓂᖅᓯᖕᖏᖅ.
- ᐊᖏᑕᐅᓂᖅ ᐃᓂᐅᐊᓂᖅ ᖃᑎᖅᕿᖅᓀᖅᑐᑦ.
- ᐊᑐᖅᑕᐅᐊᖅᓀᖅᑐᑦ ᐅᓄᖅᓂᖏᑕ ᖁᓄᐃᑕᖏᓚᓂᖏᑦ ᓀᓗᓀᐃᕆᔨᓗᑎᑦ.
- ᐊᑐᖅᑕᐅᐊᖅᓀᖅᑐᑦ ᐅᓄᖅᓂᖏᑦ ᓀᓪᓕᐅᑦᑕᕆᔨᓗᑎᑦ.
- ᐊᑐᖅᑕᐅᐊᖅᓀᖅᑐᑦ ᑎᑎᕆᖅᑕᐅᓗᑎᒃ ᐅᓄᖅᑐᑦ ᓀᒻᓂᓂᖏᑕ ᐊᓯᔪᖅᓴᖅᑕᓪᓕᐊᓂᖏᓐᓂᓂ.


ᖁᑦᓯᐅᓂᖏᓂᓂᑦ ᖁᐅᐊᓴᓴᖁᓂᓂᖅ ᐊᐱᖅᓯᖁᓂᖏᑕ

- ᐊᑭᑐᐊᖅᓀᖅᑐᖅ.
- ᐊᑐᖅᑕᐅᐊᖅᓀᖅᑐᑦ ᓀᓗᓀᐃᒃᑯᑕᐅᓗᓂ ᐅᓄᖅᓂᖏᑕ ᖁᓄᐃᑕᖏᓚᓂᖏᓐᓂᒃ.
- ᐊᑐᖅᑕᐅᐊᖅᓀᖅᑐᑦ ᐊᐅᓪᓚᒃᑎᑦᑎᓂᖅᒥ ᐊᑭᑐᓂᖅᑲᓂᒃ ᖁᓄᐃᑕᖏᓚᓂᖏᓐᓂᒃ ᐊᖅᓀᐅᒪᓂᖅᒧᑦ ᖁᐅᐊᓴᓴᖁᖅᑐᑦ.
- ᐊᖅᓀᑦᓗᐃᑦ ᐊᖏᔪᖅᑦᓗᐃᑦᓗ ᐊᓯᔪᖅᓴᖅᑕᓪᓕᐊᓂᖏᓐᓄᓐ ᐊᖅᓀᐅᐅᒪᓂᖅᒧᑦ ᖁᐅᐊᓴᓴᖁᖅᑐᑦ.
- ᐃᓗᓀᐃᒃᑯᑕᐅᐊᖅᓀᖅᓴᑦ ᐅᓄᖅᓯᓚᑦᓕᐊᓂᖏᓐᓄᓐ.


ᓂᐱᓕᐅᓗᓂᕐᒧᑦ ᖃᐅᔨᓴᖅᓂᖅ


- ᐊᑭᖑᓂᓐ ᐊᑐᕆᐊᖅᓂᖅ.
- ᖃᐅᖑᑕᓯᒃᑯᑦ ᐊᒻᒪᓗ ᖃᐅᔨᓯᐅᓂᖑᓂᓐᓂᑦ ᖃᐅᔨᓴᖅᓂᐅᕐᓂ ᐊᑭᖑᓂᓐᓂ ᐊᑎᕆᔨᕆᑎᒪᓕᕆᔨ.
- ᐊᑐᖅᑕᐅᔪᓐᓇᖅᑐᑦ ᓇᓗᓇᐃᕆᔨᓴᓕᕆᓂ ᐊᖅᖃᔪᑕᒧᑦ ᑎᑭᑦᑕᖅᖑᑕ ᐃᓂᖑᓂᓐᓂ.
- ᐊᑐᖅᑕᐅᔪᓐᓇᖅᑐᑦ ᓇᓗᓇᐃᕆᔨᓴᓕᕆᓂ ᐊᖅᖃᔪᑦ ᐃᓗᐊᓂᑐᑦ ᐃᓂᖑᓂᓐᓂ.
- ᐊᑐᖅᑕᐅᔪᓐᓇᖅᑐᑦ ᐃᑲᔪᖅᓭᐃᓗᑎᒃ ᖃᐅᔨᓴᖅᓂᖅᓂᒥ ᐱᑕᕆᐊᖑᔪᑦ.
- ᐊᑐᖅᑕᐅᔪᓐᓇᖅᑐᑦ ᓇᓗᓇᐃᕆᔨᓴᓕᕆᓂ ᐊᑐᕆᐊᖅᑲᒪᕆᔨᒃᑐᑦ ᐃᓂᕐᒥᐅᑐᓂ.
- ᐊᑐᖅᑕᐅᔪᓐᓇᖅᑐᑦ ᓴᐳᑎᓇᓕᕆᓂ ᐊᑐᕆᐊᖅᑲᒪᕆᔨᒃᑐᑦ ᐃᓂᕐᒥᐅᑐᓂ.






ᖃᒃᓗᓂᒃ ᓂᐱᒋᐅᕆᓚᓂᒃ ᐊᑐᖅᑕᐅᓂᖅ



Winter
(Dec 16 - Apr 14)
Spring Migration
(Apr 15 - Jun 8)
Calving
(Jun 9 - 22)
Post-calving
(Jun 23 - Jul 3)
Summer
(Jul 4 - Aug 22)
Late Summer
(Aug 23 - Sep 16)
Fall Migration, pre-breeding
(Sep 17 - Oct 18)
Rut/Breeding
(Oct 19 - Nov 6)
Fall Migration, post-breeding
(Nov 7 - Dec 15)
Average movement rate (km/day)
Date
01-Jan
31-Jan
02-Mar
01-Apr
01-May
31-May
30-Jun
30-Jul
29-Aug
28-Sep
28-Oct
27-Nov
27-Dec


Winter
(Dec 16 - Apr 14)
Spring Migration
(Apr 15 - Jun 8)
Calving
(Jun 9 - 22)
Post-calving
(Jun 23 - Jul 3)
Summer
(Jul 4 - Aug 22)
Late Summer
(Aug 23 - Sep 16)
Fall Migration, pre-breeding
(Sep 17 - Oct 18)
Rut/Breeding
(Oct 19 - Nov 6)
Fall Migration, post-breeding
(Nov 7 - Dec 15)
Average movement rate (km/day)
Date
01-Jan
31-Jan
02-Mar
01-Apr
01-May
31-May
30-Jun
30-Jul
29-Aug
28-Sep
28-Oct
27-Nov
27-Dec

ᓇᓗᓇᐃᖅᓯᓂᖅ ᐊᕐᕋᔪᑦ ᐃᓗᐊᓂᑉᑐᑦ ᐃᓂᖏᓐᓂ




ᓄᓇᒥᒃ ᐊᑐᕐᓂᖅ ᐊᕐᕋᔪᑦ ᐃᓕᓴᕆᓂ – ᕿᑐᕐᖏᐅᑦᓂᖅ




ᓄᓇᒥᐅᑦ ᐊᑐᕐᓂᖅ ᐊᖏᖅᔪᓂ ᐃᓕᓴᕐᓄᑦ – ᐃᓐᓕᕐᕋᓂᖓᓕᒃ ᐃᓚᒋᓂᖅ

ᑲᑎᓐᓴᒪᐅᖅᑐᓂᒃ ᓇᓗᓇᐃᔭᐃᓕᑎᐊᕐᓂᖅ




ᓄᓇᕗᑦᓂᐅᓕᖅ ᖃᐅᔨᓴᖅ




ᓄᓇᕗᑦ ᑐᖓᓕᖅ ᑎᒥᖅᑎᑦ (NTI)–ᓂᑦ ᐅᒪᔨᓪᓗᓵᕕᑎᖅᑐᑦ ᑕᐃᖃᑎ ᓱᓕ ᓂᕆᔨᓗ ᓂᕆᔨᑎᓄᑦ ᐱᓪᓗᓵᓕ ᐊᒥᓱᓄ ᐊᑐᕐᓂᖅ ᐅᓂᒃᑳᖅ ᐊᒥᓱᓄᑦ ᐃᓕᓴᕐᓂᖅᑐᑦ ᐱᓕᕆᔨᒃᑯᑦ ᐅᖃᐅᔨᓴᓪᓚᓂᖅᓴᓕᖅᑐᑦ/ᐃᖃᖅᓄᑦ ᐃᓂᐅᔨᒥ.

ᓇᓱᐊᑦ ᖃᖅᖠᓂᕆᑐᖅᑎᑕᐅᓂᒪᔨᒥᒃ ᐊᓘᐃᓇᒍᐊᑦ ᐅᓴᖅᑕᕐᕕᐅᔨᑦ ᑭᓇᐅᔭᖅ ᐊᖃᓕᑕᖅᓂᒪᔨᒥᑦ ᓄᓇᕗᒻᒥ.

ᐅᓴᖅᑕᐃᓂᖅ ᐅᓴᖅᑲᓂᖅ ᓴᖅᑭᐱᖅᑕᐃᓂᖅ ᖃᖅᑲᓂᕐᒪᑐᖅ ᖃᓄᐃᓂᖅ.





ᑐᖅᑐᐃᑦ ᓴᐳᑎᑐᒋᑦ ᓴᓇᔭᐅᓚᓂᑦᑕᐊᔪᓂᒃ




ᖃᑐᐊᖅᑕᐅᔪᓐᓇᒪᓗᒃ




ᐅᐱᐊᖅᓯᓚᑦ ᐃᖅᑎᕋᓂᖅᑎᑦ ᐃᓚᓂᓂᑦ




ᖃᑐᐊᖏᑎᐅᖅᓯᒪᓕᓚᖅᑎᑦᓴᑎᑦ



2018


Telemetry Dates: Jul 20, 2018 to Jul 21, 2018
Baker Lake
Fall 2018
Legend

2016


Telemetry Dates: Jul 20, 2016 to Jul 21, 2016
Baker Lake
Fall 2016
Legend



ᑎᑎᕆᖅᑕᐅᕈᐊᖏᓚᐅᔪᑦ ᐊᐅᓪᓚᑎᓂᕐᒧᑦ ᐸᖅᓇᐅᑎᓄᑦ

ᐊᖏᔪᓇᓕᔪᑦ ᒥᑭᕈᐊᖅᓂᐊᖅᑭᓪᓗ ᑲᑐᔾᔨᖅᑎᑎᑭᖏᔪᓄᑦ (HTO)–ᑲᑯᓐᓄᑦ ᐊᐅᓪᓚᖅᑎᑕᐅᔪᑦ ᓯᑎᐱᓗ 2018–ᒥᑦ

ᖃᐅᐅᔨᓴᒪᔪᑦ ᓄᓇᓕᖏᓐᓂ ᐃᓕᒧᑦᑐᑕᐅᔨᓂᖅ, ᐊᐱᖅᑯᑎᓂᖅ ᐊᓱᒪᓗ ᐅᖃᐅᓕᓕᓚᓂᖅ


GOVERNMENT OF NUNAVUT: DEPARTMENT OF ENVIRONMENT


Baffin Island Caribou Management Plan

Working together to ensure Baffin Island caribou harvest is sustainable

GN DOE

July 2018

DRAFT


ᓄᓇᕗᑦ ᓴᖅᓯᒪᓕ ᐊᖏᓚᐅᔪᑦ


ᖃᐅᖅᑕᒪᓕᒃ ᑐᑭᑕᓱᓂᕐᒥᑦ ᐊᐅᓪᓚᔨᓴᒪᓂᖅᓂᑦ ᐸᖅᓇᐅᑎᓄᑦ


ᐃᓕᓴᖅᑎᑎᕐᓂᖅ ᓇᓗᓇᐃᔨᓴᒪᓂᑦ ᖃᐅᖅᑕᒪᓕᒃ ᑐᑭᑕᓱᓂᑦ ᐊᖏᔪᓇᓕᖏᓐᓂ ᑲᔪᓯᔨᐅᖏᓐᓇᓴᒪᓂᑦ ᓯᑎᐱᓗ

GN DOE

ᐅᓇ 2018

DRAFT


ᐊᐢᑎᓕᓱᓴᒃᑯᑦ ᐱᓕᓱᐊᖅᑎᑦ
ᓄᓇᕗᑦ ᐅᒪᔨᓕᓱᐃᓐᓇᖅᑐᑦ ᑲᑎᒪᔨᖅᑎᑦ
Nunavut Wildlife Management Board
QIKIQTAALUK WILDLIFE BOARD
ᖃᐅᔨᒪᔭᕐᓗᒃᑐᑦ
ᐊᖏᓐᓇᓕᒃᑯᑦ
ᒥᑭᑎᐊᖅᑐᓂᒃᑲ
ᑲᑐᔾᔨᖃᑎᒌᖏᓐᓂᑦ
(HTO)–ᖅᑎᑦ
ᓄᓇᕗᒻᒥᐅᑦ
NUNAVUT TUNNGAVIK
Nunavut


ᐊᐊᐅᓪᓗᒍᑦ
ᐱᖓᓱᒃᑐᖅ
ᓄᓇᕗᑦ ᐅᒪᔨᓕᓱᐃᓐᓇᖅᑐᑦ ᑲᑎᒪᔨᖅᑎᑦ
Nunavut Wildlife Management Board
QIKIQTAALUK WILDLIFE BOARD
ᖃᐅᔨᒪᔭᕐᓗᒃᑐᑦ
ᐊᖏᓐᓇᓕᒃᑯᑦ
ᒥᑭᑎᐊᖅᑐᓂᒃᑲ
ᑲᑐᔾᔨᖃᑎᒌᖏᓐᓂᑦ
(HTO)–ᖅᑎᑦ
ᓄᓇᕗᒻᒥᐅᑦ
NUNAVUT TUNNGAVIK
Nunavut

ᑭᓯᓂᒃ ᐊᓯᔨᕆᕐᑐᖅᑕᕐᓂᓕᕐᓂᒃ?

- ᐸᖅᓭᐅᑎ ᓄᑦᖏᓴᖅᑎᑕᐅᓂᐊᖅᑐᖅ ᓄᑦᒥᒃ ᑐᐱᐅᒪᓵᕆᑎᓂᒃ ᐱᑕᖅᑐᖅᓂᑦ
  - ᑐᒃᑐᐃᑦ ᖃᒃᓄᐃᑕᖏᓂᖓᑦ
  - ᐊᖏᔪᓚᵐᓂᐅᑦ ᐊᐅᑕᕆᑎᓂᖅᓯᑦ ᖃᒃᓄᐃᑕᖏᓂᖓᑦ
  - ᖃᒃᐅᔨᕐᓂᐅᑦ ᖃᒃᓄᐃᑕᖏᓂᖓᑦ
- ᑕᒪᐃᵒᓂ ᖃᑎᖅᑭᐅᖃᒃᑯᑦᓂᐊᖅᑐᑦ ᐊᖅᑦᔪ ᖃᓄᑕ ᓇᑦᑕᖅᖏᓐᓂᒃ

  - ᑭᖏᑦᓕᖅᑎ ᐊᓯᔨᕆᑕᐅᖅᓯᐊᓂᖓᑦ (2028)–ᒥᑦ



ᓄᓇᕗᑦ ᐅᒪᔨᓕᒪᔪᖅᑐᑦ ᑲᑎᒪᔨᖏᑦ
Nunavunmi Anngutighatigut Aulapkaijitkut Katimajiat
Nunavut Wildlife Management Board

ᐱᓕᕆᑦᑕᐃᓕᓚᖅ ᐅᒪᔨᖏᓪ ᐊᖏᔪᑎᑎᓕᔪ ᐊᑐᖃᑎᒌᓚᖅ ᐃᓄᐃᑦ ᖃᐅᔨᒪᓂᖃᖅᖢᑎᑦ ᐊᓪᓚᓕ ᖃᐅᔨᓴᖃᖅᖢᑎᑦ
Tammaqtailinahuarniriit anngutighat atuqhugit Inuit qaujimajatuqangillu ilihimaniillu ilitquhiannin
Conserving wildlife through the application of Inuit Qaujimajatuqangit and scientific knowledge

October 08, 2019

Honourable Joe Savikataaq
Minister of Environment
Government of Nunavut

Dear Minister Savikataaq:

Re: Proposal to Approve the Baffin Island Caribou Management Plan

At the Nunavut Wildlife Management Board's (NWMB or Board) Regular Meeting in Iqaluit on September 11, 2019 (RM003-2019), your department asked the NWMB to approve the *Baffin Island Caribou Management Plan* (management plan). The goal of the management plan is to serve as "an adaptive guide to harvest and monitoring of the Baffin Island caribou population."

At the NWMB's In-Camera Meeting in Iqaluit on September 12, 2019 (IC003-2019), the Board considered your proposal, along with records of government consultations with affected communities in the Qikiqtaaluk Region. The NWMB raised the following concerns:

- The lack of specific management actions to address current threats to Baffin Island caribou.
- The management plan has not been updated to include recent changes to Baffin Island caribou harvest non-quota limitations
- The QWB did not provide and comment on the content of the proposed management plan. Even though your department accommodated the Qikiqtaaluk Wildlife Board's (QWB) requests for deadline extensions in three instances.

The NWMB has written to the Qikiqtaaluk Wildlife Board (QWB) to request that they provide their position on the content of the management plan to your department for consideration by December 13, 2019.

The NWMB, therefore, recommends that your department consider the points raised above and to consider the written comments from the Qikiqtaaluk Board if any.

The Board will be pleased to consider the Proposal once the points above have been addressed.

ᓄᓇᕗᑦ ᐅᒪᔨᓕᒪᓂᖅᓴᕐᔪᐊᑦ ᑲᑎᒪᔨᖏᑦ
Nunavunmi Anngutighatigut Aulapkaijitkut Katimajiat
Nunavut Wildlife Management Board

ᐱᓗᐊᑦᑕᐃᓕᓚᓂᖅ ᐅᒪᔨᖏᓪ ᐊᖏᔪᑎᑎᓗᒍ ᐊᑐᕐᑎᑕᐅᓂᖅᑐᑦ ᐃᓄᐃᑦ ᖃᐅᔨᒪᓂᕐᒥᖏᑦ ᐊᒻᒪᓗ ᖃᐅᔨᓴᕐᑎᒻᒪᕆᒃ ᖃᐅᔨᒪᓂᖅᑐᑦ
Tammaqtailinahuarniriit anngutighat atuqhugit Inuit qaujimajatuqangillu ilihimaniillu ilitquhiannin
Conserving wildlife through the application of Inuit Qaujimajatuqangit and scientific knowledge

Yours sincerely,



Daniel Shewchuk
Chairperson
Nunavut Wildlife Management Board

cc - Drikus Gissing, Director of Wildlife Management, Nunavut Department of Environment

ᑲᑎᒪᔨ ᓄᓇᕗᑦ ᐱᖃᓐᓇᐅᑉᐸᓂᖅ
Building Nunavut Together
Nunavuliuqatigiingniq
Bâtir le Nunavut ensemble

ᒥᓂᔅᑕ ᐊᓂᕆᑎᓕᕐᓂᖅᓴᖅᑯᓐᓂᖅ
Minister of Environment
Ministaat Avatiliqiyitkut
Ministre de l'Environnement

Mr. Daniel Shewchuk
Acting Chairperson
Nunavut Wildlife Management Board
P.O Box 1379
Iqaluit, NU X0A 0H0

Translation
To follow

January 6, 2020

Re: Proposal to Approve the Baffin Island Caribou Management Plan

Dear Mr. Shewchuk,

Thank you very much for your letter dated October 8, 2019 concerning the Department of Environment's (DOE) request for approval of the Baffin Island Caribou Management Plan (BICMP). Regarding the concerns raised by yourself and the Nunavut Wildlife Management Board (NWMB), I would like to provide some context and responses to your specific points.

The Board indicated that there was a *"lack of specific management actions to address current threats to Baffin Island caribou"*. Specific management recommendations such as Total Allowable Harvest (TAH) levels, which are based on scientific information, traditional knowledge, and consultation with co-management partners, were indeed not included in the management plan. This is because the wildlife management system in Nunavut requires that the NWMB and co-management partners have an opportunity to comment on management recommendations. It is our position that if specific management actions were provided within the management plan, this would be overly prescriptive and create a redundancy, or even a conflict, with the NWMB's existing approval process.

Your letter indicated, *"the management plan has not been updated to include recent changes to Baffin Island harvest non-quota limitations"*. Due to the short timeframe between the decision on the Baffin Island TAH and Non-Quota Limitation, it was overlooked as an addition to the BICMP before the submission deadline for the NWMB regular meeting in September 2019. Changes to the non-quota limitations will be added to the Baffin Island Caribou Management Plan before it is re-submitted to the NWMB.

ᑎᑎᖅᑰᑦ 2410
ᐃᖅᓗᐃᑦ, ᓄᓇᕗᑦ X0A 0H0

P.O. Box 2410
Iqaluit, Nunavut X0A 0H0

Titiqakuvia 2410
Iqaluit, Nunavut X0A 0H0

C. P. 2410
Iqaluit, Nunavut X0A 0H0

867-975-5050
867-975-5051

The final point made by the NWMB was that "*the QWB did not provide and comment on the content of the proposed management plan. Even though your department accommodated the Qikiqtaaluk Wildlife Boards (QWB) requests for extensions in three instances".* This plan has been in circulation since 2005 and has undergone multiple rounds of consultations with the more recent consultations taking place with co-management partners, including the QWB, at the beginning of 2019. We recognize the need for co-management partners to have ample opportunity to provide input. Unfortunately, QWB has not responded by December 13, 2019 and the comment period was extended to accommodate QWB's request for this extension; which has created further delays in the approval process.

The DOE appreciates the Board providing a date by which QWB should respond. Unfortunately, as noted the December 13, 2019 deadline has passed without QWB providing any input on the BICMP.

I would like to thank the Board for their consideration of this plan and for their willingness to work together with us on the conservation and management of Baffin Island caribou during this critical, recovery phase for their population.

Sincerely,

Joe Savikataaq,
Minister of Environment

Cc. Jimmy Noble Jr., Deputy Minister of Environment
Steve Pinksen, Assistant Deputy Minister of Environment
Drikus Gissing, Director of Wildlife

ᑎᑎᖅᑯᓂᐊᓕ 2410
ᐃᖅᓗᐃᑦ, ᓄᓇᕗᑦ X0A 0H0

P.O. Box 2410
Iqaluit, Nunavut X0A 0H0

Titiqakuvia 2410
Iqaluit, Nunavut X0A 0H0

C. P. 2410
Iqaluit, Nunavut X0A 0H0

867-975-5050
867-975-5051

ᓄᓇᕗᒻᒥ ᐅᒪᔨᓕᕆᔨᖅᑎᑦᑕᐅᓂᖅ ᑲᑎᒪᔨᖏᑦ
Nunavunmi Anngutighatigut Aulapkaijitkut Katimajiat
Nunavut Wildlife Management Board

ᐱᓕᕆᑦᑕᐃᓕᓚᓂᖅ ᐅᒪᔨᖏᓂ ᐊᖏᔪᑎᑦᓯᓕᕆᑦ ᐃᓚᐃᑦ ᖃᐅᔨᒪᓂᕆᖏᑦ ᐃᓚᖓ ᖃᐅᔨᓴᓕᖅᑐᑦ ᐊᒻᒪᓗ ᖃᐅᔨᓴᓂᐊᖅᓯᓗᒃᑕᒃ ᖃᐅᔨᓴᖏᑦ
Tammaqtailinahuarniriit anngutighat atuqhugit Inuit qaujimajatuqangillu ilihimaniillu ilitquhiannin
Conserving wildlife through the application of Inuit Qaujimajatuqangit and scientific knowledge

ᔪᑎᐱᓗ 18, 2019

ᓴᐃᒥᔪ ᖃᑦᑕᖅ
ᐃᓕᔪᐊᑦᑕᖅ
ᖃᑉᑭᖅᑕᓗ ᐅᒪᔨᕆᓚᓂ ᑲᑎᒪᔨᖏᑦ

ᒥᔭᑦ ᖃᑦᑕᕐᒧᑦ:

ᐱᓱᕈᓂᖅᖁᓂᖅ: ᑐᓴᔪᕐᕈᑎ ᓂᑦᑕᐅᑎᑦᖃᐃᐅᕐᔪᑎᑦᑕᓗᒍ ᖃᑉᑭᖅᑕᓗᒃ ᐅᒪᔨᕆᓚᓂ ᑲᑎᒪᔨᖏᑦᓱᓂ ᐅᓗ ᖃᑉᑭᖅᑕᓂᒃ<ᖁᓗᕈᑎ

ᓄᓇᕗᑦ ᐅᒪᔨᕆᓚᓂᔪᕐᓇᐅᑦ ᑲᑎᒪᔨᖏᑦ ᐃᓱᒪᕆᐅᑎᑐᖏᑦ

ᑲᑎᒪᓂᖅᑲᖅᑎᑦᑕᒥᑦ ᓄᓇᕗᑦ ᐅᒪᔨᕆᓚᓂᔪᕐᓇᐅᑦ ᑲᑎᒪᔨᖏᑦ ᑲᑎᒪᐃᓴᖏᓐᓄᒍᑦ ᑲᑎᒪᑎᑦᕆᑦ ᐃᖅᑲᓐᓂᒃ ᔪᑎᑐᖅ 11, 2019-ᖓᑎᑦᕆᑐ (RM003-2019), ᓄᓇᕗᑦ ᓴᖅᒪᓂᖏᑦ ᐊᐱᓚᒋᐅᔪᒪᒍ ᓄᓇᕗᑦ ᐅᒪᔨᕆᓚᓂᔪᕐᓇᐅᑦ ᑲᑎᒪᔨᖏᑐᓄᑦ ᐊᖏᕐᕈᑕᐅᔪᓂᒃᑐᑦᑕᐃᒪᑐᒃ ᖃᑉᑭᖅᑕᓗᒃᖁᓕ ᑐᖅᑐᕈᓚᕆᔪᖓᓄᑦ ᖃᐅᔨᓴᖏᑦ (ᒥᐊᓂᖅᓯᕆᔨᑎᓄᑦ <ᖁᓗᐅᑎᑦ). ᑐᖃᓐᓇᐅᓚᖏᓐᓂᑦ <ᖁᓗᐅᑎᐅᐳ ᐱᔨᑦᔪᕐᑕᐅᓂᐊᖅᓯᒪᔪᑦ “ᐊᐅᑕᖅᑎᑕᐅᕈᑎᒃᑯᑦᑕᖅᑐᑐᓄᑦ ᑐᑭᒧᐊᒍᑕᐅᓗᓂ ᑐᓴᔪᕐᕈᑎᑦᑎᓂᒃᑯᑦ ᐊᒻᒪᓗ ᓇᐅᑦᔪᓂᖅᖁᓂᖅᑕᐅᓂᒃᑯᑦ ᖃᑉᑭᖅᑕᓗᒃ, ᑐᓴᔪᕐᕈᑎ ᐃᓱᒪᕆᐅᑎᑐᖏᑦ ᑐᓴᔪᕐᕈᑎᑦᑎᓂᒃᑯᑦ ᖃᑉᑭᖅᑕᓗᒃᖁᓕ ᐅᓇᓱᐊᓂᖅᑐᑦ ᑐᓴᔪᕐᕈᑎᑦᑎᓂᒃᑯᑦ ᖃᑉᑭᖅᑕᓗᒃᖁᓕ ᐅᒪᔨᕆᓚᓂᔪᕐᓇᐅᑦ ᑲᑎᒪᔨᖏᑐᓄᑦ ᐅᒪᔨᕆᓚᓂᔪᕐᓇᐅᑦ ᑲᑎᒪᔨᖏᑦ ᐃᓱᒪᕆᐅᑎᑐᖏᑦᓱᓂ ᐅᓇᓱᐊᓂᖅᑐᑦ ᑲᑎᒪᔨᖏᑐᓄᑦ ᑐᓴᔪᕐᕈᑎᑦᑎᓂᒃᑯᑦ ᐃᓱᒪᕆᐅᑎᑐᖏᑦᓱᓂ ᐅᓇᓱᐊᓂᖅᑐᑦ ᑲᑎᒪᔨᖏᑐᓄᑦ ᐅᒪᔨᕆᓚᓂᔪᕐᓇᐅᑦ ᑲᑎᒪᔨᖏ

ᓄᓇᕗᑦ ᐅᒪᔨᓕᕆᔨᖅᑎᑦ ᑲᑎᒪᔨᖏᑦ
Nunavunmi Anngutighatigut Aulapkaijitkut Katimajiat
Nunavut Wildlife Management Board

ᐱᓗᐊᑦᑕᐃᓕᓚᖅ ᐅᒪᔨᖅᓂ ᐊᖅᑯᑎᒋᓗᐊ ᐊᑐᕆᑕᐅᓂᖏᑦ ᐃᓄᐃᑦ ᖃᐅᔨᒪᓂᑐᖅᖏᑦ ᐊᓪᓚᓗ ᖃᐅᔨᓴᕆᔨᒪᔪᓕᒃ ᖃᐅᔨᒪᓂᖏᑦ
Tammaqtailinahuarniriit anngutighat atuqhugit Inuit qaujimajatuqangillu ilihimaniillu ilitquhiannin
Conserving wildlife through the application of Inuit Qaujimajatuqangit and scientific knowledge

ᓄᓇᕗᑦ ᐅᒪᔨᓕᕆᔨᖅᑎᑦ ᑲᑎᒪᔨᖏᑦ ᑐᓴᕆᓕᖅᑐᑦ ᖃᐅᔨᓴᕆᔨᒃ ᐅᒪᔨᓕᕆᔨᑦ ᑲᑎᒪᔨᖏᓐᓄᑦ ᑐᓂᓯᔪᓐᓇᖅᑐᑦ
ᐃᓱᒪᒍᖏᑦ ᖃᐅᔨᐃᓐᓇᕐᓂᖏᓐᓄᑦ ᐱᓕᔭᐃ ᖃᐅᔨᓴᕆᔭᖓᒥ ᑐᐃᑐᑦ ᒥᐊᓂᕆᐅᔨᕐᓂᐊᕐᓇᖅᑐᖏᑦ <ᖃᓄᐅᑎᖏᑦ ᓄᓇᕗᑦ
ᑲᙵᒪᔪᖏᓐᓄᑦ ᑐᓂᔭᐅᔭᑎᑉ ᑎᑭᓕᐅᖏᓐᓄᓪᓚᑐ **ᑕᓕᑦᑎᐊᖅ, ᑎᓯᐱᓗ 13, 2019**, ᐅᓐᓄᓴᔪᑦ **5 ᖃᓄᖅᓯᒥᐅᑦ ᓯᕿᓐᓄᔪᖅᓱᖅ ᐊᑐᕐᔭᐃᔭ**.

ᐃᖏᕐᕋᑦ ᐃᖅᑲᓇᐃᔪᖅᑎᑎᓕᔅᓴᓕᑦ ᐊᐱᖅᑯᓂᖅᓴᐅᔪᓐ ᐃᓯᒪᔭᐃᓚᐅᖅᓴᐅᔪᓐᓇᔭ ᐅᑯᐊ ᑎᑎᖅᑲᑦ ᐃᓗᓕᖅᐸᓂ,
ᑐᙵᖅᑎᑕᐅᕙᓪᓕᐅᑎᓯᓇᖅᑐᑦ ᓄᓇᕗᑦ ᐅᒪᔨᓕᕆᔨᖅᑎᑦ ᑲᑎᒪᔨᖏᑦ.

ᑎᑎᕋᖅᑐᖅ,



Ċᓂᐅᑦ ᓯᐅᓴᒃ
ᐃᒃᓯᕈᐅᑕᖅ
ᓄᓇᕗᑦ ᐅᒪᔨᓕᕆᔨᖅᑎᑦ ᑲᑎᒪᔨᖏᑦ

ᑎᑎᕋᐅᓯᐊᕐᒥᔪᑦ:
- ᑐᕋᐃᑲᓴ ᑭᓯᖏ, ᑐᑭᒧᐊᖅᑎᑦᓯᔨ ᐅᒪᔨᓕᕆᓂᕐᒧᑦ ᐱᑕᕆᐃᐊᐅᓂ, ᓄᓇᕗᑦ ᒪᕈᒪᔨᖏᑦ
  ᐊᕈᑎᑕᕆᔨᒃᑯᖏᓐᓂᓂ;
- <ᐅᑦ ᐃᖏᓴᐅᑦ, ᑐᑭᒧᐊᖅᑎᑦᓯᔨ ᐅᒪᔨᓕᕆᓂᕐᒧᑦ, ᓄᓇᕗᑦ ᑐᓂᓴᐃᒃ ᑎᒥᓴᑦ; ᐊᓪᓚᓗ
- ᐱᓕᓯᐅᒋᖅ ᑯᓗᒋᖅ, ᐊᐅᑕᓕᓯᔨᓄᑦ ᑐᑭᒧᐊᖅᑎᑦᓯᔨ, ᖃᐅᔨᒋᓗᒃ ᐅᒪᔨᓕᕆᔨᑦ ᑲᑎᒪᔨᖏᑦ.

ⵊⵓⵏⵜⵓⵙⵜ

ⵓⵓⵔⵔ ⵜⵍⵔⵜⵙⵏⵔⵜⵓⵔⵜ ⵓⵎⵍⵏⵓⵔⵓⵓⵔ

ⵄⵍⵙⵄⵊⵔⵜⵓⵔⵜ

ⵊⵍⵙⵜⵓⵔⵜ: X **ⵄⵔⵍⵔⵜⵣⵎⵔ:**

ⵔⵜⵓⵣⵜⵔⵜⵏⵍⵔⵓ: ⵊⵣⵔⵔⵍⵣⵣⵎⵔ ⵔⵜⵣⵎⵔⵙⵏⵔⵜⵓⵔ ⵔⵜⵣⵎⵔⵜⵓⵔ ⵔⵜⵣⵎⵔⵓⵏⵓ ⵓⵜⵜⵔⵜⵓⵔⵜ ⵔⵍⵔⵙⵏⵓⵔⵜ ⵍⵔⵙⵏⵔⵍⵜ

ⵜⵓⵜⵜⵔⵣⵓⵓⵔⵜ:

- ⵓⵓⵔⵔ ⵏⵜⵍⵓⵔ ⵔⵜⵣⵎⵔⵙⵏⵓⵔⵓⵔ (DOE) ⵏⵜⵣⵣⵎⵔⵎⵔⵍⵔⵜⵜⵔ ⵔⵜⵣⵎⵔⵜⵔⵍⵔ ⵔⵜⵣⵎⵔⵓⵏⵓ ⵓⵜⵜⵔⵜⵓⵔⵜ ⵍⵔⵙⵏⵔⵍⵜ (LTEMP) 2012-ⵍⵔ ⵓⵜⵜⵔⵜⵏⵜⵔⵍⵎ ⵔⵜⵣⵎⵔⵓⵏⵓ-ⵜⵔⵜⵜⵔⵙⵜⵓⵓ ⵔⵜⵙⵜⵜⵔⵔ ⵔⵜⵔⵔⵍⵔⵜⵙⵜⵓⵔ ⵙⵍⵜⵓⵓⵔ ⵔⵜⵍ ⵏⵜⵜⵔⵍⵎⵔⵎⵓⵜⵔⵍⵔ, ⵔⵜⵍ ⵓⵓⵔ ⵛⵜⵙⵔⵔ ⵔⵜⵙⵜⵜⵔⵔ ⵔⵜⵔⵍⵜⵜⵔⵔ ⵙⵍⵜⵓⵓⵔ ⵊⵍⵜⵣⵓⵔⵜⵓⵔⵔ ⵊⵍⵜⵣⵓⵔⵜ. ⵔⵜⵣⵎⵔⵜⵔⵍⵔ ⵔⵜⵔⵍⵜⵔⵔ ⵏⵜⵍⵙⵏⵔⵍⵔⵜ, ⵔⵜⵍ ⵓⵓⵔ ⵛⵜⵙⵔⵔ ⵔⵜⵙⵜⵜⵔⵔ ⵔⵜⵔⵍⵜⵜⵔⵔ ⵔⵜⵣⵎⵔⵓⵏⵓ ⵔⵜⵣⵎⵔⵜⵔⵍⵔ, ⵔⵜⵣⵎⵔⵓⵏⵓ ⵔⵜⵣⵎⵔⵜⵔⵍⵔ ⵔⵜⵣⵎⵔⵓⵏⵓ ⵔⵜⵣⵎⵔⵜⵔⵍⵔ, ⵏⵜⵣⵎⵔⵜⵔⵍⵔ ⵛⵜⵙⵔⵔ ⵔⵜⵣⵎⵔⵜⵔⵍⵔ, ⵏⵜⵣⵎⵔⵜⵔⵍⵔ ⵔⵜⵣⵎⵔⵜⵔⵍⵔ ⵔⵜⵣⵎⵔⵜⵔⵍⵔ ⵔ

ᐱᑭᖅᑐᓂᑦ ᐊᒻᒪ ᐊᓇᖏᓂᑦ ᓇᓴᖅᑕᐅᓂᖓᑦ. ᒧᐟᓇᐅᔨᒥᑦ ᐃᓕᔪᑕᖓᑦ ᖃᐅᔨᓴᓂᖏᑦ ᖃᓄᐹᖏᓐᓄ ᑲᖅᓯᓪᓂᖅ ᐱᖃᓯᐅᔨᔨᑦ ᓴᐱᐱᖅᑐᐃᑦ, ᖃᐄᐱᖅᑐᐃᑦ, ᖃᐄᕐᓇᐊᑦ, ᒥᑭᔨᓂᑦ ᐅᒪᔨᐊᐃᑦ, ᐊᒻᒪ ᐃᓕᖓᓂᓂ ᑎᒥᒋᐊᓂᑦ ᓂᖅᑐᑕᖅᑕᑐᑦ.

- ᑐᑭᓯᐊᕈᑎᓕᒫᑦ ᓄᐊᑕᐅᔨᑦ ᐊᑐᖅᑐᑎᑦ ᐊᖓᖅᑕᐅᓯᒪᔨᒥᑦ ᓇᓗᓴᐃᖅᓯᒪᔨᒧᑦ ᐊᑐᖅᑕᐅᑭᐊᓂᖅ, ᐱᖅᓯᐅᑎᒃᓗᑎ ᓴᖅᑭᖅᑎᑕᐅᔨᑦ ᐅᑭᐅᑕᖅᑐᒥ ᓄᓇᖓᓂ ᐊᐊᖏᓂᕆᔨᑦ ᐱᖅᑕᐅᔨᑦ ᑭᓯᔨᐊᒥ ᐅᑭᐅᑕᖅᑐᒥ ᓄᓇᖓᓂ ᖃᐅᔨᓴᖅᓂᖅ.

- ᓄᓇᕗᒻᒥ ᖃᐅᔨᓴᖅᑕᒋᓂᖏᓐᓂᑦ ᐱᓕᕆᐊᖅ ᐊᖏᓯᐅᒃᓯᓚᐅᔨᖅ, ᐊᑯᓂᐅᔨᒧᑦ ᑭᓇᐅᔨᖅᖃᖅᑎᑕᑎᓂᖏᓐᓂᑦ ᐱᓕᕆᖃᑎᐅᔨᖅ (2019/20-2021/22) ᐅᑯᓇᓂ ᐊᑯᓂᐅᔨᒧᑦ ᐊᐊᖏᖓᓂ ᖃᐅᔨᓴᓂᖏᓐᓂᑦ ᐱᓕᕆᐊᖅ.

- ᐱᒋᐊᖅᑐᓂ 2019, ᐃᑲᔪᖅᑕᐅᓵᓗᓂ ᓄᓇᕗᒻᒥ ᓯᓚᑦᑐᓴᖏᓐᓇᖅ (NAC) ᐊᒻᒪ ᑭᓴᓕᓂᖅ ᐃᓄᐃᑦ ᑲᑐᔾᖃᖅᑎᑕᖅᑐᑦ, ᐃᓕᓴᓂᐊᖅᑎᑦ ᐊᑎᓕᐅᖅᓯᒪᔨᑦ ᓄᓇᕗᒻᒥ ᓯᓚᑦᑐᓴᖏᓐᓇᖅᓯᒪᓂᖓᓂᑦ ᐱᓕᕆᐊᒃᑲᓐᓂᑦ ᐃᖅᑲᓇᐃᔨᖅᑎᑕᐅᓚᐅᖅᑐᑦ ᐅᓂᒃᑳᓇ ᐊᑯᓂᐅᔨᒧᑦ ᐊᖓᖅᑕᐅᓂᓂᑦ ᐱᓕᕆᐊᒥᑦ, ᓴᓇᓯᒪᓚᐅᖅᑐᒍᖅ ᐱᑕᖅᓂᕐᑎᓗᓂ ᖃᐅᔨᓴᖅᑎᓂᒃᑯᓐᓂᑦ, ᐃᖅᑲᓇᐃᔨᖅᑎᑕᑎᓂᑦ ᐃᓕᓴᐃᓯᔪᓐᓇᑦ.

- ᒪᓕᒐᒃ ᓄᓇᕗᒻᒥ ᐃᓄᐃᑦ ᐃᖅᑲᓇᐃᔨᖅᑎᑕᐅᖏᓪᓗᑎᒃᑐᑦ ᐊᑐᓂ ᐊᖏᓪᓕᐊᓂᑦ, ᐱᖅᑕᐅᑎᑕᐅᔨᑦ ᐊᑕᓯᐅᖅᑐᓂᑦ ᖃᐅᔨᓴᖅᓂᖅᑐᒥ ᐊᓂᒪᔨᐅᔪᓂᖅ, ᑭᓯᐊᓂ ᐊᐅᓚᑦᑎᓐᓂᑦ ᓄᓇᕗᒻᒥ ᖃᐅᔨᓴᖅᑐᑦ (ᓯᑦᓗ ᐊᑦᑕᓇᖏᑦᑐᒥ ᐊᐅᓪᓚᓯᓂᖅ ᓄᓇᖃᖅ ᐊᒻᒪ ᐃᒪᓐᓇᑦ) ᐊᒻᒪ ᐃᓄᐃᑦ ᖃᐅᔨᓴᒪᔪᑐᖅᑐᑦ (ᓯᑦᓗ, ᐃᖅᑲᓇᐃᔨᖅᑎᑕᐅᓵᓂᖅ, ᖃᓄᑐᐃᓐᓇᖅ ᐃᓯᒪᓕᕐᓂᔪᓐᓇᖅ, ᖃᐅᔨᒪᓂᐅᔨᓂᑦ ᐱᓇᓱᓕᐅᑎᓯᓂᖅ, ᐃᓪᓕᐱᒍᓯᓯᑎᓂᖅ ᓄᓇᕗᑦ ᐊᒻᒪ ᐃᒪᓐᓇᑦ).

- ᐊᑯᓂᐅᔨᒧᑦ ᐊᖏᓪᓕᐊᓂᖅ ᖃᐅᔨᓴᓂᖏᓐᓂᑦ ᐱᓕᕆᐊᖅ ᐱᔪᒥᓇᖅᑐᖅ ᐱᓕᕆᐊᖅ ᐃᓕᓴᐅᔨᔪᐅᓂᖓᓇᓄ ᓇᓕᒪᑦ ᐊᐅᓕᑎᑐᑦ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ, ᐊᒻᒪ ᐊᖏᓪᓕᐊᓂᖅᓯᒪᓂᖓᓂᑦ ᐃᓕᓴᖅᑎᑎᓂᖅᑐᑦ ᐱᓕᕆᖃᑎᖅᑐᓂ ᐊᐅᓕᑎᑐᑦ ᐱᓕᕆᐊᖅ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ ᓄᓇᕗᒻᒥ ᐃᓄᐃᑦ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ ᓄᓇᕗᑦ ᐊᒻᒪ ᐅᒃᑯᓂᖅᑐᑦ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ ᓄᓇᕗᒻᒥ ᐃᓄᐃᑦ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ ᓄᓇᕗᑦ ᐊᒻᒪ ᐅᒃᑯᓂᖅᑐᑦ).

- ᐊᑯᓂᐅᔨᒧᑦ ᐊᖏᓪᓕᐊᓂᖅ ᖃᐅᔨᓴᓂᖏᓐᓂᑦ ᐱᓕᕆᐊᖅ ᐱᔪᒥᓇᖅ ᐱᓕᕆᐊᖅ ᐃᓕᓐᓂᐅᔨᕐᓂᖓᓇᓄ ᓇᓕᓪᓕᑦ ᐊᐅᓕᑎᑐᑦ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ, ᐊᒻᒪ ᐊᖏᓪᓕᐊᓂᖅᓴᓂᑦ ᒪᑐᐃᓐᓇᔨᑦ ᐱᓕᕆᖃᑎᖅᑐᓂ ᐊᐅᓕᑎᑐᑦ ᐱᓕᕆᐊᖅ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ ᐱᓕᕆᖃᑎᖅᑐᓂ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ ᐱᓕᕆᐊᖅ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ ᓄᓇᕗᑦ ᐊᒻᒪ ᐅᒃᑯᓂᖅᑐᑦ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ ᓄᓇᕗᒻᒥ ᐃᓄᐃᑦ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ ᓄᓇᕗᑦ ᐊᒻᒪ ᐅᒃᑯᓂᖅᑐᑦ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ ᓄᓇᕗᒻᒥ ᐃᓄᐃᑦ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ ᓄᓇᕗᑦ ᐊᒻᒪ ᐅᒃᑯᓂᖅᑐᑦ).

- ᐊᑯᓂᐅᔨᒧᑦ ᐊᖏᓪᓕᐊᓂᖅ ᖃᐅᔨᓴᓂᖏᓐᓂᑦ ᐱᓕᕆᐊᖅ ᐱᔪᒥᓇᖅ ᐱᓕᕆᐊᖅ ᐃᓕᓴᐅᔨᔪᐅᓂᖓᓇᓄ ᓇᓕᓪᓕᑦ ᐊᐅᓕᑎᑐᑦ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ ᐱᓕᕆᐊᖅ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ ᐱᓕᕆᖃᑎᖅᑐᓂ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ ᐱᓕᕆᐊᖅ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ ᓄᓇᕗᑦ ᐊᒻᒪ ᐅᒃᑯᓂᖅᑐᑦ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ ᓄᓇᕗᒻᒥ ᐃᓄᐃᑦ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ ᓄᓇᕗᑦ ᐊᒻᒪ ᐅᒃᑯᓂᖅᑐᑦ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ ᓄᓇᕗᒻᒥ ᐃᓄᐃᑦ ᐃᓕᓴᖅᑕᐅᑎᖃᖅᑐᑦ ᓄᓇᕗᑦ ᐊᒻᒪ ᐅᒃᑯᓂᖅᑐᑦ).

ᐊᐱᖅᓱᖏᓪᓗᓂ:

- ᑲᑎᒪᖅᑎᖅᑎᓂᖅ ᐅᑭᐅᔨᓂᐊᑎᒃᓴᑦ ᑲᑐᔾᖃᖅᑎᑕᐅᑎᖃᖅᑐᓂᑦ ᑲᖏᖅᑕᐅᓂᖅᑎᑦ ᐊᒻᒪ ᖃᒪᓂᑕᐃᒋᑦ ᑲᑎᒪᑎᑎᓂᒋᑕᖅᑐᑦ ᐊᐊᖏᓂᕆᔨᖅᑎᑦ ᒪᐃᑎ 2019.
- ᑲᑎᒪᑎᑎᓂᐅᒃᓴᓂᕐᓂᐅᔨᑦ ᐸᖅᓇᑕᐅᓱᒪᔨᑦ ᐊᑎᑐᐃᓂᖃᖅᑎᓕᒫᑦ ᓄᓇᕗᓂᓂᑦ ᓇᖅᑲᐊᒪᑦ 2020.

ᐊᑐᓕᖁᓕᐅᔨᑦ:


- *ᐱᑕᖃᖅᑐᑐᖅ*

<u>ᐊᑯᓂᐅᔨᒧᑦ ᐊᓂᐲᖅᓯᓂᖅᓴᑦ ᐱᓕᕆᐊᖅ: ᐊᐅᓪᓚᑎᓂᖅᓴᑦ ᓇᐃᓐᓇᕐᓂᒃ</u>


ᓴᖅᑭᖅᑎᑕᐅᓕᕆᓂᐊᖅᑐᖅ ᐅᓇᓵᓂᕆᑦ ᐅᑭᐅᑕᖅᑐᒥ ᖃᐅᖅᓯᒪᓕᕆᖅᓕᓕᕐᓂ ᒪᕐᕉᐊᖅᑎᓪᓗᓯᑐᑦᑐᓂ ᓄᓕᓵᓂᖅᓯᓂ ᐊᓕᕆᓂᓵᓂᑦ ᓇᒥᑐᐃᓂᖅ. ᑭᓯᐊᓂᓕ, ᖃᐅᔨᓴᖅᑕᐅᓂᕆᓂᑦ ᐊᑦᑐᖅᑕᐅᔨᑦ ᓯᓚᑕᐅᔨᐸᑦ ᐊᓕᕆᔨᓂᖅᓴᓂᒃ ᐅᒪᔨᓂᓕ, ᐊᔨᑎᕆᑕᖅᓴᑎᑐᑦ ᐱᓂᓕᐊᓂᖅᓴ ᖃᓄᐃᓕᓂᐅᐅᔨᑦ ᖃᐃᓂᓚᐊᓂᖅᓴᓂᒃᓴᑦ ᐅᓇᓵᓂᖅᓴ ᒥᓴᑎᐅᓴᓴᐃᔨᓂᓕ (ᓄᑦᓗ, ᖃᐃᐊᖅᔪᐅᓇᖅᑐᑐᖅ, ᐊᐅᔨᐊᐃᑦᑐᑦ ᓄᖅᓱᑎᓂᖅᑎᑦ ᐊᒪᓕ ᑕᖅᐅᖅᑭᑦᓴᖅᓴᓂᒃ ᐅᓪᓚᕆᑕᖅᓴᓂᖅ ᐊᒪᓕ ᐊᖅᑭᔨᓂᖅ), ᐅᒪᔨᐊᐃᑦ ᓂᖅᐅᐅᓇᖅᑭᑦᑐᓂᑦ ᖃᓄᐃᓕᓂᐊᕐᓇᖅᑐᑦ, ᐊᒪᓕ ᖃᐅᔨᓴᖅᑕᐅᓂᖅᓴ ᐊᓕᕆᖅᓴ ᐊᓕᕆᓂᖅᓴᓂᖅ ᐃᓚᒋᑦᑐᖅᓴᓂᖅ ᐊᒪᓕ ᐊᖅᑭᔨᓂᖅᓴᓂᖅ, ᐃᓕᓴᐃᓐ ᓂᐲᑦᑐᖅᑐᑦᑐᓂᑦ ᐊᒪᓕ ᐊᓕᕆᓂᖅᓴᓂᖅ ᐊᒪᓕ ᐊᐅᓪᓚᑎᓂᖅᓴᓂᒃᓴᑦ ᐊᓕᕆᓂᖅᓴ ᐱᔭᐅᓂᖅᓴᓂᖅ ᐊᒪᓕ ᐊᓕᕆᓂᖅᓴᓂᖅ ᐊᐅᓪᓚᑎᓂᖅᓴᓂᒃᓴᑦ ᐊᓕᕆᓂᖅᓴ ᐱᔭᐅᓂᖅᓴᓂᖅ ᐊᒪᓕ

ᖃᐅᔨᓴᓂᐊᕐᖃᖅᑕᕐᓴᓂᖅ ᓄᓇᖓᓂ ᖃᐅᔨᓴᓴᓂᖅ ᐱᓕᕆᐊᖓᓪᓚᐅᖅᑐᖅ ᐊᐅᖏᖓᓂ ᖃᓄᐅᕐᖓᓂ ᓄᓇᓕᖓᑕ ᑲᖏᓴᓪᓗᓂ (ᐊᑦᔨᓐᖓᐊᖅ 1) 2019-ᒥᑦ. ᖃᐅᔨᓴᖃᐅᔨᖅ ᑭᖓᖅᑐᐃᔪᖅ ᖃᓄᐅᕐᓚᐊᓂ 2500 km^2 ᐃᒪᖅᓚᓱᒥ ᓄᓇ ᐊᒻᒪ ᐃᒪᖏᑎ ᓇᔨᖅᑕᐅᔭᕆᑐᑦ ᐱᑕᖃᖓᓂᓂᑦ ᐊᑎᑎᓄᖓᓂ ᐅᑭᐅᑕᖅᑐᒥ, ᐃᓗᓕᖃᖅᑐᖅ ᐊᑦᔨᑦᑭᓐᓂᑎᓄᑦ ᒪᓴᓄᑦ ᐃᖃᓗᓐᓄᑦ, <ᓄᖅᑐᑦ ᑯᐊᓂᓪᓚᕆᐊᖃ, ᐅᑭᐅᖅᑕᖅ ᑐᓴᓂᖅ, ᐅᓕᓐᓂᖅ ᑎᓄᓐᓄᖅ ᐊᒻᒪ ᐃᒪᖏᑎ ᐃᓐᓇᑕᐃᑦ. ᓄᓇᒋᔭ ᐊᑎᑎᓵᓕᕆᓵᐊᓪᓚᕆᓴ ᖃᖅᖃᖅᑎ, ᐊᒻᒪ ᐱᑕᖃᐅᔪᖅᑕᑐᑦ ᐊᒻᒪ ᑕᓪᓕᓂᑦ. ᐅᔭᖅᖃᐃᑦ ᑕᑯᓴᐅᔾᔨᑦ ᑕᑯᓴᐅᓴᔨᓕᑐᑦ ᑕᒧᓄ, ᐊᒻᒪ ᑯᖅᑭᔪᓪᓚᑎᑎᐊᖅᑐᑦ ᐊᐅᑕᑎᐅᓵᓪᓗᑎ ᕿᑭᕐᕆᑦ. ᐅᔭᖅᖃᐃᑦ ᑕᑯᓴᐅᔾᔨᑦ ᑕᑯᓴᐅᓴᔨᓕᑐᑦ ᑕᒧᓄ, ᐊᒻᒪ ᑯᖅᑭᔪᓪᓚᑎᑎᐊᖅᑐᑦ ᓴᕐᓚᖓᔨᓂᑦ ᐃᐅᖅᓴᑦ, ᐃᓐᓇᑦ, ᐊᒻᒪ ᐃᓐᓇᐃᓕᓴᐃᓐᓇᖅᑐᑦ ᐊᐅᑕᑎᐅᓵᓪᓗᑎ ᕿᑭᕐᕆᑦ. ᐱᓕᕆᐊᖅᑐᑦ ᖃᓄᐃᑦᑐᖅᑐᑦ ᐊᑐᐅᓂᖅᑐᑦ, ᐃᓐᓇᑦ ᐃᓐᓇᐃᓕᓴᐃᓐᓇᖅᑐᑦ ᓴᓇᖓᓂᔨᓂᑦ ᐃᑲᖓᓂᑦ, ᐊᒻᒪ ᐃᓐᓇᐃᓕᓴᐃᓐᓇᖅᑐᑦ. ᐱᓕᕆᐊᖅᑐᑦ ᐊᑐᐅᓂᖅᑐᑦ ᐊᑐᐅᓂᖅᑐᑦ ᐊᑐᐅᓂ

ጋᖁძርᐅᑦ⊃ᒪᑎ, ᐊᖏᒪ ᖃᐅᔨᕋᐃᓂᖅ ᐊᓇᖏᑐᓂ ᓄᓴᐃᓂᕐᒥᑦ ⊃ᑭᓯᓕᐊᑭᑎᓂᑦ (ᓯᑦᔪ, ᓇᔪᓇᐃᖅᓯᒪᔨᒧᑦ ᓇᒫᑐᓂᖓᓐ ᓄᓴᖅᑕᐅᔨᑦ ᐊᑐᓂ ᐅᒪᔨᓂᑦ ᐊᑐᓂ ᓇᔨᖅᑕᐅᔨᓄᑦ ᖃᓄᐃᑦᑐᓂᖏᑦᑕ) ᐱᕆᐸᖅᑕᐅᓯᒪᔨᑦ.



ᐊᑲᓇᓐᖓᐊᖅ 1 ᖃᐅᔨᓴᖅᓱᓕᐅᔪᑦ ᓇᒻᒥᓂᖅᖢᓂᕐᒥᐅᓇᖅᑐᖅ (2019 – ᑲᔪᓯᔾᔨᓕᕐᒥᐅᓇᖅᑐᓇᖅᑐᒃ) ᓯᓚᓗ ᖃᓄᑦᑎᓂᖅᓯᒪᓕᑦ ᖃᐅᔨᓴᐅᑎᑦ, ᓄᓕᓕᒃ ᖃᐅᔨᓴᖅᓱᓕᐅᔪᖅ ᐃᓂᖏᑦ (PFP), ᐱᑭᖅᑐᐃᓐᓇᑦ ᐊᔾᔨᒌᒍᑎᑦ, ᒥᑭᓕᐊᑦ ᐅᓪᓗᓂ, ᖁᔭᓕᐊᑦ (ᐃᓯᕆᓕᓕᑦ ᓂᕐᐱᓕᐅᑭᑦ ᑎᑎᕋᐊᑯᓯᓯᒪᔪᓂ ᖃᐅᔨᓴᖅᓂᖅ), ᖁᔨᓴᖅᑐᐃᓐᓇᑦ (ᐅᓂᖅᓱᒍᑦ), ᐊᓯᒪ ᓂᕈᑐᖅᑐᑦ ᑎᑎᕋᑦ ᐱᑭᐅᖃᖅᓱᒍᑦ ᖃᐅᔨᓴᖅᑕᐅᔪᑦ ᖃᓂᒋᔭᖏᓂ ᑲᖏᖅᓯᓂᖅ.



ᐊᑦᓴᓐᖓᐊᖅ 2 ᑎᓴᒪᑦ ᖃᐅᔨᓴᖏᐅᑉᐳᓐᓇ ᖅᑐᑦ ᓇᒻᓐᓂᑦ ᓴᖅᑭᖅᑎᑕᐅᓕᐅᖅᑐᑦ ᑭᑎᑦᓗᐊᑕᖏᓐᓂᓪ, ᐊᓱᒪ 12 ᖃᐅᔨᓴᖏᐅᑉᐳᓐᓇᓂᐊᖅᑐᑦ ᓴᖅᑭᖅᑎᑕᐅᓕᐅᖅᑐᑦ ᐅᖓᓯᒻᒥᐅ ᐅᐊᓐᓇᒪᐊᒍᑦ (N=6) ᐊᓱᒪ ᓂᒋᔭᒍᑦ (N=6) ᑭᑎᑦᓗᐊᑕᖏᓐᓂᓪ ᓄᑦᕋᓕᐅᔪᖏᑉᐳᓐ ᓄᓇ ᖏᑉᐳᓐ ᓴᖅᑭᑎᑎᒋᑦ ᑲᑎᑦᑐᒋᑦ 16 ᖃᐅᔨᓴᖏᐅᑉᐳᓐᓇ ᖅᑐᑦ ᓂᒋᖏᓂ ᐅᐊᓐᓇ ᖏᓂᓂ ᐃᓗᐊᓂ ᖃᓪᓗᓂᑦᓯᐊᖅ ᑐᑦᑐᖏᑦ ᓄᑦᕋᓕᐅᔪᖏᑦ ᓄᓇᐃᑦ. ᓂᒋᖅᓯᓯᒋᓐ ᖏᓂᓂ ᖃᓄᒋᔨᓯᓕᐊᓂᑦᑐᖅ 260 ᑭᓯᒋᑕᐃᑦ ᐅᐊᓐᓇᒋᓐ ᖏᓂᓂ.





LTEMP

Long-term Ecological Monitoring Program

Nunavut Wildlife Cooperative Research Unit

Government of Nunavut
University of Alberta

2019 - Nov
Annual Report


Erik Hedlin
Alastair Franke
Kevin Hawkshaw
Rebekah McKinnon
Nick Gulotta


Disclaimer: This draft report has been prepared for the Government of Nunavut and presents preliminary findings that may change with further analysis. The views and interpretations in this report article are those of the authors and do not necessarily represent the views of the Government of Nunavut. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the Government of Nunavut.


ᐅᖃᐅᓯᖃᖅᓯᒪᓂᖅᐸᑦ

- ᓄᓇᕗᑦ ᑲᓚᒪᖅᓯᑦ ᓴᖅᑭᑎᑦᑎᓪᓗᐅᕐᓂᖅᑐᑦ ᐊᔪᓂᐅᔨᒧᑦ ᐊᓚᑎᖏᓐᓄᑦ ᖃᐅᔨᓴᓂᖅᒧᑦ ᐱᓪᓕᐊᖅ (LTEMP) 2012–ᒥ ᖃᐅᔨᓴᖅᓯᐊᑎ ᐊᓚᑎᖏᓐᓄᑦ ᖃᓄᐅᓚᓂᖅᑐᖏᓐᓄ ᐊᓯᔾᔨᖅᑐᑦ ᐊᑦᑐᐊᓂᓪᑦ ᓯᓪᓚᖏᓄ ᐊᒻᒪ ᐱᓚᕐᓪᐊᑎᑦᑎᓂᖅᒧᑦ.
- ᖃᐅᔨᓴᖅᓱᐅᔨᒥᑦ ᖃᐅᔨᓴᖅᓂᖃᓪᓚᐅᕐᖅᑐᑦ 2012-2013 ᐊᒻᒪ 2015-2016. ᖃᓄᐊᓕᓯᓂᖏᓪᓗᓄᑦ ᓄᖅᑲᖃᑦᑕᖅᑐᑦ ᐃᖅᑲᓇᐃᔪᖅᑎᓂᑦ, ᐊᐊᖏᓕᓪᓚᔨᑦᓯᑦ ᐃᓯᒪᓕᐅᓪᓚᐅᕐᖅᑐᑦ ᐱᓕᓪᖃᓄᖅᑲᖅᓯᒪᓄ ᐃᓕᓴᓂᐊᖅᓂᓕᓂᖅᓯᒧᓐᓇᓂᑦ ᐱᓕᓪᖃᓄᖏᓐᓂᓴᓂ ᓴᓇᓪᓗᓄ ᐊᔪᓂᐅᔨᒧᑦ ᐱᔾᓴᓇᓂᖏᓐᓂᓴᓄᓪᓗ ᐱᓕᓪᖃᓄᖏᓐᓂᓴᓂ ᐱᓕᕆᔨᒃᑯᑦᑐᑦ ᐱᓕᓪᖃᓄᖅᑲᐅᔪᓴᓪᓗᑎᑦ ᐱᓕᕆᔨᒥᑦ.
- ᑐᑭᓯᐅᒪᖃᖅᑎᓂᖅᑐᑦ ᐊᖏᑕᑦ ᓯᓚᑦᑐᑦᓴᖅᓴᐅᐊᒧᑦ ᐃᐅᓗᑦ, ᐱᓕᓪᖃᓄᖏᓐᓂᓴᓂ ᐃᓕᓴᓂᐊᖅᓂᓕᓂᖅᓯᒧᓐᓇᓂᑦ ᐱᓕᓪᖃᓄᖅᑲᐅᔪᓴᓪᓗᑎᑦ ᐱᓕᕆᔨᒥᑦ ᐊᓐᓇᑦ ᐃᓗᓐᓂᑦ ᐊᔪᓂᐅᔨᒧᑦ ᐱᓕᕆᔨᒥᑦ ᐱᓕᓪᖃᓄᖅᑲᐅᔪᓴᓪᓗᑎ

- ᑐᖏᓐᓄ: ᖃᐅᔨᓴᖅᓯᒍ ᐊᓂᒪᓐᓄ ᐊᐅᓴᵇᑯᖏᕕᓂᑕᖅᑕᖅᑎᑦ ᖃᒪᓕᕆᔨᐊᕐᖃᖅ ᑐᑦᑐᖅᓵᓄᑦ.
- ᐃᓱᒪᓯᑎᓴᓯᐊᑕᐅᔨᔪᑦ: 1) ᑎᑎᕆᖃᑕᐅᓕᕆᖅᑎᑦ ᐊᑯᓕᐅᔨᔭᒧᑦ ᐊᓱᓴᖃᑐᑦ ᐱᔨᔪᑎᑦᓯᓯᒍ ᑐᓱᓴᕆᓕᕆᓄᑦ, ᐊᒪᒪ 2) ᓴᖃᑭᑎᑦᑎᓕᖅ ᑐᓂᖅᓄᓕᖅᑎᓕᕆᓂᖅᓕ ᖃᓄᐃᓂᕆᓕᕆᓂᖅᓕ (ᐅᓴᕆᓂᓕᐊᕆᓴᖃᖅᐅᖅᑐᑦ ᐊᕐᓇᑦᑐᖃᓕᔨᖅ).
  - ᖃᐅᔨᓴᑦᑎᐊᖃᖅᑕᑦᓕᖅᖅ ᐱᓕᓀᐊᖅᓯᖅ ᐊᑦᑕᔪᑕᒧᑦ ᖃᓕᑭᖅᖅᓯᐊᓕ ᓄᓇᕗᖅᓄᓕ ᑲᖅᑎᓂᖅᓕ
  - ᖃᐅᔨᓴᑦᑎᐊᖃᖅᑕᑦᓕᖅᖅ ᓄᑦᑎᓕᐅᑦᓇᖅᑎᓕ ᓄᓇᖅᑎᓕ ᖃᒪᓕᕆᔨᐊᕐᖃᖅ ᑐᑦᑐᖅᓵᓄᑦ
- ᓄᓇᕗᒻᒥ ᑕᒪᐃᓕᓕᑲ ᖃᐅᔨᓴᕆᑭᑎᓄᑦ <ᕐᓇᐅᑎᑦ (NGMP) ᐊᖏᕐᕋᑲᓪᓚᐅᔨᖅᑎᑦ, ᐊᖏᕐᕋᑲᓪᓚᐅᔨᖅᑎᑦ, ᐊᑯᓂᐅᔨᔪᒧᑦ ᑭᓇᐅᔭᖃᖅᑎᑦᑎᓂᕆᔪᒧᑦ ᐱᓕᐅᔭᖅᑎᑦ (2019/20-2021/22).
- ᓄᓇᕗᒻᒥ ᓯᓚᑦᑐᓴᕐᕿᑲᑦ ᐊᒻᒪ ᑭᓴᓪᓚᕐᒥ ᐃᓄᐃᑦ ᑐᐸᓴᖃᑎᒋᑦ ᐃᑲᔪᖅᑐᑦ ᐊᓴᖏᓪᓗᓂᔪᒧᑦ ᐱᓚᕆᒥ ᐃᓚᓲᓂᐊᓕᖅᑎᓂᑦ ᐃᓕᓴᕐᕋᐅᓂᖅᓯᒪᓕᖅᑎᓂᑦ ᐃᓕᓴᕐᕋᐃᓕᓂᔪᒧᑦ ᐱᓚᕆᐊᒥᑦ.
- ᒪᓕᒃᑲᖅ ᓄᓇᕗᒻᒥ ᐃᓄᖅᑲ ᐃᓕᓴᕐᕋᐃᓕᔨᖅᑎᑕᐅᔨᔪᑦ ᐊᑐᓂ ᐊᓕᖅᑐᒍᓂ
  - ᐱᖅᑕᐅᔨᔪᖅ ᖃᐅᔨᓴᕆᔨᓂᐅᔨᒥᑦ ᐱᓚᕆᐊᓚᒋᔭᖅᑐᑦ
  - ᐊᐅᓚᓯᖅᑕᐅᓂᖅᑎᑦ ᓄᓇᓚᒻᒥ ᖃᐅᔨᒪᓕᐅᔨᔪᑦ ᐊᒻᒪ ᐃᓄᐃᑦ ᖃᐅᔨᒪᓕᕐᑐᖅᑎᑦ
- ᐊᔨᓂᐅᔨᔪᑦ ᐊᓂᒪᓐᓄᓂᓂᔪᒧᑦ ᐱᓚᕆᐊᖅ ᐱᔨᒥᓇᖅᑐᖅ ᓄᓇᓚᒻᒪᑦ ᐊᐅᓚᑕᐅᔨᔪᑦ ᖃᐅᔨᓴᕐᓂᒥᑦ ᐊᑕᖏᕐᕋᓂ ᐃᓕᓴᕐᕋᐅᓂᖅᓯᒪᓕᖅᑎᓂᑦ.


Kitikmeot
Cambridge Bay
Taloyoak
Kugaaruk
Pelly Bay
Igloolik
Hall Beach
King William Island
Gjoa Haven
Queen Maud Gulf MBS
Naujaat
204
Naujaat
FOXE BASIN
Kivalliq
Amaruq
301
Area A (M)
Meadow River
330
Meadowbank Mine
Greyhound
Area B (M)
St. Tropez
Area C (M)
Baker Lake
Kiggavik
White Hills
Cone Hill
314
Gibson MacQuoid
Kahuna Gold
Chesterfield Inlet
Parker Lake
316
320
Fox Lake
315
Kuulu
317
329
327
202
Kahuna Diamonds
Peter Lake
334
Meliadine
Rankin Inlet
Angilak
Quartzite
353
335
Pistol Bay
Whale Cove
Yandle
344
352
Noomut
Hard Cash
Arviat
Kiyuk
328
HUDSON BAY
BAIE D' HUDSON
MANITOBA
Churchill
SOUTHAMPTON ISLAND
Coral Harbour
Coats Island
MELVILLE PENINSULA
Ukkusiksalik NP
Beverly CPMA
Qamanirjuaq CPMA
McConnell River MBS
Hiry Gibbons MBS
LEGEND
0 50 100 200 300 400
Kilometres


ipcc
INTERGOVERNMENTAL PANEL ON climate change
CLIMATE CHANGE 2014
Synthesis Report
CLIMATE CHANGE 2014 Synthesis Report
A REPORT OF THE
INTERGOVERNMENTAL PANEL ON CLIMATE CHANGE
WMO
UNEP


Figure 4.9.6
QamanirJuaq
Summer Density
(Jul 4 - Aug 22)
Legend
Community
Tree Line
River/Stream
All-season Road
Winter Road
Lake
Annual Home Range*
Core Seasonal Range
Utilization Distribution (%)
50
80
90
95
100
Increasing Density
Area of Detail
0 50 100 150
Kilometres
Projection:
Canada Lambert Conformal Conic
Data Sources:
Government of Nunavut, Government of Northwest Territories, Natural Resources Canada, GeoBase®, National Topographic Database
Nunavut
Northwest Territories
CASLYS CONSULTING
September17, 2014
Northwest Territories
Nunavut
Saskatchewan
Manitoba
Hudson Bay
Aberdeen Lake
Baker Lake
Chesterfield Inlet
Rankin Inlet
Whale Cove
Arviat
Churchill
Artillery Lake
Dubawnt Lake
Whitefish Lake
Nonacho Lake
Clinton-Colden Lake
Qamanirjuaq Lake
South Henik Lake
Kasba Lake
Nueltin Lake
Selwyn Lake
Fond-du-Lac
Black Lake
Wollaston Lake
Wollaston Lake
Brochet
Cree Lake
Reindeer Lake
Lynn Lake
South Indian Lake
Leaf Rapids
Lake Eliza
Gillam
DRAFT
* Annual home ranges are from the following publication: Nagy, J. A., D. L. Johnson, N. C. Larter, M. W. Campbell, A. E. Derocher, A. Kelly, M. Dumond, D. Allaire, and B. Croft. 2011. Subpopulation structure of caribou (Rangifertarandus L.) in arctic and subarctic Canada. Ecological Applications [doi:10.1890/10-1410.1].



nature climate change

Perspective | Published: 14 May 2018

Future climate risk from compound events

Jakob Zscheischler, Seth Westra, Bart J. J. M. van den Hurk, Sonia I. Seneviratne, Philip J. Ward, Andy Pitman, Amir AghaKouchak, David N. Bresch, Michael Leonard, Thomas Wahl & Xuebin Zhang

Nature Climate Change **8**, 469–477(2018) | Cite this article

4454 Accesses | **76** Citations | **316** Altmetric | Metrics


An Author Correction to this article was published on 20 June 2018

This article has been updated

ᑭᓯᐊᓂ ᖃᓄᐃᓗᒃᑐᑦᑕᐅᑎᒧᓂᑦ ᐱᒃᓯᑕᐅᔨᑦ, ᖃᐅᔨᓴᖅᓂᖅ ᐊᐢᓂᐊᓐᓱᓂ ᐊᓯᔾᔨᖅᑐᑦ ᐱᖅᓇᖅᑐᖅ.

Abstract

Floods, wildfires, heatwaves and droughts often result from a combination of interacting physical processes across multiple spatial and temporal scales. The combination of processes (climate drivers and hazards) leading to a significant impact is referred to as a 'compound event'. Traditional risk assessment methods typically only consider one driver and/or hazard at a time, potentially leading to underestimation of risk, as the processes that cause extreme events often interact and are spatially and/or temporally dependent. Here we show how a better understanding of compound events may improve projections of potential high-impact events, and can provide a bridge between climate scientists, engineers, social scientists, impact modellers and decision-makers, who need to work closely together to understand these complex events.



Global Change Biology

Global Change Biology (2015) **21**, 4651–4661, doi: 10.1111/gcb.13051

Contrasting effects of warming and increased snowfall on Arctic tundra plant phenology over the past two decades

ANNE D. BJORKMAN[1,2], SARAH C. ELMENDORF[3,4], ALISON L. BEAMISH[1,5], MARK VELLEND[6] and GREGORY H. R. HENRY[1]

[1]*Department of Geography and Biodiversity Research Centre, University of British Columbia, Vancouver, BC V6T 1Z4, Canada,* [2]*German Centre for Integrative Biodiversity Research and University of Leipzig, Leipzig 04103, Germany,* [3]*National Ecological Observatory Network, Boulder, CO 80301, USA,* [4]*Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, CO 80309, USA,* [5]*Periglacial Research Unit, Alfred Wegener Institute, Potsdam 14473, Germany,* [6]*Département de biologie, Université de Sherbrooke, Sherbrooke, QC J1K2R1, Canada*

ᖁᐅᓴᓴᕐᓂᖅ ᐅᑎᖁᑕᖅᑐᓂᓕᕐᓂ ᓯᓚᒪᓕᐊᖏᑦᑐᑦ

Abstract

Recent changes in climate have led to significant shifts in phenology, with many studies demonstrating advanced phenology in response to warming temperatures. The rate of temperature change is especially high in the Arctic, but this is also where we have relatively little data on phenological changes and the processes driving these changes. In order to understand how Arctic plant species are likely to respond to future changes in climate, we monitored flowering phenology in response to both experimental and ambient warming for four widespread species in two habitat types over 21 years. We additionally used long-term environmental records to disentangle the effects of temperature increase and changes in snowmelt date on phenological patterns. While flowering occurred earlier in response to experimental warming, plants in unmanipulated plots showed no change or a delay in flowering over the 21-year period, despite more than 1 °C of ambient warming during that time. This counterintuitive result was likely due to significantly delayed snowmelt over the study period (0.05–0.2 days/yr) due to increased winter snowfall. The timing of snowmelt was a strong driver of flowering phenology for all species – especially for early-flowering species – while spring temperature was significantly related to flowering time only for later-flowering species. Despite significantly delayed flowering phenology, the timing of seed maturation showed no significant change over time, suggesting that warmer temperatures may promote more rapid seed development. The results of this study highlight the importance of understanding the specific environmental cues that drive species' phenological responses as well as the complex interactions between temperature and precipitation when forecasting phenology over the coming decades. As demonstrated here, the effects of altered snowmelt patterns can counter the effects of warmer temperatures, even to the point of generating phenological responses opposite to those predicted by warming alone.

Keywords: Arctic tundra, Bayesian hierarchical modeling, climate change, flowering time, interval censoring, plant phenology, seed maturation

Community-level phenological response to climate change

Otso Ovaskainen, Svetlana Skorokhodova, Marina Yakovleva, Alexander Sukhov, Anatoliy Kutenkov, Nadezhda Kutenkova, Anatoliy Shcherbakov, Evegeniy Meyke, and Maria del Mar Delgado

PNAS August 13, 2013 110 (33) 13434-13439; https://doi.org/10.1073/pnas.1305533110

Edited by William H. Schlesinger, Cary Institute of Ecosystem Studies, Millbrook, NY, and approved July 9, 2013 (received for review March 25, 2013)

Article | Figures & SI | Info & Metrics | PDF

ᖁᐅᓴᓴᕐᓂᖅ ᐅᑎᖁᑕᖅᑐᓂᓕᕐᓂ ᓯᓚᒪᓕᐊᖏᑦᑐᑦ

Abstract

Climate change may disrupt interactions among species and ecosystem synchrony with adverse consequences to ecosystem functioning. We present here a 40-y-long time series of 10,425 dates that were systematically collected in a single Russian locality for 97 plant, 78 bird, 10 herptile, 19 insect, and 9 fungal phenological events, as well as for 77 climatic events related to temperature, precipitation, snow, ice, and frost. We show that species are shifting their phenologies at dissimilar rates, partly because they respond to different climatic factors, which in turn are shifting at dissimilar rates. Plants have advanced their spring phenology even faster than average temperature has increased, whereas migratory birds have shown more divergent responses and shifted, on average, less than plants. Phenological events of birds and insects were mainly triggered by climate cues (variation in temperature and snow and ice cover) occurring over the course of short periods, whereas many plants, herptiles, and fungi were affected by long-term climatic averages. Year-to-year variation in plants, herptiles, and insects showed a high degree of synchrony, whereas the phenological timing of fungi did not correlate with any other taxonomic group. In many cases, species that are synchronous in their year-to-year dynamics have also shifted in congruence, suggesting that climate change may have disrupted phenological synchrony less than has been previously assumed. Our results illustrate how a multidimensional change in the physical environment has translated into a community-level change in phenology.

nature climate change

LETTERS

PUBLISHED ONLINE: 6 JULY 2015 | DOI: 10.1038/NCLIMATE2697

Climate sensitivity of shrub growth across the tundra biome

Isla H. Myers-Smith[1]*, Sarah C. Elmendorf[2,3], Pieter S. A. Beck[4,5], Martin Wilmking[6], Martin Hallinger[6,7], Daan Blok[8], Ken D. Tape[9], Shelly A. Rayback[10], Marc Macias-Fauria[11], Bruce C. Forbes[12], James D. M. Speed[13], Noémie Boulanger-Lapointe[14,15], Christian Rixen[16], Esther Lévesque[14], Niels Martin Schmidt[17], Claudia Baittinger[18], Andrew J. Trant[19,20], Luise Hermanutz[20], Laura Siegwart Collier[20], Melissa A. Dawes[16], Trevor C. Lantz[19], Stef Weijers[21], Rasmus Halfdan Jørgensen[22], Agata Buchwal[23], Allan Buras[6], Adam T. Naito[24], Virve Ravolainen[25], Gabriela Schaepman-Strub[26], Julia A. Wheeler[16,27], Sonja Wipf[16], Kevin C. Guay[5], David S. Hik[28] and Mark Vellend[29]

ᐃᓕᓴᕆᔨᐃᑦ ᓇᓗᓇᐃᔨᖅᓯᒪᓂᖅᐳᑦ

Rapid climate warming in the tundra biome has been linked to increasing shrub dominance[1–4]. Shrub expansion can modify climate by altering surface albedo, energy and water balance, and permafrost[2,5–8], yet the drivers of shrub growth remain poorly understood. Dendroecological data consisting of multi-decadal time series of annual shrub growth provide an underused resource to explore climate–growth relationships. Here, we analyse circumpolar data from 37 Arctic and alpine sites in 9 countries, including 25 species, and ~42,000 annual growth records from 1,821 individuals. Our analyses demonstrate that the sensitivity of shrub growth to climate was: (1) heterogeneous, with European sites showing greater summer temperature sensitivity than North American sites, and (2) higher at sites with greater soil moisture and for taller shrubs (for example, alders and willows) growing at their northern or upper elevational range edges. Across latitude, climate sensitivity of growth was greatest at the boundary between the Low and High Arctic, where permafrost is thawing[4] and most of the global permafrost soil carbon pool is stored[9]. The observed variation in climate–shrub growth relationships should be incorporated into Earth system models to improve future projections of climate change impacts across the tundra biome.

The Arctic is warming more rapidly than lower latitudes owing to climate amplification involving temperature, water vapour, albedo and sea ice feedbacks[5,7]. Tundra ecosystems are thus predicted to respond more rapidly to climate change than other terrestrial ecosystems[4]. The tundra biome spans Arctic and alpine regions that have similar plant species pools and mean climates, yet vary in topography, seasonality, land cover and glaciation history. Concurrent with the recent high-latitude warming trend[7], repeat photography and vegetation surveys have shown widespread expansion of shrubs[1–3], characterized by increased canopy cover, height and abundance. However, climate warming[7] and shrub increase[2,10] have not occurred at all sites. Models predict that warming of 2–10 °C (ref. 11) could convert as much as half of current

[1]School of GeoSciences, University of Edinburgh, Edinburgh EH9 3FF, UK. [2]National Ecological Observatory Network, 1685 38th Street, Suite 100, Boulder, Colorado 80301, USA. [3]Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, Colorado 80309, USA. [4]European Commission, Joint Research Centre (JRC), Institute for Environment and Sustainability (IES), Forest Resources and Climate Unit, Via Enrico Fermi 2749, 21027 Ispra VA, Italy. [5]Woods Hole Research Center, 149 Woods Hole Road, Falmouth, Massachusetts 02540, USA. [6]Institute of Botany and Landscape Ecology, Ernst-Moritz-Arndt University Greifswald, D-17487 Greifswald, Germany. [7]Department of Ecology, Swedish University of Agricultural Sciences,

Global Change Biology

Global Change Biology (2009) 15, 2681–2693, doi: 10.1111/j.1365-2486.2009.01935.x

Herbivores inhibit climate-driven shrub expansion on the tundra

JOHAN OLOFSSON*, LAURI OKSANEN†‡, TERRY CALLAGHAN§¶, PHILIP E. HULME‖, TARJA OKSANEN* and OTSO SUOMINEN†


*Department of Ecology and Environmental Science, Umeå University, S-901 87 Umeå, Sweden, †Department of Biology, University of Turku, FI-20014 Turku, Finland, ‡Department of Natural Sciences, Finnmark University College N-9509 Alta, Norway, §Abisko Scientific Research Station, Abisko, Sweden, ¶Department of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2TN, UK, ‖National Centre for Advanced Bio-Protection Technologies, Lincoln University, PO Box 84, Canterbury, New Zealand


ᐃᓕᓴᕆᔨᐃᑦ ᓇᓗᓇᐃᔨᖅᓯᒪᓂᖅᐳᑦ ᐱᑭᖅᑐᓂᑦ ᖃᐅᔨᓴᓂᑦ ᖁᐅᓴᐃᓂᖅᐳᑦ

Recent Pan-Arctic shrub expansion has been interpreted as a response to a warmer climate. However, [illegible] to which the abundance of shrubs in arctic ecosystems. We addressed [illegible] alternative explanations by following the changes in plant community composition during the last 10 years in permanent plots inside and outside exclosures with different mesh sizes that exclude either only reindeer or all mammalian herbivores including voles and lemmings. The exclosures were replicated at three forest and tundra sites at four different locations along a climatic gradient (oceanic to continental) in northern Fennoscandia. Since the last 10 years have been exceptionally warm, we could study how warming has influenced the vegetation in different grazing treatments. Our results show that the abundance of the dominant shrub, *Betula nana*, has increased during the last decade, but that the increase was more pronounced when herbivores were excluded. Reindeer have the largest effect on shrubs in tundra, while voles and lemmings have a larger effect in the forest. The positive relationship between annual mean temperature and shrub growth in the absence of herbivores and the lack of relationships in grazed controls is another indication that shrub abundance is controlled by an interaction between herbivores and climate. In addition to their effects on taller shrubs (>0.3 m), reindeer reduced the abundance of lichens, whereas microtine rodents reduced the abundance of dwarf shrubs (<0.3 m) and mosses. In contrast to short-term responses, competitive interactions between dwarf shrubs and lichens were evident in the long term. These results show that herbivores have to be considered in order to understand how a changing climate will influence tundra ecosystems.

Keywords: *Betula nana*, exclosures, global warming, herbivores, lemmings, reindeer, shrubs, snow, tundra, voles

Received 15 October 2008 and accepted 16 February 2009



R1Z1
R1Z2
R1Z4
R1Z5
R1Z6
R1Z7
R2Z7
R4Z7
R3Z7
R3Z6
R3Z5
R3Z4
R3Z3
R3Z2
R3Z1
Rankin Inlet
Whale Cove


Banks Lake JB
Snug Lake JB
Rankin Inlet
Whale Cove

ᑐᑦᑐᓄᑦ ᐊᑐᖅᑕᖅ

ᖃᐅᔨᓴᖅᓯᐅᔪᖅ ᓇᒻᓂᓕᖓ


ᐱᑭᖅᑐᓕᑦ ᐃᓕᐅᔪᖅ




ᐱᑭᖅᑐᑦ



ᐱᓯᖅᑐᖅᓴᐅᓚᓂᓕᖅ














ᐃᓐᓇᓱᒪᓂᑦ

ᐳᓴᓕᑎᓐᓄᑦ ᐱᖃᓱᐅᓴᓇᓕᖅᑐᖅ

ᐅᒪᓕᐊᓂᑦ ᐱᑕᖃᖅᑐᑦ/ᐱᑕᖃᖅᑎᑎᐊᖅᓯᓐᓂᑦ


Banks Lake JB
Rankin Inlet
Snug Lake JB
Whale Cove



15 m
0.5 m
ᑕᒪᐅᓇᑐᐃᓐᓂᖅ
ᑕᒪᐅᓇᑐᐃᓐᓂᖅ
ᑕᒪᐅᓇᑐᐃᓐᓂᖅ
ᑕᒪᐅᓇᑐᐃᓐᓂᖅ



50cm
15cm
ᖃᐅᔨᓴᕕᖏᓐᓂᔪᓐᓇᖅᑐᖅ ᐱᒋᖅᑐᓄᑦ
ᖃᐅᔨᓴᕕᖏᓐᓂᔪᓐᓇᖅᑐᖅ ᐊᑦᓴᑦᓄᑦ


ᑐᑦᑐᓄᑦ ᐊᑐᖅᑕᖅ

ᖃᐅᔨᓴᕐᓂᐅᔪᖅ ᓇᒫᓂᓂᖓ

ᐱᑭᖅᑐᓂᑦ ᐃᓂᐅᔪᖅ





R1Z1
R1Z2
R1Z4
R1Z5
R1Z6
R1Z7
R2Z7
R4Z7
R3Z7
R3Z6
R3Z5
R3Z4
R3Z3
R3Z2
R3Z1
Rankin Inlet
Whale Cove


Banks Lake JB
Snug Lake JB
Rankin Inlet
Whale Cove

ᖃᐅᓂᖅᐸᐊᖅ ᓄᖅᕿᓕᐅᖅᐃᖏᑦ ᓄᓇᐃᑦ

	2012	2013	2014	2015	2016	2017	2018	2019
ᑐᑭᒧᐊᖅᑎᑦᑎᐸᖅ	N. Lecomte	N. Lecomte	/	M. Wilson	M. Wilson	/	NuWCRU	NuWCRU
ᐱᑭᐅᑐᑦ ᓄᓇᒥ	ᐃ	ᐃ	/	ᐃ	ᐃ	/	ᐃ	ᐃ
ᖃᐅᒪᔪᑦ	ᐊᖅᑕᐅᓂᖓ	ᐊᖅᑕᐅᓂᖓ / ᓇᓴᖅᑕᖅ	/	ᐊᖅᑕᐅᓂᖓ	No	/	ᐊᖅᑕᐅᓂᖓ	ᐊᖅᑕᐅᓂᖓ / ᓇᓴᖅᑕᖅ
ᐊᐃᔨᓕᐅᑭᑦ	ᐊᑦᖏ	ᐊᑦᖏ	/	ᐃ	ᐃ	/	ᐊᑦᖏ	l ᐊᑦᖏ
ᓂᐱᓄᑦ ᓂᐱᓕᐅᑭᖅ	ᐊᑦᖏ	ᐊᑦᖏ	/	ᐃ	ᐃ	/	ᐊᑦᖏ	No



RANKIN INLET









ᖃᓂᒃᑯᑦ ᓂᕆᔨᒋᑦ
ᓂᕆᔨᒋᑦᓴᐃᓕᓐᓂᒃ
ᓂᕆᔨᒋᑦᓴᐃᓕᓐᓂᒃᑲᒪᒋᔭᖅᑎᓪᓗᒍ
ᓂᕆᔨᒋᑦᓴᐃᓕᓐᓂᒃ
ᐱᓕᕆᔨᒃᑐᖅᓴᐃᓕᓐᓄᑦ



ᐊᒋᑦᑎᓂ
ᐅᓂᒃᑕᖅᑐᑦ
ᐱᓕᕆᔨᖅ
ᐃᓚᖏᑦ
ᐊᑐᕐᓂᖅ
ᐅᖃᓕᒫᑦ
ᐊᑐᕐᓂᖅ




ᐱᑭᖅᑐᓂᑦ ᖃᐅᔨᓴᓂᖅ ᐅᑎᖃᒋᖅᑐᓂᑦ






ᐳᔪᐃᑦ ᐱᑭᖅᑐᑦ





ᐱᑭᖅᑐᓂᑦ ᖃᐅᔨᓴᕐᓂᖅ ᐅᑎᖃᒋᖅᑐᓂᑦ



ᐳᔾᐃᑦ ᐱᑭᖅᑐᑦ


1. ᐅᓪᓗᖓ ᐊᐳᑎᖃᑭᓚᕐᓂᖓ
2. ᐅᓪᓗᖓ ᓯᕗᓪᓕᖅᑕᒥᑦ ᓴᖅᑭᖅᑐᑦ ᐱᕐᔪᖅᓯᐊᑦ
3. ᐅᓪᓗᖓ ᓯᕗᓪᓕᖅᑕᒥᑦ ᐱᑭᑭᓇᐅᑎᑦ ᑕᔪᐃᓴᐅᔾᔪᑦ (ᕐᔪᓯᖅᑐᑦ ᐱᑭᖅᑐᓕᕆᐅᑭᑎᖓ)


ᐱᑕᖃᓂᖅᑐᑦ ᖁᓇᐃᓕᓴᐅᓂᖅᑐᑦ






ᐱᑕᖅᓂᖅᑐᑦ ᖃᓄᐃᓕᓚᖅᑐᑦ











ᐱᑕᖃᕐᓂᖅᑕ ᖃᓄᐃᓕᓐᖓᓂᖅᑕ








ᐱᑕᖃᓂᓕᕆᓂ ᖃᓄᐃᓕᓚᖅᓂᓕᕆᓂ


ᓄᓇᖅᑕ ᕿᓕᓚᖅ



ᐅᒪᔨᖃᑕᐅᓂᖅ


- Andromeda polifolia
- bare ground
- Bryoria nitidula
- detritus
- Empetrum nigrum
- Flavocetraria cucullata
- Rhododendron tomentosum
- Lycopodium annotinum
- Masonhalea richardsonii
- Rhododendron lapponicum
- Rock
- Thamnolia subuliformis sp.
- Vaccinium uliginosum
- Vaccinium vitis-idaea

ᓄᕐᓂ ᖃᐅᔨᓴᖃᐅᔨᓪᓗ





ᐱᑕᖃᕐᓂᖅᑐᑦ ᖃᓄᐃᓕᓴᖃᓂᖅᑐᑦ




ᐅᒪᔨᖃᑕᐅᓂᖓ

Betula glandulosa


no vegetation

ᓄᓇᓕᓂ ᖃᐅᔨᓴᕐᕕᐅᔪᑦ






ᐱᑕᖅᓂᖅᑐᑦ ᖃᓄᐃᓕᓴᖅᑐᑦ




ᓄᓇᓕ ᖃᐅᔨᓴᓇᐅᔪᑦ





ᐃᓕᓴᕐᓂᖅ ᐅᓂᖅᓴᓂᐅᕙᓪᓕᐊᓂᖅ



ᐃᓱᒪᖅᑐᓄᑦ ᐅᓂᖅᓴᑦ











ᒪᑭᔪᓂᖅ ᐅᒪᔪᓂᖅ
ᐊᖏᕐᕋᓱᒃ
ᐅᑎᖅᑕᖅᑐᑦ
ᖃᓄᐃᓕᓴᐃᓂᖅ
ᐱᔪᓐᓇᖅᑐᑦ
ᓄᓇᕗᑦ ᓂᕆᑐᖅᑐᖅ
ᑕᒪᓐᓇᑦ ᐅᓂᒃᑎᑎᑐᓂᖅ
ᓴᓇᔨᓕᐅᕐᓂᖅ
ᓄᓇᕗᑦ ᓂᕆᑐᖅᑐᖅ
ᐃᓚᖏᑦ ᐅᓂᒃᑎᑎᑐᓂᖅ
ᐊᑐᖅᑐᑦ ᓂᕆᑐᖅᑐᖅ
ᐃᓚᖏᑦ ᐅᓂᒃᑎᑎᑐᓂᖅ
ᐊᑐᖅᑐᑦ ᓂᕆᑐᖅᑐᖅ
ᐃᓚᖏᑦ ᐅᓂᒃᑎᑎᑐᓂᖅ
ᐊᑐᖅᑐᑦ ᓂᕆᑐᖅᑐᖅ
ᐃᓚᖏᑦ ᐅᓂᒃᑎᑎᑐᓂᖅ
ᐊᑐᖅᑐᑦ ᓂᕆᑐᖅᑐᖅ







20
19
18
17
16
15
14
13
12
11
10
9
8
7
6
5
4
3
2
1
A
B
C
D
300m


15
14
13
15m



1m

ᓄᓇᒥᐅᑉ ᓂᕆᔨᒋᑦᓯᒪᓂᖅ
ᐊᒥᓱᓕᕆᓂᖅ
ᐃᓕᓴᕐᓂᐊᕐᓂᖅ
ᑲᑎᒪᔨᖅ ᑕᒪᐃᓐᓇᖅ
ᐊᑐᕐᓂᖅ
ᐊᒥᓱᓕᕆᓂᖅ
ᐃᓚᖓᑕᐅᓂᖅ
ᐊᑐᕐᓂᖅ
ᐊᑐᕐᓂᖅ
ᐊᑐᕐ











12 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 21 22 23 24






12 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 21 22 23 24











File Edit Multitrack Clip Effects Favorites View Window Help
Waveform
Multitrack
Workspace: Default
Search Help
Files
Favorites
Name
Status
Duration
XC43259... gambelii.mp3 *
0:50.909
Markers
Name
Start
End
Durati
Media Browser
Effects Rack
Properties
XC43259...w - Zonotrichia leucophrys gambelii.mp3 *
Info
Duration: 0:50.909
Sample Rate: 24000 Hz
Channels: Stereo
Bit Depth: 32-bit (float)
Format: MP3 160 Kbps CBR
File Path: C:\User...onotrichia leucophrys gambelii.mp3
History
Video
Editor: XC432599 - White-crowned Sparrow - Zonotrichia leucophrys gambelii.mp3 *
0:26.287
Levels
Selection/View
Start
End
Duration
Selection 0:26.287 0:26.287 0:00.000
View 0:20.115 0:50.909 0:30.794







Ecology and Evolution

Open Access

Using bioacoustics to examine shifts in songbird phenology

Rachel T. Buxton[1], Emma Brown[2], Lewis Sharman[3], Christine M. Gabriele[3] & Megan F. McKenna[2]

[1]Department of Fish, Wildlife and Conservation Biology, Colorado State University, 1474 Campus Delivery, Fort Collins, Colorado 80523
[2]Natural Sounds and Night Skies Division, National Park Service, 1201 Oakridge Drive, Fort Collins, Colorado 80525
[3]Glacier Bay National Park and Preserve, PO Box 140, Gustavus, Alaska 99826

Keywords
Acoustic Complexity Index, Alaska, climate change, Glacier Bay, monitoring, songbirds, soundscape.

Correspondence
Rachel T. Buxton, Department of Fish, Wildlife and Conservation Biology, Colorado State University, 1474 Campus Delivery, Fort Collins, CO 80523, USA.
Tel: 1 970 491 8740;
Fax: 1 970 491 5091;
E-mail: Rachel.Buxton@colostate.edu

Funding Information

Abstract

Monitoring patterns in biodiversity and phenology have become increasingly important given accelerating levels of anthropogenic change. Long-term monitoring programs have reported earlier occurrence of spring activity, reflecting species response to climate change. Although tracking shifts in spring migration represents a valuable approach to monitoring community-level consequences of climate change, robust long-term observations are challenging and costly. Audio recordings and metrics of bioacoustic activity could provide an effective method for monitoring changes in songbird activity and broader biotic interactions. We used 3 years of spring and fall recordings at six sites in Glacier Bay National Park, Alaska, an area experiencing rapid warming and glacial retreat, to examine the utility of bioacoustics to detect changes in songbird phenology. We calculated the Acoustic Complexity Index (ACI), an algorithm representing an index of bird


ᐊᓯᖏᑲᓐᓂᕐᑐᑦ ᓇᓗᓇᖅᓂᓚᕆᖏᓐᓂ ᖃᐅᔨᓴᕐᓂ ᖃᓄᐃᓗᒃᑖᑦ ᐅᑭᐅᖅᑯᑦ ᐅᐱᕙᓪᓕᐊᓂ ᐊᑐᖅᑕᐅᓂᐅᖅᑐᑦ
ᖃᐅᔨᔪᒪᐅᓪᓗᑎ ᑎᑭᓂᓂᖏᓐᓄᑦ ᐅᑎᖅᑕᖃᕐᑕᖅᑐᑦ ᖃᐅᔨᓴᕐᓂ

Ecology and Evolution 2016; 6(14): 4697–4710

doi: 10.1002/ece3.2242


abrupt in the fall due to weather events, suggesting spring recordings are better suited to indicate phenology. To ensure changes in ACI values were detecting real changes in songbird activity, we explored the relationship between ACI and song of three species: varied thrush (*Ixoreus naevius*), Pacific wren (*Troglodytes pacificus*), and ruby-crowned kinglet (*Regulus calendula*). ACI was positively related to counts of all species, but most markedly with song of the varied thrush, the most common species in our recordings and a known indicator of forest ecosystem health. We conclude that acoustic recordings paired with bioacoustic indices may be a useful method of monitoring shifts in songbird communities due to climate change and other sources of anthropogenic disturbance.




ACI
R1Z7 - 2015
21000
20500
20000
19500
19000
18500
18000
Jun 01
Jul 01
Aug 01


ᑎᒃᓚᐊᖅ ᓂᕆᑐᖅᑐᖅ








Avian top predators









ᓄᕐᕋᓕᐅᕐᐸᖏᑦ ᓄᓇᐃᑦ – ᖃᐅᔨᓴᕐᓂᖅᑕᐊᕐᓂᖅ

2015-2016 < 50 ᐊᐱᓵᖏᐊᑦ ᐊᖅᑯᐸᖅᑐᓂ ᖃᐅᔨᓴᓂᕐᒥᑦ



2015-2016 ≥ 181, 960 ᒥᓂᑦᔪ ᐊᖅᑯᐸᖅᑐᓂ ᖃᐅᔨᓴᓂᕐᒥᑦ
2015-2016 > 48, 000 ᐊᐱᓵᖏᐊᑦ ᐊᖅᑯᐸᖅᑐᓂ ᖃᐅᔨᓴᓂᕐᒥᑦ



2012-2015 ≥ 192 ᐃᓂᐅᔪᖅ*ᐊᕐᕿᒍᐃᑦ, 695 ᑐᑦᑐᓄᑦ ᐊᓇᖏᑦ
2012-2015 ≥ 192 ᐃᓂᐅᔪᖅ*ᐊᕐᕿᒍᐃᑦ, 294 ᒥᑭᔪᓄᑦ ᐅᒪᔪᓄᑦ ᐊᓇᖏᑦ
2012-2015 ≥ 192 ᐃᓂᐅᔪᖅ*ᐊᕐᕿᒍᐃᑦ, 294 ᐊᖏᕐᓂᓄᑦ ᐊᓇᖏᑦ





2012-2019 ≥ 192 ᐃᓂᐅᔪᖅ*ᐊᕐᕿᒍᐃᑦ, 3,840 ᐅᒪᔪᐃᑦ ᓇᓗᓇᐃᖅᑕᐅᔪᑦ
2012-2019 ≥ 192 ᐃᓂᐅᔪᖅ*ᐊᕐᕿᒍᐃᑦ, 1920 ᒥᑭᔪᑦ ᖃᐅᔨᓴᖅᐳᔪᑦ ᐊᐱᓵᖏᐊᑦ



ᑲᖏᕐᓀᓂᖅ – ᖃᐅᔨᓴᓂᕐᑎᐊᕐᓂᖅ

1980-2019 ≥ 1200 ᐱᑭᖅ*ᐊᕐᕿᒍᐃᑦ

2015-2019 ≥ 550 1-ᑭᓪᒋᑕᐃᑦ ᖃᐅᔨᓴᕐᕕᒃᐳᔪᑦ ᒧᓐᓇᐅᔪᒥᑦ ᖃᐅᔨᓴᐃᓂᕐᒥᑦ
2019 ≥ 73, 130 ᒥᓂᑦ ᐊᖅᑯᐸᖅᑐᓄᑦ ᖃᐅᔨᓴᐃᓂᖅ

2008-2019 ≥ 720 ᐊᔪᓕᐅᖅᑲᔪᑦ*ᐅᑦᓄᐃᑦ/ᐊᕐᕿᒍ

2019 ≥ 17, 377 ᐊᔪᓕᐅᖅᑲᔪᑦ *ᐃᑲᕐᕿᐃᖅ, 376 ᐅᒪᔪᐃᑦ ᓇᓗᓇᐃᖅᑕᐅᔪᑦ

2019 ≥ 10 ᓄᓀᓕᒃ ᖃᐅᔨᓴᕐᕕᐅᔪᖅ ᐃᓂᖏᑦ, 1500 ᐅᒪᔪᐃᑦ ᓇᓗᓇᐃᖅᑕᐅᔪᑦ

ᐊᐱᕐᓂᑦ





LTEMP

Long-term Ecosystem Monitoring Program

Nunavut Wildlife Cooperative Research Unit
Government of Nunavut
University of Alberta

2019 - Nov
Annual Report

Erik Hedlin
Alastair Franke
Kevin Hawkshaw
Rebekah McKinnon
Nick Gulotta

Disclaimer: This draft report has been prepared for the Government of Nunavut and presents preliminary findings that may change with further analysis. The views and interpretations in this report article are those of the authors and do not necessarily represent the views of the Government of Nunavut. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the Government of Nunavut.

Long-term Ecological Monitoring Program: Executive Summary It is well established that temperatures in the Arctic have risen at almost twice the rate of temperatures elsewhere. However, studying the e[U+FB00]ects of climate change on wildlife is challenging; unlike physical processes that generally respond to increased temperature in predictable ways (e.g., permafrost, glacial erosion and oceanic winds, tides and densities), animals can respond in surprising ways, and assessing environmental change that results from natural and anthropogenic drivers in ecosystems that are inherently variable exacerbates these challenges. Ecosystem change is expected to be signi[U+FB01]cant and ongoing. Shifts in the distribution and demography of tundra-breeding wildlife, such as barren-ground caribou, are expected to occur with climate-induced increases in vegetation growth and range expansion, potentially bene[U+FB01]tting species associated with shrub-dominated habitats, and harming species more reliant on tundra habitat, through climate-induced habitat loss. Similarly, pathogens and parasites currently excluded from the Arctic are expected to become established through range expansion and warming. Novel relationships among hosts and parasites can increase the prevalence of infectious diseases, increase host morbidity and mortality, and have cascading impacts on food webs and ecosystem functioning. Similarly, over the long-term, the frequency of extreme weather events, outside of historical range in variation, is likely (e.g., rain on snow, summer storms, temperature highs and lows, snow cover and depth). The ecological consequences of these events include changes to food webs and timing of key ecological relationships, shifts in species distribution and community composition, and changes in demography.

To address two primary concerns (documenting long-term changes associated with climate change, and documenting baseline conditions in a region of high interest for economic development), the Government of Nunavut, Department of Environment (DOE) established the Long-Term Ecological Monitoring Program (LTEMP) in 2012. The goal of the program is to undertake annual monitoring at spatial and temporal scales relevant to monitoring ecosystem components within the summer range of the Qamanirjuaq caribou herd. Within the calving grounds of the Qamanirjuaq herd, extensive monitoring has been conducted periodically at 16 permanent locations since 2012, and starting in 2019, intensive annual monitoring was initiated near the community of Rankin Inlet. The intensive monitoring near Rankin Inlet incorporates multiple ecosystem components across three trophic levels (primary producers to tertiary consumers), while the extensive monitoring in the summer range of the Qamanirjuaq herd focuses on vegetation and pellet counts.

To address research and staffing capacity issues associated with the LTEMP, and to ensure the continuity of the project, the DOE decided to collaborate with academic partners. A Memorandum of Understanding with the University of Alberta, Department of Biological Sciences, and Dr. Alastair Franke was developed to facilitate the partnering of academic researchers with Government of Nunavut-led research projects, such as LTEMP. The main purpose of the LTEMP is to maintain a long-term (i.e., decades), field-based study to monitor ecosystem changes, and incorporate the results into wildlife management practices. Specifically, this involves site-specific, standardized data collection for terrestrial (scat counts, bird surveys, small mammal surveys, invertebrate sampling, and vegetation surveys) and physical (temperature, precipitation, land cover) ecosystem components.

Intensive research site monitoring was conducted during the summer near the community of Rankin Inlet (Figure 1) in 2019. The study area represents approximately 2500 km2 of coastal tundra and marine habitat typical of the low Arctic, containing a diversity of wet meadows, dwarf shrubs, dry eskers, rocky ridges, tidal flats and sea cliffs. The landscape consists of low rolling hills, interspersed with ponds and lakes. Rock outcrops are common throughout the area, and are well-developed as barrier islands. Ridge tops, upland areas, and well-drained slopes support lichens, moss, and low shrubs. Dominant vascular plants are Labrador tea (Ledum palustre), mountain cranberry (Vaccinium vitisidaea), and crowberry (Empetrum nigrum). Bell heather (Cassiope tetragona) occurs in low-lying areas and rock crevices. Vegetation on most slopes and hillsides consists of heath patches comprising Dryas integrifolia, Carex spp., Cassiope, and lichens.

We have structured our study design to incorporate monitoring of multiple ecosystem components across three trophic levels (primary producers to tertiary consumers). Monitoring of vegeta-

tion, including a warming experiment (staring in 2020), is central to the overall study design. To monitor plant species composition and vegetation structure, five stations each comprising four replicates (two control and two treatment) were selected based on distribution and proximity of three vegetation types, shrub, heath, and graminoid. For each replicate, three 1m X 1m point-framing-plots (PFPs) have been permanently established within uniform patches of three vegetation types (shrub, heath, and graminoid), resulting in a total of 60 PFPs (30 control and 30 treatment). To monitor plant phenology, we have established a 50-m transect to monitor three species, Dryas integrifolia (mountain avens), Eriophorum vaginatum (cottongrass), and Salix arctica (Arctic willow), respectively. Other ecosystem components (small mammals, arthropods, avian community, and raptors) are monitored simultaneously, along with abiotic variables associated with weather, snow-cover, freeze-up and thaw. For arthropods, one pitfall trap was placed in each vegetation type (graminoid, shrub, and heath) at each control and treatment plot (4 plots total) at each PFP site (5 sites total), for a total of 60 pitfall traps. In addition, we documented pollinator diversity using pan traps. Two small mammal trapping stations were established each comprising effort no less than 720 trap-nights per trapping session.

Extensive, short-term monitoring was also conducted within the calving grounds of the Qamanirjuaq herd (Figure 2) in the summer of 2019. Sixteen permanent sampling sites have been established along a gradient of historical intensity of use by caribou, as determined by collar data collected between 1993 and 2012. Each of the 16 sampling locations contain three vegetation plots, one each for shrub, graminoid, and heath dominated micro-environments. Vegetation sampling occurs at all plots, and can be broken down into two main categories: (i) visual estimation of percent cover, and, (ii) species presence/richness. In addition, we record species-specific pellet counts annually as an index of animal use.

Data collection, back-up and storage, and preliminary data analysis has been conducted for vegetation community sampling (n = 15 locations), carabid and spider sampling (n = 15 locations), passive songbird sampling (n = 13 locations), active songbird sampling (n = 60 locations), historical shrub growth sampling (n = 7 locations), rainfall measurements (n = 4 locations), and raptor monitoring (study site wide) for the Rankin Inlet study area. Historical data collected (i.e., prior to 2018) in the calving grounds has similarly been backed-up and stored, and basic analysis of the pellet count data (e.g., site-specific counts per species for each habitat type) has been completed.

Contents

1 Introduction **6**
1.1 Study Objectives . . . 6
1.2 Program History . . . 7
1.2.1 Qamanirjuaq Calving Grounds . . . 7
1.2.2 Rankin Inlet . . . 8
1.3 Study Area . . . 9
1.3.1 Rankin Inlet . . . 9
1.3.2 Qamanirjuaq Caribou Herd Calving Grounds . . . 9

2 Methods **12**
2.1 Rankin Inlet . . . 12
2.1.1 Primary Producers . . . 12
2.1.2 Primary and Secondary Consumers . . . 15
2.1.3 Avian Herbivores and Insectivores . . . 16
2.1.4 Tertiary Consumers . . . 18
2.1.5 Physical . . . 18
2.2 Qamanirjuaq Calving Grounds . . . 19
2.2.1 Vegetation monitoring . . . 19
2.2.2 Pellet Counts . . . 20
2.2.3 Trail Cameras, ARUs and Weather Monitoring . . . 21

3 Results **23**
3.1 Rankin Inlet . . . 23
3.1.1 Primary Producers . . . 23
3.1.2 Primary and Secondary Consumers . . . 25
3.1.3 Avian Herbivores and Insectivores . . . 27
3.1.4 Tertiary Consumers . . . 28
3.1.5 Physical . . . 29
3.2 Qamanirjuaq Calving Grounds . . . 33
3.2.1 Vegetation Monitoring . . . 33
3.2.2 Pellet Counts . . . 34
3.2.3 Motion Sensitive Cameras . . . 36
3.2.4 ARUs . . . 38

4 Discussion **40**

5 Qamanirjuaq - Yearly Range **50**

6 Data Management Plan **51**

7 README Template in Markdown **54**

8 C3 Plant and Lichen Collecting Protocols **56**

9 Autonomous Recording Unit Transcription Protocol **63**
9.1 Converting File Formats . . . 63
9.2 Listening Software . . . 64
9.2.1 Software Settings . . . 64
9.3 Entering Data . . . 65
9.3.1 Site Info Data Entry . . . 65

9.3.2 Species Detections Entry 65
9.3.3 Additional Resources for Confirming Species ID 66

1 Introduction

It is well established that temperatures in the Arctic have risen at almost twice the rate of temperatures elsewhere [35]. However, studying the effects of climate change on wildlife is challenging [6]; unlike physical processes that generally respond to increased temperature in predictable ways (e.g., permafrost, glacial erosion and oceanic winds, tides and densities), animals can respond in surprising ways, and assessing environmental change that results from natural and anthropogenic drivers is challenging. In the Arctic, ecosystems show considerable annual variability, which exacerbates the challenge to understand the effects of natural and anthropogenic perturbations [7].

Ecosystem change is expected to be significant and ongoing [35, 86]. Shifts in distribution and demography of tundra-breeding wildlife, such as caribou, are expected to occur with climate-induced increases in vegetation growth and range expansion [75], potentially benefitting species associated with shrubs, and harming tundra obligates through climate-induced habitat loss [78, 54]. Similarly, pathogens and parasites currently excluded from the Arctic are expected to become established through range expansion and warming [43, 79]. For example, warmer conditions are thought to be associated with increasing prevalence of a parasitic nematode in muskoxen *Ovibos moschatus* [43]. These novel relationships among hosts and parasites can increase the prevalence of infectious diseases [27, 2, 79], increase host morbidity and mortality, and have cascading impact on food webs and ecosystem functioning [32]. Over the long-term, Collins et al. [1] indicated that the frequency of extreme weather (i.e., outside of historical range in variation) events is likely (e.g., rain on snow, summer storms, temperature highs and lows, snow cover and depth). The ecological consequences of these events include trophic mismatch [67], shifts in species distribution [82, 44], and species assemblages [67], and changes in demography [3]. Furthermore, changes in inputs from precipitation and outputs from evaporation and plant transpiration [34] can have greater impact on ecosystems than the progression of climate change *per se* [64], potentially leading to episodic tundra drying, altered hydrology, reduced permafrost and deeper active layer in summer [32]. In Nunavut, climate-sensitive factors are exacerbated by the potential for cumulative effects of economic development [19].


In order to effectively investigate questions of ecosystem function and cumulative effects, assessments should be considered at spatial and temporal scales relevant to species' of interest. Johnson et al. [36] argued that monitoring of multiple organism groups reduced uncertainty when conducting studies focused on detection of ecological change. In a separate study, Johnson et al. [37] compared response time of indicators of ecological change, and argued that nonlinear responses among organism groups can be used to identify early warning indicators. Ims and Fuglei [34] pointed out that many tundra species exhibit cycles, and argued that cycles result from trophic interactions within plant-based food webs, where microtine rodents play a critical role as prey, or as consumers of vegetation. Periodicity among species varies considerably; lemmings for example are thought to cycle approximately every four years [65, 8], ptarmigan species every ten years [4, 59], and caribou are thought to cycle over several decades [53]. Thus, distinguishing between natural cycles and declines that may cause local extinction [49] is extremely challenging, particularly for species that cycle over very long periods.

1.1 Study Objectives

The Government of Nunavut, Department of Environment recognizes the need to increase research capacity and productivity, including work related to wildlife population trends, species interactions, and research relating to the effects of anthropogenic (e.g., economic development) and natural disturbance regimes (e.g., climate change) on wildlife species and their habitat. The Department recognizes that a major challenge for management of wildlife in Nunavut will be to mitigate cumulative landscape impacts. This will be particularly important for critical areas, such as calving grounds, den sites, feeding grounds, migration routes and molting areas, and implies conserving important habitats at scales that preserve functional connectivity, ecosystem resilience and facilitates adaptation to climate change.

To expand research capacity and productivity that will support management of wildlife species in Nunavut, the Department of Environment signed a Memorandum of Understanding with Alastair Franke, the University of Alberta, Department of Biological Sciences. Alastair Franke and Caryn Smith established the Nunavut Wildlife Cooperative Research Unit (NuWCRU) as the basis for collaboration among four main stakeholder groups: (i) the department; (ii) community-based organizations; (iii) university-based researchers at the University of Alberta, and; (iv) proponents of economic development.

Within the context of NuWCRU goals and objectives, the Department directed Alastair Franke to build on the Long-term Ecosystem Program (L-TEMP).



1.2 Program History

The purpose of the Long-Term Ecological Monitoring Program is to assess ecosystem change within the summer range of the Qamanirjuaq caribou herd. The Qamanirjuaq Caribou Herd is the largest caribou herd in the western arctic occupying 300,000 km^2 [10]. Each year, Kivalliq Inuit and Manitoba Dene harvest approximately 8,000 and 5,000 animals, respectively. Aboriginal hunters in Saskatchewan and Northwest Territories harvest an additional 1,000 to 5,000 animals annually [17]. The estimated value of the harvest is 15 to 25 million dollars annually. A decline in herd size, first reported in the early 1950's, resulted in follow-up surveys dedicated to estimating herd size. Although estimates vary considerably, Boulanger et al. [10] suggest that the Qamanirjuaq herd is in a slow decline. The goal of long-term monitoring outlined here is to complement existing species-specific monitoring initiatives (e.g., government-led caribou and muskox surveys) in order to describe long-term change associated with natural and anthropogenic drivers within the calving grounds of the Qamanirjuaq caribou herd. For caribou, ecosystem monitoring should be conducted at broad spatial scales, over long time periods (e.g., decades) using multiple organism groups, including those that are early warning indicators.

1.2.1 Qamanirjuaq Calving Grounds

Work was established in 2012 under the direction of the Government of Nunavut (GN) ecosystems biologist with the goal to undertake extensive monitoring at spatial and temporal scales relevant to monitoring change within the Qamanirjuaq caribou herd calving grounds. The initial study design was limited to assessing vegetation and pellet counts in the calving grounds. Monitoring with motion sensitive cameras and autonomous recording units was added in 2015. Fieldwork was not conducted in 2014 and 2017 because the Ecosystems Biologist position was vacant, and vegetation and scat

sampling were not conducted in 2016. Responsibility for program delivery was assigned to the GN-led Nunavut Wildlife Cooperative Research Unit (Table 1) in 2018, in part to enable continuity of annual monitoring.

	2012	2013	2014	2015	2016	2017	2018	2019
Lead	N. Lecomte	N. Lecomte	/	M. Wilson	M. Wilson	/	NuWCRU	NuWCRU
Veg	Yes	Yes	/	Yes	No	/	Yes	Yes
Scat	Removal	Removal/Count	/	Removal	No	/	Removal	Removal/count
Camera	No	No	/	Yes	Yes	/	No	No
ARU	No	No	/	Yes	Yes	/	No	No

Table 1: Summary of field work completed as part of the Long-term Ecosystem Monitoring Program from 2012 – 2019.

Intensive summer monitoring in the calving grounds is not possible due to the potential for disturbance to caribou during the calving period. Monitoring capable of evaluating change across multiple trophic levels requires frequent sampling [55]. For example, arthropod pitfall traps need to be emptied and recharged bi-weekly, vegetation phenology metrics have to be examined every two days, small mammal sampling has to be conducted daily, and ensuring sufficient sample size for vegetation communities requires full time work over the course of several weeks. Implementing such a program throughout the Qamanirjuaq Calving grounds is not feasible, and NuWCRU therefore established an additional study area surrounding Rankin Inlet in 2019.

All L-TEMP data were provided to NuWCRU personnel in October 2018. An external hard-drive containing all relevant data was shipped to the NuWCRU office at the University of Alberta by O. Chaikina (received 7 October 2018). Lapses in fieldwork have resulted in three years of only pellet removal (i.e., to remove multi-year accumulation of scat), and only one year (2013) where counts reflect accumulation over the course of a single calving season. Vegetation sampling and time-series photographs of quadrats were repeated in four years (2012, 2013, 2015 and 2018) from 2012 - 2018. Trail camera and autonomous recording unit (ARU) data were collected only twice (2015 and 2016) over the seven-year period.

1.2.2 Rankin Inlet

In 2019, all existing LTEMP equipment was moved to Rankin Inlet with the intent of conducting intensive monitoring. Data collection was implemented in 2019 (Table 2).

	2012	2013	2014	2015	2016	2017	2018	2019
Lead	/	/	/	/	/	/	/	NuWCRU
Veg Phenology	/	/	/	/	/	/	/	No
Veg Community	/	/	/	/	/	/	/	Yes
Dendro	/	/	/	/	/	/	/	Yes
Rain	/	/	/	/	/	/	/	Yes
ARU	/	/	/	/	/	/	/	Yes
Distance Sampling	/	/	/	/	/	/	/	Yes
Pitfall	/	/	/	/	/	/	/	Yes
Pollinator Trap	/	/	/	/	/	/	/	No
Malaise Trap	/	/	/	/	/	/	/	No
Raptor Monitoring	/	/	/	/	/	/	/	Yes
Small Mammal	/	/	/	/	/	/	/	Yes

Table 2: Summary of field work completed as part of the Long-term Ecological Monitoring Program from 2012 to 2019 in the Rankin Inlet study area.

1.3 Study Area

1.3.1 Rankin Inlet

The study area (Figure 1) surrounds the community the Rankin Inlet, Nunavut, Canada (62.81° N, 92.09° W), encompassing approximately 2500 km^2 of coastal tundra habitat. The landscape is typical of the low Arctic, containing a diversity of wet meadows, dwarf shrubs, dry eskers, rocky ridges, tidal flats and sea cliffs. Summers are short and cool. Spring thaw begins in mid-May, reaching a peak by the first week of June, when ice cover on small lakes and ponds begins to melt. Land-fast sea ice persists until the second week of July. Annual rainfall is approximately 16cm, and snowfall is approximately 118.0. The landscape consists of low rolling hills, interspersed with ponds and lakes. Rock outcrops are common throughout the area, and are well developed as barrier islands. Ridge tops, upland areas, and well-drained slopes support lichens, moss, and low shrubs. Dominant vascular plants are Labrador tea (*Rhododendron tomentosum*), mountain cranberry (*Vaccinium vitisidaea*), and crowberry (*Empetrum nigrum*). Bell heather (*Cassiope tetragona*) occurs in low-lying areas and rock crevices. Vegetation on most slopes and hillsides consists of heath patches comprising *Dryas integrifolia*, Carex spp., Cassiope, and lichens.

1.3.2 Qamanirjuaq Caribou Herd Calving Grounds

The Dubwant Lake plain/upland ecoregion is primarily used in July and August during the post calving period. The eastern and southeastern portions of Kazan River upland ecoregion are used during fall migration and rut (September and October). The Coastal Hudson Bay lowland ecoregion within the Hudson Plains ecozone is most commonly used during late winter and occasionally during late fall (Appendix 5). The study region is rich in biological resources and diversity, and is under continuing pressure for mineral development. The Kivalliq Region contains several Tier 2 and 3 mineral projects that overlap with range of the Qamanirjuaq herd.

In 2012, cumulative caribou collar relocation data spanning from 1993 to 2012 were used to delineate seven zones that varied in intensity of use (Figure 2); Zone 7 contained the highest concentration of use during the calving period, and Zone 1 containing the lowest concentration of use[a]. As a result, sixteen permanent site locations extending northward and southward from the core calving grounds were established (Figure 2). Within each site location, Inuit Qaujimajatuqangit (IQ) was

[a] see Appendix 5 for annual movement




Figure 1: Sampling locations surrounding Rankin Inlet

used to identify plots dominated by three different vegetation types (shrub, graminoid and heath) representing the habitat that caribou were most likely to use. Similarly, IQ was used to exclude locations least likely to be used (i.e., those containing more than 30% mires, areas which were prone to flooding, or those comprised primarily of rock).




Figure 2: Sampling locations within the core calving (red), and post-calving (yellow) areas. The map on the right shows the utlization distributions used to inform the study design.

2 Methods

Fieldwork in 2019 was conducted from May 15 to August 25 in Rankin inlet, and August 7 to August 13 in the calving grounds. The following section describes field methods applied across all years combined (i.e., not all methods applied in all years), for both study areas.

General Study Design

2.1 Rankin Inlet




Figure 3: Diagram of the sampling schedule.

We have structured our study design to incorporate monitoring of multiple ecosystem components across trophic levels (primary producers to tertiary consumers). Monitoring of vegetation, including future implementation of a warming experiment, is central to the overall study design. To monitor plant species composition and structure, five stations each comprising four replicates (two control and two treatment) were selected based on distribution and proximity of three vegetation types. For each replicate, three 1mX1m point-framing-plots (PFPs) have been permanently established within uniform patches of three vegetation types (shrub, heath, and graminoid), resulting in a total of 60 PFPs (30 control and 30 treatment). To monitor plant phenology, we have established a 50 m transect to monitor three species, *Dryas integrifolia* (mountain avens), *Eriophorum vaginatum* (cottongrass), and *Salix arctica* (Arctic willow), respectively. Other ecosystem components (small mammals, arthropods, avian community, and raptors) will be monitored simultaneously in time and space, including abiotic variables associated with weather, snow-cover, freeze-up and thaw. For arthropods, one pitfall trap will be placed in each vegetation type (graminoid, shrub, and heath) at each control and treatment plot (4 plots total) at each PFP site (5 sites total), for a total of 60 pitfall traps. In addition, we shall document pollinator diversity using pan traps. Two small mammal trapping stations will be established each comprising effort no less than 720 trap-nights per trapping session.

2.1.1 Primary Producers

Vegetation Community Composition

Field Methods Arctic tundra ecologists involved in the International Tundra Experiment (ITEX; [30]) developed and published standard protocols for measurements, experimental design and statistical analyses in the ITEX Manual. ITEX researchers recommend the point quadrat method for assessing vegetation changes in tundra plant communities because it is quantitative and repeatable [30]. A total of 30 permanent point framing plot (PFP) locations were established in the Rankin Inlet area in 2019. The 30 plots are grouped into 5 sampling stations (6 plots in each station), and each station contains two heath-dominated plots, two grass-dominated plots, and two shrub-dominated plots. Each plot is composed of a 10 by 10 grid that results in 100 intersections at which the canopy height, canopy vegetation species, and ground vegetation species are all identified. These plots will serve as permanent, long-term monitoring plots to evaluate changes in community composition.

Analytical Methods - Repository Because sampling plots are nested temporally and spatially, we will use hierarchical models that include random intercepts for sample location and year to estimate change in canopy height (Gaussian distribution), trends in vegetation percent cover (Binomial distribution), and species richness (Poisson distribution).




Figure 4: Diagram of the sampling scheme.

Plant Phenology

Field Methods To monitor plant phenology, we have established a 50-m transect to monitor three species, *Dryas integrifolia* (mountain avens), *Eriophorum vaginatum* (cottongrass), and *Salix herbacea* (Alpine dwarf shrub), respectively. Data shall be collected every 2 to 3 days from snow melt (mid-June) to senescence (Mid-August) each year. The long-term objective is to monitor trends in species-specific phenological milestones and realized growing season length (time between leaf emergence and leaf senescence) in conjunction with snow melt, and spring and autumn temperatures. For *Salix herbacea*, we shall monitor: 1. date snow free; 2. date of first leaf bud burst; 3. date first pollen visible on male plants; 4. date of onset of seed dispersal in female plants 5. date first yellowing of leaves 6. date last green leaf turning yellow 7. date all leaves dead
For *Dryas integrifolia*, we shall monitor:

1. date snow free; 2. date first appearance of white color on flower buds; 3. date first open flower; 4. date last petal shed, and; 5. date first twisting of filaments on seed heads.

For *Eriophorum vaginatum*, we shall monitor:

1. date snow free; 2. date first appearance of flower buds, and; 3. date first pollen visible (yellow anthers).

Analytical Methods Phenological events will be estimated with a Bayesian hierarchical model using an interval censored response variable. Given the limitations associated with a survey frequency of every 2 to 3 days, the exact time at which a phenological event occurs may be unknown (the event occurred after the last survey, but before the current survey). Resulting uncertainty can bias model estimates depending on how one chooses to deal with the uncertainty [9]. With an interval censored response variable, lower and upper boundaries of time within which the event must have occured are created to reflect the uncertainty, and parameter estimates are corrected as a result. Following Bjorkman et al. (2015), we define the upper boundary as the first day in which the phenological event was detected, and the lower boundary as the day of the previous survey. Factors potentially affecting yearly variation in phenology such as (i) winter temperature, (ii) spring temperature,





Figure 5: Diagram of the sampling scheme.

(iii) previous summer temperature, and (iv) timing of snowmelt per plot, per year, will be included as fixed effects within the model, and as the sample locations within the transect will remain the same across years, plot will be added as a random intercept to account for individual-specific variation.

Plant Growth

Field Methods Annual growth increments in woody species can provide information on trends in growth related to local growing conditions, including long-term climate change. Sample processing shall follow methods adapted for tundra shrub species [56]. Thin cross sections of either the largest stem or the root collar shall be mounted on glass slides and photographed under a dissecting microscope. Using the digital image, individual rings shall be counted and measured for each section of shrub along four radii with a resolution of 1m [70].




Analytical Methods - Repository - As one sample from a single individual will be used to generate growth metrics among multiple years, we will include a random intercept for individual, and

year to estimate change in radial growth over time. Models will be built separately for each species of shrub, and will include within-year covariates such as winter, spring, and summer conditions obtained from archived records, as well as local topography, and NDVI. Spatial correlation structures will be added to the model to investigate potential spatial relationships in radial shrub growth within the study area.

2.1.2 Primary and Secondary Consumers

Microtine Rodents

Field Methods Following the procedures outlined by [41], we propose to establish two permanent small mammal trapping sites. Trapping sites have been selected based on their inclusion of habitats suitable for brown and collared lemmings (presence of old lemming nests, runways and burrows, seed bearing plants, wet and dry tundra, and a total area that is equal to or larger than 700 m in length). We will establish four permanent line transects (GPS-located) at each trapping site. Line transects shall be 300 m long and travel from xeric to hydric conditions. Each transect contains 20 trapping stations that consist of a flagged stake and three museum special snap traps attached to the stake using string (1 m in length), resulting in a total of 240 traps. Traps shall be distributed around the stake, with extra effort taken to target areas of suspected lemming use (lemming burrows or runs). Snap traps are placed no further than 1 m from the station flag, and baited with peanut butter. All traps shall be checked once daily for three trap-nights, resulting in 720 trap-nights per trapping session. We shall record all captures, misfires, or missing bait from each trap. Lemming abundance will be expressed as the total number of lemmings caught along all transects per 100 trap nights [24].




Arthropod Diversity

Field Methods To measure epigeic arthropod diversity, one pitfall trap is placed in each vegetation type (graminoid, shrub, and heath) at each control and treatment plot (4 plots total) at each PFP site (5 sites total), for a total of 60 pitfall traps. Traps are placed adjacent to the vegetation plot so that sampling efforts are specific to the vegetation type, but sufficiently far away to ensure that the vegetation plot isn't heavily trampled.

Traps are placed in a hole in the ground such that the pitfall trap fits snugly in, with its upper rim exactly level with the soil surface. Litter and vegetation surrounding the freshly dug hole is replaced as much as possible. Each trap consists of an outer plastic container, an inner plastic container, and

a rain shelter pegged into the ground at two corners. The inner pitfall trap container is filled 1/3 to 1/2 full with a mix of 50% non-toxic monopropylene glycol, 50% water.

Pitfall traps are first deployed in early July when the surface soil layer has fully thawed, and removed roughly one week into August. Trapped specimens are removed from the pitfall traps every two weeks, and transferred into 70% ethanol. Specimens are then returned to the lab in Edmonton for pinning, identification, and archival.



Analytical Methods - Repository - Arthropod sampling adopts the same hierarchical structure as plant species diversity, and models will therefore be similar - random intercepts for plot and year will be included to account for repeated observations. However, unlike plant species diversity, repeated observations with arthropods enables the estimation of detection error, so n-mixture models will be used to estimate species richness at each location while accounting for imperfect detection. We will estimate trends in species richness among years while including within-year covariates such as vegetation type, winter conditions, spring conditions, topography, and NDVI.

2.1.3 Avian Herbivores and Insectivores

Passive Sampling

Field Methods A permanent grid of locations has been established for passive avian sampling using autonomous recording units (ARUs). Each year, ARUs are deployed and powered on in May, and sensors remain active until early August. Due to the length of recording period, units are visited every 3 weeks for maintenance and battery replacement if needed. ARUs are programmed to replicate the recommended sampling duration for human conducted point counts [48], and record in two daily sampling periods, 1) 10 minute recordings every hour from 02:00 to 12:00, and 2) 10 minute recordings every hour from 17:00 to 23:59.

Analytical Methods A permanent grid of locations has been established for autonomous recording unit deployment. A total of 13 units were deployed in Rankin Inlet in 2019 in a range of areas dominated by srhub, heath, and graminoid vegetation. The units were deployed between July 3 and July 11, and were active until August 15. All units were programmed to record daily for 10 minutes every hour between 2:00AM and 12:00PM, and 10 minutes every hour between 5:00PM and

12:00AM, resulting in 17 recordings per day per unit. Transcription protocols have been developed and refined to reduce observer bias in data extraction (Appendix 9), and a database has been created to aid data entry and ensure errors are minimized. All recordings have been converted from WAC to WAV format to allow transcription, and recordings are currently being processed using the following two methods.

1. Traditional Transcription - Repository -
    - Using the protocols outlined in Appendix 9, an observer listens to the recording and identifies all species detected.
    - Data structure enables hierarchical modeling that simultaneously estimates detection probabilities, and probability of species occurrence.
    - As data processing is slow, a subset of recordings from each site are processed.
2. Acoustic Indices - Repository -
    - Using R, audio files are analyzed to investigate sound patterns typically associated with wildlife activity. Numerical outputs that indicate certain qualities of the recording are produced, and can be used to compare the level of wildlife activity across days in the summer.
    - This process drastically reduces processing time considerably, as it doesn't require an observer to listen to the recording. Index outputs can't fully replace transcription as they can't be used to determine species occurence, however, this method increases the amount of data various aspects of songbird phenology (acoustic complexity will increase as birds arrive in the spring).
    - all recordings obtained can be processed each year.

Active Sampling

Field Methods We used distance sampling transects [12], each 1 km in length, to estimate abundance and habitat associations of avian species and guilds. We generated 400 random start points at the start of each field season, and surveyed approximately two transects per day spanning the period from spring melt to the start of southward migration for many avian species. Based on work conducted in this study area (Hawkshaw, 2019, pers. comm.), we expect the sampling period to extend from the end of May to mid-August. Sampling is classified into two periods: early summer (prior to July 11), and late summer (all surveys conducted after July 11). This date corresponds to the division between incubation and brood rearing for most avian species in our study area (Hawksahw, 2019, pers. comm.), and thus was a logical cut-off when investigating how habitat associations and abundance might change seasonally. All transects are replicated before and after this date. To avoid the confounding effects of date and habitat in our counts, all replicates were a minimum of 15 days apart. Time of day of surveys ranged from 9:00 – 20:00. In the arctic, typical activity patterns in avian species are generally relaxed and they can remain active throughout the day [72].

Analytical Methods Density surface modelling [52] will be used to relate the abundance of species to spatial or habitat covariates to create distribution maps. To estimate detection functions, we shall follow the approach recommended by Thomas et al. [76]. Candidate models (i.e., detection functions) will include covariates that potentially influence detection (e.g., Julian day, time of day, precipitation and wind speed). The top detection function used for each guild shall be selected using Akaike's Information Criterion (AIC). Detection functions shall be estimated using the package Distance (Miller et al. 2017) in the R statistical environment (R Development Core Team 2010). To model spatiotemporal variation in abundance of avian guilds, we shall use an extension of distance

sampling known as density surface modelling (DSM). DSM incorporates the probability of detection estimated using the detection function and models adjusted counts on the basis of spatial or habitat covariates in a generalized additive model framework. As a final step, the resulting model shall be used to predict abundance over a grid of each permanent site. Analyses will completed using the R package dsm (Miller et al. 2017) and mgcv (Wood 2015).

2.1.4 Tertiary Consumers

Raptors

Field Methods As falcons arrive from their wintering grounds in May of each year, all historical and potential new breeding sites shall be surveyed by snowmobile to determine the presence or absence of territorial, breeding-aged adults. All known sites (i.e. cliff or rock outcrop used by a single pair of peregrines including known alternate nest sites within a breeding territory) shall be checked until occupied or until the breeding season is sufficiently advanced to conclude that the site was vacant. In cases where sites are occupied by individuals previously fitted with unique alphanumeric color-coded leg bands, bands shall be read using a Questar Field Model telescope, hand held digital camera, or Reconyx trail camera. If territories are occupied by unmarked individuals, effort shall be made to trap and mark these birds with an alphanumeric color band on one leg and a US Fish and Wildlife band on the other.

After breakup of the sea-ice, territories will be surveyed by boat to determine occupancy and productivity at each site; inland sites shall be accessed by all-terrain vehicle for the same purpose. During summer, banded adult birds shall be identified by either trapping or visually by telescope. Unmarked individuals shall be trapped and banded where possible. Once nestlings reached a minimum of 25 days of age (first week of August), they too shall be banded. Only young that survived beyond 25 days and were banded were included in the annual nestling census.

Analytical Methods Using the mgcv package in R, general additive models (GAM) are used to estimate annual trends for occupancy and productivity. Models include random intercepts at the year level, and smoothers are applied to account for potential non-linear patterns. Typical model validation procedures were performed to ensure that no model assumptions were violated (normal residuals), and the observed data was adequately described by the model.

2.1.5 Physical

Meteorological Conditions

Field Methods Up to four Rainwise PortLog Weather Stations will be deployed approximately uniformly within the study area, and will record hourly estimates of wind speed, wind direction, temperature, humidity, dew point, barometer, rainfall, and solar radiation. In addition, one rain gauge and one phenocam will be deployed at each PFP. A snow gauge shall be deployed within the field of view of the phenocam to monitor snow depth. Analysis of these data shall include annual summaries of precipitation, temperature, snow depth and cover for the purpose of direct long-term comparisons. However, we shall also use these measures as covariates in other analyses. For example, snow depth data could be included in occupancy models that may explain annual variation in caribou or other wildlife (e.g., song birds) phenology.

Analytical Methods - Repository - General linear models (Gaussian distribution) that incorporate site and year as random intercepts will be used to estimate trends in rainfall patterns, total rainfall amounts, temperature, snow-depth, and snow melt.

2.2 Qamanirjuaq Calving Grounds

As indicated, 16 permanent sampling sites have been established within the caribou calving grounds. The sampling locations have been intentionally placed along a gradient of historical caribou intensity of use, as determined by collar data collected between 1993 and 2012 during the calving period. Each of the 16 sampling locations contain three vegetation plots, one for shrub, graminoid, and heath dominated micro-environments (Figure 6). All micro-environment plots are anchored at a fixed location using permanent stakes and flagging, and all corners of the plot are identified with a unique ID (AA, BB, CC, and DD) so that all sampling within the plot can be reliably compared among years.




Figure 6: Diagram of the sampling scheme.

2.2.1 Vegetation monitoring

Vegetation sampling occurs at all plots, and can be broken down into two main categories: (i) visual estimation of percent cover, and, (ii) species presence/richness.

Percent Cover Percent vegetation cover can be estimated in the field by technicians, however, digital photography and image processing can be used as a replacement to decrease the amount of time spent in the field [15]. Each year, a total of eight digital photographs are taken of 50cm x 50cm quadrats placed in permanent locations around the perimeter of each micro-environment plot (Figure 6). A measuring tape is extended across one edge of the quadrat, and a digital camera is placed at a consistent height of 1.2 m above the ground using a tripod. All photos are taken in jpg format, and the image is composed so that the 50cm x 50cm quadrat fills as much of the frame as possible. Placing an erasable whiteboard with custom labels that includes plot ID, location ID (ex. AA), dominant vegetation, and date, within the image makes organizing the pictures after field work much easier (Figure 7).

Percent Cover - Repository -

Estimating yearly percent cover of the entire 15m x 15m quadrats is not feasible given the low sampling effort within the quadrat (n = 16 samples per year). Alternatively, we will estimate percent cover using the 8 digital photographs taken around the quadrat (Figure 6). This will be completed by overlaying a 10 x 10 grid on every image, and identifying specimens located at each of the 100 grid intersections within the image. Each photo will therefore be reduced to 100 data points from which percent cover can easily be estimated (for example, if 20/100 intersections contained graminoid species, percent cover of gram would equal 20%). Percent cover calculations for each im-




Figure 7: Example veg cover image




Figure 8: Photo file structure

age associated with a vegetation quadrat will be averaged to provide an estimate of percent cover for the entire 15m x 15m plot.

Species Presence/Richness In an effort to build vegetation species accumulation curves, a 50cm x 50cm quadrat was randomly thrown into each of the four quadrants of all vegetation plots each year (Figure 6). Plant species touching the corners of the quadrat after random tossing were identified to species, which resulted in a total of 16 plants identified to species per vegetation plot. When technicians were unable to ID plant species, samples were collected and sent to the Canadian Museum of Nature for consultation (see Appendix C for detailed protocol on specimen collection).

Species Presence/Richness Analysis - Repository -

As many samples were shipped to Jeff Saarela in 2019, we don't have all Species Presence/Richness data. Once data is received, Bayesian linear models can be used to estimate site-level changes in species richness/evenness over time.

All photos are named with the vegetation type, corner ID, sampling location ID, and year (ie. `Heath AA - R2Z7 2018.jpg`), and file organization uses the convention depicted in Figure 8.

2.2.2 Pellet Counts

The relationship between pellet counts and the number of individuals is well established [21]. To determine the annual pellet counts for a number of species, Four 15cm x 15cm pellet quadrats are located in the corner of each dominant vegetation plot (Figure 9), and all pellets contained within each

of the quadrats are counted, identified to species, and collected for potential lab work (e.g., genetic structure, stress hormones etc.). Pellet counts/collection are conducted annually, and data therefore reflect animal use throughout the entire year.

Pellet Count Analysis - Repository -

The goal of this analysis was to investigate pellet count trends, and compare variables potentially related to pellet counts across the 16 plots in the study area. Pellets were counted at 4 locations within each of the three habitat specific quadrats (graminoid, heath, and shrub) in the years 2012, 2013, and 2015. Because pellets were not removed prior to 2012, counts from that year included all historical scat deposits that had not degraded. After counts were conducted in 2012, all pellets were then removed from the plots so that 2013 data would represent one year of herbivore presence. The same protocol was used in 2013, however biologists did not return to the plots until 2015. Counts from 2015 therefore represent herbivore presence between the time when surveys were conducted in 2013, and the time when surveys were conducted in 2015 (two years).

Pellet counts were pooled within habitat-specific quadrats at every site within each year. There are 16 sites within the study area, each with a graminoid, heath, and shrub plot, so our analysis involved a total of 48 values for every year that data was collected (i.e., 2012-graminoid = 16, 2012-heath = 16, and 2012-shrub = 16). Although pellets for all species were recorded, the number of detections were only sufficient for analysis among caribou, ptarmigan, and small mammals. We modeled pellet count using a Poisson distribution with the covariates distance north (i.e., the plot located furthest to the south = 0 km), habitat (graminoid, heath, or shrub), year (2012, 2013, or 2015), and zone (Z1 through Z7). We predicted that caribou pellet counts would be higher in the core of the calving grounds, an area located part way along the south to north gradient. We therefore expected pellet counts to have a non-linear relationship with distance north where pellet abundance was reduced at the low and high values of the variable. To test for this prediction, we compared models containing this variable with a linear-predictor, against models containing this variable with smoothing splines [85]. All other covariates were categorical, and were included as dummy variables. A total of five models containing various combinations of the above covariates were estimated for caribou, ptarmigan, and small mammals separately using uninformative priors in a Bayesian framework with Stan, and the R package brms [13] Models were then compared using leave-one-out cross-validation (LOO), and the widely applicable information criterion (WAIC; [81]).

2.2.3 Trail Cameras, ARUs and Weather Monitoring

At each of the 16 sites, one motion-sensitive camera, one acoustic monitor, one rain gauge and one temperature logger is mounted on a tripod located within the vicinity of the three 15m x 15m quadrats. Motion-sensitive cameras are programmed to capture one photograph per hour, every day for the entire duration of the calving and post calving period (June-August), and three images when triggered by motion. Acoustic monitors are programmed to record for 10 min every hour from the time of installment (late April) to retrieval (late August). Historically, a rain gauge and temperature logger are mounted on a tripod to record local conditions. NuWCRU is in the process of securing parts to construct environmental monitoring units (EMUs) composed of temperature/humidity, soil moisture, and light sensors [51].

Trail Camera Analysis - Repository -

To get a better sense of migratory timing and general patterns of occupancy across the south to north gradient, all available images were looked at, and summarized by plotting the maximum number of individuals detected on a daily basis from May 1 and August 25. Although motion sensitive cameras are common and useful tools for assessing population abundance and movement, a higher intensity of sampling design is necessary to distinguish spatial and temporal patterns in abundance. As such, cameras were not deployed in the calving grounds in 2019.




Figure 9: Permanent pellet sampling quadrats

Autonomous Recording Units To efficiently and consistently process data obtained with ARUs, we've developed protocol (Appendix 9) and have built a database. We have converted all data from WAV to WAC format to allow transcription, and are currently processing historical recordings. Here, we summarize the tally of recordings that were obtained in the summers of 2015 and 2016 (no ARUs were deployed in subsequent years). To increase the density of sampling within a given area, ARUs were deployed in the Rankin Inlet area in 2019.

Weather Monitoring Many ecological changes have been reported in the north as temperatures increase [54, 55, 63], however, site-specific differences between studies have resulted in substantial variation [22]. Sources of such variation include site-specific characteristics such as topography, permafrost presence, moisture regimes, and local environmental variables [22]. Site-specific environmental data can therefore enhance modeling beyond random grouping variables, and may provide additional explanations for observed site-level differences.

Environmental monitors that record soil moisture, temperature/humidity, and photo information are currently being constructed – no results are available to present here.

3 Results

3.1 Rankin Inlet

3.1.1 Primary Producers

Community Composition A total of 30 permanent point framing plot (PFP) locations were established in the Rankin Inlet area in 2019. The 30 plots were grouped into 5 sampling stations (6 plots in each station), and each station contained two heath dominated plots, two grass dominated plots, and two shrub dominated plots. Each plot is composed of a 10 by 10 grid that results in 100 intersections at which the canopy height, canopy vegetation species, and ground vegetation species are all identified.

In 2019, 3 quadrats (1 heath, 1 gram, 1 shrub) were sampled from each station for a total of 1500 vegetation IDs. Species occurence for each plot have been visualized in Figure 10 and Tables 3 to 6, and species richness analysis will be added to the following report on March 31, 2020.




Figure 10: All plant species detected within each vegetation type.

Graminoid Plot

spp	*n*
detritus	223
bare ground	64
Carex spp.	33
Dryas integrefolia	32
Moss spp.	28
Flavocetraria cucullata	20
Carex aquatalis	17
Grass spp.	17
spp unk.	13
Vaccinium uliginosum	12
Andromeda polifolia	11
Rhododendron lapponicum	7
Betula glandulosa	5
Polygonum viviparum	3
Cassiope tetragona	2
Salix herbacea	2
Vaccinium vitis-idaea	2
Allectoria ochroleuca	1
Cetraria ericitorum	1
Equisetum arvense	1
Equisetum variegatum	1
Lichen spp.	1
Lycopodium annotinum	1
Pedicularis spp.	1
Rock	1
Salix spp.	1

Table 3: Tally of detections for all graminoid plots

Heath Plot

spp	*n*
Bryoria nitidula	93
detritus	80
Flavocetraria cucullata	65
Dryas integrefolia	47
Vaccinium uliginosum	43
Grass spp.	22
Cassiope tetragona	20
bare ground	18
Rhododendron tomentosum	18
Allectoria ochroleuca	14
Moss spp.	13
Empetrum nigrum	10
Arctostaphylas spp.	9
Cetraria ericitorum	9
Vaccinium vitis-idaea	6
Rhododendron lapponicum	5
Thamnolia subuliformis sp.	5
Dactylina arctica	4
Andromeda polifolia	3
Masonhalea richardsonii	3
Rock	3
Carex spp.	2
Oxytropis spp.	2
Salix reticulata	2
Lycopodium annotinum	1
Oxytropis maydelliana	1
Pedicularis spp.	1
Salix herbacea	1

Table 4: Tally of detections for all heath plots

Shrub Canopy

spp	*n*
Betula glandulosa	277
no vegetation	218
Salix pulchra	5

Table 5: Tally of detections at shrub plots, exclusively at the canopy layer.

Shrub Ground

spp	*n*
detritus	198
Cassiope tetragona	56
Lycopodium annotinum	35
Vaccinium vitis-idaea	28
Vaccinium uliginosum	26
Dryas integrefolia	25
Moss spp.	23
Arctostaphylas spp.	20
Flavocetraria cucullata	19
Empetrum nigrum	16
Rhododendron lapponicum	16
Grass spp.	14
Masonhalea richardsonii	6
Oxytropis maydelliana	6
Allectoria ochroleuca	5
Cetraria ericitorum	4
Fungus unk	1
Rhododendron tomentosum	1
Salix reticulata	1

Table 6: Tally of detections at shrub plots, exclusively at the ground layer.

Plant Phenology Data collection will start in 2020.

Plant Growth A total of 7 shrub root collars were collected for analysis in 2019 in the Rankin Inlet area. This included 6 Salix samples, and one Betula glandulosa sample. The samples will be cross sectioned and sanded to maximize growth ring contrast, and yearly radial growth will be quantified from growth ring measurements. Result will be submitted March 31, 2020.

3.1.2 Primary and Secondary Consumers

Microtine Rodents Historical data going back to 2008 is available for one of the small mammal trapping locations used in 2019 (Figure 11). The second trapping location was established in the summer of 2019 to determine how microtine rodent abundance varies within the study area. Snap trapping was conducted from July 23 to 26 at the new location (grid number 2), and conducted from July 30 to August 2 at the historical location (grid number 1). A total of 8 collared lemmings were detected, with 5 detections at grid 1, and 3 detections at grid 2. Figure 11 combines the results from both grids in 2019, and standardizes them into the number of detected small mammals per 100 trap nights to enable comparison with historical records. Small mammal abundance was highest in the first year of sampling (2008), and despite a similar peak in 2013, has largely decreased over the 11 year period.




Figure 11: Lemming detections per 100 trap nights across years. Trapping efforts varied by year, and total trap nights are visualized in red.

Arthropod Diversity In 2019, 30 pitfall traps were deployed from July 17 to August 17. Traps were recharged once during this period, and resulted in a total of 60 collections and 17,377.9 trap hours. Pitfall samples are currently being sorted into the following categories: 1. carabidae, 2. spiders, and 3. non-carabidae beetles.

Specimen sorting is almost complete, and current detections per site are summarized in Table 7. So far, a total number of 108 carabidae, 189 spiders, and 12 beetles not in the carabidae family have been sorted. Under spervision of NuWCRU staff, three undergraduates at the University of Alberta showing interest in invertebrate ecology have been recruited to identify specimens to species, and undertake correct sample archival.

PFP1			
Veg Type	**Carabidae**	**Spiders**	**Other Beetle**
gram	6	18	0
heath	3	NA	2
shrub	31	44	2
total	*40*	*62*	*4*
PFP2			
Veg Type	**Carabidae**	**Spiders**	**Other Beetle**
gram	6	25	4
heath	6	7	0
shrub	3	10	0
total	*15*	*42*	*4*
PFP3			
Veg Type	**Carabidae**	**Spiders**	**Other Beetle**
gram	15	13	0
heath	9	24	0
shrub	14	14	0
total	*38*	*51*	*0*
PFP4			
Veg Type	**Carabidae**	**Spiders**	**Other Beetle**
gram	6	8	0
heath	1	0	4
shrub	11	5	0
total	*18*	*13*	*4*
PFP5			
Veg Type	**Carabidae**	**Spiders**	**Other Beetle**
gram	1	4	0
heath	9	9	0
shrub	10	13	0
total	*20*	*26*	*0*

Table 7: Current tally of detected invertebrates at all locations. Specimen sorting is not yet complete (approximately 10% of samples left to process) so these numbers will increase.

3.1.3 Avian Herbivores and Insectivores

Passive Sampling All recordings have been converted to WAV format for processing, and 5 recordings have been transcribed from each ARU station. A total of 28 species were detected across all recordings, with the tallies shown in Table 8.

spp	*n*	*spp*	*n*
TUSW	38	REDP	4
SAVS	35	UNPA	4
HEGU	28	UNSH	4
AMPI	25	COLO	3
REDP	22	RLHA	3
CANG	21	LESA	2
HOLA	20	RTLO	2
SACR	16	SESA	2
LTDU	15	CANG	2
LALO	12	ARGS	1
PALO	9	CORA	1
SEPL	7	UNDU	1
UNLO	6	UNGO	1
FOX	4	UNKN	1

Table 8: All species detected within the first round of transcritptions (n = 48 recordings) for 2019 in Rankin Inlet.

3.1.4 Tertiary Consumers

Raptors Model results (Figure 12, and Table 9) indicate that occupancy has remained stable, throughout the monitoring period. Productivity appears to have declined over the same period. Potential mechanisms for these patterns have been investigated in standalone studies ([46], [23], [3], [44]). After fitting each GAM, we validated model fit using histograms and kernel density plots to inspect the residuals for normality. We assessed homogeneity of variance by plotting residual values against fitted values, and inspected each plot to ensure that points were uniformly distributed. Model outputs for each GAM included estimates of the grand mean for each attribute of interest for each monitoring site (i.e., the intercept term).

	Coefficient		Smoothing Terms				
model	*Intercept*	*SE*	*edf*	*p-value*	*D (%)*	*GCV*	*n*
occupancy ~ s(year)	0.74	0.010	1.00	0.16	5.7	0.004	36
productivity ~ s(year)	1.07	0.085	1.23	0.043	19.3	0.28	36

Table 9: Model structure and smoothing terms for peregrine falcon occupancy and productivity at Rankin Inlet, NU. The intercept estimates the grand mean, we do not report a p-value as the intercept is expected to be significantly different from zero *a priori*. Effective degrees of freedom (edf) were estimated by generalized cross-validation, where higher values indicate less smoothing and lower values indicate more smoothing; significant p-values indicate improved model fit (%D, deviance explained) compared to a linear model. Non-significant p-values indicate highly variable data.




Figure 12: Peregrine Falcon occupancy and productivity since 1981.

3.1.5 Physical

Temperature Historical climate data from the Rankin Inlet weather station dates back to 1981. To visualize and compare annual trends, we've summarized daily temperatures for two decades on the extreme ends of the available data, 1981-1991, and 2009-2019. For both decades, we plotted the minimum, maximum, and mean daily temperature for each day of the year. We then subtracted the difference in mean daily temperature between the two decades for each day, and calculted the monthly difference in mean temperature. The results are presented in Figure 13, which shows an overall increase in mean monthly temperature by 1.7°C (0.95), a maximum increase in the month of December of 3.94°C, and a minimum increase in April of 0.510°C.




Figure 13: A comparison of daily minimum, maximum, and mean temperatures from 1981-1991, and 2009-2019.

Snow on the Ground To visualize trends in ground snow accumulation over the past 40 years, we visually compared the minimum, maximum, and mean ground snow measurements for the years 1981-1991, and 2009-2019 (14). The resulting figure reveals substantial differences in the mean amount of snow on the ground from December to May, a trend observed by others [9]. Although snow depths were greater from 2009-2019, first snow free dates do not differ.




Figure 14: A comparison of daily minimum, maximum, and mean depths of ground snow from 1981-1991, and 2009-2019.

Summer Precipitation To investigate trends in rainfall since 1981, we tallied total monthly precipitation from May through August for all years, as well as the average amount of precipitation per event. The majority of monthly precipitation occurs in July and August, but there is substantial variation between years (Figure 15). Overlaying a simple general linear model indicates little to no trend in the data. Similarly, there doesn't appear to be any pattern in the amount of precipitation falling per event (Figure 16).




Figure 15: Total precipitation in the summer months from 1981 to 2019.




Figure 16: The mean (± SD) and max rainfall amounts within the summer months for each year from 1981 to 2019.

3.2 Qamanirjuaq Calving Grounds

3.2.1 Vegetation Monitoring

Visual comparison of vegetation quadrats images (Figure 17) show clearly that plant matter in 2012 was much drier compared to 2018. In 2102, quadrat photographs were recorded between September 18th and 23rd, and in 2018 photographs were recorded approximately one month earlier (between


August 26th and 29th).

The goal of this analysis was to provide a preliminary comparative visual assessment of the vegetation quadrats over times for each habitat type at each sampling station. Using the library of images recorded in 2012, and 2018, we randomly selected one 1x1 meter quadrat in each of the three permanent 15m x 15m quadrat (heath, shrub, graminoid), per site, and constructed a panel of 96 images (2 years by 16 sites, by 3 types of vegetation communities). We have started to translate methods from Chen et al. 2010 [15] into R for analysis.


3.2.2 Pellet Counts

Here we report on predicted values for pellets counts averaged across years (mean ± 1 standard deviation). Pooled pellet count trends for caribou, small mammals and ptarmigan (Figure 18a) increased from south to north for both caribou and ptarmigan, while pooled counts for small mammals decreased from south to north. For caribou the highest predicted counts were recorded at the northernmost station (R3Z1) averaged to 9.9±7.0, 7.0±5.0, and 4.0±2.8, for heath, shrub, and graminoid quadrats respectively. Similarly, the highest pellet counts for ptarmigan were recorded at the northernmost station and averaged 3.9±3.4, 3.6±3.1, and 1.2±1.04, for heath, shrub, and graminoid quadrats respectively. Pellet counts for small mammal were highest in the southernmost station and averaged 13.2±9.4, 10.0±7.1, and 4.2±3.0, for shrub, graminoid and heath quadrats respectively. The lowest counts for caribou summed to 2.7±2.93, 1.73±1.9, and 1.0±1.1 for heath, shrub, and graminoid quadrats respectively. Although the lowest counts for small mammals occurred within R3Z6, and were slightly higher in the northernmost zone (R3Z1), confidence intervals in counts for each zone overlapped, suggesting no difference. Small mammal pellt counts in R3Z6 summed to 0.4±0.7, 0.3±0.6, and 0.1±0.2, for shrub, graminoid and heath quadrats respectively. For caribou the lowest counts recorded summed to 4.0±2.8, 2.8±2.0, and 1.6±1.2, for heath, shrub, and graminoid quadrats respectively, and lowest counts for ptarmigan summed to 1.3±1.1, 1.1±1.0, and 0.4±0.3, for heath, shrub, and graminoid, respectively. Smoothed trends of annual pellet counts for caribou, small mammals and ptarmigan (Figure 18b, Figure 18c, and Figure 18d) show the lowest counts in 2013 (i.e., the year following removal). The smoothed trends for all species in all years followed the same trend for pooled years; caribou and ptarmigan increased from south to north, and small mammals decreased from south to north. Pellet counts were highest in 2012 for caribou and for ptarmigan, but for small mammals, counts were highest in 2015.


Total predicted counts for species by year (Figure 19) indicate, unsurprisingly, that for all species counts were lowest in 2013, following removal in 2012. For caribou and ptarmigan, highest counts were observed in 2012 (the first year of the program). Predicted counts for small mammal pellets were highest in 2015 despite removal in 2012 and 2013.




(a) Pellet counts pooled across all years



(b) Caribou pellet counts



(c) Small mammal pellet counts



(d) Ptarmigan pellet counts

Figure 18: A summary of pellet counts among three veg dominated plots (heath, graminoid, and shrub), and three years




Figure 19: pellet count pooled within each year for all species

3.2.3 Motion Sensitive Cameras

A total of 31 cameras were installed over two years; 15 in 2015, and 16 in 2016 (Figure 21). In 2015, cameras were deployed from May 14 to May 16, and were retrieved at the end of August. A total of 16,893 images (excluding images captured after camera had fallen over) were captured, with an average of 1,126 images per site. The number of days over which cameras captured quality images in 2015 averaged 58 per site, and ranged from 6 (R3Z5), to 101 (R3Z7). A total of five cameras fell, or were pushed over prior to retrieval resulting in a reduced duration of operation. In 2016, 31,130 images were captured averaging 1,945 per site. The average number of days with useable images in 2016 was greater than 2015 with an average of 72 per site, and ranged from 9 days at site R1Z6 to 121 days at R4Z7, and R1Z4. A total of 6 cameras were disturbed, or pushed over by wildlife (muskox, caribou, and grizzly), and an additional three fell from unknown causes.

The timing of caribou migration, as determined by the first yearly Caribou detection in any year, ranged from May 9 at the site furthest to the south (R3Z1), to July 3 at the site furthest to the north (Figure 20).




Figure 20: A visual of daily caribou detections from remote cameras between May 1 and August 25




Figure 21: Motion sensitive cameras functioning in 2015 (blue) and 2016 (green)

3.2.4 ARUs

A total of 18,196 ten-minute recordings spanning approximately 12 weeks in 2015, and approximately 10 weeks in 2016 are available. ARUs were installed at 12 sites in 2015 with recordings beginning in late May, and continuing through August (variation in monitoring period among sites, likely due to battery power), and 8 sites in 2016 with recordings beginning in late April, and continuing through early July (Figure 22). The number of ten-minute recordings available for each day of coverage ranges from 5 to 15. The first and last recording times within each day vary, however mainly range from 12:00 AM to mid-afternoon. Files are currently being transcribed.




Figure 22: Autonomous recording units successfully recording in 2015 (blue) and 2016 (green).




Figure 17: To compare changes in vegetation between 2012 and 2018, photos of the same location within respective plot are displayed beside one another.

4 Discussion

To date, large amounts of data have been collected and archived. This report outlines efforts taken to summarize and analyze data that have been collected, describes potential problems associated with the data, compiles information relevant to ongoing work, describes the deployment of an additional field program around Rankin Inlet (for efficacy related to helicopter access), provides preliminary analysis and data summaries, and offers recommendations to ensure that ecosystem-level monitoring in the Qamanirjuaq caribou summer range continues with best practices in mind.

In the 2019 field season, many components of a new field program were rolled out near the community of Rankin Inlet. Following the general L-TEMP approach to monitoring ecosystem change across trophic levels(primary producers, and primary to tertiary consumers). Permament sampling locations were established to ensure consistent sampling across years for all trophic levels, and new equipment and sampling methods were trialed. Work completed in 2019 includes:

- Primary Producers
  - vegetation community dynamics
  - dendro ecology, and secondary growth patterns
- Primary and Secondary Consumers
  - microtine rodent sampling
  - invertebrate sampling
  - avian herbivore and insectivore sampling
- Tertiary Consumers
  - raptor monitoring
- Physical
  - rain loggers
  - work with historical data

NuWCRU continues to investigate additional data collection methods to employ in the upcoming field season, and we anticipate adding 1. vegetation phenology transects for *Dryas integrifolia* (mountain avens), *Eriophorum vaginatum* (cottongrass), and *Salix herbacea* (Alpine dwarf shrub), 2. pan traps for sampling pollinating insects, and 3. extensive work with available earth observation data.

Work continued in the Qamanirjuaq calving grounds following methods established in 2012. Concern over sampling design resulted in moving environmental sensors (ARUs and remote cameras) to Rankin Inlet, but vegetation (random sampling, images, and dendro ecology), and pellet sampling were completed. Analysis of historical pellet count data indicated that counts of caribou and ptarmigan pellets increased from north to south, while counts of small mammal pellets decrease from south to north. The general pattern of increased caribou pellet counts from south to north is consistent with observations that in recent years calving has tended to occur within the northeastern extents of the calving grounds, and with increased post-calving abundance in the same region. Our analysis indicated that caribou pellets were highest in heath habitat, which may be associated with selection of elevated heath communities that are relatively snow-free compared to shrub and graminoid-dominated vegetation. Alternately, heath communities tend to be more xeric which may contribute to lower break-down rates compared to relatively mesic vegetation. However, inconsistent deployment of a field program obscures these results. Only one year (2013) of seven characterized pellet counts for a calendar year, and although the general trend for caribou remains consistent with counts for multiple years, counts are very low. Single year accumulations of ptarmigan and small mammal pellets were essentially zero. We suggest that the current approach of recording the

number of individual pellets up to a maximum of four, and recording all other counts (i.e., five or more) within a single category (i.e., >4), may have obscured true pattern. It is also possible that the quadrat size over-which pellet counts are currently tallied is too small to capture among-year variation of pellet counts given the current herd size. Microtine rodent abundance is known to be cyclical [65], and it appears that microtine rodent abundance was high in 2014 and/or 2015; unfortunately, we cannot determine which of the three possibilities occurs because pellet removal did not occur in 2014.

Although further analysis of data collected to date is possible, inconsistency in data collection has reduced power, and many challenges are associated with the data. As such, NuWCRU prioritized the development of analytical methods, databases, protocol, analytical transparency and version control, best practices for data management, so that incoming data is organized and well-documented. The infrastructure for transparency and data management has now been established, and NuWCRU continues to work through data incoming from the 2019 field season, as well as archived data from historical work in the calving grounds (2012-2018).

Acknowledgements

A lot of hard work and investment from many people has made the collection of data to date possible. We would like to thank Page Burt for her extreme hospitality, expertise, and patience in helping with unknown plant specimens. We also thank Mitch Campbell and Keenan Lindell for their guidance, and help with data collection throughout the program deployment.

References

[1] Long-term Climate Change: Projections, Commitments and Irreversibility Pages 1029 to 1076. In Intergovernmental Panel on Climate Change, editor, *Climate Change 2013 - The Physical Science Basis*, pages 1029–1136. Cambridge University Press, Cambridge, 2014.

[2] S. Altizer, R. S. Ostfeld, P. T. J. Johnson, S. Kutz, and C. D. Harvell. Climate Change and Infectious Diseases: From Evidence to a Predictive Framework. *Science*, 341(6145):514–519, August 2013.

[3] Alexandre Anctil, Alastair Franke, and Joël Bêty. Heavy rainfall increases nestling mortality of an arctic top predator: Experimental evidence and long-term trend in peregrine falcons. *Oecologia*, 174(3):1033–1043, March 2014.

[4] A. Andreev. The ten year cycle of the willow grouse of Lower Kolyma. *Oecologia*, 76(2):261–267, July 1988.

[5] G R Beard, W A Scott, and J K Adamson. The Value of Consistent Methodology in Long-term Environmental Monitoring. *Environmental Monitoring and Assessment*, 54(3):239–258, 1999.

[6] D Berteaux, Mm Humphries, Cj Krebs, M Lima, Ag McAdam, N Pettorelli, D Réale, T Saitoh, E Tkadlec, Rb Weladji, and Nc Stenseth. Constraints to projecting the effects of climate change on mammals. *Climate Research*, 32:151–158, 2006.

[7] Dominique Berteaux and Nils Chr. Stenseth. Measuring, understanding and projecting the effects of large-scale climatic variability on mammals. *Climate Research*, 32(2):95–97, 2006.

[8] Frédéric Bilodeau, Gilles Gauthier, and Dominique Berteaux. The effect of snow cover on lemming population cycles in the Canadian High Arctic. *Oecologia*, 172(4):1007–1016, August 2013.

[9] Anne D. Bjorkman, Sarah C. Elmendorf, Alison L. Beamish, Mark Vellend, and Gregory H. R. Henry. Contrasting effects of warming and increased snowfall on Arctic tundra plant phenology over the past two decades. *Global Change Biology*, 21(12):4651–4661, 2015.

[10] John Boulanger, Mitch Campbell, and David S Lee. Estimating Abundance and Trend of the Qamanirjuaq Mainland Migratory Barren-Ground Caribou Subpopulation - June 2017. page 99, 2018.

[11] ST Buckland, DR Anderson, KP Burnham, JL Laake, DL Borchers, and L Thomas. *Introduction to Distance Sampling Estimating Abundance of Biological Populations.* Oxford University Press., 2001.

[12] Stephen T. Buckland, E. A. Rexstad, Tiago A. Marques, and C. S. Oedekoven. *Distance Sampling: Methods and Applications*. Methods in Statistical Ecology. Springer International Publishing, 2015.

[13] Paul-Christian Bürkner. Advanced Bayesian Multilevel Modeling with the R Package brms. *The R Journal*, 10(1):395, 2018.

[14] Marie-Christine Cadieux, Dominique Fauteux, and Gilles Gauthier. Technical manual for sampling small mammals in the Arctic. page 60.

[15] Zhaohua Chen, Wenjun Chen, Sylvain G. Leblanc, and Gregory H.R. Henry. Digital Photograph Analysis for Measuring Percent Plant Cover in the Arctic. *ARCTIC*, 63(3), September 2010.

[16] Lee W. Cohnstaedt, Kateryn Rochon, Adrian J. Duehl, John F. Anderson, Roberto Barrera, Nan-Yao Su, Alec C. Gerry, Peter J. Obenauer, James F. Campbell, Tim J. Lysyk, and Sandra A. Allan. Arthropod Surveillance Programs: Basic Components, Strategies, and Analysis. *Annals of the Entomological Society of America*, 105(2):135–149, March 2012.

[17] Intergroup Consultants. Economic valuation and socio-cultural perspectives of the estimated harvest of the Beverly and Qamanirjuaq caribou herds. Technical report, Winnipeg, MB, 2008.

[18] COSEWIC. Caribou (Rangifer tarandus) barren-ground population COSEWIC assessment and status report 2016. https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry/cosewic-assessments-status-reports/caribou-barren-ground-population-2016/chapter-1.html, August 2017.

[19] National Research Council, Division on Earth and Life Studies, Polar Research Board, , Board on Environmental Studies Toxicology, and Committee on Cumulative Environmental Effects of Oil and Gas Activities on Alaska's North Slope. *Cumulative Environmental Effects of Oil and Gas Activities on Alaska's North Slope*. National Academies Press, October 2003.

[20] Sasha R. X. Dall, Alasdair I. Houston, and John M. McNamara. The behavioural ecology of personality: Consistent individual differences from an adaptive perspective. *Ecology Letters*, 7(8):734–739, 2004.

[21] L. S. Eggert, J. A. Eggert, and D. S. Woodruff. Estimating population sizes for elusive animals: The forest elephants of Kakum National Park, Ghana. *Molecular Ecology*, 12(6):1389–1402, June 2003.

[22] Sarah C. Elmendorf, Gregory H. R. Henry, Robert D. Hollister, Robert G. Björk, Noémie Boulanger-Lapointe, Elisabeth J. Cooper, Johannes H. C. Cornelissen, Thomas A. Day, Ellen Dorrepaal, Tatiana G. Elumeeva, Mike Gill, William A. Gould, John Harte, David S. Hik, Annika Hofgaard, David R. Johnson, Jill F. Johnstone, Ingibjörg Svala Jónsdóttir, Janet C. Jorgenson, Kari Klanderud, Julia A. Klein, Saewan Koh, Gaku Kudo, Mark Lara, Esther Lévesque, Borgthor Magnússon, Jeremy L. May, Joel A. Mercado-Dıaz, Anders Michelsen, Ulf Molau, Isla H. Myers-Smith, Steven F. Oberbauer, Vladimir G. Onipchenko, Christian Rixen, Niels Martin Schmidt, Gaius R. Shaver, Marko J. Spasojevic, Þóra Ellen Þórhallsdóttir, Anne Tolvanen, Tiffany Troxler, Craig E. Tweedie, Sandra Villareal, Carl-Henrik Wahren, Xanthe Walker, Patrick J. Webber, Jeffrey M. Welker, and Sonja Wipf. Plot-scale evidence of tundra vegetation change and links to recent summer warming. *Nature Climate Change*, 2(6):453–457, June 2012.

[23] Alastair Franke, Jean-François Therrien, Sébastien Descamps, and Joël Bêty. Climatic conditions during outward migration affect apparent survival of an arctic top predator, the peregrine falcon Falco peregrinus. *Journal of Avian Biology*, 42(6):544–551, November 2011.

[24] Gilles Gauthier, Joël Bêty, Marie-Christine Cadieux, Pierre Legagneux, Madeleine Doiron, Clément Chevallier, Sandra Lai, Arnaud Tarroux, and Dominique Berteaux. Long-term monitoring at multiple trophic levels suggests heterogeneity in responses to climate change in the Canadian Arctic tundra. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1624), August 2013.

[25] Paola F. Giordano, Joaquín L. Navarro, and Mónica B. Martella. Building large-scale spatially explicit models to predict the distribution of suitable habitat patches for the Greater rhea (Rhea americana), a near-threatened species. *Biological Conservation*, 143(2):357–365, February 2010.

[26] P. J. Gregory, S. N. Johnson, A. C. Newton, and J. S. I. Ingram. Integrating pests and pathogens into the climate change/food security debate. *Journal of Experimental Botany*, 60(10):2827–2838, July 2009.

[27] C. D. Harvell. Climate Warming and Disease Risks for Terrestrial and Marine Biota. *Science*, 296(5576):2158–2162, June 2002.

[28] Erik Hedlin, Melanie Wilson, Keenan Lindell, Page Burt, Kevin Hawkshaw, Mitch Campbell, and Alastair Franke. Qamanirjuaq Caribou Calving Grounds Long-term Ecological Monitoring Program: 2018 Annual Report to Government of Nunavut. Technical report, Government of Nunavut, 2018.

[29] Thrine Moen Heggberget, Eldar Gaare, and John P. Ball. Reindeer (Rangifer tarandus) and climate change: Importance of winter forage. *Rangifer*, 22(1):13, June 2002.

[30] G.H.R. Henry and U. Molau. Tundra plants and climate change: The International Tundra Experiment (ITEX). *Global Change Biology*, 3(S1):1–9, December 1997.

[31] Alexandre H. Hirzel and Gwenaëlle Le Lay. Habitat suitability modelling and niche theory. *Journal of Applied Ecology*, 45(5):1372–1381, October 2008.

[32] E Hoberg and J Kutz. Chapters - Arctic biodiversity, Conservation of Arctic Flora and Fauna (CAFF). https://arcticbiodiversity.is/index.php/participants/13-the-report/chapters?start=15.

[33] Toke T. Høye, Eric Post, Hans Meltofte, Niels M. Schmidt, and Mads C. Forchhammer. Rapid advancement of spring in the High Arctic. *Current Biology*, 17(12):R449–R451, June 2007.

[34] Rolf A. Ims and Eva Fuglei. Trophic Interaction Cycles in Tundra Ecosystems and the Impact of Climate Change. *BioScience*, 55(4):311, 2005.

[35] Intergovernmental Panel on Climate Change and Ottmar Edenhofer, editors. *Climate Change 2014: Mitigation of Climate Change: Working Group III Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, New York, NY, 2014. OCLC: ocn892580682.

[36] Richard K. Johnson, Daniel Hering, Mike T. Furse, and Ralph T. Clarke. Detection of ecological change using multiple organism groups: Metrics and uncertainty. *Hydrobiologia*, 566(1):115–137, August 2006.

[37] Richard K. Johnson, Daniel Hering, Mike T. Furse, and Piet F. M. Verdonschot. Indicators of ecological change: Comparison of the early response of four organism groups to stress gradients. *Hydrobiologia*, 566(1):139–152, August 2006.

[38] T. G. F. Kittel, W. L. Steffen, and F. S. Chapin. Global and regional modelling of Arctic-boreal vegetation distribution and its sensitivity to altered forcing. *Global Change Biology*, 6(S1):1–18, December 2000.

[39] Kari Klanderud and H. J. B. Birks. Recent increases in species richness and shifts in altitudinal distributions of Norwegian mountain plants. *The Holocene*, 13(1):1–6, January 2003.

[40] Pekka Kontiainen, Hannu Pietiäinen, Kalle Huttunen, Patrik Karell, Heikki Kolunen, and Jon E. Brommer. Aggressive Ural owl mothers recruit more offspring. *Behavioral Ecology*, 20(4):789–796, July 2009.

[41] Charles Krebs, Don Reid, Alice Kenney, Doug Morris, and Scott Gilbert. Small Mammal Population Monitoring.

[42] Leif Kullman. Rapid BlackwellScienceLtd recent range-margin rise of tree and shrub species in the Swedish Scandes. *Journal of Ecology*, page 10, 2002.

[43] Susan J. Kutz, Sylvia Checkley, Guilherme G. Verocai, Mathieu Dumond, Eric P. Hoberg, Rod Peacock, Jessica P. Wu, Karin Orsel, Karin Seegers, Amy L. Warren, and Arthur Abrams. Invasion, establishment, and range expansion of two parasitic nematodes in the Canadian Arctic. *Global Change Biology*, pages n/a–n/a, September 2013.

[44] Vincent Lamarre, Pierre Legagneux, Alastair Franke, Nicolas Casajus, Douglas C. Currie, Dominique Berteaux, and Joël Bêty. Precipitation and ectoparasitism reduce reproductive success in an arctic-nesting top-predator. *Scientific Reports*, 8(1), December 2018.

[45] Hedwig Lankau, Alex MacPhail, Michelle Knaggs, and Erin Bayne. Acoustic Recording Analysis Protocol. Technical report, University of Alberta and Alberta Biodiverisyt Monitoring Institute, Edmonton, Alberta, 2015.

[46] Vincent L'Hérault, Alastair Franke, Nicolas Lecomte, Adam Alogut, and Joël Bêty. Landscape heterogeneity drives intra-population niche variation and reproduction in an arctic top predator. *Ecology and Evolution*, 3(9):2867–2879, September 2013.

[47] Tara G. Martin and Hugh P. Possingham. Predicting the impact of livestock grazing on birds using foraging height data: Bird foraging height predicts impact of grazing. *Journal of Applied Ecology*, 42(2):400–408, April 2005.

[48] Steven M. Matsuoka, C. Lisa Mahon, Colleen M. Handel, Péter Sólymos, Erin M. Bayne, Patricia C. Fontaine, and C. John Ralph. Reviving common standards in point-count surveys for broad inference across studies. *The Condor. 116(4): 599-608*, 116(4):599–608, 2014.

[49] Diethart Matthies, Ingo Bräuer, Wiebke Maibom, and Teja Tscharntke. Population size and the risk of local extinction: Empirical evidence from rare plants. *Oikos*, 105(3):481–488, June 2004.

[50] Sean M. McMahon, Sandy P. Harrison, W. Scott Armbruster, Patrick J. Bartlein, Colin M. Beale, Mary E. Edwards, Jens Kattge, Guy Midgley, Xavier Morin, and I. Colin Prentice. Improving assessment and modelling of climate change impacts on global terrestrial biodiversity. *Trends in Ecology & Evolution*, 26(5):249–259, May 2011.

[51] James G. Mickley, Timothy E. Moore, Carl D. Schlichting, Amber DeRobertis, Emilia N. Pfisterer, and Robert Bagchi. Measuring microenvironments for global change: DIY environmental microcontroller units (EMUs). *Methods in Ecology and Evolution*, December 2018.

[52] David L. Miller, M. Louise Burt, Eric A. Rexstad, and Len Thomas. Spatial models for distance sampling data: Recent developments and future directions. *Methods in Ecology and Evolution*, 4(11):1001–1010, November 2013.

[53] Claude Morneau and Serge Payette. Long-term fluctuations of a caribou population revealed by tree-ring data. 78:7, 2000.

[54] Isla H. Myers-Smith, Sarah C. Elmendorf, Pieter S. A. Beck, Martin Wilmking, Martin Hallinger, Daan Blok, Ken D. Tape, Shelly A. Rayback, Marc Macias-Fauria, Bruce C. Forbes, James D. M. Speed, Noémie Boulanger-Lapointe, Christian Rixen, Esther Lévesque, Niels Martin Schmidt, Claudia Baittinger, Andrew J. Trant, Luise Hermanutz, Laura Siegwart Collier, Melissa A. Dawes, Trevor C. Lantz, Stef Weijers, Rasmus Halfdan Jørgensen, Agata Buchwal, Allan Buras, Adam T. Naito, Virve Ravolainen, Gabriela Schaepman-Strub, Julia A. Wheeler, Sonja Wipf, Kevin C. Guay, David S. Hik, and Mark Vellend. Climate sensitivity of shrub growth across the tundra biome. *Nature Climate Change*, 5(9):887–891, September 2015.

[55] Isla H. Myers-Smith, Meagan M. Grabowski, Haydn J. D. Thomas, Sandra Angers-Blondin, Gergana N. Daskalova, Anne D. Bjorkman, Andrew M. Cunliffe, Jakob J. Assmann, Joseph S. Boyle, Edward McLeod, Samuel McLeod, Ricky Joe, Paden Lennie, Deon Arey, Richard R. Gordon, and Cameron D. Eckert. Eighteen years of ecological monitoring reveals multiple lines of evidence for tundra vegetation change. *Ecological Monographs*, page e01351, March 2019.

[56] Isla H. Myers-Smith, Martin Hallinger, Daan Blok, Ute Sass-Klaassen, Shelly A. Rayback, Stef Weijers, Andrew J. Trant, Ken D. Tape, Adam T. Naito, Sonja Wipf, Christian Rixen, Melissa A.

Dawes, Julia A. Wheeler, Agata Buchwal, Claudia Baittinger, Marc Macias-Fauria, Bruce C. Forbes, Esther Lévesque, Noémie Boulanger-Lapointe, Ilka Beil, Virve Ravolainen, and Martin Wilmking. Methods for measuring arctic and alpine shrub growth: A review. *Earth-Science Reviews*, 140:1–13, January 2015.

[57] R. B. Myneni, C. D. Keeling, C. J. Tucker, G. Asrar, and R. R. Nemani. Increased plant growth in the northern high latitudes from 1981 to 1991. *Nature*, 386(6626):698–702, April 1997.

[58] John Andrew Stephen Nagy. *Use of Space by Caribou in Northern Canada*. PhD thesis, University of Alberta, Edmonton, Alberta, 2011.

[59] Ólafur K. Nielsen and Gunnlaugur Pétursson. Population fluctuations of gyrfalcon and rock ptarmigan: Analysis of export figures from Iceland. *Wildlife Biology*, 1(1):65–71, January 1995.

[60] Pantoula Nikolakaki. A GIS site-selection process for habitat creation: Estimating connectivity of habitat patches. *Landscape and Urban Planning*, 68(1):77–94, May 2004.

[61] J. Olofsson, S. Stark, and L. Oksanen. Reindeer influence on ecosystem processes in the tundra. *Oikos*, 105(2):386–396, May 2004.

[62] Johan Olofsson, Heidi Kitti, Pirjo Rautiainen, Sari Stark, and Lauri Oksanen. Effects of summer grazing by reindeer on composition of vegetation, productivity and nitrogen cycling. *Ecography*, 24(1):13–24, February 2001.

[63] Johan Olofsson, Lauri Oksanen, Terry Callaghan, Philip E. Hulme, Tarja Oksanen, and Otso Suominen. Herbivores inhibit climate-driven shrub expansion on the tundra: HERBIVORES INHIBIT SHRUB EXPANSION. *Global Change Biology*, 15(11):2681–2693, November 2009.

[64] Camille Parmesan, Terry L. Root, and Michael R. Willig. Impacts of Extreme Weather and Climate on Terrestrial Biota *. *Bulletin of the American Meteorological Society*, 81(3):443–450, March 2000.

[65] Frank A. Pitelka and George O. Batzli. Population cycles of lemmings near Barrow, Alaska: A historical review. *Acta Theriologica*, 52(3):323–336, September 2007.

[66] Radim Plhal, Jiří Kamler, and Miloslav Homolka. Faecal pellet group counting as a promising method of wild boar population density estimation. *Acta Theriologica*, 59(4):561–569, October 2014.

[67] E. Post, M. C. Forchhammer, M. S. Bret-Harte, T. V. Callaghan, T. R. Christensen, B. Elberling, A. D. Fox, O. Gilg, D. S. Hik, T. T. Hoye, R. A. Ims, E. Jeppesen, D. R. Klein, J. Madsen, A. D. McGuire, S. Rysgaard, D. E. Schindler, I. Stirling, M. P. Tamstorf, N. J.C. Tyler, R. van der Wal, J. Welker, P. A. Wookey, N. M. Schmidt, and P. Aastrup. Ecological Dynamics Across the Arctic Associated with Recent Climate Change. *Science*, 325(5946):1355–1358, September 2009.

[68] E. Post and C. Pedersen. Opposing plant community responses to warming with and without herbivores. *Proceedings of the National Academy of Sciences*, 105(34):12353–12358, August 2008.

[69] H. Scherm, R.W. Sutherst, R. Harrington, and J.S.I. Ingram. Global networking for assessment of impacts of global change on plant pests. *Environmental Pollution*, 108(3):333–341, June 2000.

[70] Caroline A. Schneider, Wayne S. Rasband, and Kevin W. Eliceiri. NIH Image to ImageJ: 25 years of image analysis. *Nature Methods*, 9(7):671–675, July 2012.

[71] Bo Soderstrom, Tomas Part, and Erik Linnarsson. Grazing Effects on Between-Year Variation of Farmland Bird Communities. *Ecological Applications*, 11(4):11, 2001.

[72] Silke S. Steiger, Mihai Valcu, Kamiel Spoelstra, Barbara Helm, Martin Wikelski, and Bart Kempenaers. When the sun never sets: Diverse activity rhythms under continuous daylight in free-living arctic-breeding birds. *Proceedings. Biological Sciences*, 280(1764):20131016, August 2013.

[73] Ron Store and Jukka Jokimäki. A GIS-based multi-scale approach to habitat suitability modeling. *Ecological Modelling*, 169(1):1–15, November 2003.

[74] Ron Store and Jyrki Kangas. Integrating spatial multi-criteria evaluation and expert knowledge for GIS-based habitat suitability modelling. *Landscape and Urban Planning*, 55(2):79–93, July 2001.

[75] Ken Tape, Matthew Sturm, and Charles Racine. The evidence for shrub expansion in Northern Alaska and the Pan-Arctic. *Global Change Biology*, 12(4):686–702, April 2006.

[76] Len Thomas, Stephen T. Buckland, Eric A. Rexstad, Jeff L. Laake, Samantha Strindberg, Sharon L. Hedley, Jon Rb Bishop, Tiago A. Marques, and Kenneth P. Burnham. Distance software: Design and analysis of distance sampling surveys for estimating population size. *The Journal of Applied Ecology*, 47(1):5–14, February 2010.

[77] H. Tømmervik, B. Johansen, J.Å. Riseth, S.R. Karlsen, B. Solberg, and K.A. Høgda. Above ground biomass changes in the mountain birch forests and mountain heaths of Finnmarksvidda, northern Norway, in the period 1957–2006. *Forest Ecology and Management*, 257(1):244–257, January 2009.

[78] H. Tømmervik, B. Johansen, I. Tombre, D. Thannheiser, K. A. Høgda, E. Gaare, and F. E. Wielgolaski. Vegetation Changes in the Nordic Mountain Birch Forest: The Influence of Grazing and Climate Change. *Arctic, Antarctic, and Alpine Research*, 36(3):323–332, August 2004.

[79] Caroline Van Hemert, John M Pearce, and Colleen M Handel. Wildlife health in a rapidly changing North: Focus on avian disease. *Frontiers in Ecology and the Environment*, 12(10):548–556, December 2014.

[80] R. van Klink, F. van der Plas, C. G. E. Toos van Noordwijk, M. F. WallisDeVries, and H. Olff. Effects of large herbivores on grassland arthropod diversity: Large herbivores and arthropods. *Biological Reviews*, 90(2):347–366, May 2015.

[81] Aki Vehtari, Andrew Gelman, and Jonah Gabry. Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing*, 27(5):1413–1432, September 2017.

[82] Gian-Reto Walther, Eric Post, Peter Convey, Annette Menzel, Camille Parmesan, Trevor J. C. Beebee, Jean-Marc Fromentin, Ove Hoegh-Guldberg, and Franz Bairlein. Ecological responses to recent climate change. *Nature*, 416(6879):389–395, March 2002.


[83] T. N. Wasserman, S. A. Cushman, A. S. Shirk, E. L. Landguth, and J. S. Littell. Simulating the effects of climate change on population connectivity of American marten (Martes americana) in the northern Rocky Mountains, USA. *Landscape Ecology*, 27(2):211–225, February 2012.


[84] Max Wolf and Franz J. Weissing. Animal personalities: Consequences for ecology and evolution. *Trends in Ecology & Evolution*, 27(8):452–461, August 2012.


[85] Simon N. Wood. Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models: Estimation of Semiparametric Generalized Linear Models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 73(1):3–36, January 2011.


[86] Jakob Zscheischler, Seth Westra, Bart J. J. M. van den Hurk, Sonia I. Seneviratne, Philip J. Ward, Andy Pitman, Amir AghaKouchak, David N. Bresch, Michael Leonard, Thomas Wahl, and Xuebin Zhang. Future climate risk from compound events. *Nature Climate Change*, 8(6):469–477, June 2018.


5 Qamanirjuaq - Yearly Range


























6 Data Management Plan

Data Collection

File formats, and methods of collection

- tabular data will be initially collected on paper (vegetation data sheets) or in digital delimited format (ie. weather data), followed by entry into databases specific to the data
- spatial data although all collected data has a spatial component, here we refer to earth observation data (satellite imagery) as "spatial". All such data will be pulled from existing databases, most commonly in delimited format
- images are generated in .jpg format
- audio is recorded in `WAV` format, and converted to `WAC` format for processing

Data security and storage during collection

- tabular data collected on paper will be backed up in the field using a camera. Tabular data in delimited format will be copied to additional drives and cloud storage
- spatial data all spatial data is pulled from existing databases, and can therefore be copied to additional drives and cloud storage immediately
- images will be copied to additional drives and cloud storage
- audio will be copied to additional drives and cloud storage

Data amount and long-term storage

- most of the mass of data will be in audio and image files. At our current rate of sampling, we expect to collect 0.5 TB each year, and will therefore need a total of 1.0 TB of hard rive space, and 0.5 TB of cloud storage for secure back up.

Data organization and version control

- Data will be stored in databases specific to each sub-project. Data will be queried and exported for specific analysis, and as such, stored with all necessary components of the analysis. The analytical components (figures, scripts, input data, results tables) will be stored within a "Project" folder using RStudio's Project functionality. Furthermore, all analytical projects will be linked to GitHub to implement version control, and promote efficient collaboration.

Documentation and Metadata

Documentation to ensure data is read and interpreted correctly in the future

- All analytical projects will be accompanied with a README file that outlines: 1) who collected the data, 2) when it was collected, 3) how was it collected, 4) relevant study documents, 5) data queries used to pull data from database, 6) naming conventions used within data and scripts.

Metadata standard and tools used to document and describe the data

- A template README file[b] will be included within the project folder of each analysis. The template can be altered, but will provide a good starting point that ensures all necessary information is included

[b] see Appendix 7 for the template README

Insurance that the documentation is created or captured consistently throughout your project

- The template README file[c] will be a primary tool to ensure consistency, and all scripts will be heavily commented to ensure sufficient communication of rationale and methodology.

Storage and backup

Data storage and back-up during the project

- All data is backed up on at least two hard drives that are stored in separate locations. Google Drive and Github are use for cloud storage.

Sensitive data security

- Hard drives are located in secure locations, only accessible to key staff. Data stored in the cloud is encrypted and password protected.

Preservation

Selecting which data to destroy, or preserve

- All data will be preserved due to the high potential of reuse, and incorporation in an historical context.

Data life after the end of the project

- All data will be transferred to the Government of Nunavut at the conclusion of this project

Preparation of data preservation and access

- tabular data will be entered into databases, but all originals will be preserved
- because spatial data is acquired to fulfill analytical needs, storage is typically within the analytical project folder. To increase ease of sharing, all spatial data will be copied into a folder specific to spatial data as well.
- images are preserved in their original format, as well as any converted formats (ie. IHS)
- audio will be preserved and shared in `WAC` format

Methods of sharing data

- Data sharing restrictions are imposed by the Government of Nunavut, and data access requests will be processed by them

End-user agreements included with data sharing

- as data sharing requests are processed by the Government of Nunavut, all end-user agreements are supplied by them as well

Responsibilities and Resources

Data management responsibilities

- the Principal Investigator and Research Associate will be responsible for data management

[c] see Appendix 7

Personnel changes and transitioning responsibilities

- Due to the thorough and consistent documentation accompanying all data and analysis, personnel changes will not be an issue

Party responsible for data sharing and preservation after the project has concluded

- All data will be transferred to the Government of Nunavut upon the completion of the program. The Government of Nunavut will also adopt all data management responsibility at this time

Resources required for data storage

- hard drives, and cloud storage subscriptions

7 README Template in Markdown

The following is the contents of the template `README.md` file contained within each project analysis folder. An exampl can be seen here.

```
# NuWCRU README Template

## Project Description
This is a basic template to be included with all NuWCRU
    github repositories. The purpose of this template is to
    standardize th  metadata generated for each project/
    analysis to implement reproducibility and transparency,
    and efficient knowledge sharin between  personnel. This
    template should be used as a mere starting point, and can
    /should be altered to suit the specific needs o each
    project.

For basic workflow techniques using R, Hadley Wickham's
    freely available [R for Data Science](https://r4ds.had.co
    .nz/index.html is  highly recommended. See [Chapter 8:
    Workflow](https://r4ds.had.co.nz/workflow-projects.html)
    for working with R Projects.


## General Project File Structure
To avoid project folders filled with random files, it helps
    to organize files into a common structure. This is just
    an example and  both the files and their descriptions
    will be unique to each project.

‘‘‘
| --- .gitignore <- Files that should be ignored by git.
|
| --- README.md <- The top-level README including general
    project descriptions
|
| --- data
| |-- tab_data.csv <- Converted, but pre-process tab data
| |-- processed.csv <- Data ready for modeling
| |-- raw <- Raw data pulled from source
| |-- temp <- Temporary files
|
| --- docs
| |--process_doc.md <- Displaying process and some results
|
| --- figures
| |-- range.png <- figure description
| |-- eda.png <- figure description
| |-- growth.png <- figure description
|
| --- scripts
|-- d_wrangle.R <- wrangling pre-processed tab data, output:
```

```
    processed.csv
|-- src.R <- cleaned and standardized covariates
|-- model.R <- model building/runs
|-- vis.R <- model results, tables, and visualizations
```

Data
* Description of the data here. All relevant information
    needed to understand the data, and subsequently the
    analysis should b documented here.
* This includes covariate abbreviations, specific data
    collection methodology, years data was collected, and
    possibly eve contacts in the case of questions.
*If data was pulled from a database, notes about the queries
    used should be documented here

Relevant Study Documents
If methods are adopted from others (articles or research
    groups), links to the original documents are placed here.

Authors
Contact information for the most recent personnel working on
    the project.

8 C3 Plant and Lichen Collecting Protocols

Questions: email Jeff Saarela (jsaarela@mus-nature.ca) or Paul Sokoloff (psokoloff@mus-nature.ca)

Botanists at the Canadian Museum of Nature are building a record of all the plants, lichens, mosses and liverworts that live i Canada's Arctic. Species occurrences are documented with specimens housed permanently in the national collections of the Canadia Museum of Nature. During the Canada C3 voyage, the science team will be collecting plants whenever possible to build pan-Canadian collection as a record of the voyage and to contribute to documentation of Canada's botanical biodiversity.

Things you'll need:

Collecting

- collecting satchel
- GPS
- camera
- digger
- pruning shears
- field notebook
- ziploc collecting bags
- paper collecting bags
- sharpies
- pencils
- field guide
- hammer and chisel

Pressing

- plant press ends (x2)
- plant press straps (x2)
- cardboard
- newsprint
- silica gel envelopes
- jewelers tags
- sharpies
- national herbarium specimen sheets
- cheesecloth

Collection

Site selection The main goal at each landing is to document, with collections, as much plant and lichen diversity as possible. Ideally you wil be able to collect at a few sites per landing, time permitting. When you arrive on land, you will need to locate an area in whic to collect. If there is a distinctive coastal community in the vicinity of the landing area, start there. This habitat will no usually extend too far beyond the wrack line.

Once you have collected examples of all plants, lichens and algae in the coastal community (including macroalgae in the intertida zone), move to another community type further inland.
Here are some guidelines to keep in mind:

- If there are multiple inland habitats in the general landing area, start with a site that appears to have the greates plant and lichen diversity.
- If there is an area that looks to have different or uncommon species compared with what you have collected at othe sites on the trip, and collecting time is limited, start there. We want to capture diversity at each site, but also across th Arctic – and species that are uncommon either locally or regionally are always of interest and collecting those species shoul be a priority.

- If the whole general area looks similar in terms of plant and habitat diversity, any spot will work.
- A single collection site can be fairly large if the plant community is fairly homogeneous (up to about 300 m radius fro the recorded GPS coordinates), but if you move far away from your original site, or enter a dramatically new habitat with different suite of species (e.g., a wetter area, a drier area, a south-facing slope) it is best to start a new site.

Data collection For each collection site, fill out a site record; below that fill out a species record for eac specimen collected in the field. Each record should be filled out as completely as possible while you are the field makin collections. You won't be able to remember all the details to fill this in later. The fields necessary are defined below. All thi data will appear on the herbarium labels.

For each *collection site*, fill out the following fields:

1. Site Number: This is a unique identification number for a collection site. Use the date, followed by th site visited in sequential order: i.e. the first site visited on July 1, 2017 would be 2017-July-1-1, the second sid 2017-July-1-2, etc.
2. Site Information: Record site information, including country, province or territory, island (if applicable) region and/or lower level political division (leave these blank if you don't know; we can figure out the correct informatio later), a general description of the locality, GPS co-ordinates (these are critical) and elevation. The locality should includ named landmarks or geographical features to assist in relocating the sites, i.e. "10 km north of Cambridge Bay, along road t Mount Pelly" or "western shore of Lady Gaga Bay, just south of the mouth of the Beyoncé River". Do your best to describ where you are; we can edit this info later to improve accuracy. Record coordinates in degrees, minutes, seconds, and ensur the datum on your GPS is set to WGS84 (check this when you get the GPS on your leg, and then don't change it). Record both th uncertainty displayed on the GPS, and the radius (estimated by the collector) of the collection site from the location wher you recorded the coordinates. If you are saving waypoints in the GPS unit record their names/numbers here.
3. Habitat Information: Describe the habitat of the site, noting moisture levels and makeup of the habitat i.e. dry tundra, wet meadow, gravelly scree, coastal area, along river, prostrate shrub tundra, etc. Take some photographs o the general habitat of the collecting site and record the photo IDs in this field so we can later associate the photos wit sites.
4. Collector(s): Record the names of all collectors in your party, starting with the collector whose numbe series (see Collector Number) is being used.
5. Associates: List the dominant plant species growing in the general habitat. If you don't know the species leave this blank and we can fill it in later from the habitat photos.

For each *specimen* collected, fill out the following fields:

1. Collection Number: This is a unique identification number assigned to each specimen, allowing the fiel notes to be linked to the sample. Numbers must not be repeated. Best practice is for an individual collector to start at 1. For C3, append your initials to the collection number (`JS-1, JS-2`, etc.). If a collection is later found to be mixture of more than one species, we will subdivide the collection with letters (e.g., `JS-720a, JS-720b`).
2. Species: The species name of the specimen (if known). If you don't know the species, identify it as closel as possible (i.e, to family or to genus, or with a description such as "flower with 4 purple petals" or "evergree shrub"). Be sure to circle the type of specimen you've collected

(vascular plant, moss, lichen, macroalgae, or other), an whether or not a silica gel sample and/or photograph were taken (and the photo's filename from the camera). Silica gel sample won't be taken until you are back in the ship; it is at that point, when you are processing your samples, when you should not in the field book if a silica sample was taken.

3. Specimen Notes: Record flower and fruit colour as observed in the field, as these often change while drying Describe anything in particular that you notice about how the plant was growing (e.g., rhizomatous, tufted/cespitose, cushio plant), its reproductive stage (e.g., in flower, in fruit), its size (e.g. height, especially if referring to a larger wood plant from which you are only collecting one or more branches, or to the typical height, if the sample you are collecting i not representative; stem diameter for trees or shrubs).

4. Substrate: For mosses and lichens record the moisture level (e.g., wet, dry, mesic), light level (e.g. exposed, shaded), and type of substrate (e.g., rock, soil, old shingles) exactly where the specimen was growing. This ca differ from the general conditions at the site (e.g., dry rocks above the water level in a stream or pond). If it's raining try to guess how sunny or dry the substrate usually is.

5. Species Abundance at Site: Record a rough estimate of plant abundance at the collected site, e.g., dominant common (the plant is frequently encountered throughout the site), rare (only a few plants in the area; in these cases it i appropriate to count and record the number of individuals present, if time permits), locally common (common in a localize area rather than all over the site).

NOTE:

- A collection is a gathering of one species in one spot. Ideally each gathering will consist of multiple individuals o the species. You may end up collecting the same species more than once at a site, if you think the plants are differen species, or if one gathering of a species is in a fruiting stage and another in a flowering stage. This is normal, and in suc cases each collection gets its own number. Even though the collections are the same species, they represent differen collecting events.

- If you make more than nine collections from a site (in most cases you will), start a new page with the collection sit number noted on top. There is no need to fill out the location information again if all specimens are collected from the sam location.

Specimen Collection At each site, collect as many different plant, lichen, and algae (if applicable) species. The mor you wander around a site, exploring as many microhabitats as possible, the more species you will encounter. If you are not sure i plants represent different species (often it will be obvious, but in some cases it will not be, particularly for small species) use your best judgment to decide whether or not to make a collection: if the plants or lichens look different to you, it is bes to collect.

Collecting Vascular Plants

1. Take photographs of the plant you're collecting (time permitting), aiming for a photo that shows the whole plant in it local habitat. These photos will be useful for helping us describe the habitat and associated species. Feel free to tak close-ups of flowers, etc., but it would be better to do this if you have time after you have collected all the plan diversity in each area.

2. Dig herbaceous (non–woody) plants out by the roots (try your best to keep the roots intact), and shake off any exces dirt. However, if you don't get the roots or rhizomes, still keep the specimens. If you're collecting a woody plant (e.g., willow or birch shrub), use the pruning

shears to cut off one or more branches with flowers and/or fruits where possible Branches should be cut a bit shorter than the size of a herbarium sheet, on which they need to fit later.

3. Try to collect enough individual plants (or clippings) from a population to fill a standard 11' by 18' herbarium sheet In general, collecting many individuals is always better than collecting a few. Individual Arctic plants often make poo specimens on their own. The exceptions to this rule are rare plant species, or extremely small populations. NEVER collect all or most) of the specimens growing in an area.

4. Fill out the collection information for this specimen in your field notebook (see Data Collection, above).

5. Place your specimen in a Ziploc bag and write the collection site number and the collection number for that specimen o the outside of the bag with a Sharpie. Large specimens can be folded, but otherwise plants should be handled carefully so the don't get mangled. You may put multiple collections in the same bag, but make sure (i) you don't mix species that have any chance of being confused later when pressing the collections, and (ii) you write th collection numbers of each collection on the bag, so you can associate the correct specimens with the correct numbers.

6. Place this bag in your collecting satchel, and take care not to crush the specimens in the bag.

Collecting Bryophytes

1. Take a photograph of the collection site, and the plant you're collecting (time permitting).

2. Collect groups of entire plants. All parts of a bryophyte help in identification. Use an implement such as your digge to lift plants from rocky or woody substrates without leaving their bases behind. Collect a small sample of substrate with th plants if they are too firmly attached to their substrates to lift them in one piece (but generally avoid collecting larg quantities of soil or litter with the plants). Plants with sporophytes (capsules) – even old ones – are more useful fo identification than those without.

3. Each sample should ideally be about the size of your palm. Never collect a sample if it will remove most or all of th species from the site.

4. Each sample should have a single species dominant. Bryophytes often grow intermixed, but try to make separat collections for each species. For example, if species A is mixed with B, try to collect a sample that is mostly species A an another that is mostly species B.

5. Each sample should be collected in a separate paper bag. On drying, bryophyte samples become brittle and difficult t separate from one another without inflicting damage. Some samples that form solid cushions when moist can fall apart when dr and mix together with other species in the same bag. Furthermore, mosses in over-stuffed bags may dry too slowly to preven mold from rendering the samples useless. Paper is used instead of plastic because bryophytes are preserved best in a dr state, and no further processing is necessary once the sample is placed in a bag. The top of each bag should be folded ove tightly so that the sample doesn't fall out, then place the bag in your collecting satchel, ensuring the specimen is no crushed.

6. Each bag should be legibly labeled in permanent ink (use your Sharpie). Include at least the collection number. The res of the specimen information can be recorded in your collection book. Alternatively, if you prefer to transcribe th information later, on the ship, record it on the bag and make sure to record the collection numbers for the site in the book

Collecting Lichens

Lichens are collected in much the same way that bryophytes are. Of the three main types of lichen (*fruticose*, *foliose*, and *crustose*), only some of the larger *fruticose* species grow free from a substrate. The rest ofte grow attached to tree branches, rocks, or soil. It is therefore often necessary to collect the lichen with a part of it substrate. This includes chipping *crustose* lichens off rock (try to get as much of the lichen as possible, including th center, and the margins), cutting away a branch from a larger shrub with a lichen attached to it (be sure to record the shru species), or carefully digging out a tiny bit of soil with a soil crust lichen on it. Once you make a collection, follow steps through 8 noted above under bryophytes.

Collecting Macroalgae

Macroalgae (kelp) can be found in the intertidal zone both as complete organisms (species that grow in this zone and that ar adapted to tidal life), and as fragments (or whole organisms) that have washed ashore from deeper waters.

Collecting both types of kelp are important contributions to marine biodiversity science. When collecting either type be sure t collect as intact a specimen as possible, ideally one possessing a hold-fast, stipe, and blade (or part thereof).

1. Select a specimen, and place this specimen in a Ziploc collection bag.
2. Fill out a collection record for the site (if necessary) and specimen, and write the collection number on the Ziploc bag.
3. Place this Ziploc in your collecting satchel, and protect the specimen from being crushed.

Processing

Vascular Plants Ideally, plants should be pressed as soon as possible after collection, but this is not alway practical (it's late in the day; you are tired and hungry; there's something more interesting going on, etc.). It is okay t process your collections the morning after they were collected. Some species, such as poppies, will lose their petals if no processed quickly. If you don't have time to process all your collections upon return to the ship, try to at least process th most sensitive ones that are losing their petals, or that are really shrivelling up, etc. Prior to processing keep the collectin bag containing the plant specimens in a cool, shaded place to prevent wilting. Ensuring the Ziploc bags are closed will also slo down excessive wilting (some wilting is okay and makes it easier to press plants; excessive wilting, in some cases, will result i poorer specimens).

1. Cover the workspace in newsprint (this can be re-used) to minimize the mess, or work on a large tarp that will captur all the extra dirt and debris from the specimens.
2. Organize your bagged samples by site - it's often easier to press in the order you collected the plants in. However since you should have assigned collection numbers to all your collections, ordering them is not strictly necessary, especiall if time is limited.
3. If necessary, clean excessive dirt and especially mud from the roots of the plants using a bowl of water, and shake th excess water off. If pressed for time, just shake the roots out onto your tarp. Specimens with lots of dirt attached to the dry slower and are messy when the specimens are back in the museum.
4. Place a plant press end flat on your workspace, place a cardboard on top, and open a blank newsprint onto it. Arrang the plants from a single collection on the newsprint so that they all fit on the sheet. For plants with large flowers, arrang the specimen so the flowers are fully

visible when the plant is dry. For small plants and grass-like plants it is sufficien to just lay them on the sheets in any orientation. Space the specimens out, as much as possible, so they are not laying ato one another, as this slows down drying. Fold specimens that are larger than the newsprint sheets. If a specimen is folded t fit in the newsprint, it will fit on a herbarium sheet. Do not simply fold a large plant in half; rather, fold over the part or parts) of the plant that extend beyond the sheet margins. If you have lots of material, you can use multiple newsprin sheets. Be sure the collection number is marked on each sheet.

5. Write the collector name and collection number (required; e.g. JS 25 or Saarela 25) on the inside and outside of th newsprint, along the bottom or side edges, and the species name on the outside of the newsprint (time permitting; this speed up sorting back in the herbarium) Make sure we can read the collection numbers please!

6. As you're processing your plants, go through your camera, and write the filename of any photos of that plant on th collection form. Be sure to transfer these photos to the lab computer.

7. DNA Samples: Collect leaf material from one plant per collection, and put it in a silica gel envelope labeled with you name, the collection number, the date, and the location. For large specimens with lots of leaf material, sample several leave (again, more is always better). For small plants with few leaves, sample one or more leaves, but don't remove all the leave from the specimen, rendering it useless. For very small individuals, it will be difficult to sample one or more leaves, an these will be so small it will be difficult to get DNA out of them. In these cases, sample a whole plant into the silica ge envelope. Seal the envelope with a bit of water (don't lick it…remember what happened to George Costanza's fiancé?) and put i back in the silica gel, which will rapidly remove the moisture from the leaf tissue, preserving the DNA for later extraction Write the range of the collection numbers of the silica gel envelopes on the outside of the silica gel Ziploc (e.g. JS-1 t JS-25)

8. Tag the plant you removed the leaf material from with a jewelry tag marked with "DNA" and your collection number. If yo sampled a whole plant, note this in the fieldbook; in such cases there will be no plant to tag. Close the newsprint, place new cardboard on top of the closed sheet, and repeat the arranging and pressing process, alternating cardboards and fille newsprints, until you have all your specimens processed.

9. Place a plant press end on the top and bottom of the stack, and cinch the stack together as tight as you can with tw plant press straps. It often helps to have one person stand on the press while another person tightens it.

10. Place the tightened press in front of a space heater, set to low, to speed the drying process. To dry, warm air needs t move through the press to remove moisture from the specimens. The press can also be dried in a windy and sunny area. Do no get the press wet.

11. Check the press for dry plants when you are getting close to running out of cardboard. Plants that are stiffl 2-dimensional, and dry to the touch, are dry (the newsprint in which they are pressed in should also be dry), and can b removed from the press. Small plants and most grasses, sedges and rushes will dry in one day, if dried with a heater; larg plants, wet plants and woody plants usually take several days to dry. If you run out of cardboard and find that specimens i the press are not yet dry, you may double- or triple-up sheets of newsprint between cardboards. The specimens will dry slower but will turn out okay. Once removed from the press, specimens must be packaged for transport by wrapping a stack of specimen tightly in several sheets of paper. Do not leave loose sheets of newsprint with dried plants laying around. If you don't hav enough dry specimens to prepare a package, put a stack of specimens between two pieces of cardboard and securely tighten th pile with string. Ensure each specimen remains in its newsprint, so that the collection remains associated with its number. I the

plants fall out of their sheets they become useless because they cannot be associated with their collection data.

Lichens and Bryophytes

Dry samples as soon as possible upon return to the ship. Moss and lichen bags need only be labeled with the collection number which should have been done in the field), and dried in a single layer in front of the heater. Be sure that there are measures i place to prevent samples on the floor from being stepped on, or kicked/blown out of reach or out of sight. It is not necessary t take samples for later DNA extraction from bryophytes and lichens. Drying samples should be shifted periodically until thei contents are completely dry to the touch (be sure to test the middle of large clumps as well as the outside). Once dry the sample can be stored almost indefinitely in their paper bags, in a mesh sack in a dry place. Do not put the paper bags into a mesh sac if they are wet or moist, or fungi will colonize them, ruining the specimens.

Macroalgae

Macroalgae are processed in the same way as vascular plants (including sampling of tissue for DNA analyses - blot the tissue firs to speed up drying) except for one modification designed to prevent the wet specimens from sticking to the newsprint. When yo open the newsprint on the cardboard, place a National Herbarium of Canada plant specimen sheet on the stack, on top of the ope newsprint. Lay the wet macroalgae specimen on the plant specimen sheet, and then place a sheet of cheesecloth on top of th specimen. Close the newsprint, and continue pressing plants. The specimen will be stuck to the herbarium sheet when dry, and th cheesecloth will prevent the top side of the newsprint from sticking to the specimen.

9 Autonomous Recording Unit Transcription Protocol

This protocol describes the methodology used to identify and count species detected within the recordings [45].

9.1 Converting File Formats

The autonomous recording units are set to record in `WAC`, a proprietary format created by the manufacturers of the cording units, Wildlife Acoustics. To open the recordings in software necessary for anscription, the files first need to be converted into `WAV` format. This can be done using a command line converter veloped by Peter Solymos[d], or with software developed by Wildlife Acoustics available for free - download Kaleidoscope here.

Once installed, opening Kaleidoscope brings you to the main screen (Figure 23).




Figure 23: A screen capture of the main window in Kaleidoscope.

The kaleidoscope window is composed of two main parts:

- the `INPUTS` section (red) that specifies the file format and location of files to be converted,
- the `OUTPUTS` section (blue) that specifies the format files should be converted to, and their future location.

[d] See the github repository for the converter here

9.2 Listening Software

Many software programs can be used to help identify, and count the species detected in recordings accurately. The programs best ited for this purpose all visually display the audio information in a spectrograms. Spectrograms display sound frequencies ross time, and given the unique aural signatures at the species and individual level, a visual representation of sound enables a stener to both "see" and hear vocalizing animals and thus more easily identify all individuals (Figure **??**).




Figure 24: A screen capture from Adobe Audition's spectrogram display. We can clearly see the Sandhill Crane's vocalizing.

9.2.1 Software Settings

In this protocol, we focus on the programs Adobe Audition and Audacity. With a few adjustments to the default settings in these programs, we can create a much tter environment for detecting bird species/individuals in the recordings.

Spectral Display The settings responsible for tailoring the spectral display to specific monitor resolutions are cated in the top menu: `Edit` ⇒ `Preferences` ⇒ `General`, with recommended ttings found in Table 10. These settings ensure that vocalizations appear sharp and constrasted ainst background noise.

Size	Window Type	Scale	Decibel Range (dB)	Length Min	Length x
2048	Blackmann-Harris	Logarithmic	132	15	30 to

Table 10: Spectral display settings used to improve the appearance of vocalizations in recordings

Frequency Range Songbirds vocalize within a certain range of sound frequencies, and identifying calls is made easier adjusting the range of the spectrogram accordingly.

- create screen cap of spectrogram with inappropriate range
- compare with proper frequency range

Timeframe When the audio recording is first opened, the spectrogram spans the entire recording (10 minutes in our se). This resolution is too low to reliably detect individuals, and the spectrogram should be adjusted so that 20-40 seconds are splayed, depending on the monitor size. This can be achieved in Adobe Audition by resizing the scroll bar above the spectrogram, d the zoom buttons located above the spectrogram can be used in Audacity.

9.3 Entering Data

All data entry is completed in the `Data Entry` form within the LTEMP ARU Database (Figure 25).




Figure 25: Screen capture of the data entry form within the ARU database used while transcribing recordings.

9.3.1 Site Info Data Entry

- Site - unique ID associated with the sampling site.
- StationID - unique ID of the ARU location within the site. This was added to prevent overlapping data tries if multiple ARUs are deployed within each Site.
- Method
  - 1 = listen to the first 5 minutes, visually scan the last 5 minutes
  - 2 = listen to the first 5 minutes with no additional visual scan
  - 3 = visually scan all 10 minutes with listening to confirm species/individual IDs
- Rain - on a scale of 1 → 5 of intensity, with 5 being the highest
- Wind - same scale as rain

9.3.2 Species Detections Entry

While listening to the recording, each individual detected should be recorded on a seperate row, with the following information lled in:

- Species - Entered using the four letter AOU standardized code.

- Ind_ID - The default value for this field is '1'. It corresponds to uniquely identified individuals of a species on the recording. In other words, if you hear two Snow buntings in the recording, you would have one row with NBU 1' and second row with 'SNBU 2'. As many individuals as you can identify can be added. Evidence for multiple individuals eds to be strong (see TMTC below). Fill in the field with '0' if the species is 'NONE'.
- O min → 9 min - Enter the second when each individual was detected within each nute. In other words, every box should either be empty, signifying that individual was not heard during that minute or have a value between 1 d 59. Detections made within the first 1 second (e.g. <1) are labeled as '1'. Each audio program contains a digital timer to lp pinpoint the second the species was heard.
- TMTC - The TMTC (too many to count) column is used to indicate whether one individual was clearly detected, multiple individuals with simultaneous and overlapping vocalizations were detected. A value of "1" indicates just one dividual was heard, while a "2" indicates there are too many individuals to count seperately. A value of 2 is often tributed to migrating flocks.
- VT - This field identifies what type of vocalization was heard. Songs are unique and repetitive territorial calizations, and are thus easier to identify. Calls are often less unique and therefore more difficult to identify to ecies.
- Confidence - Confidence in species identification, as well as whether or not this is a new individual of the me species (ie. high confidence that the detected individual is a second SNBU for example)
- Comments - additional comments associated with the detected individual.

Recording Unknowns If a song or call cannot be identified, the species code should begin with "UN", followed by the first two letters of the bird oup (UNPA, UNSH, UNGU for unknown passerine, shorebird and gull respectively). A clip of the song should be saved into a folder tled "unknown", using the following naming scheme: siteID-stationID-date-time of recording-time of clip-species code.wav. An ample of this would be: `R1Z1-A-20150515-063000-0150-unpa.wav`. This procedure should also be used for species signments with low confidence.

9.3.3 Additional Resources for Confirming Species ID

Many individuals will be difficult to identify due to sub-optimal recording, or unusual vocalizations. In cases where individuals an unknown species are detected, online libraries of bird vocalizations can be used to compare similar sounding species:

- Xeno-Canto - a vast library of user submitted bird vocalizations from around the rld. Recordings include both song and calls, and each recording is accompanied by a spectrogram.
- Environment Canada's Dendroica - fewer recordings than Xeno-Canto for ch species, but good user-interface, and spectrograms can be viewed here as well.

ᑐᓂᓴᐅᔨᓪ ᓄᓇᕗᑦ ᐅᒪᔪᓕᕆᔨᖅᑯᓂᓪᓗᓂ ᑲᑎᒪᔨᖏᓐᓄᓂ

ᐃᒪᐃᓕᒪᔨᐅᓪ

ᑐᕿᓚᖏᖅ: **ᐊᖏᑭᒐᔨᓪ: X**

ᐱᓱᒋᐅᔨᖅ: ᐃᓗᒪᓕᐊᖏᒃ ᐱᓴᒪᐊᖏᖅᕿᑦ ᐅᓪᒪᑭᓂᐊᔪ/ᐅᓪᒪᑭᓴᐅᓂᐊᖏᑦᓕᓂᓂ ᖃᓄᓕᕐᓄᓂ - ᐊᖏᕕᑦ ᐃᓕᖏᑦ (Blue Whale) (ᐅᐊᖏᓇᖅᑯᓗᖏᓕᓂ-ᐱᖏᕐᓇᖅᑯᓗᖏᓕᓂ ᐊᑦᓯᖓᒃ ᑕᒪᐅᖏᑦ ᖃᓄᓕᕐᓂᐃᑦ) ᖁᓇᐃᓕᐅᒪᐊᖏᓂᔪᑦ ᑯᓇᐅᑎ-ᔪᑦ.

ᖁᓇᐃᓕᓐᓄᓂᖏᓐᓄ:
ᐊᖏᕕᑦ Blue Whale (Northwest Atlantic population) ᖁᓇᐅᔨᖏᑕᐅᓕᖏᓪᓚᑕ ᓄᑎᒪᔨᖏᓐᓄᓂᓪ ᖁᓇᐃᓕᓐᓄᓚᖏᓐᓄᑕ ᐅᒪᔨᐃᓪ ᐊᑦᑕᕆᓭᑐᑦᑐᑦ ᓇᓭᑕᒥ Committee of the Status of Endangered Wildlife in Canada (COSEWIC) ᐅᓗᓕᐊᓭᑦᑐᒧᓐᖏᓐᓂᓂᓇᓂᓂᓇ 2002 ᓭᓗᓭᐃᖏᑕᐅᓕᖏᓪᓚᑕ ᐊᓕᒪᓗ ᑎᑎᖏᑦᑕᐅᓪᓗᑎᓇ ᐅᒪᔨᐃᓂ ᐊᑦᑕᕆᓭᑐᑦᑐᑦᑐᓗᓂ 2005. ᓄᓇᕗᒻᒥ ᐅᒪᔨᑦᓕᕆᔨᖅᑯᓂᓪᓗᓂ ᖁᓇᐅᔨᑎᑕᐅᓕᖏᓐᓄᑦ ᓇᓕᒪᔨᓄᓂ ᐊᑦᑕᕆᓭᑐᑦᑐᑦᑐᓗᓂ ᐅᒪᔨᐃᓂ ᖁᓇᐅᔨᖏᑦᑕᐅᓂᖏᓐᓄᓇᓂ ᐃᖃᓗᓕᕕᓴᐅᒪᔨᖏᓐᓄᓇᓗ ᑎᑎᖏᑦᑕᐅᓂᐊᖏᓪᓚᑕ 2004. ᓄᓇᕗᒻᒥ ᐅᒪᒋᑦᓕᕆᔨᖅᑯᓂᓪᓗᓂ ᓇᓕᒪᔨᖏᓐᓂᓂ ᐊᐱᓕᔪᐅᓕᖏᑦᒥᓂ ᐊᖏᑎᔪᓂᖏᓕᖏᓪᓚᑕ ᐅᓇᓱᐊᓂᐊᖏᑦᑕᐅᓕᓴᐃᖅᑐᓂᓪ ᖁᓇᐅᑎᖏᓐᓄᓂᓇᓂᓇ 2007 ᓇᔨᓕᑎᑕᐅᓕᐅᑦᓴᓂᓪᓗ 2010. ᐅᓗᓕᐊᓭᑦᑐᓂᖏᓐᓄᓂᓇ ᐊᑦᑕᕆᓭᑦᑐᓂᖏᓐᓄᓂᓇ ᐃᒪᔨᐃᓂ, ᐱᔪᓕᐊᖃᓂᖏᓐᓄ ᒪᑕᓪᓗᔪ ᑭᔨᔨᐃᓂ ᐊᑦᑕᕆᓭᑦᑐᒧᓐᖏᓐᓂᓂ ᐱᒪᕐᔨᐃᓂ ᓴᓐᕿᑕᒪᐊᖏᖅᖁᓪᓚᑕ ᐊᐅᓪᔪᕿᑎᐊᖏᓂᓪᓕᔪᑦ ᖁᓇᐅᑎᓂ ᐊᖃᔪ ᑕᓪᓕᒪᐃᓂ ᐃᓗᓐᓂᐊᑦ ᐊᑎᖏᑦᑕᐅᓂᐊᖏᓪᓚᑕᓗᓂᓪᓗ.

ᑎᑎᖏᑦᔪᒪᔨᑦ 5.2.34 (d) (i) ᓄᓇᕗᑦ ᓄᓇᖅᓱ ᐱᑎᔪᐅᓕᓂᖅᓱᓄᓂ ᐊᖅᑎᖏᕆᑎᑐᓂ ᐅᖏᖏᔪᒪᓕᒪᓚᑕ ᓄᓇᕗᒻᒥ ᐅᒪᔨᐃᓪᓚᑕᐅᓕᖏᓪᓚᑕ, ᐃᓗᒪᖏᔪᓗᑎᒃ, ᐊᖅᑎᖏᔪᓄᓐᓇᖏᓂᖅᓱᓂᖅᓱᓂ ᖁᓇᐅᑎᐅᔨᓂᓂᓪ ᐊᐅᓕᑕᐅᓂᖅᓱᓐᓂᓄᑦ, ᐅᓗᓇᐃᓕᐅᔪᓕᓂᖅᓱᓐᓂᓄᑦ, ᓴᓪᔪᐅᔪᓕᓂᖅᓱᓐᓂᓄᑦ, ᖁᓇᐅᑎᐅᓂᖅᓱᓐᓂᓄᑦ ᐅᒪᔨᐃᓂ ᐱᓗᐊᖏᑕᐅᓂ ᐅᓗᓕᐊᓭᑦᑐᓂᖏᓐᓄᓂ ᐃᒪᔨᐃᓂ. ᐃᓗᓕᓐᓄᓂ ᖁᓇᓂ ᐊᖅᑎᖏᑕᐅᓕᔪᑲᓴᖏᓐᓄᑕ ᐊᐅᓕᑕᐅᓂᖅᓱᓐᓂᓄᑦ ᖁᓇᐅᑎᓂ, ᐅᓗᓕᐊᓭᑦᑐᓂᖏᓐᓄᓂᓇᓗ ᐊᑎᓚᓗ ᐊᐅᓕᓴᐃᖅᑐᓂᖏᓐᓄᓂᓪ ᐊᖅᑎᖏᕆᑎᑐᓂ ᐅᖏᖏᔪᓄᓐᓇᖏᓂᖅᓱᓂ ᐃᓗᓐᓂᐊᑦ 2008 “ᓄᓐᖏᖏᑎᑎᔪᑕᐅᓕᑦ ᑐᑭᔪᐅᓴᖏᖏᑎᑎᓂᖅᓱᓂᓪᓗᑦ ᑲᑎᓴᓐᑕᐅᓂᖅᓱᓐᓂ ᐊᓵᐅᓄᖏᓂᑐᑦ, ᐅᓗᓇᐃᓕᐅᔪᓕᓂᖅᓱᓐᓂᓄᑦ ᐃᒪᔨᐃᓂ ᑕᙳᓂ ᓇᓭᑕᒥ ᐱᑎᔪᐅᓕᓂᖅᓱᓐᓂᓄᑦ ᐊᖅᑎᖏᕆᑎᑐᓂ ᐊᖏᑎᔪᓂᖏᓕᖏᓪᓚᑕ ᐊᐅᓕᑕᐅᓂᖅᓱᓐᓂᓄᑦ ᐊᖅᑎᖏᕆᑎᑐᓂ ᐃᒪᔨᐃᓂ ᐊᑦᑕᕆᓭᑦᑐᓂᖏᓐᓄᓂ ᐱᖏᓐᓄᓂᓪ”.

ᐊᖏᕕᑦ Blue Whale (Northwest Atlantic population) ᐊᐅᓕᑕᐅᓕᖏᓪᓚᑕ ᐸᓇᐅᑎᓂ (ᓭᐃᐅᓴᖏᓐᓂᔪᑦ ᐅᖏᑲᓐᓄᓕᐃᓂ ᑕᙳᐅᖏᑦᑕᓂ ᐊᑎᓚᓗ ᐃᓇᓄᑐᖏᓐᓂᖏᓕᐅᑕᐅᓐᓂᓕᐅᓐᓄᓄᑦ) ᐅᖏᑲᓐᓂᔪᑦ ᖁᓇᐃᓕᐅᒪᐅᖏᓪᓚᑕᐅᓂᖅᓱᓐᓂᓄᑦ ᐱᕐᔪᐅᑎᑐᓐᓂᔪ ᐅᙳᑐᓴᐅᓂ ᐃᑲᔪᑕᐅᓗᓐᓄᑦ ᑐᓂᓯᖏᓐᓄᓗᓂ ᐸᓇᐅᑎᓂᖏᓐᓄᓂ ᐊᖅᑎᖏᕆᑎᑐᓂ ᐅᖏᑲᓐᓂᔪᑦ ᐊᖅᑎᖏᕆᑎᑐᓂ ᐃᒪᔨᐃᓂ ᐊᑦᑕᕆᓭᑦᑐᓂᖏᓐᓄᓂ ᐱᖏᓐᓄᓂᓪ

<ᖁᓇᐅᑎᓕᐅᓚᓂᖅᓗᒃ, ᐊᐢᖏᑎᓕᓚᐳᓴᓯᑦ ᖃᑭᖁᖃᓯᓴᐅᐳᓱᑦ ᑕᑦᓯᓕᖅᓴᓴᓯᑦ ᑎᑎᖅᑲᓕᐊᒥᑦ, ᓇᓗᓇᐃᖁᓗᑎᑐᓗ ᐃᓚᒪᑭᖅᓚᐃᓪᓚᓴ ᑕᑕᓇ ᖃᓇᐃᓕᓕᐅᓚᐊᖁᓂᓗᒃ ᐸᖁᓇᐅᑎ.

ᑕᑕᓇ ᑎᑎᖅᑲᖅ ᑲᑎᒪᔨᓇᐊᖅᑕᐅᖅᑲᐅᑕᐅᓕᐅᖅᑐᖅ ᐃᑦᓯ 2016-ᒥᑦ, ᑕᐃᑲᓂ ᑲᑎᒪᔨᑦ ᖃᑎᖅᑭᐅᖁᔭᓕᐅᖅᑎᑕᖅᓯᑦ ᑭᓯᐊᓂ ᑎᑎᖅᑲᖅ ᑭᖅᑦᓕᖅᐸᓕᐊᖅᓯᒪᓕᖅᐸᑦ. ᑎᑎᖅᑲᖅ ᑭᖅᑦᓕᖅᐸᓕᐊᖅᓯᒪᓕᖅᑐᖅ.

ᓯᐸᑕᒪᐸ 11, 2019, ᑐᖅᐳᖅᑐᑦ ᑭᓚᓕᓴᐃᑦ ᐱᓕᓕᐊᖅᓂᕆᔨᓂᒃ <ᖁᓇᐅᑦ ᑲᑎᒪᔨᓂᒃ ᐅᖃᐅᓯᐅᓕᐅᖏᒪᑦ. ᑲᑎᒪᔨᑦ ᑭᓯᐊᓵᓂ ᐊᖏᓕᐅᖏᑕᖅᓯᑦ ᐱᓕᓕᐊᖅᓂᕆᔨᓂᒃ <ᖁᓇᐅᑦ ᐃᓯᒪᓗᓐᓇᖅᑐᖃᐅᓕᖏᒪᑦ ᐃᓕᖏᑦ ᐊᖅᖅᓯᑎᑦ ᑲᑎᒪᔨᖏᑦ ᖃᓄᑕᖏᓕᓂᓯᓯᐅᖅᑐᑦ ᑐᖅᐳᖅᑐᑦ ᑭᓚᓕᓴᐃᑦᐃᓯᒪᔭᑎᖃᖏᓐᓇᖅᓂᐊᖏᒪᒃᑕ. ᑲᓇᑕᐅᐸ ᐃᒪᖏᐅᑕᓕᓚᔨᖏᑦ ᑐᓂᓯᒪᓕᐅᖏᒪᒃᑕ ᐊᔨᖏᓕᓂᓱ ᐱᓕᓕᐊᖅᓂᕆᔨᓂᒃ <ᖁᓇᐅᑎᐸ ᓄᓇᓕᓕᓂᒃ 6-ᓄᑦ ᐅᖃᐅᓯᓴᖃᖅᑎᓯᐸᔪᑦ ᐃᓕᐅᔪᒪᓂᖏᓕᓂᓱ ᑐᖅᐳᖅᑐᑦ ᑭᓚᓕᓴᐃᑦᐅᖃᐅᓯᓴᖏᓕᓂᓱᓗᔪᓐᓇ ᓂᓚᐅᓗᔪᑎᑎᒃ <ᖁᓇᐅᓗᒍᑦ ᑎᑭᓕᐅᓂᖏᓐᓄᓂ ᓄᐃᐱᓚ 25, 2019. ᑐᓴᓯᕋᓕᐅᕋᔪᐊᖏᒪᑦ, ᒪᑭᐃᓇᓴᑎᑦ ᐊᖅᓄᓇᓴᑎᑦ ᑲᑎᒪᔨᖏᑦ ᑭᐅᓕᐅᖅᐳᒃ ᑕᑯᓕᐅᖏᓕᓂᓱᖅᑐᓂᒃ ᐅᖃᐅᓯᓴᖃᖅᑎᓯᐸᔪᑦ ᑐᖅᐳᖅᑐᑦ ᑭᓚᓕᓴᐃᑦᐅᖃᐅᓯᓴᖏᓕᓂᓱᓗᔪᓐᓇ ᐅᖃᐅᓯᓴᖃᓕᐅᖅᓯᒪᓕᐅᖏᓕᓂᓱ, ᐃᓯᓇᓕᐅᖅᐳᒍᑦ ᖃᐅᔨᓴᖅᑐᓂᒃ ᐅᖃᐅᓯᓴᖏᓕᓂᓱᓗᔪᓐᓇ ᑐᖅᐳᖅᑐᑦ ᑭᓚᓕᓴᐃᑦᐱᓕᓕᐊᖅᓂᕆᔨᓂᒃ ᐅᖃᐅᓯᓴᖃᓕᐅᖅᓯᒪᓕᒃ, ᐃᓯᓇᓕᓕᓂᓱᓗᔪᓐᓇ ᐃᒍᓗᔪᑕᐅᖅᓯᒪᓕᖅᑐᑦ ᑕᒪᐃᓕᓐᓄᑦ.

ᐊᑐᓕᖁᔭᐅᓪᓚᑦ:
ᐊᐢᖏᑎᓕᓚᐳᓴᓯᑦ ᖃᑎᖅᑭᓗᓄ ᐅᐸᓗᒥᒍᓕᖅᑐᖅᓯᓯᒪᓚᖅᓴᖅ ᖃᖅᑲᐃᓕᓕᐅᖅᓯᒪᔪᑦ <ᖁᓇᐅᑎ ᖃᐅᓚᓂᖅᓂᒃ - ᐊᖁᐃᑦ ᐃᓕᖅᑎᑦ (Blue Whale) (ᐅᐊᖅᓇᖅᐸᓯᖅᓂᓕᓂᒃ-ᐱᖅᓂᖅᓇᖅᐸᓯᖅᓂᓕ ᐊᑦᓭᖅᑎᒃ ᑕᓚᐅᖁᒥᑦ ᖃᐅᓇᓚᐃᑦ), ᓂᐸᓕᐅᑎᖃᖁᓗᑎᒃ ᖃᖅᑲᑐᐃᖅᓇᖅ ᐃᖁᖅᓗᓕᓚᐳᓴᓯᒪᓐᓄᑦ ᐊᓯᒪᓗ ᐃᓯᒪᓕᓕᐊᖅᓂᖅᓴᖅᑐᒃ ᑐᓂᓯᓗᑎᒃ.

ᐃᖁᐅᓯᖁᑎᓕᐅᔪᖅ ᐅᒧᖅᓂ:
ᓴᒻ ᓯᑎᐅᖅᓴᖅ, ᐅᒪᔨᐃᑦ ᐊᑦᑕᖁᓇᖁᑎᒋᑦᑐᑦ ᖃᐅᐅᔨᓴᖁᑎ, ᐃᒪᖁᑎᐅᑕᓕᐅᓚᐳᓴᓯᑦ ᑲᓇᑕ, ᖃᐅᑎᐊᓄᑦ ᐅᑭᐅᖁᑦᑕᖁᑐᒧᑦ , ᕿᐃᓂᐱᖅ

ᐅᑦᓗᖅᓂ:
ᓴᓂᐊᓇ 16, 2020

ᐃᓕᓴᐸᐧᐦᓂᖅ 1 – ᖃᐅᔨᓴᐸᖅᑕᐅᓂᓄᑦ ᐊᖏᓐᓇᓕᒃᑎᑦ ᑲᑎᒪᔨᖅᓯᓂᑦ ᐱᓕᕆᕐᔪᓂᖅᓄᑦ ᐱᖁᔭᐅᑎᖏᓐᓂᑦ ᑐᖏᓐᓂᒃ ᖃᓄᐃᓕᓴᕐᓂᑦ

ᑐᓴᖅᑎᑦᑎᓂᖅ ᐊᖏᓐᓇᓕᒃᑎᑦ ᑲᑎᒪᔨᖅᓯᓂᖅ:
ᐊᒃᑐᐱᓚᓗ 9, 2019 – ᐱᐊᓕᔪᓄᑦ ᑎᑎᖅᑎᒍ ᑐᕈᓕᒃᓂᕐᑎᑦ ᑐᖏᖅᑐᒧᑦ ᖃᓄᐃᓕᓴᕐᓂᑦ ᐸᖏᓐᓂᑦ, ᐃᖅᓗᓐᓂᑦ,, ᖃᑭᑕᔪᐊᖏᓐᓇᑦ, ᑲᓛᓕᓴᐱᐊᓐᓇᑦ ᒥᑎᒪᑕᓕᓐᓇᓪᓗ ᐊᖏᓐᓇᓕᒃᑎᑦ ᑲᑎᒪᔨᖅᓯᓂᖅ, ᑎᑎᖅᓂᐊᖏᓐᖁᖅ ᐅᖅᓯᒪᑦ ᑕᒪᖏᑐᒃ ᑐᓂᔭᐅᔾᔨᑦ ᐊᒃᑐᐸ 17th.

ᐊᒃᑐᐱᓚᓗ 28, 2019 – ᖃᓚᑕᐅᔨᔪᑦ ᑐᓂᔭᐅᔪᖅ, ᐃᓗᐊᓂᔪᐅᖅᑐᖅ ᑎᑎᖅᑲᖅ, ᐊᖏᓐᓇᓕᒃᑎᑦ ᑲᑎᒪᔨᓕᒪᓕᕐᓂᓄᑦ ᐃᒪᖏᑕᑕᓕᕐᓂᓄᑦ ᑐᓴᕐᓯᖅᑕᖏᓐᓄᑦ ᐊᖏᓐᓇᓕᒃᑎᑦ ᑐᖏᖅᑐᒧᑦ ᖃᓄᐃᓕᓴᕐᓂᑦ.

ᓄᖕᐱᓚᐃᐅᓐᓇᑦ 20-21 – ᐅᖃᓗᓴᑐᑦ ᐊᖏᓐᓇᓕᒃᑎᑦ ᑲᑎᒪᔨᖅᓯᓂᖅ ᐅᑯᓄᖓ ᐃᖃᓗᐃᑦ, ᑭᒥᑦ ᒥᑎᒪᑕᓕᒃ, ᑭᓯᐊᓂ ᑐᓂᓯᖅᑐᑦ ᐅᖃᐅᓯᐸᓴᖅᓯᓂᒃ.

ᖃᖅᑲᓇᐱᐅᓐᓇᑦ ᐊᖏᓐᓇᓕᒃᑎᑦ ᑲᑎᒪᔨᖅᓯᓂᖅ: ᑕᑯᓯᐃᓕᑦᑐᑦ ᑐᖏᖅᑐᓂᒃ ᖃᓄᐃᓕᓴᕐᓂᒃ, ᐅᖃᐅᓯᐸᓴᐃᑐᑦ.

ᐸᖅᓂᑐᐊᓐᓇᑦ ᐊᖏᓐᓇᓕᒃᑎᑦ ᑲᑎᒪᔨᖅᓯᓂᖅ: ᐅᖃᓯᐸᓴᐃᓕᑦᑐᑦ ᑐᖏᖅᑐᓄᓂᒃ ᖃᓄᐃᓕᓴᕐᓂᒃ .

ᖃᑭᖅᑕᖅᓱᐸᑦ ᐊᖏᓐᓇᓕᒃᑎᑦ ᑲᑎᒪᔨᖅᓯᓂᖅ: ᓄᐃᐱᓚ 21, ᑭᐅᓕᐅᖅᑐᖅ ᖃᐅᔨᓚᑐᐊᖅᑎᑐᖅ ᑎᑎᖅᑲᓂᒃ ᐱᓕᐅᖅᓯᒪᓕᒃᑎᓕᐅᖅᑐᒃ, ᒪᑭᐃᖅᑎᒃ ᐅᖃᓗᓕᒃᑲ ᑭᐅᖅᑎᑐᑦ.

ᓇᐃᓐᓇᕐᓂᒃ ᐅᖃᐅᓯᖏᓐᓄᑦ – ᐊᐅᓪᓚᕆᐊᖅᓂᖅᓗᑦ ᐸᖅᓇᐅᑎᑦ ᐊᖅᓴᑦ Northwest
Blue Whale (ᑕᓪᐅᕐᓚᓂ Atlantic ᐱᑕᖃᖅᓂᕐᕆᑦ)

ᑕᓪᐅᕐᓚᓂ Atlantic ᐊᖅᓴᑦ blue whale ᑎᑎᕖᖅᑕᐅᓯᒪᖏᒪᑕ ᐅᓗᓪᓕᐊᓇᖅᑐᑦᓂᕐᕆᓂᕐᓂᖅ ᒪᑕᕐᓗᓄᔪ ᐅᓪᓚᕐᔪᑦ ᐅᓗᓪᓕᐊᓇᖅᑐᖅᑐᑦ ᐱᖅᘻᔪᑦ ᐊᖏᔪᓚᓐᓂ 2005. ᐱᓕᕆᑕᐅᑦᓗᐊᕆᖅᑐᑦ ᐅᓗᓪᓕᐊᓇᖅᑐᑎᕐᓂᓄᑦ ᓂᐸᖏᖅᑕᖅᓂᐅᔪᑦ ᐃᒧᓂ, ᓂᖏᕐᕄᖅᑕᖅᑎᐊᖅᓂᕐᓂᕐᓂ, ᓯᑐᖅᓇᖅᑐᑦ, ᑐᓗᖅᑐᓂᕐᒥᔪᑦ, ᐅᐃᒪᓴᖅᑕᐅᓂᕐᓂᓄᑦ, ᐊᓚᒪᓗ ᓂᑖᕐᕄᓂᕐᒥᔪᓄᑦ ᓄᓇᐊᓄᑦ ᒪᖅᑎᑕᐅᑎᓄᑦ. ᑎᑎᕖᖅᑕᐅᓂᔪ 2009, ᐅᓄᖅᓯᒡᓕᐊᑎᑕᐅᓇᓯᓂᕐᓂᓄᑦ ᑐᖅᑖᐅᔪᑦ ᐱᖅᑎᓯᒡᓗᑎᓄᑦ ᐱᖏᐅᓇᓯᑐᖅ ᖃᐅᔨᖏᐅᖅᓯᒡᓕᓪᓗᑎᓄᑦ ᐋᖅᓴᑦ ᒥᖏᓐᓄᑦ, ᐃᓂᒋᔭᖅᓂᕐᒥᑦ, ᐊᓚᒪᓗ ᐅᓗᓪᓕᐊᓇᑭᑕᐅᔪᑦ ᐊᓚᒪᓗ ᐊᑐᓕᖅᑎᑕᐅᓗᑎᓄᑦ ᐋᖅᑭᒡᓕᐊᑭᑕᐅᔪᖓᓂᑦ ᐅᓗᓪᓕᐊᖅᑎᓕᖅᑎᓄᑦ.

ᐊᐅᓚᕆᐊᖅᓂᕐᓗᑦ ᐸᖅᓇᐅᑎᑦ ᐊᖅᓴᖅᓄᑦ Northwest Atlantic blue whale ᐊᑐᓕᖅᑎᑕᐅᓂᐊᖅᑐᑦ ᖃᓄᓕᒥᐅᔪᖅᑐᑦ ᐊᓚᒪᓗ ᐊᑯᓂᕐᓯᔭᖅᑐᑦ ᐃᑲᔪᑕᐅᓇᓯᓗᑎᓄᑦ ᐅᓄᖅᓯᒡᓕᓪᓗᐊᖅᓇᖏᓚᐅᑎᓄᑦ ᐊᓗᓕᒥᐅᑕᓕᓚᖅᑯᓄᑦ ᐃᓚᒥᐅᑕᓕᖅᑎᑕᐅᓗᑎᓄᑦ ᐃᓚᒥᐅᑕᓕᖅᑎᑕᐅᓗᑎᓄᑦ ᐃᓕᓴᕐᓂᖅ. ᐅᓄᖅᓯᒡᓕᓪᓗᐊᖅᓱᔪᑕᐅᓇᓯᔭᖅᑐᑦ ᓯᔭᑕᐅᔪᑦ ᐱᑕᓚᐊᖓᓂᐊᖅᑐᑦ ᐃᒪᖅᒥᐅᑕᓕᓚᖅᑯᑦᓄᑦ ᐃᓚᒥᐅᑕᓕᖅᑎᑕᐅᓗᑎ

Species at Risk Act
Action Plan Series

Action Plan for the Blue Whale (*Balaenoptera musculus*), Northwest Atlantic Population, in Canada

Blue Whale, Northwest Atlantic Population



2019

Canada

Recommended citation:

Fisheries and Oceans Canada. 2019. Action Plan for the Blue Whale (*Balaenoptera musculus*), Northwest Atlantic Population, in Canada. *Species at Risk Act* Action Plan Series. Fisheries and Oceans Canada, Ottawa. iv + 21 pp.

For copies of the Action Plan, or for additional information on species at risk, including COSEWIC Status Reports, residence descriptions, recovery strategies, and other related recovery documents, please visit the Species at Risk Public Registry.

Cover illustration: Y. Morin, DFO

Également disponible en français sous le titre
«Plan d'action pour le rorqual bleu (*Balaenoptera musculus*), population de l'Atlantique Nord-Ouest, au Canada. »

© Her Majesty the Queen in Right of Canada, represented by the Minister of Fisheries and Oceans, 2019. All rights reserved.
ISBN
Catalogue no.

Content (excluding the illustrations) may be used without permission, with appropriate credit to the source.

Preface

The federal, provincial, and territorial government signatories under the Accord for the Protection of Species at Risk (1996) agreed to establish complementary legislation and programs that provide for the effective protection of species at risk throughout Canada. Under the *Species at Risk Act* (S.C. 2002, c.29) (SARA), the federal competent ministers are responsible for preparing action plans for species listed as extirpated, endangered, and threatened, and for which recovery has been deemed feasible. They are also required to report on progress five years after the publication of the final document on the Species at Risk Public Registry.

The Minister of Fisheries and Oceans is the competent minister under SARA for the Northwest Atlantic Blue Whale and has prepared this Action Plan to implement the Recovery Strategy, as per Section 47 of SARA. The minister responsible for the Parks Canada Agency is the competent minister for individuals in the waters of Forillon National Park. In preparing this Action Plan, the competent ministers have considered, as per Section 38 of SARA, the commitment of the Government of Canada to conserving biological diversity and to adhering to the principle that, if there are threats of serious or irreversible damage to the listed species, cost-effective measures to prevent the reduction or loss of the species should not be postponed for a lack of full scientific certainty. To the extent possible, this Action Plan has been prepared in cooperation with Parks Canada Agency and the Mingan Island Cetacean Study, as per Section 48(1) of SARA.

As stated in the preamble to SARA, success in the recovery of this population depends on the commitment and cooperation of many different constituencies that will be involved in implementing the directions and actions set out in this Action Plan and will not be achieved by Fisheries and Oceans Canada or any other jurisdiction alone. The cost of conserving species at risk is shared between different constituencies. All Canadians are invited to join in supporting and implementing this Action Plan for the benefit of the Northwest Atlantic Blue Whale and Canadian society as a whole.

Under SARA, an action plan provides the detailed recovery planning that supports the strategic direction set out in the recovery strategy for the species. The plan outlines recovery measures to be taken by Fisheries and Oceans Canada, Environment and Climate Change Canada and other jurisdictions or organizations to help achieve the population and distribution objectives identified in the Recovery Strategy. Implementation of this Action Plan is subject to the appropriations, priorities, and budgetary constraints of the participating jurisdictions and organizations.

Acknowledgments

Fisheries and Oceans Canada would like to thank Andréanne Demers, Hugues Bouchard, Sarah Larochelle and Charline Le Mer, who drafted this Action Plan, as well as all those in the Department who supported its development. The Department also wishes to acknowledge the contributions of Parks Canada, as well as Richard Sears and Christian Ramp of the Mingan Island Cetacean Study.

Executive summary

The Northwest Atlantic Blue Whale was listed as Endangered under the *Species at Risk Act* in 2005. The main threats to its recovery are anthropogenic noise, lack of food availability, contaminants, collisions, disturbances and entanglements. Published in 2009, the Recovery Strategy proposed three recovery objectives intended to increase knowledge of the population, its habitat, and threats, and implement measures to mitigate threats.

The Action Plan for the Northwest Atlantic Blue Whale presents measures that will be implemented in the short and medium term to assist in meeting the recovery objectives. The first set of measures will be taken by Fisheries and Oceans Canada. They consist primarily of research aimed at estimating the size of the Northwest Atlantic Blue Whale population and its use of Canadian waters. The measures also include adopting or enforcing legislation or policies to protect Blue Whale habitat and mitigate threats. The second set of measures will be implemented by the Department in partnership with the various stakeholders involved in the Blue Whale's recovery. They include research on krill and the use of hydroacoustics to document the presence of these whales. The third set of measures consists of actions that concerned stakeholders will be able to take voluntarily. They include observations and photo-identification of Blue Whales and outreach and awareness initiatives.

The measures set out in the Action Plan could affect some stakeholders such as non-governmental organizations or the shipping industry. However, their implementation would not necessarily result in incremental costs to these stakeholders. Canadian society as a whole would benefit from the implementation of the Action Plan, given the economic value that Canadians attach to the recovery of the species and the protection of its habitat.

Table of Contents

Preface i
Acknowledgments ii
Executive summary iii
1. Recovery actions 1
1.1 Context and scope of the Action Plan 1
1.2 Measures to be taken and Implementation Schedule 3
1.3 Critical habitat 17
1.3.1 Critical habitat identification 177
2. Socioeconomic assessment 177
2.1 Stakeholder profile 177
2.2 Socioeconomic costs of implementing the Action Plan 188
2.3 Benefits of implementing the Action Plan 199
2.4 Distributional impacts 20
3. Measuring progress 20
4. References 21
Appendix A: Effects on the environment and other species 222
Appendix B: Record of cooperation and consultation 233

1. Recovery actions

1.1 Context and scope of the Action Plan

The Blue Whale (*Balaenoptera musculus*) is a baleen whale that uses coastal and offshore Atlantic Canadian waters mainly in the summer (Figure 1), to feed primarily on euphausiids, commonly known as krill. The Blue Whale population in the Northwest Atlantic was designated as Endangered by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) in May 2002. This population was reassessed by COSEWIC in 2012 and its status confirmed. It was listed as Endangered under the *Species at Risk Act* (SARA) in January 2005. Commercial whaling historically carried out in the Atlantic Ocean reduced the population by about 70%; at least 11,000 Blue Whales were killed before the 1960s including at least 1,500 animals in eastern Canadian waters. The size of the Northwest Atlantic population is currently unknown, but experts estimate that the number of mature animals is unlikely to exceed 250 individuals.

The Recovery Strategy for the Northwest Atlantic Blue Whale (Beauchamp et al. 2009) presents the various threats facing the population. In addition to historical whaling and natural sources of mortality such as ice entrapment and predation, several threats have been identified and their general level of concern assessed. This level of concern depends on the potential impact of the threat on the population, knowledge on the threat, and its extent within the distribution range. The main threats to the recovery of the Northwest Atlantic Blue Whale population were determined by experts to be anthropogenic noise, which causes a degraded underwater acoustic environment and alters behaviour, and the lack of food resources, which could result from ecosystem changes caused in particular by climate change.

Contaminants, vessel collisions, disturbances caused by whale watching activities, entanglements in fishing gear, epizootics, toxic algal blooms and toxic spills also pose a threat to the Blue Whale. Based on the best available information when the recovery strategy was published in 2009, these threats were assessed as presenting a low level of concern, either because their impact was considered low, very localized or difficult to quantify or because the threats were considered potential. Based on new data from photo-identification (Gaspard et al. 2017), the impact of the threat of entanglement in fishing gear needs to be reassessed, as it may be more significant than previously thought. Given the small size of the Blue Whale population, even activities that affect a small number of individuals can have a significant impact on the species' survival in the Atlantic Ocean.

The long-term goal of this Recovery Strategy is to reach a total of 1,000 mature individuals. To reach this recovery goal, three objectives were set for the Canadian range:

Objective 1: Define and conduct a long-term assessment of the size, structure and trends of the Northwest Atlantic Blue Whale population, and determine their range and critical habitat within Canadian waters;

Objective 2: Implement control and monitoring measures for activities that could hinder the recovery of the Blue Whale in its Canadian range;

Objective 3: Increase knowledge of the main threats to the recovery of the Blue Whale in Canadian waters both to determine their true impact and to identify effective measures to mitigate the negative consequences for the population's recovery.

The Recovery Strategy is proposing recovery approaches based on three broad strategies: research and monitoring, conservation, awareness and education. The purpose of this Action Plan is to outline priority actions to meet the above recovery objectives, following the same broad strategies. These actions are related to all the threats described in the Recovery Strategy and the full range of the population in Atlantic Canada's waters. The Recovery Strategy provides more details on the strategies and approaches for recovering Northwest Atlantic Blue Whales, on studies to identify their critical habitat, and on their biology (Beauchamp et al. 2009).

Under Section 47 of SARA, the competent minister must prepare one or more action plans based on the Recovery Strategy. Action planning for species at risk recovery is therefore an iterative process. The Implementation Schedule in this Action Plan may be modified in the future depending on the level of progress made towards recovery.




Figure 1. Geographic range and concentration areas (darker blue) of the Blue Whale in the Northwest Atlantic Ocean.

1.2 Measures to be taken and Implementation Schedule

Success in the recovery of this population is not dependent on the actions of any single jurisdiction; rather, it requires the commitment and cooperation of the many different constituencies involved in implementing the directions and actions set out in this Action Plan.

This Action Plan outlines measures that maximize the likelihood of achieving the population and distribution objectives for the Northwest Atlantic Blue Whale. It also includes measures to address threats to the population and monitor its recovery, in order to guide not only activities to be undertaken by Fisheries and Oceans Canada (DFO), but also those in which other jurisdictions, organizations and individuals have a role to play. DFO strongly encourages all Canadians to participate in the conservation of the Blue Whale by supporting and implementing the priority recovery measures outlined in this Action Plan. DFO recognizes the important role of key players in the field in the implementation of recovery measures for this population.

The following three tables are not intended to present all the measures that could be taken to foster the Blue Whale's recovery; instead, they target measures likely to be taken

in the medium term (within the next 5 to 10 years). These measures are technically and financially feasible and will help identify measures that could be implemented in the longer term based on the knowledge that will be acquired. Given that the Northwest Atlantic Blue Whale has a vast range, the proposed research focuses on targeted areas to optimize results and efficiency. Much of this research could help support future critical habitat identification.

Table 1. Implementation Schedule: Measures to be undertaken by Fisheries and Oceans Canada

Table 1 identifies measures to be taken by DFO to support the recovery of the Blue Whale. This table primarily presents research and monitoring measures that will have to be led by DFO, particularly because of the complex logistics involved in monitoring a Blue Whale population with such a vast range. This research is a continuation or logical extension of studies already performed by DFO as part of the Recovery Strategy and as part of its mandate for marine mammal conservation. For an overview of the research activities and conservation and awareness measures carried out from 2009 to 2014, see the Report on the Progress of Recovery Strategy Implementation for the Blue Whale, Northwest Atlantic Population (Fisheries and Oceans Canada 2016). The measures listed in Table 1 as well as Tables 2 and 3 address the three recovery objectives:

Objective 1: Increase knowledge of the population and its size (Measures 1 to 3): A better understanding of the high-concentration areas of Blue Whales will make it possible to carry out targeted surveys to assess the population size more effectively; this research will also make it possible to understand the reasons why Blue Whales are attracted to certain locations. It is being conducted to meet both Objective 1 of the Recovery Strategy and the Schedule of Studies to identify critical habitat included in the Recovery Strategy. The studies on high-concentration areas and their characteristics, such as high concentrations of krill, could lead to the identification of Blue Whale critical habitat, at least partially, in an update of the Recovery Strategy.

Objective 2: Mitigate threats (Measures 7 to 10): Fisheries and Oceans Canada will also implement conservation measures to protect the Blue Whale's food resources, mitigate the risk of collisions and protect the Blue Whale from disturbances through policies and regulations.

Objective 3: Develop a better understanding of threats (Measures 4 to 6): The impact of threats, such as noise, disturbance or entanglement, which can alter the behaviour of Blue Whales and thereby adversely affect their recovery, are still poorly understood. Studies on noise sources and the impact of disturbance and entanglement on the recovery of the population will make it possible to implement suitable mitigation measures and improve on existing mitigation measures.

The measures presented in Table 1 are in addition to the implementation of DFO's mandate and enforcement of laws in effect. The Blue Whale is protected under SARA, which prohibits anyone from harming or harassing an endangered species. The Blue Whale’s needs must also be taken into account in the environmental assessments of various projects under the *Fisheries Act,* the *Canadian Environmental Assessment Act,*

2012 or the *National Energy Board Act*. Mitigation measures can also be included in authorization conditions issued under various types of legislation enforced by the Department. Environmental assessments of projects submitted to federal-provincial offshore petroleum boards are also reviewed by DFO to ensure that species at risk are taken into account. Moreover, scientific research protocols on the Blue Whale are reviewed to minimize disturbances. By enforcing legislation, other government departments and agencies also contribute to Blue Whale recovery. For example, Parks Canada is implementing the *Marine Activities in the Saguenay–St. Lawrence Marine Park Regulations*.

Table 1. Measures to be undertaken by Fisheries and Oceans Canada

#	Recovery Measures	Priority[1]	Threats or Objectives Addressed[2]	Timeline
Broad Strategy: *Research and monitoring*				
1	Continue delineating high-density seasonal areas in Canadian waters using satellite transmitters, especially in southwestern Newfoundland and on the Scotian Shelf. Transmitters have been put on several Blue Whales since 2010 and data have been used to define feeding areas and migratory routes. This research will identify locations where the Blue Whale is found in order to assess their numbers through targeted surveys.	High	Objective 1	5 years
2	Conduct targeted surveys in high-density areas to assess the population size. Once high-density areas have been identified, aerial or vessel-based surveys can focus on these areas and thereby be more effective in estimating the number of Blue Whales.	High	Objective 1	5 years
3	Assess the extent to which the biological processes (krill aggregations) and physical processes (currents, tides) affect the distribution, behaviour, and migrations of the Blue Whale.	High	Objective 1	5 years

[1] **Priority**: Reflects the degree to which the measure contributes directly to the recovery of the population or is an essential precursor to a measure that contributes to the recovery of the population.

- "High" priority measures are considered those most likely to have an immediate or direct impact on the achievement of the recovery objective for the population.
- "Medium" priority measures may have a less immediate or direct impact on the achievement of the population and distribution recovery objectives, but are still important for the recovery of the population.
- "Low" priority recovery measures will likely have an indirect or gradual impact on the achievement of the recovery objectives. They are nonetheless considered important contributions to the knowledge base or public involvement and to the value the public ascribes to the species.

[2] **Threats or objectives addressed**: Indicates which recovery objective the measure addresses or which threat it mitigates. Summary of the objectives: Objective 1: Increase knowledge of the population and its size; Objective 2: Mitigate threats; Objective 3: Develop a better understanding of threats.

#	Recovery Measures	Priority[1]	Threats or Objectives Addressed[2]	Timeline
4	Study the Blue Whale's behavioural responses to various noise sources in various contexts. The Blue Whale's tolerance to noise must be better understood in order to develop appropriate mitigation measures for noise-generating human activities.	Medium	Objective 3 Noise	10 years
5	Study the behavioural response of Blue Whales to various approach distances of boats (whale watching, research, and recreational boats). This knowledge will help to establish acceptable approach distances for the Blue Whale.	Medium	Objective 3 Disturbance	2 years
6	Assess the fishing effort and characterize the fishing gear used in areas occupied by the Blue Whale to determine the potential impact of the threat of gear entanglement on the population.	Low	Objective 3 Entanglement	10 years
Broad Strategy: *Conservation*				
7	Assess whether any commercial fishery of krill would affect the integrity of the ecosystem or the energy needs of a recovering or a recovered Blue Whale population, in accordance with the Policy on New Fisheries on Forage Species.	High	Objective 2 Availability of food	Ongoing
8	Adopt and implement the amendments to the *Marine Mammal Regulations* to help protect the Blue Whale from anthropogenic disturbances in all of the areas that they occupy. The amended regulations include approach distances for species at risk like the Blue Whale in order to reduce disturbance.	High	Objective 2 Disturbance Collisions	Ongoing
9	Determine the best management tools for achieving the St. Lawrence Estuary Area of Interest's conservation objectives and implement them, including the designation of a marine protected area.	High	Objective 2	5 years
10	By enacting regulations, designate a marine protected area in the American Bank area located off the Gaspé Peninsula. This site is, among other things, considered as a high-density Blue Whale area. Examples of potential conservation measures would be voluntary measures and rules of ethics to regulate marine observation activities and reduce disturbances, as well as	Medium	Objective 2 Disturbance Entanglement	2 years

#	Recovery Measures	Priority[1]	Threats or Objectives Addressed[2]	Timeline
	apply fishery measures aimed at protecting forage species, such as krill, the Blue Whale's main prey.		Collisions	
Broad Strategy: *Outreach*				
11	Encourage the whale watching public to report their observations. For example, the "Tell Jack" outreach campaign encourages the whale watching public in Newfoundland and Labrador to play a role in DFO marine mammal science and promote DFO's research on marine mammals including the Blue Whale. Anecdotal reports and pictures sent to the marine mammal research group can provide useful information. The campaign uses social media, proactive media relations, and public outreach activities to solicit observation reports and pictures from the public via Twitter and email.	Medium	Objective 1	Ongoing
12	Carry out outreach and educational activities intended for whale watching operators, the whale watching public, and other mariners about best practices for observing marine mammals including the Blue Whale. For example, in Newfoundland and Labrador, school visits and public events such as Oceans Day are organized using life-size Blue Whale tails made of fabric. The tails could be used to educate people on the Blue Whale's biology, behaviour, and lifecycle, as well as its SARA status.	Medium	Objective 2 Disturbance	Ongoing

Table 2. Measures to be undertaken collaboratively between Fisheries and Oceans Canada and its partners

Table 2 shows the actions that will be taken by Fisheries and Oceans Canada and its partners working in collaboration. The order in which partners are listed in the table is not indicative of their contribution or degree of involvement. Implementation of these measures will be dependent on a collaborative approach, in which Fisheries and Oceans Canada is a partner in recovery efforts, but cannot implement the measures alone. Table 2 presents research and monitoring measures that DFO is implementing in close partnership with universities, other government agencies, the private sector and research organizations. The purpose of these measures is to:

1. Continue the characterization of noise sources and levels and the Blue Whale's exposure to noise in the areas they use the most (measure 13).
2. Study the Blue Whale's primary food source, krill (Measures 14 to 16). DFO intends to continue its research on krill production in the Estuary and Gulf of St. Lawrence, and on krill interaction with Blue Whales, in collaboration with universities.
3. Increase knowledge of the Blue whale distribution in areas outside the Estuary and Gulf of St. Lawrence, for which few observational data are available (Measures 17 to 22). DFO and several partners intend to use hydroacoustics and photo-identification for this purpose. These measures also support the schedule of studies intended to identify critical habitat set out in the Recovery Strategy.
4. Better assess the effect of the threat of entanglement on the health and recovery of the Blue Whale (measure 23).

This table also proposes several partnerships to implement conservation measures and thereby mitigate threats to Blue Whale recovery. Certain initiatives are underway or are planned for the near future to mitigate threats, such as noise (Measure 24), toxic spills (Measure 25), vessel collisions (Measures 26 and 27) and entanglements (Measure 28). Two awareness campaigns are also underway to improve the data collected by observers (Measure 29) and to raise awareness among pleasure boaters around the Saguenay–St. Lawrence Marine Park in order to reduce disturbances in this important area for the Blue Whale (Measure 30).

Table 2. Collaborative measures between Fisheries and Oceans Canada and its partners.

Acronyms: DFO (Fisheries and Oceans Canada), NGO (Non-Governmental Organizations), NOAA (National Oceanographic and Atmospheric Administration, US).

#	Recovery Measures	Priority[3]	Threats or Objectives Addressed[4]	Timeline	Partnerships[5]
Broad Strategy: *Research and monitoring*					
13	Continue characterizing of noise sources and levels, and analyzing the exposure of Blue Whales to noise in the areas they use most often in Canadian Atlantic waters.	Medium	Objective 3 Noise	Ten years	DFO Transport Canada Industry
14	Study krill distribution, population dynamics and production processes. Tides, currents, and krill behaviour will determine the aggregation areas essential for effective Blue Whale feeding. A better understanding of these factors could be integrated into critical habitat identification.	High	Objective 3 Availability of food	3 years	DFO Universities Research institutes
15	Study the Blue Whale's energy needs to estimate the krill biomass necessary to support the current population and eventually a recovering population.	High	Objective 3 Availability of food	3 years	DFO Universities Research institutes
16	Study the trophic interactions between the Blue Whale and krill to try to explain the high inter-annual and inter-regional variability of Blue Whale occurrences and residence time.	High	Objective 3 Availability of food	3 years	DFO Universities Research institutes

[3] See footnote 1.
[4] See footnote 2.
[5] **Potential partners**: Université du Québec à Rimouski, Dalhousie University, Institut des sciences de la mer à Rimouski, Group for Research and Education on Marine Mammals (GREMM), Mingan Island Cetacean Study, Réseau d'observation des mammifères marins, Indigenous groups, Shipping Federation of Canada, whale watching tour operators, emergency response networks, etc.

#	Recovery Measures	Priority[3]	Threats or Objectives Addressed[4]	Timeline	Partnerships[5]
17	Use hydroacoustic techniques to monitor the occurrence and number of Blue Whales in Atlantic Canada's waters, especially on the Scotian Shelf and south of Newfoundland. Bottom-moored recorders will make it possible to collect data on all marine mammal species that vocalize in a specific location.	High	Objective 1	2 to 5 years	DFO Universities Research institutes
18	Acquire data on Blue Whale distribution and abundance outside the Gulf of St. Lawrence (Cabot Strait and Scotian Shelf) where very little information is available, using photo-identification and the installation of satellite transmitters.	High	Objective 1	5 years	NGO DFO Universities
19	Conduct necropsy on dead Blue Whales whenever possible and follow protocols for data collection and sharing.	High	Objective 1 Objective 3	Ongoing	NGO DFO Universities
20	Continue collecting tissue samples and conducting biopsies to assess the population structure, pregnancy rates, and level of contaminants, especially in the Gulf of St. Lawrence, the Cabot Strait and the Scotian Shelf.	Medium	Objective 1 Objective 3	5 years	NGO Universities DFO
21	Better integrate all Blue Whale sightings data collected by various sources.	Low	Objective 1	5 years	NGO Universities Research institutes DFO
22	Establish international research partnerships to increase understanding of the Blue Whale's distribution and migration routes. Sharing hydroacoustic data is an example of collaboration.	Low	Objective 1	10 years	DFO NOAA Universities

#	Recovery Measures	Priority[3]	Threats or Objectives Addressed[4]	Timeline	Partnerships[5]
23	Use photo-identification to monitor entanglement rates in Blue Whales in order to better assess the impact of this threat on the health and recovery of the population.	Medium	Objective 2 Entanglement	In progress	NGO DFO
Broad Strategy: *Conservation*					
24	Study and implement measures to reduce the negative impact of noise caused by human activities, such as shipping, construction, and seismic exploration.	High	Objective 2 Noise	Ongoing	DFO Transport Canada Industry Parks Canada Universities NGO
25	Develop and implement response plans to reduce impacts to Blue Whales and their habitat likely to be caused by toxic spills. The government's initiative to strengthen the tanker safety system includes improving response planning in the event of toxic spills in targeted areas such as the Gulf of St. Lawrence, a high-risk area for spill impacts.	Medium	Objective 2 Spills	5 years	Provincial and federal departments Industry NGO
26	Maintain the Marine Mammal Response Program in the various regions of Canada's Atlantic coast. This program supports organizations that maintain call centres and databases, and that intervene when a marine mammal is in distress. It also supports training for emergency responders.	Medium	Objective 2 Entanglement Collisions	Ongoing	NGO DFO Indigenous groups
27	Study how to reduce the risk of vessel collisions with Blue Whales in the St. Lawrence Estuary and Gulf of St. Lawrence beyond existing voluntary measures for speed reduction, with the goal of identifying measures to implement (e.g., moving shipping lanes, establishing no-go areas).	Medium	Objective 2 Collisions	5 years	DFO Transport Canada Industry NGO

#	Recovery Measures	Priority[3]	Threats or Objectives Addressed[4]	Timeline	Partnerships[5]
28	Develop projects to promote the recovery of ghost or lost fishing gear in the Estuary and Gulf of St. Lawrence	Low	Objective 2 Entanglement	5 years	DFO Industry NGO
Broad Strategy: *Outreach*					
29	Continue training observers to improve marine mammal identification and information gathering. Training is intended for all those who have the opportunity to gather observational information (researchers, bird watchers, observers on platforms).	Medium	Objective 1	Ongoing	DFO NGO Industry
30	Educate pleasure boaters and captains of whale watching excursions on the impacts their activities have on Blue Whales near the Saguenay-St. Lawrence Marine Park. In collaboration with fisheries officers, the goal of this project is to educate users on the appropriate behaviour to adopt outside the marine park. It could be extended to other areas where there is a high incidence of disturbance by pleasure boaters and whale watchers.	Medium	Objective 2	5 years	DFO Parks Canada NGO

Table 3. Measures that represent opportunities for other jurisdictions, organizations or individuals to lead

As all Canadians are invited to join in supporting and implementing this Action Plan for the benefit of the Blue Whale and Canadian society as a whole, Table 3 identifies measures likely to support the recovery of the Northwest Atlantic Blue Whale that could be taken voluntarily by other jurisdictions, institutions, groups, and individuals interested in supporting the species' recovery. The order in which partners are listed in Table 3 is not indicative of their contribution or degree of involvement. The research and monitoring measures presented in Table 3 are activities that could be implemented by non-governmental organizations, particularly the Mingan Island Cetacean Study (MICS). These measures involve continuing Blue Whale monitoring activities through photo-identification, hydroacoustics, tissue sample collection, and biopsies (Measures 31 to 36) in Canadian and international waters. Several organizations, including Parks Canada, are already involved in efforts to raise marine user awareness of the impact their activities have on the Blue Whale, and these efforts are expected to continue (Measures 37 and 38).

If your organization is interested in participating in one of these measures, please contact the Species at Risk Management Division in Quebec at lep-sara-qc@dfo-mpo.gc.ca.

Table 3. Measures that represent opportunities for other jurisdictions, organizations or individuals to lead.

#	Recovery Measures	Priority[6]	Threats or objectives addressed[7]	Partnerships[8]
Broad Strategy: *Research and monitoring*				
31	Continue photo-identification activities, especially in various areas of the Gulf of St. Lawrence, and analyze photos to continue identifying individuals. These analyses help to better understand the Blue Whale's annual use of Canadian waters and assess its abundance. The continuation of these activities implies maintaining a database of observations and photos.	High	Objective 1	NGO Indigenous groups
32	Conduct acoustic monitoring (recording vocals using moored recorders) in various sectors of the Gulf of St. Lawrence and Atlantic Canada that are difficult to access in order to better characterize their use by the Blue Whale.	High	Objective 1	NGO Universities Research institutes
33	Conduct an analysis of the Northwest and Northeast Atlantic photo-identification catalogues to verify whether there are connections between these two areas of the Atlantic Ocean.	Medium	Objective 1	NGO Universities Research institutes
34	Biopsy Blue Whales to monitor their contaminants levels. The accumulation of contaminants in the Blue Whale is barely known and biopsies could make it possible to identify contaminants found in tissues and monitor their evolution over time.	Medium	Objective 3 Contaminants	NGO Universities Research institutes
35	Conduct genetic analyses of tissues collected from individuals that died in spring 2014 in southern Newfoundland. The comparison of these analyses with biopsies carried out across the North Atlantic will allow for a better understanding of the extent of exchange among the whales in the various regions.	Low	Objective 1	NGO Universities Research institutes

[6] See footnote 1.
[7] See footnote 2.
[8] See footnote 5.

#	Recovery Measures	Priority[6]	Threats or objectives addressed[7]	Partnerships[8]
36	Conduct a survey off Mauritania in the winter to determine whether the Northwest Atlantic Blue Whale uses the waters off West Africa to breed.	Low	Objective 1	NGO Universities Research institutes
Broad Strategy: *Outreach*				
37	Continue raising awareness among marine users of the impact their activities have on the Blue Whale (marine observation activities, commercial shipping, and pleasure boaters).	Medium	Objective 2	Parks Canada Transport Canada NGO Industry Indigenous groups
38	Continue mandatory training for all captains and kayaking guides who conduct their activities in the Saguenay-St. Lawrence Marine Park to familiarize them with the best practices for observing marine mammals (marine park regulations, biology, and ways to diversify excursions).	Medium	Objective 2	Parks Canada

1.3 Critical habitat

1.3.1 Critical habitat identification

The Act requires an action plan which or that includes an identification of critical habitat to the extent possible. When published in 2009, the Recovery Strategy for the Northwest Atlantic Blue Whale included a schedule of studies to identify critical habitat. Several such studies have been carried out since the publication of the Recovery Strategy, and others are ongoing. These studies are summarized in the Report on the Progress of Recovery Strategy Implementation for the Blue Whale, Northwest Atlantic population. The results were presented and peer-reviewed at a DFO Science Advisory Meeting in winter 2016. A Science Advisory Report was published recently (Fisheries and Oceans Canada 2018) and will be used to support the identification of Blue Whale critical habitat, to the extent possible. This identification will be presented in an update of the Recovery Strategy, rather than in this Action Plan.

2. Socioeconomic assessment

The *Species at Risk Act* requires that an Action Plan include an assessment of the socioeconomic costs associated with its implementation and of the benefits to be derived from the implementation (SARA 49(1)(*e*), 2002). This assessment addresses only the incremental (new) socioeconomic costs associated with the implementation of this Action Plan at the national level as well as the social and environmental benefits of implementing it in its entirety, recognizing that not all aspects of its implementation are under the jurisdiction of the federal government. It is intended to inform the public and guide partners in their decision-making on the implementation of the Action Plan.

This assessment will first identify the main stakeholders that could be affected by or involved in the implementation of the recovery measures listed in Tables 1 to 3 of the Action Plan. Section 2.2 then examines whether these measures could involve incremental costs to stakeholders. Lastly, Section 2.3 presents an overview of the benefits of implementing the Action Plan.

2.1 Stakeholder profile

The Blue Whale recovery measures set out in Tables 1 to 3 are grouped into three types of broad strategies: research and monitoring, conservation, and outreach. The types of stakeholders that would take part in the implementation of the Action Plan are also identified in these tables.

Research and monitoring

DFO's main partners in carrying out the research and monitoring activities are universities (e.g., University of Quebec at Rimouski, Dalhousie University), governmental (e.g., Transport Canada) and non-governmental (e. g., Mingan Island Cetacean Study) organizations.

Conservation

The implementation of conservation measures would involve a number of stakeholders, including the federal and provincial governments, non-governmental organizations, and the private sector. Several private sector industries could be affected by the implementation of conservation measures. In particular, noise reduction measures could have repercussions for industries whose activities involve shipping, marine construction, and seismic exploration (Table 2 – Measure 24). Initiatives to enhance the security systems of tankers and reduce the risk of collision between Blue Whales and ships may also have an impact on the shipping industry (Table 2 – Measures 25 and 27).

Outreach

Outreach activities would be conducted primarily by the federal government in collaboration with non-governmental organizations and the whale watching industry.

2.2 Socioeconomic costs of implementing the Action Plan

Many of the measures identified in the Action Plan are initiatives underway within the federal government and its partners, and these measures are expected to continue even in the absence of the Action Plan. Although the measures set out in the Action Plan could affect some stakeholders identified above, their implementation would not systematically result in incremental socioeconomic costs to these stakeholders.

There is not enough information to quantify the incremental socioeconomic costs that could result from implementing the Action Plan for the Blue Whale. Therefore, the potential costs of the Action Plan are evaluated qualitatively because most of the available information is qualitative.

Research and monitoring

Of the 38 recovery measures in the action plan, 23 measures involve research and monitoring activities. Many of these research projects are an extension of projects already being carried out by DFO and its partners. Certain projects go beyond the scope of the Blue Whale Recovery Strategy and include the acquisition of knowledge that can be applied to several species. It is therefore realistic to think that many of the research activities listed in Tables 1 to 3 would be carried out by DFO and its partners, even in the absence of the Action Plan.

DFO-led research and monitoring would be funded through the Department's regular programs and would not mean incremental costs to DFO. Measures undertaken by other organizations (universities, NGOs, research institutes) could be funded in part by existing federal government programs. However, additional costs could be incurred by local and regional stakeholders who chose to become involved in Blue Whale recovery efforts.

There is not enough information at this stage to quantify these costs, but they would vary depending on the extent of the research activities undertaken.

Conservation

The Action Plan identifies eight conservation measures to implement mitigation and monitoring measures for activities that could disrupt the recovery of the Blue Whale in its Canadian range. Most of these measures fall within the framework of initiatives that are already underway within the federal government; the implementation of these measures therefore will not involve any incremental costs to the government.

One of the conservation measures aimed at reducing the risk of collisions between the Blue Whale and ships in the St. Lawrence Estuary (Table 2 – Measure 27) could lead to additional costs for the shipping industry. As details on the implementation of these measures are not known, it is not possible to estimate the incremental costs, if any, to the shipping industry.

The implementation of conservation measures is not expected to generate any additional costs to the other stakeholders.

Outreach

The awareness activities included in the Action Plan are all activities that are currently underway and are intended to protect several species of marine mammals. Therefore, realistically, these activities would be carried out even in the absence of the Action Plan for the Blue Whale. Consequently, awareness activities are not expected to generate incremental costs to the federal government or any of the stakeholders.

2.3 Benefits of implementing the Action Plan

The implementation of the measures outlined in this Action Plan will contribute positively to the achievement of the long-term goal of the Northwest Atlantic Blue Whale Recovery Strategy, which is to reach a total of 1,000 mature individuals in the population.

The benefits of the recovery of the Blue Whale are difficult to quantify. However, the Act recognizes that “wildlife, in all its forms, has value in and of itself and is valued by Canadians for aesthetic, cultural, spiritual, recreational, educational, historical, economic, medical, ecological and scientific reasons” (SARA 2002). Self-sustaining and healthy ecosystems with their various elements in place, including species at risk, contribute positively to the livelihoods and the quality of life of all Canadians. A review of the literature confirms that Canadians value the preservation and conservation of species in and of themselves. Measures taken to preserve a species, such as habitat protection and restoration, are also valued. In addition, the more measures contribute to the recovery of a species, the higher the value the public ascribes to such measures (Loomis and White 1996; Fisheries and Oceans Canada 2008).

Specifically, a study estimating the economic benefits of marine mammal recovery in the St. Lawrence Estuary reveals that Canadians would be willing to pay $229 annually per household for a multi-species recovery strategy resulting in a measurable improvement in the status of species at risk for a number of marine mammals, including the Blue Whale (Boxall et al. 2012).

The implementation of the Action Plan should also generate benefits beyond the recovery of the Blue Whale. The acquisition of knowledge and the development of conservation measures should benefit several other marine mammal species.

2.4 Distributional impacts

Many different stakeholders will be involved in implementing the recommendations set out in this Action Plan. Given that most of the measures set out in the plan relate to existing programs and are a continuation of activities already underway, the incremental costs to DFO and its partners should be minimal.

The benefits of implementing the Blue Whale Action Plan will be enjoyed by the Canadian society as a whole, given the economic value that Canadians attach to the recovery of the species and the protection of its habitat.

3. Measuring progress

The recovery objectives presented in the Recovery Strategy propose a method for defining and measuring progress made toward achieving population and distribution objectives.

A report on the implementation of the Action Plan (under s. 55 of SARA) will be prepared to assess the progress made towards the implementation of the recovery measures.

A report on the ecological and socioeconomic impacts of the Action Plan (under s. 55 of SARA) will be prepared to provide information on the monitoring of the species' recovery and its long term viability, and on the implementation of the Action Plan.

4. References

Beauchamp, J., Bouchard, H., de Margerie, P., Otis, N., and J.-Y. Savaria. 2009. Recovery strategy for the Blue Whale (*Balaenoptera musculus*), Northwest Atlantic population, in Canada. *Species at Risk Act* Recovery Strategy Series. Fisheries and Oceans Canada, Ottawa.

Boxall, P.C., Adamowicz, W.L., Olar, M., West, G.E., and G. Cantin. 2012. Analysis of the Economic Benefits Associated with the Recovery of Threatened Marine Mammal Species in the Canadian St. Lawrence Estuary. Marine Policy. 36 (1):189-197.

Fisheries and Oceans Canada. 2008. Estimation of the Economic Benefits of Marine Mammal Recovery in the St. Lawrence Estuary. Policy and Economics Regional Branch, Quebec 2008.

Fisheries and Oceans Canada. 2016. Report on the Progress of Recovery Strategy Implementation for the Blue Whale (*Balaenoptera musculus*), Northwest Atlantic population, in Canada for the Period 2009 – 2014. *Species at Risk Act* Recovery Strategy Report Series. Fisheries and Oceans Canada, Ottawa.

Fisheries and Oceans Canada. 2018. Identification of important habitats for the Blue Whale in the western North Atlantic. DFO Can. Sci. Advis. Sec., Sci. Advis. Rep. 2018/003.

Gaspard, D., C. Ramp, J. Delarue, S. Landry and R. Sears. 2017. Entanglement threat has been underestimated for three rorqual species in the Gulf of St. Lawrence, Canada. Poster presented at the 22nd Biennial Conference on the Biology of Marine Mammals, October 22-27, 2017, Halifax, NS, Canada.

Loomis, J.B., and D.S. White. 1996. Economic Benefits of Rare and Endangered Species: Summary and Meta-Analysis. Ecological Economics. 18: 197-206.

Appendix A: Effects on the environment and other species

A Strategic Environmental Assessment (SEA) is conducted on all SARA recovery planning documents under SARA and in accordance with the Cabinet Directive on the Environmental Assessment of Policy, Plan and Program Proposals. The purpose of an SEA is to incorporate environmental considerations into the development of public policies, plans, and program proposals to support environmentally sound decision-making and to evaluate whether the outcomes of a recovery planning document could affect the environment in any way or the achievement of any of the Federal Sustainable Development Strategy's goals or targets.

Recovery planning is intended to benefit species at risk and biodiversity in general. However, it is recognized that implementation of action plans may inadvertently have environmental impacts beyond the intended benefits. The planning process based on national guidelines directly incorporates consideration of all environmental impacts, with a particular focus on the potential impacts on non-target species or habitats. The results of the SEA are incorporated directly into the action plan itself, but are also summarized below in this statement.

The threats affecting the Blue Whale also weigh on several marine mammal species that share its range. Measures aimed at reducing the impact of threats to the Blue Whale should therefore also benefit these species. Research intended to better understand krill production and behaviour and could lead to conservation measures will also be positive for all levels of the food chain.

Appendix B: Record of cooperation and consultation

This document was prepared with the participation of DFO researchers working on the Blue Whale. Representatives of the Saguenay-St. Lawrence Marine Park also participated, because the species is often present within its boundaries. The expertise of the Mingan Island Cetacean Study, which has been collecting observational data on whales for over three decades and which acts as curator of the Blue Whale photo-identification catalogue, was also solicited.

The document has been reviewed by the different divisions in the five relevant DFO administrative regions and by national headquarters. It was also sent for comment to provincial governments and Indigenous groups present within the range of this Blue Whale population. The non-governmental organizations, the shipping and fishing industries and the commercial marine life observations activities were also consulted.


ᐃᓕᓴᕐᓂᐊᕐᓂᖅ ᑐᓂᔨᐅᓂᒃᑯᑦ

ᓄᓇᕗᒻᒥ ᐅᒪᔨᓕᓇᖅᑐᖃᐃᓐᓇᑦ ᑲᑎᒪᔨᖅᑭᓐᓄᑦ ᐊᒻᒪᓗ ᓄᓇᕕᒻᒥ ᐃᒪᖃᐅᑉᑕᖅᓯᒃ ᐅᒪᒪᔨᖅᑐᖃᐃᓐᓇᑦ ᑲᑎᒪᔨᖅᑭᓐᓄᑦ ᐅᓄᓇᓕᒃ

ᑐᙵᕆᔭᖅ: ᐊᖅᑭᑦᑕᐅᔪᖅ: X ᐃᓱᒪᓕᐅᕐᑕᐅᔪᖅ: X

ᐃᓱᒪᓗᐊᑎ:2020/21 ᑲᑎᒪᓂᖅ ᑭᖓᔪᓕᑦᑕᐅᔪᓐᓇᖅᑐᑦ ᐅᐊᓐᓇᖅᑕᓱᒻᒥ (*Pandalus borealis*) ᐊᒻᒪᓗ Striped (*Pandalus montagui*) ᑭᖓᔪᓕᐊᐃᑦ ᑲᓇᓐᓇᖅᑕᓱᒻᒥ ᐱᖓᓐᓇᖅᑕᓱᒻᒥᓗ ᖃᐅᔨᓴᖅᑕᐅᔭᐃᐅᔪᓂ.

<u>ᓄᓇᖅᑐᐊᖅ:</u>
ᑐᖓᔪᖅᑕᐅᓂᖓᑦ – ᑲᓇᓐᓇᖅᑕᓱᒻᒥ ᖃᐅᔨᓴᖅᑕᐅᔭᐃᓐ.
ᐅᓴᐅᓴᐅᓂᖓᑦ – ᐱᖓᓐ.



<u>ᐃᓗᓕᖅᑭᑦ</u>

ᒥᓂᑦᑕᐅᐃᐅᓴᖅ ᑐᓂᓯᔪᓴᖅᑎᑦ ᓄᓇᕗᓴᒥ ᐅᒪᔪᓕᓚᓕᔪᐊᕐᓂᖅ ᑲᑎᒪᔨᖅᑭᓐᓄᑦ ᐊᒪᓕᓚᓕᔪᐊᕐᓂᖅ (ᑲᑎᒪᔨᖅᑭᑦ) ᐃᕗᐊᓚ 2020-ᒥ ᑐᖓᓕᐃᕐᓂᓯᓂ ᑲᑉᐸᔨᓗᑎᑦ ᐊᖏᕈᓂᕐᒥᒃᓴᓂᖅ ᐃᓯᒪᓕᐅᕕᑕᐅᓯᒪᔪᓄᓗᓗ ᒪᔭᕐᓄᑦ ᐊᑦᔨᑦᖁᓂᑐᓄᓂᑦ ᑭᖓᔪᐸᓂᓄᑦ ᐱᖓᓱᓇᖅᐸᓯᒪᒥ ᖃᐅᔨᔪᖅᑕᐅᕐᓇᒪᒥ (WAZ) ᐊᒪᓚᓗ ᑲᓇᓄᓇᖅᐸᓯᒪᒥ ᖃᐅᔨᔪᖅᑕᐅᕐᓇᒪᒥ (EAZ). ᑐᑭᓯᓂᐊᖏᑎᓂᖓᓪᓚᖁᓯᒪᔪᓂ 2019-ᒥ ᐊᑦᔨᑦᖁᓂᑐᓂᓗᓂᑦ ᐅᒪᔪᓕᓂᓂᑦ ᖃᐅᔨᔪᓂᕐᒥᑦ ᑐᓴᐅᒪᑦᐸᐅᑕᐅᓂᐊᖏᖅᑐᓂᖅ 2020/2021–ᒥ ᑭᖑᑦᑕᑕᐅᔪᓄᓂᐊᖏᖅᑐᓂᑐᓂᑦ ᐅᑐᐃᓄᓇᐅᓯᒪᓕᕐᓂᑐᓂᑦ ᐅᑯᐊ ᑐᓂᓗᐅᑎᓗᓚᓂᑦ. ᒥᓂᑦᑕᐅᐃᐅᓴᖅ ᓂᓚᐅᓄᓂᖅᓱᓂᖅᑐᓂᑦ ᐊᑐᐃᓄᓇᐅᓯᒪᓕᕐᓂᑐᓂᑦ ᐱᖓᓱᓇᖅᐸᓯᒪᒥ ᐊᓯᔨᓄᓂᖅᓱᓂᖅᑐᓂᑦ ᐊᑕᐅᓯᒪᔪᓄᑦ (ᐅᖓᓗᑕᓂ 25%) ᑭᖑᑦᑕᓄᓂᐊᕆᐅᓄᑦ ᐊᑕᐅᓯᒪᔪᓄᑦ ᐅᖓᓗᑕᓄᓕᔭᓂᓄᑦ ᐃᓗᐊᖅᑐᓄᓂᑦ ᑕᑯᓄᓕᕐᒪᑐᐃᓐᓇᖅᑐᓂᑦ, ᐃᓕᒍᓯᐅᑎᒥᑦ ᐊᐅᓪᓚᕆᐊᓕᕐᓂᑦ, ᐃᓕᒍᓯᐅᑎᒥᑦ ᐊᐅᓪᓚᕆᐊᓕᕐᓂᑦ ᐃᓗᐊᖅᑐᓄᓂᑦ ᑕᑯᓄᓕᕐᒪᑐᐃᓐᓇᖅᑐᓂᑦ ᐃᓗᐊᖅᑐᓄᓂᑦ ᐃᓗᐊᖅᑐᓄᓂᑦ ᐃᓗᐊᖅᑐᓄ

- 2019–Γ ᖃᐅᔨᓴᖅ ᓇᓗᓇᐃᖅᓯᖅ 19.5%–Γᑦ ᐃᓄᐃᓴᓕᕆᓕᖅᑐᑦ ᑭᖓᔪᓕᐃᑦ ᑲᑎᒪᔨᖅᑐᑦ, ᑭᖓᔪᖏᒍᑦ ᐊᓪᓚᑦᑐᒪᓚᒥᑦ ᐊᒥᓯᖓᔪᓕᕆᓕᖅᑐᖅᓯᓂ (77.7%) 2018–Γ ᖃᐅᔨᓴᖅᓂᒥ.
- ᑲᔪᓯᑎᑕᐅᓂᖅᑐᑦ Ḷᓪᓇ ᑲᑎᑦᓯᑎᑦ ᑭᖓᔪᔪᑦᑕᐅᔪᓇᖅᑐᑦ ᐃᒪᐃᓕᖏᒪᑲᖅᑐᖅ ᓄᖓᔪᔪᓕᕆᓂᖅᑐᑦ ᐃᖏᕐᕋᓂᖅᑎᓇᑎᑦ 18.6%–Γᑦ.
- ᑲᔪᓯᑎᑕᐅᔪᔪᑕ 15% ᓄᖓᔪᔪᓕᕆᓂᖅᑐᑕ ᐃᖏᕐᕋᓂᖓ ᐃᒪᐃᓕᖏᒪᑲᖅᑐᖅ ᑲᑎᑦᓯᑎᑦ ᑭᖓᔪᔪᑦᑕᐅᔪᓕᓇᖅᑐᑦ 9,640t, ᐃᓄᐃᓴᓕᕆᓕᖅᑐᑦ 19.5%–Γᑦ.

ᐃᓱᒪᕐᓴᖅᓱᐅᕈᑎᓕᕆᑕᐅᔪ ᑲᑎᒪᓕᑦ ᐊᐠᖅᑭᑕᖏᓂᖅᑐᓄᑦ 2020/21 ᐃᖃᓗᓕᕆᓂᕐᒧᑦ

- ᑐᑭᓯᓂᐊᖅᑎᑦ ᖃᐅᔨᔪᓇᖏᑕᑦ ᐊᐅᓪᓂᖅᓂᖅ ᓇᓕᓂᖏᑕ ᑲᑎᒪᐅᖅᑐᖅ Ḷᓪᓇᐅᑎᓕᓄᑦ. ᐅᓴᖅᑐᑦᓂᖅᑯᓕᓕᓄ ᐊᐅᓪᓂᖅᑕᖃᕐᓂ ᓇᓕᓄᑐᐃᓇᖅᑐᓄᑦ ᑭᖓᔪᓯᐊᑦ ᐱᖓᓱᓇᖅᑐᖅ ᖃᓄᐃᓕᓴᕐᓂᖅ, ᓱᕐᓗ ᑲᑎᒪᐅᖅᑐᑦ ᖃᓄᐃᓕᓴᕐᓂᖅᑐᑦ ᐊᐅᓪᓂᖅᑕᖃᕐᓂ ᐃᖏᕐᕋᓂᖅᑎᓇᑎᒍᑦ ᐅᑭᐅᖅᑕᐅᓂᖅᑐᑦ.
- ᐱᖓᓱᓇᖅᑐᖅᓯᓂ ᐱᒋᐊᖃᖅᓯᓂ, ᐊᑕᐅᓱᑲᓇᖅᑐᖅᓯᓂᖅᑐᓄᑦ ᖃᓄᐃᓕᓴᕐᓂᖅᑐᑦ ᐃᓕᓴᕆᔨᐅᔪᓂᖅᑐᑦ ᐊᐅᓪᓂᖅᑕᖃᕐᓂᖅᑐᑦ ᑲᑎᒪᐅᖅᑐᖅ ᓇᓗᓇᐃᔾᔨᐃᔪᓂᖅᑐᑦ ᐃᓄᐃᓴᓕᕆᓂᕐᒥᓂᖅᑐᑦ ᑭᖓᔪᓯᐊᑦ ᐃᓄᐃᓴᓕᕆᓂᕐᒥᓂᖅᑐᑦ ᐃᓄᐊᓂ ᐃᓄᐅᓱᓪᓚᐊᓂᖅᑐᑦ 2013–Γᓂᑦ ᑕᒪᐸᐅᓂᑦ ᑭᖓᔪᓯᓂᑦ.
- ᐱᖓᓱᓇᖅᑐᖅᓯᓂ ᖃᐅᔨᓴᖅᓱᐅᔪᖅ ᐊᐅᓪᓚᑕᐅᖅᑕᑦᑕᖅᓱᓚᖅ ᑕᐃᓂ 10%–Γᑦ ᓄᖓᑎᖅᓯᓕᕆᓂᖅᑐᑦ ᐃᖏᕐᕋᓂᖅᑎᖅᓯᓂᑦ, ᐱᓱᐊᓂᓕ ᓄᖓᑎᖅᓯᓕᕆᓂᕐᒥᓂᖅᑐᑕ ᑕᒪᐅᐸᑦ ᖃᓄᐃᓕᓴᕐᓂᖅᑐᑦ ᑕᒪᐅᐸᑦ ᖃᓄᐃᓕᓴᕐᓂᖅᑐᑦ ᑕᓪᓕᓱᓚᖅᑐᑦ 2017/`18–Γ ᐃᖃᓗᓕᕆᓂᖅᓱᓚᐅᔪᕐᓚᓂᓕᓂᑦ.
- 20% ᓄᖓᑎᖅᓯᓕᕆᓂᖅᓯᓂᑦ ᐃᖏᕐᕋᓂᖅ ᑲᑎᒪᖅᑕᐅᖅᑕᑦ ᐊᒥᑎᖅᓱᐃᓕᑦᑕᓂᑦ ᓇᓚᒪᓕᔪᐅᕐᖅ ᐅᖅᓱᒋᓪᓕᓴᕐᑕᐃᓕᓕᓗᓂ ᐃᖏᕐᕋᓂᖅᓱᓂ ᐃᓄᐃᓴᓕᕆᓂᕐᓂᖅᑐᑦ ᑲᑎᒪᔨᖅᑕᐅᖅᑕᖃᕐᓂᖅᑐᑦ ᓇᓗᓇᐃᖅᓯᐅᓴ

- ᑲᑎᒪᓕᑦ ᓄᓇᕗᒻᒥ ᓄᓇᕕᖓᒥᓗ ᐊᓐᓚᓗ ᐃᑭᖏᑦ ᐅᒥᐊᖅᐊᑦ ᐃᓱᒪᓕᐅᕆᓱᒪᔪᑦ ᑲᔪᓱᖏᓪᓗᓂᑎᑦ ᑲᑎᑦᓱᑎᑦ ᑭᖑᓪᓕᐊᑦᑕᐅᔭᓇᖅᑐᓂᒃ 840t 2014–ᒥᓂᑦ.

2020/21 ᑲᑎᑦᓱᑎᑦ ᑭᖑᓪᓕᐊᑦᑕᐅᔭᓇᖅᑐᑦ ᓇᓕᖏᑦᑎᐊᑦ ᑲᓇᓂᓇᖅᑐᒥ ᑭᖑᓪᓕᐊᑦᑕᐅᔭᓇᖅᑐᑦ P. montagui:

ᓇᓕᖏᑦᑎᐊᖅ 1: ᑲᑎᒪᓕᑦ ᐃᓱᒪᑦᓴᓱᐅᑕᑎᖅᑦᓂᑎᑦ ᐃᓱᒪᓕᐅᕆᑕᑎᖅᑦᓂᒥᑦᒃ 15% ᐊᒥᓱᓯᒪᔪᐊᕆᓂᒥᑦᒃ ᑭᖑᓪᓕᐊᑦᑕᐅᔭᓇᖅᑐᑦ *P. montagui* ᑲᑎᑦᓱᑎᑦ, ᐃᒪᐃᓕᓐᓄᑦᓗᓂᑎᑦ ᑲᑎᑦᑐᑦ 966t ᓄᖑᓱᑦᓕᓕᐊᓂᖓᑦ 11.4%

ᓇᓕᖏᑦᑎᐊᖅ 2 (ᐃᓱᒪᓕᐅᕈᑕᐅᔪᖅ): ᑲᔪᓱᑎᑕᐅᓗᓂᑎᑦ ᑲᑎᑦᓱᑎᑦ ᑭᖑᓪᓕᐊᑦᑕᐅᔭᓇᖅᑐᑦ 840t (9.9% ER)

ᒥᓂᔨᑕᐅᕕᑦ ᑐᑦᓱᕋᖅᑐᖅ ᐋᖅᑭᑦᑕᐅᔪᓂᑦ ᑲᑎᑦᓱᑎᑦ ᑭᖑᓪᓕᐊᑦᑕᐅᔭᓇᖅᑐᑦ ᓄᓇᕗᒻᒥ ᓄᓇᕕᖓᒥᓗᓂᓗ.

ᑲᓇᐅᓇᖅᓯᓱᒥ ᖃᐅᔨᓴᖏᐅᔪᒥ – ***ᑭᖑᓪᓕᐃᑦP. borealis***

ᑐᑭᓱᓂᐊᖅᑎᓂᑦ ᑐᓴᕋᓴᐃᑦ ᓇᓗᓇᐃᖅᓱᔪᑦ ᐊᓪᓚᑎᐃᑦᑐᓴᒪᓕᒪᓕᒃ ᐊᖏᔪᓪᓚᑎᐊᖅᑐᑦ (102.8%) ᑭᖑᓪᓕᓴᓱᐊᕋᕋᓴᓂ ᑲᑎᒪᐅᖅᑐᑦ 95,238t–ᓄᑦ, ᑭᖑᕐᓴᒍᒧᑦ 19.6%-ᒥᒃ ᐊᖏᔪᓪᓚᑎᐊᓕᐅᖅᑎᓪᓗᓂᑦ ᑭᖑᓪᓕᐊᑦᑕᐅᔭᓇᖅᑐᑦ ᑲᑎᒪᐅᖅᑐᑦ ᐊᖏᕐᕋᓂᑦ. ᑲᑎᒪᐅᖅᑐᑦ ᐊᒥᓴᖏᓱᐊᖅᑐᑦ ᐅᔨᓴᖅᑐᓴᓕᓐᓂᑦ ᑐᙵᓴᒪᓂᒃ. ᑲᑎᒪᐅᖅᑐᑦ ᓇᓗᓇᐃᖅᓱᔪᑦ ᐃᓚᖓᔪᓐᓇᖅᑐᑦ ᑲᑎᒪᐅᖅᑐᑦ ᐃᓚᐃᓕᓐᓄᑦᓴᓗᑦ ᑲᑎᒪᐅᖅᑐᑦ (MSC).

- 2019/20 ᑭᖑᓪᓕᐊᑦᑕᐅᔭᓇᖅᑐᑦ 8,610t ᓄᖑᓱᑦᓕᓕᐊᓂᖓᑦᓗᓂᒃ 18.4%, ᐊᒥᓱᓯᒪᓕᒃᓕᓕᖅᑐᑦ 7,840t (20% ᓄᖑᓱᑦᓕᓕᐊᓂᖓᑦ) 2018/19–ᒥ.
- ᑲᔪᓱᑎᑦᓱᓂᖅ ᒪᔭᓇ ᑭᖑᓪᓕᐊᑦᑕᐅᔭᓇᖅᑐᓂᑦ ᐃᓚᐃᓕᓐᓄᒪᓕᒃᖅᑐᑦ ᓄᖑᓱᑦᓕᓕᐊᓂᖓᑦ 9.1%.
- 15%–ᒥᑦ ᐊᖏᑦᓕᓕᒥᐊᓂᖅ ᐃᓚᐃᓕᓐᓄᒪᓕᒃᖅᑐᖅ 9,901t (+1,292t) 10.4% ᓄᖑᓱᑦᓕᓕᐊᓂᖓᑦ.
- 20%–ᒥᑦ ᐊᖏᑦᓕᓕᒥᐊᓂᖅ (+1,722t) ᐃᓚᐃᓕᓐᓄᒪᓕᒃᖅᑐᑦ ᑭᖑᓪᓕᐊᑦᑕᐃᑦ 10,332t (10.9% ᓄᖑᓱᑦᓕᓕᐊᓂᑦ).
- 25%–ᒥᑦ ᐊᖏᑦᓕᓕᒥᐊᓂᖅ (2,153t) ᐃᓚᓕᐃᓄᒪᓕᒃᖅᑐᑦ ᑭᖑᓪᓕᐊᑦᑕᐃᑦ 10,763t (11.3% ᓄᖑᓱᑦᓕᓕᐊᓂᑦ).
- 15%–ᒥᑦ ᓄᖑᓱᑦᓕᓕᐊᓂᑦ ᑭᖑᓪᓕᐊᑦᑕᐅᔭᓂᑦ 14,271t, 65.7%–ᒥᑦ ᐊᖏᑦᓕᓕᐊᓂᒃᑕᑦ.

2020/21 ᑲᑎᑦᓱᑎᑦ ᑭᖑᓪᓕᐊᑦᑕᐅᔭᓇᖅᑐᑦ ᓇᓕᖏᑦᑎᐊᑦ ᑲᓇᓂᓇᖅᑐᒥ ᑭᖑᓪᓕᐊᑦᑕᐅᔭᓇᖅᑐᑦ P. borealis

ᓇᓕᖏᑦᑎᐊᖅ 1: ᐊᖏᑦᓕᓕᒥᐃᓗᓂᑦ ᑭᖑᓪᓕᐊᑦᑕᐅᔭᓇᖅᑐᑦ ᑲᑎᑦᓱᑎᑦ 15% (+1,292t) to 9,901t (10.4% ᓄᖑᓱᑦᓕᓕᐊᓂᖓᑦ). ᓇᓕᖏᑦᑎᐊᖅ 2: ᐊᖏᑦᓕᓕᒥᐃᓗᓂᑦ ᑭᖑᓪᓕᐊᑦᑕᐅᔭᓇᖅᑐᑦ ᑲᑎᑦᓱᑎᑦ 20% ᐅᓄᖓ 10,332t (10.9% ᓄᖑᑎᖅᑦᓕᓕᐊᓂᑦ), ᑕᑯᓐᓇᖅᓱᑦ ᐊᓪᓚᑎᐃᑦᑐᓴᒪᓕᒪᓕᒃ ᐊᒥᓱᓯᒪᔪᓂᖓᑦ ᑭᖑᓪᓕᐊᑦᑕᐅᔭᓇᖅᑐᑦ ᑲᑎᒪᐅᖅᑐᑦ ᐊᓪᓚᑎᓗ ᐊᑦᓕᓱᑦᑐᒥ ᓄᖑᓱᑦᓕᓕᐊᓂᑦ.

ᒥᓂᔨᑕᐅᕕᑦ ᑐᑦᓱᕋᖅᑐᖅ ᐋᖅᑭᑦᑕᐅᔭᓇᖅᑐᑦ ᑲᑎᒪᐅᖅᑐᑦ ᐊᖏᕐᕋᓂᑦ ᓄᓇᕗᒻᒥ ᓄᓇᕕᖓᒥᓗᓂᑎᑦ.

ᐅᑭᐅᖅᑕᖅᑐᒥ ᑭᖑᓚᓱᓂᕐᓗᑦ ᑲᑎᒪᐅᕋᓂᑦ

ᐅᑭᐅᖅᑕᖅᑐᒥ ᑭᖑᓚᓱᓂᕐᓗᑦ ᑲᑎᒪᐅᕋᓂᑦ ᑲᑎᒪᓂᖅᑐᑦ ᒧᑦᑭ 5, 2020. ᑕᐃᑲᓂ ᑲᑎᒪᓂᕐᒥ, ᒥᓂᔨᑕᐅᕕᐅᔪᑦ ᑐᑭᓱᓂᐊᕐᓂᖅᑕᖅᑐᑦ ᐃᑭᖑᑦ ᑭᖑᔪᓂᐊᖅᑕᖅᑎᓂᑦ ᐃᖅᑲᓇᐅᓂᐊᖅᑎᓂᑦ ᑲᑎᒪᐅᖅᑐᑦ ᑭᖑᓪᓕᐊᑦᑕᐅᔭᓇᖅᑐᓂᑦ ᑲᓇᓂᓇᖅᑐᒥ. ᑲᑎᑦᓱᑎᑦ ᑭᖑᓪᓕᐊᑦᑕᐅᔭᓇᖅᑐᑦ ᑲᓇᓂᓇᖅᑐᒥ ᐱᖓᓱᓇᖅᑐᒥᓗ ᐅᖃᖅᑎᑦᑎᓂᒃᑯᑦᑕᐅᓯᖅᑐᑦ ᓄᓇᕗᒻᒥ ᓄᓇᕗᒻᒥᓗ ᐅᖃᖅᓯᓚᑦᓴᓄᑦ ᓄᓇᖃᖅᑲᓇᕐᓂᖅᑎᑦᑎᓂᒃᑯᑦ ᑲᑎᒪᐅᖅᑐᑦ ᐊᓪᓚᑎᓗ ᓄᓇᕗᒻᒥ ᑕᒪᓐᓇᖅᑐᖅ 4-ᒥ. ᓇᓕᖏᑦᑎᐊᑦ ᐃᓱᒪᓕᐅᕈᑕᐅᔪᓐᓇ ᑲᑎᒪᓕᑦ ᐊᖏᔪᓐᓇᖅᑐᑦ ᑲᑎᒪᐅᖅᑐᑦ ᑕᒪᓐᓇᖅᑐᖅ

ᓱᖏᓇᐅᔨᓴᕐᓂ: ᐊᐸᑎᒥᖅ **ᐃᖃᓗᓕᕆᓂᖏᑦᑐᓪᓗ ᐊᐅᒃᓚᖏᓂᖅᐲᑦ**, ᐃᖃᓗᓕᕆᓂᕐᒧᑦ ᐃᓚᖏᑕᓕᕆᓂᕐᓪᓗ ᑲᓇᑕᒥ

ᐅᓪᓗᖓ: ᕿᓂᐊᕆ 27, 2020

[ᑐᑭᓯᓂᐊᖅᑎᓂᒃ ᑭᐅᔭᕐᓂᒃ 2020/014: ᑐᓴᕋᔭᖅ ᒪᑎᒪᐅᖅᑐᑦ ᒥᑦᓴᓄᑦ ᓇᓗᓇᐃᖅᓯᕐᓂᒃ ᐅᑭᐅᖅᑕᖅᑐᑦ ᑭᓯᐊᓂᖅᑐᑦ, ᑕᐃᔨᐅᓂᕐᓂ Pandalus Borealis, ᐊᓱᒪᓗ, Pandalus montagui, ᐱᖓᓱᓇᖏᑦ ᒪᓇᓱᓇᖏᓐᓂ ᖃᐅᔨᓴᕐᕕᐅᓂ, ᓴᓱᓂᐊᕆᓗ 2020]

ᖃᐅᔨᓴᖅᑎᓄᑦ ᑭᐅᔭᕆᐊᕆᔨᖅ 2020/014

ᓇᐃᑦᑐᓕᕈᑦ: ᓄᑦᑭᕈᓐᓇᖅᑕᐅᔭᕐᓂᑦ ᐅᒪᔨᖏᑦ ᖃᓄᐃᓕᖏᓚᓂᐊᒃ ᓇᓗᓇᐃᒃᑯᑦᖏᓐᓄᑦ ᐅᑭᐅᖅᑕᖅᑐᑦ ᑭᒧᒧᔨᖏᓄᑦ ᔨᓐᓂᒋᔭᖅᓯᓗᒃᑐᑦ, ᔨᓐᓂᒋᔭᖅᓯᓗᒃᑐᑦ, ᔨᓐᓂᒋᔭᖅᓯᓗᒃᑐᑦ *ᐊᒥᑕᒍ*, ᐃᓗᐊᓂ ᐱᓕᕆᔨᒃᑲᓐᓂᑦ ᐊᑭᑐᖅᑕᐅᓂᖅ ᐃᓂᓪᓗᐊᓂ 2020

ᓄᑦᑭᕈᓐᓇᖅᑐᑦ ᓇᓗᓇᐃᒃᑯᑦᖏᓐᓂᒃ

ᖃᓇᓐᓂᒃ ᖃᒥᖅᑭᕐᕕᖏᑦᑕ ᐃᓂᓪᓚ – ᔨᓐᑎᓕᖅ ᔪᓐᓇᕐᓂᖅ

ᐃᖃᓗᓕᕆᓂᖅ

ᑲᑎᒪᒫᓐᓇᒍ ᐃᖃᓗᐊᑕᐅᔪᑦ ᑭᖑᓪᓕᓂᒃ ᔨᓐᓂᒋᔭᖅᓯᓗᒃᑐᑦ ᔪᓐᓇᕐᓂᖅ ᐃᓗᐊᓂ EAZ ᐅᓂᒃᑲᓚᐅᔪᒪᓕᒪᓕᖅ ᖃᓇᑕᐅᔨ ᐊᑦᓯᓐᓄᒥᒥ ᑯᑕᖓᒃᑕ ᐅᓂᒃᑲᖏᑦ (CAQR), ᐅᓪᓗᐊᑕ 7 ᐅᓐᓄᐊᓚ 2020, ᑎᑭᐅᒪᓪᓗᓂ 4,687 ᑕᔪᓄᑦ, ᑖᓐᓇ 56% ᐱᒃᓴᑕ ᑭᓯᓚᐊᖑᒪᔨᔨ TAC (ᖃᒥᖅᑕᕆᓴᖅ 1, ᑕᑯᓴᓴᖅ 2). ᐊᖅᑭᒧᖓᓄᑦ 2019/20 ᐃᖃᓗᓕᕆᓂᖅ ᐊᐅᓚᓂᖅ ᑎᑭᐅᓐᓇᔪᓂᒃ ᑲᒪᒃᑯᓗᒍ 31 ᐃᓕᓴ 2020, ᑕᒪᐃᓪᒪᓕᑦ ᑖᓱᐊᐃ ᐱᓇᔪᒃᑕᓄᑦ ᐅᓂᒃᑲᓕᑦ ᐱᒋᐊᕈᑎᑐᐃᓐᓇᑦ ᐊᖅᑭᒧᖓᓄᑦ 2019/20.

ᐅᒪᔨᖏᑕ ᐊᑲᑎᑎᓂᔭᖅ

ᐃᖃᓗᓕᕆᔨᒃᑕᐅᔪᓇᖅᑐᒃᑐᖅ ᐅᒪᔨᖏᑕ ᐊᑲᑎᑎᓂᔭᖅᑕ ᓇᐃᓴᐅᑦ ᐊᖏᑕᓕᐊᓕᐅᖅᑐᒃᑐᖅ 102.8% ᐱᒋᐊᖅᓯᒍ 2018 ᐊᖅᑭᒧᖓᓄᑦ 2019 ᐃᓪᓇᓗ ᖃᓄᑎᓯᒧᓕᓚᒃᑐᒋᑕᖅᑐᒃᑐᖅ (95,138 ᑕᔪᓄᑦ) ᑕᐃᑲᖓᓂᑕ ᐅᓪᓗᖅᑎᑕᑦ ᐃᖃᓗᓕᕆᔨᑕᑦ ᐱᒋᐊᖅᑎᑕᐅᓪᓚᕆᒃ (ᖃᒥᖅᑕᕆᓴᖅ 2, ᑕᑯᓴᓴᖅ 3a). ᐊᖅᓇᓪᓗᓂ SSB ᓇᐃᓴᐅᑦ ᑕᑯᓴᖅᑕᐃᓪᓗᑎᒃ ᐊᖅᑭᒪᔭᑎᓂᔭᖅᑐᒃᑐᖅ 74.0% ᐅᓪᓗᒥᐅᔪᖅ ᖃᓄᑎᖅᓯᓂᖅᓴᑦ ᑐᓴᓂᓚᖅᑕᖅᓯᒃ (57,143 ᑕᔪᓄᑦ) (ᖃᒥᖅᑕᕆᓴᖅ 2, ᑕᑯᓴᓴᖅ 3b).

ᐱᓴᓚᐅᑦᑕᕐᓂᖅᓯᒃ

ᐅᓂᒃᑲᓚᐅᔪᒪᓕᔨᑦ ᐱᓴᓚᐅᑦᑕᕐᓂᖅᓯᒃᑕ ᓇᐃᓴᐅᑕ (ER) ᓇᓗᓇᐃᕈᑎᖓᑦ ᐊᖅᑭᒧᒍᑦ 2019/20, ᐅᓪᓗᐊᑕᒍᑦ 7 ᐅᓐᓄᐊᓚ 2020, ᑎᑭᐅᒪᓪᓗᓂ 4.9% (ᑕᑯᓴᓴᖅ 4a). ᑕᐃᒃᓯᒃᑲᑦ ᑖᓐᓇ ᐃᖃᓗᓕᕆᓂᖅ ᒪᑐᐃᖓᓪᓗᓂᖅᓯᒃᑕ ᑲᑎᒪᑎᓪᓗᑦᑕ, ᐅᓂᒃᑲᓚᐅᔪᒪᓕᔨᖅ ᐱᓴᓚᐅᓂᖅᓯᒃᑕ ER ᖃᓄᑎᓯᒧᓕᓚᒃᑐᒋᑕᖅᑐᒃᑐᖅ ᐃᓕᓴᐅᓂᖅᓯᒃᑐᖅ ᐃᖃᓗᐊᖅᓯᒪᓂᖅᓯᒃ. ᑕᐃᒪᓕ ᐃᓗᐃᑦᑐᐃᔨᖅ ᑭᒃᓕᓕᐊᖑᔨᖅ TAC ᐱᓴᓚᐅᑐᐃᓐᓇᓚᖅᑕᒃᖏᑦ ᐃᖃᓗᓕᕆᓂᖅᓯᓚᐅᑦ ᑕᖅᑭᖑᔪᑦ, ᓴᖅᑭᑐᐃᓐᓇᓚᖅᑕᒃᑐᖅ ER ᓇᐃᓴᐅᑦ ᐊᖅᑭᒧᖓᓄᑦ 2019/20 ᑎᑭᑦᓗᓂᖅ 8.8% (ᑕᑯᓴᓴᖅ 4b). ᐊᖅᑭᒧᐊᓴᓂᑦ ᖃᑎᖅᑭᓗᐊᖅᑐᒥᐅᒃ ᓇᐃᓴᐅᑦ ᓴᖅᑭᑦᑐᓕᓇᖅᑐᒃᑯᑦ ᐱᓴᓚᐅᑐᓂᖑᓂᓕᖅᑐᒃᑐᖅ ER ᑭᓚᖅᑐᓂᖅ 15.4%, ᑖᓐᓇ ᖃᓄᐅᓴᓇᒃ ᑎᓵᑯᐊᖅᑕᐅᔪᒪᓕᔨᖅ ᑎᑭᐅᓐᓇᔪᓂᒃᑲᓐᓂᒃ ER 15% ᖃᓄᐃᙵᑦᑎᐊᖅᑐᑦ ᐃᓄᒃᑎᓄᑦ ᐃᓗᐊᓄᑦ EAZ (DFO, 2018).

ᐅᓪᓗᒥ ᖃᓄᐃᓪᓗᐊᑭᑦ

ᐊᖅᓇᓪᓗᓄᑦ SSB ᓇᐃᓴᐅᑦ ᔨᓐᑎᓕᖅ ᔪᓐᓇᕐᓂᖅ ᐃᓗᐊᓂ ᑖᓱᒪᒪ EAZ ᓱᓕ ᓱᓕᑕᓄᓕᓱᒪᙵᑦᑐᑦ ᑖᓱᒪᒪ ᖃᓄᐃᓕᖏᓐᓄᑦᑐᐊᖅᑐᑦ ᐃᓄᒃᑎᓄᑦ ᐃᓗᐊᓄᑦ PA ᐊᓂᔪᓕᐊᓱᒪᓕᕆᑕ (ᑕᑯᓴᓴᖅ 5).

ᖃᓇᓐᓂᒃ ᖃᒥᖅᑭᕐᕕᖏᑦᑕ ᐃᓂᓪᓚ – ᔨᓐᑎᓕᖅ *ᐊᒥᑕᒍ*

ᐃᖃᓗᓕᕆᓂᖅ

ᑲᑎᒪᒫᓂᑦ ᐱᓴᓚᐅᓂᖓᓂᑦ ᑭᖑᓪᓕᓂᖓᓂᑕ ᔨᓐᑎᓕᖅ *ᐊᒥᑕᒍ* ᐃᓗᐊᓂ EAZ, ᑕᐃᒃᓯᒃᑲᑦ ᐅᓪᓗᖅᑎᑕᑦ 7 ᐅᓐᓄᐊᓚ 2020, 113 ᑕᔪᓄᑦ (ᖃᒥᖅᑕᕆᓴᖅ 1, ᑕᑯᓴᓴᖅ 6) ᐊᒪᓚᓗ ᑲᑕᒃᓯᓪᓕᐊᖑᓪᓕᖅᓴᓂᒃ ᑕᐃᒪᓐᓇᑦ ᐊᖅᑭᒧᖓᓄᑦ 2001. ᑖᓐᓇ 2019/20 ᐃᖃᓗᓕᕆᓂᖅ ᐊᐅᓚᓂᐊᕐᓇᖅᑐᒃᑯᑦ ᑎᑭᐅᓐᓇᔪᓂᒃ ᑲᒪᒃᑯᓗᒍ 31 ᐃᓕᓴ 2020, ᑕᐃᒪᓐᓇ ᐱᓇᔪᒃᑕᐅᓂᖓᓂᑦ ᓇᓗᓇᐃᖅᑎᑦ ᐱᒋᐊᕈᑎᑐᐃᓐᓇᕐᓇᐅᔭᓕ ᐊᖅᑭᒧᖓᓄᑦ 2019/20.

ᐅᒪᔪᖏᑦ ᐊᐳᑎᓂᓗᒃ

ᐅᒪᔪᖏᑦ ᐊᐳᑎᓂᓗᒃᑕᑦ ᓇᐋᕆᐅᑎᐊᒥᖏᑦ ᑭᖓᓴᕐᑦ <ᓵᑦᑕᓪᓗ ᐃᑕᑐᐃᐦ ᐃᓗᓐᓂᐊᓂ EAZ ᐅᑎᖅᑕᕋᐃᓵᓇᖅᑐᖅ ᐊᖏᕐᕕᒃᓴᖏᑦ ᐊᐳᑎᓂᓗᓴᐳᑦᑕᑦ, ᑭᓯᐊᓂ ᐃᓕᓵᓂᐳᓕᕆᓂᖓ ᐱᐊᖅᖢᓴᓕᕆᓂᖓ ᐅᒪᔪᖏᑦ ᐊᐳᑎᓂᓗᒃᑕᑦ ᐊᐳᑎᓂᓗᒃᑕᑦ ᐅᒪᔪᖏᑦ ᐊᐳᑎᓂᓗᒃᑕᑦ ᓇᐋᕆᐅᑎᖅᓯᓂᑦ ᐅᓂᒃᑲᐃᓂᖅ 2012 (ᑕᒻᒥᐅᕐᖅ 7). ᑕᒪᐃᓵᓂᑦ ᐃᖅᑲᓚᑲᐅᕐᖅᑕᑦ ᐅᒪᔪᖏᑦ ᐊᐳᑎᓂᓗᒃᑕᑦ ᐊᖏᕐᕕᒃᓴᖏᑦ ᐊᐳᑎᓂᓗᒃ ᐊᖏᒪᑦᑕᐅᖅ ᐊᖏᕐᕕᒃᓴᖏᑦ SSB ᓇᐋᕆᐅᑎᖅᓯᓂᑦ ᑲᑎᑕᐊᖅᓯᓂᖅᑐᑦ ᐊᖏᕐᕕᒃᓴᖏᑦ ᐊᐳᑎᓂᓗᒃᑕᑦ 2019 ᑎᑭᓪᓗᒍ 8,503 ᑕᓐᓂᑦ ᐊᖏᒪᓗ 4,415 ᑕᓐᓂᑦ, ᐊᑐᓂᖓᓕᕆᔨᑎᒍᑦ, ᐊᖏᒪᓗ ᐊᑎᖓᔪᖓᓂᓂᖅᑐᑦ ᐊᖏᕐᕕᒃᓴᖏᑦ ᐊᐳᑎᓂᓗᒃᑕ

ᐱᖓᓐᓄᓕᒃ ᖃᑎᖅᑕᕐᕕᖏᑦ ᐃᓕᓴᖅ – ᓴᓇᓕᓐ ᐊᒥᓱᐃᓕᓴᕐᓂᖅ

ᑕᐃᒪᓕᒪ ᑕᒪᐃᓇᓂᒃ ᖃᐅᔨᓴᖓᔪᓄᑦ ᐊᖏᕐᕋᕐᓂ ᐅᓂᒃᑲᓇᐅᑦᑐᓕᕆᔨᖅᑐᑦᓯᓂᑦ, ᑕᒃᔪᐊᕐ ᑭᖑᓂᖓᓂᒃᑯᑦ ᐃᓕᓴᕐᕕᖏᓂᒃ ᐊᐅᓚᑦᑎᓂᒃ ᐃᓕᓴᕐᕕᖏᓂᒃ ᐊᐅᓚᑦᑎᓂᒃ ᐃᓚᖓᑕᒥ

- ᐸᓐᓂᒃ ᐊᒥᐱᐃᐧᐃᐊᓂᒃ ᐅᒪᔨᓕᒃ ᐊᑦᑎᒋᓴᖅᑕ ᓇᐋᔭᐅᑎᓐᓂᒃ ᐄᐋᔨᐅᑦᑎᓐᓂᒃ ᐱᒻᓂᕆᑎᓐ ᐃᑭᖅᑐᒪᓴᑦᑐᖅ, ᑕᒪᓐᓇ ᓇᓗᓇᖏᑎᖅᓯᐅᑎᐳᖅ ᖃᓇᐃᓕᑐᓂᒃᑕ ᐅᒪᔨᓕᒃ ᐊᑦᑎᒋᓴᐊ ᐃᓗᐊᓂ EAZ.
- ᒪᓕᒃᓴᐃᒍ ᑕᒪᑐᒪ ᐊᖏᕐᕋᔪᑦ ᑲᑕᒋᐊᓂᓐᖓ ᐃᓗᐊᒍᑦ ᐅᒪᔨᓕᒃ ᐊᑦᑎᒋᓴᖅᑕ, ᑭᖓᑦᐸᑦ ᐸᓐᓂᒃ/ᓯᑕᒍᐃ ᐃᖃᓗᓀᓴᖏᑦ ᐃᓂᖃᑎᑕᑦ ᐅᓴᓐᓄᓂᒃ ᑕᑯᓀᓯᒪ ᓂᐲᔪᓐᓇᐅᑉᐳᓂᒃ ᐊᑐᖃᑕᐅᒪᔨᓂᖅ ᑐᖓᔪᓐᓇᒪᓐᖓ.
- ᓄᖃᑐᐃᐃᓇᐊᓕᓂᒃ ᐱᓴᓗᐅᑕᑦᑎᓂᖓᓐᓄᑦ ᓇᐃᓴᐅᑕ ᑕᑯ ᐃᓗᐃᑦᑐᓗᓂ ᐃᖃᓗᐃᑦᑐᓗᓂ TAC ᓄᖓᑕᐅᐳᓂᑭᓂ ᐊᑦᑕᒍᐊᓂ 2019/20 ᑎᑭᐅᑎᑯᓂᖃᖅ 9.9%.

ᐱᖓᓂᓀᑕ ᖃᒥᑦᑭᐟᖓᓕᑕ ᐃᓂᖓ (WAZ)

ᐸᓐᓂᒃ ᐳᐊᓂᓱᒃ ᑦ

- ᖃᓄᐃᑦᑐᓂᓗᒃ ᐅᒪᔨᓴᖃᑎᒋᑦ ᑭᖓᔪᐸᑦ ᐅᓪᓗᒥ, ᐱᓪᓗᒍ ᐱᑕᖃᖅᖁᑦᓯᒪᑦ ᓯᓂᔅᑎᓴᖃᓕᕐᓂᒍᓄᑦ ᐊᑐᖃᑕᐅᒪᔨᓯᓂᖃᖅ ᑐᓴᕐᓂᒍᓄᑦ ᐸᓐᓂᒃ ᐳᐊᓂᓱᒃ ᑭᖓᔪᐸᑦᑎᓂᒃ ᐅᓪᓗᐊᓂ WAZ.
- ᑕᒪᐃᓪᓗᒃ ᐃᖃᓄᓕᕆᑕᐅᔨᓇᖅᑐᒃ ᐅᒪᔨᓕᒃ ᐊᑦᑎᒋᓴᖅᑕ ᐊᓖᒪᓗ ᐊᖏᓇᓪᓗᑦ ᑭᖓᔪᓂᒡᐅᓯᑦ ᐊᒥᓯᓂᖅᑕ ᐅᒪᔨᓕᒃ ᐊᑦᑎᒋᓴᖅᑕ ᐃᓗᐊᑎᓐᓂᒃ ᑲᑕᒋᐊᓖᑦᓯᓯᒪᔨᑦ ᐊᒋᓴᐃᓂᒃ 2018 ᑎᑭᓪᓗᒍ 2019.
- ᐅᓂᖅᑲᓱᐅᓯᒪᔨᑦ ᐱᓴᓗᐅᑕᑦᑎᓂᖓᓐᓄᑦ ᐃᓗᐊᑐᑦ ᐊᖏᕐᕋᔪᓐᓇᑦ 2019/20 ᑎᑭᑦᑐᖃᖅ 3.4%. ᐅᓪᓗᒥᐅᔨᖅ ᐃᖃᓄᓕᕆᑕᐅᔨᓇᖅᑐᒃ TAC ᓇᓕᒧᓯᖃᑎᒌᐅᔨᔨᖅ ᓄᖓᑕᐅᒍᐊᓕᓚᒧᑦ ᐱᓴᓗᐅᑕᑦᑎᓂᖓᓐᓄᑦ ᓇᓗᓇᐃᕈᑦᓄᑦ ᑎᑭᑦᑐᖃᖅ 15.5%.

ᐸᓐᓂᒃ ᐊᒥᐱᐃᐧᐃᐊᓂᒃ

- ᖃᓄᐃᑦᑐᓂᓗᒃ ᐅᒪᔨᓴᖃᑎᒋᑦ ᑭᖓᔪᐸᑦ ᐅᓪᓗᒥ, ᐱᓪᓗᒍ ᐱᑕᖃᖅᖁᑦᓯᒪᑦ ᓯᓂᔅᑎᓴᖃᓕᕐᓂᒍᓄᑦ ᐊᑐᖃᑕᐅᒪᔨᓯᓂᖃᖅ ᑐᓴᕐᓂᒍᓄᑦ ᐸᓐᓂᒃ/ᓯᑕᒍᐃ ᑭᖓᔪᐸᑦᑎᓂᒃ ᐅᓪᓗᐊᓂ WAZ.
- ᑕᒪᐃᓪᓗᒃ ᐃᖃᓄᓕᕆᑕᐅᔨᓇᖅᑐᒃ ᐅᒪᔨᓕᒃ ᐊᑦᑎᒋᓴᖅᑕ ᐊᑦᓀᓪᓗᑦ ᑭᖓᔪᓂᒡᐅᓯᑦ ᐊᒥᓯᓂᖅᑕ ᐅᒪᔨᓕᒃ ᐊᑦᑎᒋᓴᖅᑕ ᐃᓗᐊᑎᓐᓂᒃ ᑲᑕᒋᐊᓖᐅᖅᑐᒃ 2018 ᑎᑭᓪᓗᒍ 2019.
- ᐅᓂᖅᑲᓱᐅᓯᒪᔨᑦ ᐱᓴᓗᐅᑕᑦᑎᓂᖓᓐᓄᑦ ᐃᓗᐊᑐᑦ ᐊᖏᕐᕋᔪᓐᓇᑦ 2019/20 ᑎᑭᑦᑐᖃᖅ 11.3%; ᓄᖃᑐᐃᐃᓇᐊᓕᓚᒧᑦ ᐱᓴᓗᐅᑕᑦᑎᓂᖓᓐᓄᑦ ᐃᓗᐊᑐᑕᑦ ᐊᖏᕐᕋᔪᒧᑦ 2019/20 ᑎᑭᑦᑕᓂᒃ 18.6%.

ᑕᐃᒃᓂ ᑲᑎᒪᔨᖅ ᐱᓕᕆᖏᑕᖅᓱᒪᔪᑦ, ᑭᖓᔪᒻᒪᓴᐅᔪᑦ ᑲᑎᒪᒪᕆᑐᐃᓂ ᐊᒡᒪᓗ ᓄᖓᔪᑎᓚᕐᓂᐊᓂᕐᒧᑐᓴᑦ ᐃᖏᕐᕋᓂᖏᑦ ᐱᖓᓱᓚᒥ ᖃᕆᓕᓴᐃᓕᕆᔨᒥ ᐊᒍᕐᔪᒍᑎᓂᑦ ᖃᕆᓕᓴᐃᓚᕆᕐᑐᖓᓂᖃᖅᓯᒪᓂᑦ

ᐱᖓᓱᓚᖅᑎ WAZ		2019/20	2018/19	2017/18	2016/17	2015/16
Montagui	FB (t)	79,834	44,915	31,724	55,194	77,078
	TAC (t)	11,975	6,138	6,138	6,138	6,138
	ER	15%	13.7%	19.3%	11.1%	8%
Borealis	FB (t)	21,088	10,487	13,116	28,532	21,713
	TAC (t)	3,163	2,080	2,080	2,080	2,080
	ER	15%	19.8%	15.9%	7.3%	9.6%

FB – ᑭᖓᔪᒪᓕᕆᐊᕐᓂᓴᐃᑦ ᑲᑎᒪᕆᖅᑐᑦ

TAC – ᑲᑎᒪᔨᓕᑎᑦ ᑭᖓᔪᒻᒪᑕᐅᔭᓚᖅᑐᑦ

ER – ᓄᖓᑎᖅᓯᒻᓗᐊᓂᕐᓂᒪᑕ ᐃᓕᕐᕋᓂᕐᓂᑦ

<u>ᐊᑐᑎᑐᐃᓴᓚᖅᓯᒻᒥᑦ / ᐃᓂᐅᔨᑦ 2020/21–ᒧᑦ</u>

	Montagui				Borealis			
	ᑲᑎᒪᔨᒪᑎᑦ ᑭᖓᔪᒻᒪᑕ ᐅᔭᓚᖅᑐᑦ	ᑯᑕ ᐊᓯᔾᔨᖅᑐᑦ	ᐊᓯᔾᔨᖅᑐᑦ ᑲᑎᒪᔨᒪᑎᑦ ᑭᖓᔪᒻᒪᑕ ᐅᔭᓚᖅᑐᑦ	ᓄᖓᔪᒻᒪᓕᓴᐃᓚᖅᑐᖓᓂᒃ	ᑲᑎᒪᔨᒪᑎ ᑭᖓᔪᒻᒪᑕ ᑕᐅᔭᓚᖅᑐᑦ	ᑯᑕ ᐊᓯᔾᔨᖅᑐᑦ	ᐊᖏᕐᓂᒥᒃ ᐊᖅᑐᑦ ᑲᑎᒪᔨᒪᑎᑦ ᑭᖓᔪᒻᒪᑕ ᐅᔭᓚᖅᑐᑦ	ᓄᖓᔪᒻᒪᓕᓴᐃᓚᖅᑐᖓᓂᒃ
ᑲᔪᓯᔪᑦ ᑭᖓᔪᒻᒪᑕᐅᔭᓚᖅᑐᑦ	**11,975t**	**0**	**0**	18.6%	**3,168t**	**0**	**0**	15.5%
ᑲᔪᓯᔪᑦ ᓄᖓᔪᒻᒪᓕᓴᐃᓚᖅᑐᑦ	9,640t	-2,335t	-19.5%	15%	3,057t	-106t	-3.4%	15%
ᐅᖃᓕᒍᓐᓇᖏᑦ ᑐᖓ ᓄᖓᔪᒻᒪᓕᓴᐃᓚᖅᑐᑦ*	12,404t	+429t	+3.6%	19.3%	4,035t	+872t	+27.6%	19.8%
ᐊᒋᓂᐅᔾᔨᖏᑦᑐᖓ ᓄᖓᔪᒻᒪᓕᓴᐃᓚᖅᑐᑦ*	**5,141t**	-6,834t	-57%	8%	**1,488t**	-1,675t	-53%	7.3%

*ᑐᓴᓴᓕᓂᒃ ᑕᐃᒃᓂᓕᕆᓐᖓ ᐃᓂᓕᔾᐅᓕᓴᓂᒃ





ᐃᓕᓴ 27, 2015

ᓄᓕᖅᑲᓇᕐᓂᖅ ᑭᐅᓂ ᓴᐃ
ᒥᓂᑦᑐᓕ ᐃᒪᖅᐳᑕᓕᕆᔨᓂ
15 ᓇᑎᖑᓕ, ᓄᓐᑎᓂᐅᓂ ᐃᓐᓇᖅᐸᐃᓂ
200 ᑭᓐᓂ ᐊᑐᕆᐊᖅ
ᐊᑐᖓ, ᐊᓐᑎᐅᓕᐅ
K1A 0E6

ᒥᓂᑦ ᓴᐃ:

ᐱᓐᓇᔪ: ᓄᓇᕗᖅᒥ ᐅᒪᔨᓕᕆᔨᖅᐸᐃᓂ ᑲᑎᒪᔨᖅᑎᑦ ᐊᒪᓚᓗ ᓄᓇᕐᒪᖅ ᑕᓕᐅᖅᒥᐅᑕᓕᕆᔨᓐᓂᖅᓂ ᑲᑎᒪᔨᖅᑎᑦ ᐃᓄᒃᑎᑐᑦᑎᓂᑦ ᐊᒪᓚᓗ ᐊᐃᑕᖅᑯᓂᖅᓂ ᐱᓐᓇᑦ ᒥᐅᔨᒃᑕᐅᓂᖅᓂ ᐊᒪᓚᓗ ᐊᓂᔨᒃᑕᐅᓂᖅᓂ ᑭᒍᓪᓕᐊᑦ ᑲᑎᒪᓂᑦ ᐊᓯᖏᓂᖅ ᐊᐅᓂᖅᑎᖅᑐᑦ ᐊᒪᓚᓗ ᐱᓕᕆᖅ ᑲᓇᐃᔨᖅᑐᑦ ᐱᓐᓇᑦ ᐱᓕᕆᔨᒃᑯᑦ ᐊᒪᓚᓗ ᐱᓕᕆᔨᒃᑯᑦ ᑭᒍᓪᓕᐊᑦ ᐱᓕᕆᔨᓐᓂᖅᓂ ᐊᐅᓂᖅᑎᖅᑐᑦ ᓇᓕ

ᐊᖏᕐᕕᐅᔨᖅ ᐃᓚᒃ, ᒪᓕᕐᑐᒍ ᐃᓕᓴᒃ 5.6.4 ᐅᑯᐊ ᓄᓇᔨᖏᑦ ᓄᓇᖁᑎᑦᓂᓕᕐᓂᑦ ᐊᖑᕐᕋᑎᑎᓕᑎᑦ, ᐅᒪᔨᓕᓚᓴᖅᔪᐊᑦ ᐊᖑᕐᓂᑐᑎᒃ ᐅᑯᓂᖓ ᑭᒧᐊᕐᓂᖑᓂᒃ ᓄᓇᔨᑦ ᐱᖓᓱᓴᖓᓂ ᐊᒻᒪᓗ ᓄᓇᔨᑦ ᑲᓇᓱᓴᖓᓂ ᐊᐅᓚᕆᒪᕕᐅᔪᓂ ᒪᓕᒐᓱᑦᓄᑦ ᐊᒻᒪᓕᒐᓱᑦᓄᑦ ᐊᑯᓂᐅᑎᓕᕐᒥᒃ (2015-16 ᐅᐃᖓᓱ 2016-17 ᐃᓚᐅᑐᓂᓂ):

1. *Lᑯᓂᖓ Pandalus montagui* ᑭᖓᔪᐊᕐᓇᑦ:
    a. ᐱᓕᕆᔨ ᐃᒪᑎᑎᒋ 3,069 ᑕᐅᔭ ᓄᓇᔨᑦ ᐱᖓᓱᓴᖓᓂ ᐊᐅᓚᕆᒪᕕᐅᔪᑦ ᓇᓗᓂ ᐃᑐᐊᓂ ᓄᓇᔨᑦ ᓄᓇᖁᑎᑦᓂᖅᓯᒪᔪᓂᒃ ᓇᓗᓂ; ᐊᒻᒪᑐ
    b. ᐱᓕᕆᔨ ᐃᒪᑎᑎᒋ 301 ᑕᐅᔭ ᓄᓇᔨᑦ ᑲᐅᓇᖓᓂ ᐊᐅᓇᕆᒪᕕᐅᔪᑦ ᓇᓗᓂ ᐃᑐᐊᓂ ᓄᓇᔨᑦ ᓄᓇᖁᑎᑦᓂᖅᓯᒪᔪᓂᒃ ᓇᓗᓂ; ᐊᒻᒪᑐ
2. *Lᑯᓂᖓ Pandalus borealis* ᑭᖓᔪᐊᕐᓇᑦ:
    a. ᐱᓕᕆᔨ ᐃᒪᑎᑎᒋ 1,040 ᑕᐅᔭ ᓄᓇᔨᑦ ᐱᖓᓱᓴᖓᓂ ᐊᐅᓚᕆᒪᕕᐅᔪᑦ ᓇᓗᓂ ᐃᑐᐊᓂ ᓄᓇᔨᑦ ᓄᓇᖁᑎᑦᓂᖅᓯᒪᔪᓂᒃ ᓇᓗᓂ; ᐊᒻᒪᑐ
    b. ᐱᓕᐅᖏᓚᓴᕐᓗᑎᓇᓕᒋᑦ ᐃᒪᑎᑎᒋ 183 ᑕᐅᔭ ᓄᓇᔨᑦ ᑲᐅᓇᖓᓂ ᐊᐅᓚᕆᒪᕕᐅᔪᑦ ᓇᓗᓂ ᐃᑐᐊᓂ ᓄᓇᔨᑦ ᓄᓇᖁᑎᑦᓂᖅᓯᒪᔪᓂᒃ ᓇᓗᓂ.

ᐊᖏᕐᕕᐅᔨᖅ ᐃᓚᒃ, ᒪᓕᕐᑐᒍ ᐃᓕᓴᒃ 5.2.3 ᐅᑯᐊ ᓄᓇᓕᒃᑲ ᐃᓄᐃᑦ ᓄᓇᖁᑎᑦᓂᖅᓯᒪᔪᓂᒃ ᐊᖑᕐᕋᑎᑎᒋᓕᒃᑲᓂᖓ, ᓄᓇᓕᒃᑲ ᑕᓚᐅᖅᑎᐅᑕᓚᓴᓕᑦ ᐊᖑᕐᓂᖅᓱᑦ ᐅᑯᓂᖓ ᑲᑎᑎᑕᑦ ᐊᔪᖑᑕᑦ ᐱᓕᐅᓂᖅᑐᒃ ᑭᖓᔪᐊᕐᓇᑦ ᓄᓇᓕᒃᑲ ᐊᐅᓚᕆᒪᕕᐅᔪᑦ ᓇᓗᓂ ᐃᑐᐊᓂ ᓄᓇᓕᒃᑲ ᐃᓚᒋᔭᐅᓂᖓ ᐱᓕᕆᔨ ᐃᒪᑎᑎᒋ 3,069 ᑕᐅᔭ ᓄᓇᓕᒃᑲ ᐱᖓᓱᓴᖓᓂ ᐊᐅᓚᕆᒪᕕᐅᔪᑦ ᓇᓗᓂ ᐃᑐᐊᓂ ᓄᓇᓕᒃᑲ ᐃᓚᒋᔭᐅᓂᖓ; ᐊᒻᒪᑐ
    b. ᐱᓕᕆᔨ ᐃᒪᑎᑎᒋ 129 ᑕᐅᔭ ᓄᓇᓕᒃᑲ ᐊᐅᓇᖓᓂ ᐊᐅᓇᕆᒪᕕᐅᔪᑦ ᓇᓗᓂ ᐃᑐᐊᓂ ᓄᓇᓕᒃᑲ ᐃᓚᒋᔭᐅᓂᖓ; ᐊᒻᒪᑐ
2. *Lᑯᓂᖓ Pandalus borealis* ᑭᖓᔪᐊᕐᓇᑦ:
    a. ᐱᓕᕆᔨ ᐃᒪᑎᑎᒋ 1,040 ᑕᐅᔭ ᓄᓇᓕᒃᑲ ᐱᖓᓱᓴᖓᓂ ᐊᐅᓚᕆᒪᕕᐅᔪᑦ ᓇᓗᓂ ᐃᑐᐊᓂ ᓄᓇᓕᒃᑲ ᐃᓚᒋᔭᐅᓂᖓ; ᐊᒻᒪᑐ
    b. ᐱᓕᐅᖏᓚᓴᕐᓗᑎᓇᓕᒋᑦ ᐃᒪᑎᑎᒋ 46 ᑕᐅᔭ ᓄᓇᓕᒃᑲ ᐊᐅᓚᕆᒪᕕᐅᔪᑦ ᓇᓗᓂ ᐃᑐᐊᓂ ᓄᓇᓕᒃᑲ ᐃᓚᒋᔭᐅᓂᖓ.

1.2 ᑲᑎᑎᑕᑦ ᐊᔪ

ᐱᑎᐊᖑᑎᐅᔨᖅ ᐃᓚᒃ, ᒪᓕᕐᑐᑎᑦ ᐃᓕᓴᑦ 15.3.4 ᐊᒻᒪᑐ 15.3.7 ᐅᑯᐊ ᓄᓇᔨᑦ ᓄᓇᖁᑎᑦᓂᖅᓯᒪᔪᓂᒃ ᐊᖑᕐᕋᑎᑎᒋᓕᒃᑲᓂᖓ ᐊᒻᒪᑐ ᐃᓕᓴᒃ 5.4.4 ᐅᑯ ᓄᓇᓕᒃᑲ ᐃᓄᐃᑦ ᓄᓇᖁᑎᑦᓂᖅᓯᒪᔪᓂᒃ ᐊᖑᕐᕋᑎᑎᒋᓕᒃᑲᓂᖓ, ᐅᒪᔨᓕᓚᓴᖅᔪᐊᑦ ᐊᒻᒪᑐ ᓄᓇᓕᒃᑲ ᑕᓚᐅᖅᑎᐅᑕᓚᓴᓕᑦ ᐊᑐᓕᖁᓴᓗᑎᒃ ᐅᑯᓂᖓ ᑲᑎᑎᑕᑦ ᐊᔪᖑᑕᑦ ᐱᖓᓱᓴᖓᓂ ᐊᐅᓚᕆᒪᕕᐅᔪᑦ ᓇᓗᓂ ᐃᑐᐊᓂ ᓄᓇᓕᒃᑲ

*ᓯᓚᖅᑕᐅᓂᖅ ᐱᑎᐊᖏᑎᑕᐅᔾᔨᖅ ᐃᒧᑦ, ᒪᓕᒃᑐᑦ ᐃᓕᓴᖅ 15.3.4 ᐊᓪᓚᓗ 15.3.7 ᐅᑯᐊ ᓄᓇᔾᔨᒥ ᓄᓇᖁᓂᑦᑕᓂᖏᑦ ᐊᓴᖅᑲᑎᑭᓱᓕᑦ ᐊᓪᓚᓗ ᐃᓕᓴᖅ 5.4.4 ᐅᑯᐊ ᓄᓇᓕᕆᒥ ᐃᓄᐃᑦ ᓄᓇᖁᓂᒋᑦᑕᓂᖏᑦ ᐊᓴᖅᑲᑎᑭᓱᓕᑦᑕᓂᖏᑦ, ᐅᒪᔨᓯᓚᓵᖅᖢᑦ ᐊᓪᓚᓗ ᓄᓇᓕᕆᒥ ᑕᓚᐅᖅᑎᐅᒋᓯᓚᐅᑉᑯᑦ ᐊᐃᓕᖁᓚᐅᑎᒃ ᑐᑭᓯᓚᖅ “ᐊᓯᔾᔨᑲᓪᓚᕐᓂᐊᖅᑕ ᐅᒪᔨᓂᖅᓂ”, ᐊᓯᔾᔨᑦ ᐱᑕᖅᑭᑕᐅᓂᐊᖅᑐᑎᓂᖅᓄᑦ ᐊᓯᔾᔨᑦ ᐊᓯᓚᐅᑉᑯᑦ ᐊᐃᓕᖁᓚᐅᑎᒃ ᐊᐃᓕᓯᐅᓚᐅᑉᑯᑦ ᐊᓯᔾᔨᑦ ᐊᓂᓕᖅᑐᐃᓐᓇᖅᑐ

1. ᑐᖓᓚᐃᖅᑎᑦ ᒪᓕᒐᓗ ᐃᓕᖓ 5.3.3(a) ᐅᑯᐊ *ᓄᓇᕗᒻᒥ ᓄᓇᖅᑐᑦᓯᓂᖅᓯᒪᓂ ᐊᖏᕐᕋᐅᓕᕆᔨᓂ* ᐊᓚᓐᓇᑕ ᐃᓕᖓ 5.5.3(a) ᐅᑯᐊ *ᓄᓇᓚᒃᒥ ᐃᓄᐃᑦ ᓄᓇᖅᑐᑦᓯᓂᖅᓯᒪᓂ ᐊᖏᕐᕋᐅᓕᕆᔨᓄᖏᑦ* - ᐊᖑᓚᐊᕈᑎᑦ ᑭᓪᓕᕋᖅᑎᑦ ᐃᓄᐃᑦ ᐊᖏᔪᓂᕐᓂᐅᓂᒃ ᑭᖑᓪᓕᐊᓂᒃ *ᓄᓇᕗᒻᒥ ᓄᓇᖅᑐᑦᓯᓂᖅᓯᒪᓂ ᐊᖏᕐᕋᐅᓕᕆᔨᓂ* ᓇᓗᓂ ᐊᓚᒪᓗ *ᓄᓇᓚᒃᒥ* ᐃᓚᖓᓱᒪᓕ ᓇᓗᓂ ᑎᑭᐅᓕᒃᑐᐊᓗᒃ ᐊᑐᖅᑎᓪᓗᑎᑦ ᓴᓪᓕᐊᓗᒃᑐᑦ ᐱᔪᓐᓇᓪᓕ ᐃᒪᓐᓕᒃ:
    a. ᐱᓯᒪᔭᐃᓐᓇᕐᓂᒃ ᐊᑐᖅᑐᒪᓚᓐᓂᒃ ᖃᓄᐃᓐᓇᕐᓂᓐᓂᕐᓂᒃ ᑭᖑᑦᑕᐃᑦ ᐱᑕᖃᖅᑐᑦ ᐊᓯᖏᓐᓂᖅᑐᑦ ᓄᓕᒻᒥᐅᑦᑕᐅᓂᖅᑎᓪᓗᒍ ᐊᖏᕐᕋᐅᓕᕆᔨᒥᑦᑐᒃ ᐱᑕᖃᖅᑐᒥᑦ ᑕᐃᒪᒃ *ᓄᓇᕗᒻᒥ ᓄᓇᖅᑐᑦᓯᓂᖅᓯᒪᓂ ᐊᖏᕐᕋᐅᓕᕆᔨᓂ* ᑎᑎᕋᖏᓐᓂᒃ 5 (ᐃᓕᓐᓄ 5.1.5(c)) ᐊᒪᓗ *ᓄᓇᓚᒃᒥ ᐃᓄᐃᑦ ᓄᓇᖅᑐᑦᓯᓂᖅᓯᒪᓂ ᐊᖏᕐᕋᐅᓕᕆᔨᓄᖏᑦ* ᑎᑎᕋᖏᓐᓂᒃ 5 (ᐃᓕᓐᓄ 5.1.5(b)); ᐊᒪᓗ
    b. ᐱᓯᒪᔭᐃᓐᓇᕐᓂᒃ ᐱᐅᓯᖏᑦᓯᒃ ᐃᑦᓗᑉᓂᓐᓂᕐᓂᒃ ᐊᐅᒋᑕ ᖃᓄᐃᓐᓇᕐᓂᑦ ᐃᓗᐊᓂ *ᓄᓇᕗᒻᒥ* ᓄᓇᖅᑐᑦᓯᓂᖅᓯᒪᓂ ᐊᖏᕐᕋᐅᓕᕆᔨᓂ ᓇᓗᓂ ᐊᓚᒪᓗ *ᓄᓇᓚᒃᒥ* ᐃᓚᖓᓱᒪᓕ ᓇᓗᓂ (*ᓄᓇᕗᒻᒥ ᓄᓇᖅᑐᑦᓯᓂᖅᓯᒪᓂ ᐊᖏᕐᕋᐅᓕᕆᔨᓂ* ᐊᓚᒪᓗ *ᓄᓇᓚᒃᒥ ᐃᓄᐃᑦ ᓄᓇᖅᑐᑦᓯᓂᖅᓯᒪᓂ ᐊᖏᕐᕋᐅᓕᕆᔨᓄᖏᑦ* ᐃᓕᓐᓄ 5.1.5(a));

2. ᐱᓱᓴᓐᓂᒋᑦ ᐱᑕᖃᖅᑎᓪᓗᓕᒃ ᐅᐊᓂᒃ *ᓄᓇᕗᒻᒥ ᓄᓇᖅᑐᑦᓯᓂᖅᓯᒪᓂ ᐊᖏᕐᕋᐅᓕᕆᔨᓂ* ᐃᓕᓐᓄ 5.3.4 ᐊᒪᓗ *ᓄᓇᓚᒃᒥ ᐃᓄᐃᑦ ᓄᓇᖅᑐᑦᓯᓂᖅᓯᒪᓂ ᐊᖏᕐᕋᐅᓕᕆᔨᓄᖏᑦ* ᐃᓕᓐᓄ 5.5.4.1 ᐃᓗᐊᒃ (i) ᐅᒪᔨᓪᓚᐅᓴᖅᑕᐃᑦ ᐊᒪᓗ ᒥᓂᑦ ᐃᓱᒪᑎᓴᖅᑐᓴᖅᑐᒃ ᑭᖑᑦᑕᐅᓂᐊᕐᓂᒃᓴᒃ ᓱᓕᑦᓂ *ᓄᓇᕗᒻᒥ ᓄᓇᖅᑐᑦᓯᓂᖅᓯᒪᓂ* ᐊᓚᒪᓗ (ii) *ᓄᓇᓚᒃᒥ* ᑕᓪᓕᐅᒥᐅᑕᓪᓚᕆᔭᐃᑦ ᐊᒪᓗ ᒥᓂᑦ ᐃᓱᒪᑎᓴᖅᑐᓴᖅᑐᒃ ᑭᖑᑦᑕᐅᓂᐊᕐᓂᒃᓴᒃ ᓱᓕᑦᓂ *ᓄᓇᓚᒃᒥ* ᐃᓚᖓᓱᒪᓕ ᓇᓗᓂ, ᑭᒃᔨᑐᐃᓐᓇᕐᓄᑦ *ᓄᓇᖅᑎᑐᓄᑦ* ᐊᓱᐊᓂ;

3. ᐃᓯᐱᑎᓗᒍ ᐃᓗᐊᒃ ᑭᖑᑦᑕᐅᓂᐊᕐᓇᖅᑐᑦ ᐃᓗᐊᓂ ᑕᒪᑲᒃᑭᑦ *ᓄᓇᕗᒻᒥ ᓄᓇᖅᑐᑦᓯᓂᖅᓯᒪᓂ* ᐃᓚᖓᓱᒪᓕ ᓇᓗᓂ ᐃᓗᐊᓂᕐᓂᐅᓇᓗᓂ *ᓄᓇᓚᒃᒥ* ᐃᓚᖓᓱᒪᓕ ᐱᔪᓐᓇᖅᑐᑦ ᐱᐅᓯᖏᑦᓯᒃ ᐊᐱᖅᑯᑦᑐᒃ ᐅᓴᖃᑦᓯᐊᕐᓇᖅᑐᓂ, ᐊᓕᓴᐅᑕᐅᓕᕐᓂᐊᕐᓕᖅᓂᒪᓕ ᐱᓗᐊᔪᓐᓇᕐᓂᒃᓴᒃ ᐃᓕᓴᕐᓗᒍ ᐊᐱᖅᑯᑦᑐᒃ ᐅᓴᖃᑦᓯᐊᕐᓇᖅᑐᓂ *ᓄᓇᓚᒃᒥ* ᐃᓚᖓᓱᒪᓕ ᐃᓗᐊᓂᕐᓂᐅᓇᓗᓂ ᓄᓇᓕᖕᒥᓂᖅᓴᓂᒃ ᐊᐱᖅᑯᑦᑐᒃ ᐅᓴᖃᑦᓯᐊᕐᓇᖅᑐᓂ ᐃᓚᖓᓱᒪᓕ ᐃᓗᐊᓂᕐᓂᐅᓇᓗᓂ

Fisheries and Oceans Canada | Pêches et Océans Canada
Ecosystems and Oceans Science | Sciences des écosystèmes et des océans

Canadian Science Advisory Secretariat
Central and Arctic Region **Science Response 2020/014**

UPDATE OF STOCK STATUS INDICATORS FOR NORTHERN SHRIMP, *PANDALUS BOREALIS,* AND STRIPED SHRIMP, *PANDALUS MONTAGUI,* IN THE WESTERN AND EASTERN ASSESSMENT ZONES, JANUARY 2020

Context

Fisheries and Oceans Canada (DFO) Resource Management has requested an update on the stock status of the two species of shrimp, Northern Shrimp (*Pandalus borealis*) and Striped Shrimp (*P. montagui*), in the Western Assessment Zone (WAZ) and Eastern Assessment Zone (EAZ). The last Zonal Peer Review (ZPR) for the WAZ and EAZ, which fully assessed the stock status, took place in February 2019 (DFO 2019). The full assessments and the updates follow the framework developed in 2007 for Northern Shrimp off Labrador and the northeastern coast of Newfoundland (DFO 2007a). The basis of this update is a series of previously reported fishery-independent surveys and fishery data, and new survey and catch data from the 2019/20 fishing season.

This Science Response Report results from the Science Response Process (SRP) of January 28, 2020 for the Northern and Striped Shrimp Update for the Eastern and Western Assessment Zones, and Striped Shrimp Update for Shrimp Fishing Area (SFA) 4.

Background

The EAZ and WAZ (Figure 1) were adopted in 2011 as the basis for assessing the status of shrimp in SFA 2 and SFA 3 (DFO 2011). The combined boundary of the EAZ and WAZ is equivalent to the combined areas of SFAs 2 and 3 (subsequently redefined as SFAs Davis Strait, Nunavut and Nunavik) and their corresponding Management Units (Figure 1) implemented in the 2013/14 fishing season.

Two exploitation rates are presented for each assessment zone and species because the total allowable catch (TAC) is not generally taken. Exploitation rate refers to the realized rate based on reported catch, while the potential rate is calculated as though the TAC had been fully taken.

In the EAZ, the first two years of survey data (2006–2007) are not considered comparable with the rest of the time series because of poor trawl performance and incomplete sampling coverage in the Resolution Island survey area. These years are not considered when assessing trends in the indices from the EAZ.

In 2014, the Central and Arctic Region reorganized the surveys conducted within the region. For shrimp, DFO stopped conducting the WAZ survey; since then, the area has been surveyed during the joint DFO-Northern Shrimp Research Foundation survey so that the WAZ, EAZ and SFA 4 (NAFO 2G) survey areas are all sampled with the same ship, gear and at the same time of year. While this resulted in restarting the time series in the WAZ, the future benefits to the assessment of shrimp outweighs this shortcoming. It should be noted that while all historical data are presented in this report, the two surveys are not directly comparable, as the surveys for the two zones were conducted with different ships, gear and at a different time of the year.

Resource status in the EAZ was evaluated within a Precautionary Approach (PA) framework (DFO 2006). Reference points (RP) were based on the geometric mean of female spawning stock biomass (SSB) during a seemingly productive time period. The Limit Reference Point (LRP) is 30% of the mean and the Upper Stock Reference (USR) is 80% of the mean. RPs for SFA 2 were developed based on estimates from available surveys (2006-2008) at that time (DFO 2009) and implemented in the Integrated Fisheries Management Plan (IFMP; DFO 2007b). These RPs were subsequently used in the EAZ when stock areas changed. Currently, there are no RPs for the WAZ.

Buffered random sampling was applied to allocation of sampling stations within the depth strata (Kingsley et al. 2004).

Upper and lower confidence intervals used in biomass and exploitation rate calculations, were estimated by resampling statistics (bootstrapping; Bruce et al. 2000).

Analysis and Response

Update of Indicators

Eastern Assessment Zone – *P. borealis*

Fishery

The total catch of *P. borealis* in the EAZ reported in the Canadian Atlantic Quota Report (CAQR), as of 7 January 2020, was 4,687 t, which is 56% of the TAC (Table 1, Figure 2). The 2019/20 fishery runs until 31 March 2020, thus catch records should be considered preliminary for 2019/20.

Biomass

The fishable biomass index increased by 102.8% from 2018 to 2019 and is now at the highest level (95,138 t) since the time series began (Table 2, Figure 3a). The female SSB index showed an increase of 74.0% and is currently at the second highest level (57,143 t) (Table 2, Figure 3b).

Exploitation

The reported exploitation rate (ER) index for 2019/20, as of 7January 2020, was 4.9% (Figure 4a). Since the fishery was still open at the time of the meeting, the reported ER may be higher at the end of the season. Should the entire TAC be taken this fishing season, the potential ER index for 2019/20 would be 8.8% (Figure 4b). The long term average of the potential ER is 15.4%, which is near the base target ER of 15% in the Healthy Zone for the EAZ (DFO 2018).

Current Outlook

The female SSB index for *P. borealis* in the EAZ is well within the Healthy Zone of the PA Framework (Figure 5).

Eastern Assessment Zone – *P. montagui*

Fishery

The total catch of *P. montagui* in the EAZ, as of 7 January 2020, was about 113 t (Table 1, Figure 6) and has been declining since 2001. The 2019/20 fishery runs until 31 March 2020, thus catch records are preliminary for 2019/20.

Biomass

Biomass indices for *P. montagui* in the EAZ have oscillated around the long term mean, with the exception of a particularly high biomass reported in 2012 (Figure 7). Both the fishable biomass and the female SSB indices decreased in 2019 to 8,503 t and 4,415 t, respectively, and were below the long term mean (Table 3, Figure 7a,b).

Exploitation

The reported ER index for 2019/20 was very low, 1.0%, due to low catches reported in the CAQR as of 7 January 2020 (Figure 8a). The potential ER index for this stock would be 9.9% if the entire TAC is taken (Figure 8b).

Current Outlook

The female SSB index in the EAZ has declined in 2019 placing the stock in the Cautious Zone (Figure 9). Given the wide fluctuations in biomass indices for *P. montagui* observed in the past (e.g., between years 2011, 2012 and 2013, and also last year) the status of this resource is considered uncertain.

Western Assessment Zone – *P. borealis*

While all survey years are presented, only the last six years of survey data can be considered in the WAZ assessment because of the change in the surveys, resulting in the start of a new time series.

Fishery

As of 7 January 2020, the CAQR records show that about 620 t, which equals to 19.6% of the TAC, have been taken (Table 1, Figure 10).

Biomass

The fishable biomass and female SSB indices decreased from 2018 to 2019 by 3.4% and 8.1%, respectively (Table 4, Figure 11a,b). The fishable biomass index in 2019 was 20,378 t, which was in the close proximity of the time series mean. The female SSB index was 11,845 t, which was above the time series mean.

Exploitation

The reported ER index for 2019/20 was relatively low, 3.4%, due to low catches reported in the CAQR as of 7 January 2020 (Figure 12a). As a consequence of the decline in the fishable biomass in 2019, the potential ER index has increased to about 15.5% (Figure 12b).

Current Outlook

Currently, there is no PA framework for *P. borealis* in the WAZ. The establishment of reference points for this resource is planned for 2020, to be implemented in 2021 stock assessment. Historical records show that the TAC for this stock has rarely been fully taken.

Western Assessment Zone – *P. montagui*

While all survey years are presented, only the last six years of survey data can be considered in the WAZ assessment because of the change in the surveys, resulting in the start of a new time series.

Fishery

The total catch of *P. montagui* was 6,884 t, which is 57.5% of the TAC as per 7 January 2020 CAQR (Table 1, Figure 13). The 2019/20 fishery runs until 31 March 2020 thus catch records should be considered preliminary for the 2019/20 season.

Biomass

The fishable biomass index decreased by 19.5%[1] from 2018 to 2019, with 64,268 t observed in 2019 (Table 5, Figure 14a), which was above the time series mean. A more pronounced biomass decrease (37.2%) was observed for the female SSB index, with 29,079 t estimated in 2019, which was in the close proximity of the time series mean (Figure 14b).

Exploitation

Along with decreasing fishable biomass, the reported ER index in 2019/20 increased to 11.3% (Figure 15a). If the entire TAC was taken the potential ER index would be 18.6% (Figure15b).

Current Outlook

Currently, there is no PA framework for *P. montagui* in the WAZ. The establishment of reference points for this resource is planned for 2020, to be implemented in 2021 stock assessment. Historical records show that the TAC for this stock has often been nearly fully taken. With the recently increased TAC it is unlikely that it will be fully taken in 2019.

Conclusions

Eastern Assessment Zone (EAZ)

Pandalus borealis

- Currently, the *Pandalus borealis* resource is in the Healthy Zone of the Precautionary Approach Framework.
- In 2019, the fishable biomass and female spawning stock biomass indices recorded relatively large increases; the fishable biomass is at the highest level while the female spawning stock biomass is at the second highest level since the time series began.
- The potential Exploitation Rate (ER) index for 2019/20 is 8.8%. That is below both the long term mean of the potential ER (15.4%) and the base target ER of 15% in the Healthy Zone for the EAZ.

Pandalus montagui

- *Pandalus montagui* biomass indices have fluctuated widely in the past, adding to the uncertainty about the status of the stock in the EAZ.
- Following this year's decrease in the biomass, the *Pandalus montagui* resource is currently in the Cautious Zone of the Precautionary Approach Framework.

[1] Erratum February 2020 – 17.5% now reads 19.5%

- The potential exploitation rate index if the entire TAC is taken in 2019/20 would be 9.9%.

Western Assessment Zone (WAZ)

Pandalus borealis

- The status of the stock is currently uncertain, as there is no Precautionary Approach Framework for *Pandalus borealis* in the WAZ.
- Both the fishable biomass and female spawning stock biomass indices declined slightly from 2018 to 2019.
- The reported exploitation rate index for 2019/20 is 3.4%. The current TAC equates to a potential exploitation rate index of 15.5%.

Pandalus montagui

- The status of the stock is currently uncertain, as there is no Precautionary Approach Framework for *Pandalus montagui* in the WAZ.
- Both the fishable biomass and female spawning stock biomass indices decreased from 2018 to 2019.
- The reported exploitation rate index for 2019/20 is 11.3%; the potential exploitation rate index for 2019/20 is 18.6%.

Contributors

- Wojciech Walkusz, DFO Science, Central and Arctic Region
- Sheila Atchison, DFO Science, Central and Arctic Region
- Chantelle Sawatzky, DFO Science, Central and Arctic Region
- Katherine Skanes, DFO Science, Newfoundland and Labrador Region
- Krista Baker, DFO Science, Newfoundland and Labrador Region
- Will Coffey, DFO Science, Newfoundland and Labrador Region
- Erika Parrill, DFO Centre for Science Advice, Newfoundland and Labrador Region
- Martin Henri, DFO Resource Management, Newfoundland and Labrador Region
- Courtney D'Aoust, DFO Resource Management, National Capital Region
- Brittany Beauchamp, DFO Fish Population Science, National Capital Region

Approved by

Sen Wang, Regional Director of Science, Central and Arctic Region

Lianne Postma, A/Division Manager, Arctic Aquatic Research Division, Central and Arctic Region

(Approved January 30, 2020)

(Erratum: February 19, 2020)

Sources of Information

Bruce, P., Simon, J.L., and Oswald, T. 2000. Resampling Stats User's Guide. Resampling Stats, Inc. Arlington, VA. 127 p.

DFO. 2006. A Harvest Strategy Compliant with the Precautionary Approach. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2006/023.

DFO. 2007a. Assessment Framework for Northern Shrimp (*Pandalus borealis*) off Labrador and the northeastern coast of Newfoundland; 28-30 May 2007. DFO Can. Sci. Advis. Sec. Proceed. Ser. 2007/034.

DFO. 2007b. Integrated Fisheries Management Plan: Northern Shrimp - Shrimp Fishing Areas (SFAs) 0-7 and the Flemish Cap.

DFO. 2008. Assessment of Northern Shrimp (*Pandalus borealis*) and Striped Shrimp (*Pandalus montagui*) in shrimp fishing areas 0, 2 and 3. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2008/018.

DFO. 2009. Proceedings of the Precautionary Approach Workshop on Shrimp and Prawn Stocks and Fisheries; November 26-27, 2008. DFO Can. Sci. Advis. Sec. Proceed. Ser. 2008/031.

DFO. 2010. Assessment of Northern Shrimp (*Pandalus borealis*) in SFA 0, 2, 3 and Striped Shrimp (*Pandalus montagui*) in SFA 2, 3 and 4 west of 63°W. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2010/024.

DFO. 2011. Assessment of Northern Shrimp (*Pandalus borealis*) and Striped Shrimp (*Pandalus montagui*) in Western and Eastern assessment zones (SFA 2 and 3). DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2011/010.

DFO. 2012. Monitoring update for Northern Shrimp (*Pandalus borealis*) and Striped Shrimp (*Pandalus montagui*) in the western and eastern assessment zones (SFA 2 and 3). DFO Can. Sci. Advis. Sec. Sci. Resp. 2012/001.

DFO. 2013. Assessment of Northern Shrimp (*Pandalus borealis*) and Striped Shrimp (*Pandalus montagui*) in the eastern and western assessment zones (Shrimp Fishing Areas 2 and 3). DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2013/031.

DFO. 2014. Update of stock status indicators for Northern Shrimp, *Pandalus borealis*, and Striped Shrimp, *Pandalus montagui,* in the western and eastern assessment zones. DFO Can. Sci. Advis. Sec. Sci. Resp. 2014/003.

DFO. 2015. 2015 Assessment of Northern Shrimp, *Pandalus borealis,* and Striped Shrimp, *Pandalus montagui,* in the eastern and western assessment zones. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2015/017.

DFO. 2016. Update of stock status indicators for Northern Shrimp, Pandalus borealis, and Striped Shrimp, *Pandalus montagui*, in the Western and Eastern Assessment Zones for 2016. DFO Can. Sci. Advis. Sec. Sci. Resp. 2016/006.

DFO. 2017. Assessment of Northern Shrimp, *Pandalus borealis*, and Striped Shrimp, Pandalus montagui, in the Eastern and Western Assessment Zones, February 2017. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2017/010. (Erratum: April 2017)

DFO. 2018. Integrated Fisheries Management Plan: Northern shrimp and striped shrimp – Shrimp fishing areas 0, 1, 4-7, the Eastern and Western Assessment Zones and North Atlantic Fisheries Organization (NAFO) Division 3M.

DFO. 2019. Assessment of Northern Shrimp, *Pandalus borealis*, and Striped Shrimp, *Pandalus montagui*, in the Eastern and Western Assessment Zones, February 2019. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2019/011.

Kingsley, M.C.S., Kanneworff, P., and Carlsson, D.M. 2004. Buffered random sampling: a sequential inhibited spatial point process applied to sampling in a trawl survey for northern shrimp *Pandalus borealis* in west Greenland waters. ICES Journal of Marine Science 61:12-24.

Siferd, T.D. 2014. An Assessment of Northern Shrimp and Striped Shrimp in the Eastern Assessment Zone and Western Assessment Zone (Shrimp Fishing Areas 2 and 3). DFO Can. Sci. Advis. Sec. Res. Doc. 2014/028. vi + 63 p.

Siferd, T.D. 2015. 2015 Assessment of Northern Shrimp *(Pandalus borealis)* and Striped Shrimp *(Pandalus montagui)* in the Eastern and Western Assessment Zones (SFAs Nunavut, Nunavik and Davis Strait). DFO Can. Sci. Advis. Sec. Res. Doc. 2015/010. v + 70 p.

Appendix 1. Tables

Table 1. Nominal reported catches (t) for the Eastern Assessment Zone and Western Assessment Zone for Pandalus borealis *and* P. montagui*.*

Year	Eastern Assessment Zone		Western Assessment Zone	
	P. borealis	***P. montagui***	***P. borealis***	***P. montagui***
2019*	4687	113	620	6884
2018	6198	234	1307	5531
2017	6488	233	918	5609
2016	6667	358	643	5660
2015	4816	59	353	4616
2014	4972	401	847	5836
2013	6793	1075	973	4775
2012	5555	1173	13	1105
2011	7687	135	0	857
2010	6908	483	57	345
2009	5159	564	0	0
2008	5184	808	0	0
2007	6359	1832	0	0
2006	6028	925	0	0
2005	6387	1427	-	0
2004	5842	2301	-	0
2003	5617	1217	-	0
2002	5695	3081	-	0
2001	6275	3867	-	0
2000	5718	4238	-	0
1999	5465	3780	-	0
1998	5372	3360	-	0
1997	5870	3050	-	0
1996	33467	3058	-	0
1995	2489	3192	-	0
1994	456	154	-	0
1993	68	0	-	0
1992	1210	1	-	0
1991	1150	623	-	0
1990	1634	174	-	5
1989	3133	1265	-	10
1988	2873	603	-	13
1987	0	0	-	0

Table 1. continued

Year	Eastern Assessment Zone		Western Assessment Zone	
	P. borealis	***P. montagui***	***P. borealis***	***P. montagui***
1986	50	483	-	0
1985	0	0	-	0
1984	0	0	-	0
1983	21	0	-	0
1982	46	0	-	0
1981	1	9	-	2
1980	487	103	-	3
1979	1	58	-	25

** Catch based on CAQR as of 7 January 2020. Since the fishery is still open the catch is preliminary for 2019.*

Table 2. Fishable and female spawning stock biomass estimates for Pandalus borealis *in the Eastern Assessment Zone for the 2006-2019 surveys. LCL and UCL are the lower and upper 95% confidence limits. Year over year (YOY) change indicates the relative change in comparison to the previous year.*

Year	Biomass	YOY change (%)	Weight (tonne)		
			Mean	LCL	UCL
2019	Fishable	102.8	95138	48333	146788
2018	Fishable	19.6	46900	36344	58928
2017	Fishable	-40.2	39198	30225	48907
2016	Fishable	-17.0	65570	42137	93569
2015	Fishable	56.5	78984	50852	106962
2014	Fishable	1.5	50458	38914	62340
2013	Fishable	-17.9	49697	38427	60631
2012	Fishable	-22.9	60534	43074	79960
2011	Fishable	10.5	78530	23900	135037
2010	Fishable	-9.8	71065	40234	108703
2009	Fishable	54.3	78755	48850	110115
2008	Fishable	17.9	51053	37117	66708
2007	Fishable	32.0	43306	31015	58346
2006	Fishable	-	32816	21969	44152
2019	Female SS	74.0	57143	28420	87654
2018	Female SS	32.4	32842	23548	44126
2017	Female SS	-28.8	24800	19888	30252
2016	Female SS	-42.8	34827	24220	46979
2015	Female SS	78.7	60869	33379	88386
2014	Female SS	6.3	34069	25157	43000
2013	Female SS	-22.2	32049	26762	37607
2012	Female SS	-13.8	41190	29498	54383
2011	Female SS	9.1	47807	13470	82926
2010	Female SS	12.7	43800	19025	79665
2009	Female SS	40.5	38856	23122	56820
2008	Female SS	-0.2	27653	22507	39368
2007	Female SS	64.8	27698	19249	39007
2006	Female SS	-	16805	10523	23026

Table 3. Fishable and female spawning stock biomass estimates for Pandalus montagui *in the Eastern Assessment Zone for the 2006-2019 surveys. LCL and UCL are the lower and upper 95% confidence limits. Year over year (YOY) change indicates the relative change in comparison to the previous year.*

Year	Biomass	YOY change (%)	Weight (tonne)		
			Mean	LCL	UCL
2019	Fishable	-59.3	8503	3930	13948
2018	Fishable	-16.3	20895	12617	29450
2017	Fishable	81.0	24957	17246	32311
2016	Fishable	124.7	13792	6452	21126
2015	Fishable	-63.0	6137	3445	8629
2014	Fishable	371.0	16600	11203	22084
2013	Fishable	-87.8	3524	1738	6208
2012	Fishable	272.7	28845	8582	48946
2011	Fishable	4.3	7740	2871	14285
2010	Fishable	-52.7	7423	5714	9290
2009	Fishable	6.9	15679	6190	29774
2008	Fishable	203.8	14667	7287	21973
2007	Fishable	81.0	4828	3389	6673
2006	Fishable	-	2667	210	5122
2019	Female SS	-68.0	4415	1742	7275
2018	Female SS	-19.8	13806	9362	20052
2017	Female SS	64.4	16537	9866	23250
2016	Female SS	159.4	10056	2986	17280
2015	Female SS	-69.5	3877	2085	5452
2014	Female SS	357.1	12696	8834	16622
2013	Female SS	-88.2	2778	1301	4949
2012	Female SS	653.8	23552	6218	40985
2011	Female SS	-46.3	3124	1599	4721
2010	Female SS	-33.7	5819	4509	7136
2009	Female SS	-17.7	8776	4205	13955
2008	Female SS	440.9	10660	4269	17047
2007	Female SS	-7.7	1971	903	3490
2006	Female SS	-	2134	50	4219

Table 4. Fishable and female spawning stock biomass estimates for Pandalus borealis *in the Western Assessment Zone. Year over year (YOY) change indicates the relative change in comparison to the previous year. Note 2014 represents a start of the new time series and values are not directly comparable to previous years. LCL and UCL are the lower and upper 95% confidence limits.*

Year	Biomass	YOY change (%)	Weight (tonnes)		
			Mean	LCL	UCL
2019	Fishable	-3.4	20378	12852	29080
2018	Fishable	101.0	21088	12627	33452
2017	Fishable	-20.0	10487	5073	17185
2016	Fishable	-54.0	13116	7867	18868
2015	Fishable	31.4	28532	18531	39501
2014	Fishable	-1.3	21713	14353	31046
2013	Fishable	11.7	21999	15906	28518
2011	Fishable	26.7	19692	12468	27961
2009	Fishable	6.4	15544	7613	25529
2007	Fishable	-	14615	4907	28872
2019	Female SS	-8.1	11845	7529	16299
2018	Female SS	147.0	12884	7121	19203
2017	Female SS	-34.9	5216	3045	7676
2016	Female SS	-45.5	8015	4780	11590
2015	Female SS	19.5	14710	9270	20379
2014	Female SS	25.8	12309	8792	16398
2013	Female SS	53.5	9785	7106	12829
2011	Female SS	66.1	6377	4182	8909
2009	Female SS	18.8	3840	1154	7479
2007	Female SS	-	3231	1687	5361

Table 5. Fishable and female spawning stock biomass estimates for Pandalus montagui *in the Western Assessment Zone. Year over year (YOY) change indicates the relative change in comparison to the previous year. Note 2014 represents a start of the new time series and values are not directly comparable to previous years. LCL and UCL are the lower and upper 95% confidence limits.*

Year	Biomass	YOY change (%)	Weight (tonnes)		
			Mean	LCL	UCL
2019	Fishable	-19.5	64268	29711	112173
2018	Fishable	77.7	79835	34057	132111
2017	Fishable	41.6	44915	29179	63381
2016	Fishable	-42.5	31724	19507	44908
2015	Fishable	-28.4	55194	35769	76429
2014	Fishable	68.9	77078	44854	111562
2013	Fishable	-36.2	45647	32899	59438
2011	Fishable	53.3	71558	40264	108612
2009	Fishable	-13.6	46673	25756	73342
2007	Fishable	-	54044	17007	99461
2019	Female SS	-39.2	29079	14930	45581
2018	Female SS	57.8	47834	19926	81534
2017	Female SS	62.1	30305	18830	43434
2016	Female SS	-31.6	18691	11090	27334
2015	Female SS	-29.7	27324	18282	37041
2014	Female SS	44.2	38875	23553	55849
2013	Female SS	-17.2	26955	18616	35736
2011	Female SS	80.8	32549	20296	46119
2009	Female SS	-6.6	17999	9775	28160
2007	Female SS	-	19277	5668	36606

Appendix 2. Figures




Figure 1. Location of the Western and Eastern assessment zones (left panel) and corresponding Shrimp Fishing Area (SFA) management units (right panel). Boundaries of the Nunavut (NU), Nunavik (NK) and Nunatsiavut Land Claims Areas are identified with red lines. Abbreviations for Davis Strait (DS), East (E) and West (W) are used.




Figure 2. Eastern Assessment Zone Pandalus borealis TAC and catch recorded in the CAQR. Catch based on CAQR as of January 7, 2020; since fishery is still open the catch is not complete for 2019/20. Quota for 1999-2012/13 renamed to correspond to new management areas.




Figure 3. Eastern Assessment Zone Pandalus borealis a) fishable and b) female spawning stock biomass indices for the survey years 2006-2019. Error bars are 95% confidence ranges.




Figure 4. Eastern Assessment Zone Pandalus borealis a) reported and b) potential exploitation rate indices for the period of 2006/07-2019/20. Error bars are 95% confidence ranges.




Figure 5. Eastern Assessment Zone trajectory of Pandalus borealis female spawning stock biomass and exploitation rate indices in relation to reference points. USR=Upper Stock Reference and LRP=Limit Reference Point. Error bars are 95% confidence ranges.




Figure 6. Eastern Assessment Zone Pandalus montagui TAC and catch recorded in the CAQR. Catch based on CAQR as of January 7, 2020. Since fishery is still open the catch is not complete for 2019/20. New management units were implemented for the 2013/14 season.




Figure 7. Eastern Assessment Zone Pandalus montagui a) fishable and b) female spawning stock biomass indices for the survey years 2006-2019. Error bars are 95% confidence ranges.




Figure 8. Eastern Assessment Zone Pandalus montagui a) reported and b) potential exploitation rate indices for 2006/07-2019/20. Error bars are 95% confidence ranges.




Figure 9. Eastern Assessment Zone trajectory of Pandalus montagui female spawning stock biomass and exploitation rate indices in relation to reference points. USR=Upper Stock Reference and LRP=Limit Reference Point.




Figure 10. Western Assessment Zone Pandalus borealis TAC and catch recorded in the CAQR. Catch based on CAQR as of January 7, 2020; since the fishery is still open the catch is not complete for 2019/20. New management units were implemented for the 2013/14 season.




Figure 11. Western Assessment Zone Pandalus borealis, a) fishable biomass and b) female spawning stock biomass indices for the four years of DFO/Cosmos surveys and six years conducted by the NSRF-DFO/Campelen. Biomass indices from the two surveys are not directly comparable. 2014 represents the start of a new time series for the WAZ. Error bars are 95% confidence ranges.




Figure 12. Western Assessment Zone Pandalus borealis a) reported and b) potential exploitation rate indices for the four years of DFO/Cosmos surveys and six years conducted by the NSRF-DFO/Campelen. Exploitation rate indices from the two surveys are not directly comparable. 2014 represents the start of a new time series for the WAZ. Error bars are 95% confidence ranges.




Figure 13. Western Assessment Zone Pandalus montagui TAC and catch recorded in the CAQR. Catch based on CAQR as of January 7, 2020; since fishery is still open the catch is not complete for 2019/20. New management units implemented for the 2013/14 season.




Figure 14. Western Assessment Zone Pandalus montagui, a) fishable biomass and b) female spawning stock biomass indices for the four years of DFO/Cosmos surveys and six years conducted by the NSRF-DFO/Campelen. Biomass indices from the two surveys are not directly comparable. 2014 represents the start of a new time series for the WAZ. Error bars are 95% confidence ranges.




Figure 15. Western Assessment Zone Pandalus montagui a) reported and b) potential exploitation rate indices for the four years of DFO/Cosmos surveys and six years conducted by the NSRF-DFO/Campelen. Exploitation rate indices from the two surveys are not directly comparable. 2014 represents the start of a new time series for the WAZ. Error bars are 95% confidence ranges.

This Report is Available from the:

Center for Science Advice
Central and Arctic Region
Fisheries and Oceans Canada
501 University Crescent
Winnipeg, Manitoba
R3T 2N6

Telephone: (204) 983-5131
E-Mail: xcna-csa-cas@dfo-mpo.gc.ca
Internet address: www.dfo-mpo.gc.ca/csas-sccs/

ISSN 1919-3769
© Her Majesty the Queen in Right of Canada, 2020



Correct Citation for this Publication:

DFO. 2020. Update of stock status indicators for Northern Shrimp, *Pandalus borealis,* and Striped Shrimp, *Pandalus montagui,* in the Western and Eastern Assessment Zones, January 2020. DFO Can. Sci. Advis. Sec. Sci. Resp. 2020/014. (Erratum: February 2020)

Aussi disponible en français :

MPO. 2020. Mise à jour des indicateurs de l'état des stocks des crevettes nordiques, Pandalus borealis*, et les crevettes ésopes,* Pandalus montagui*, dans les zones d'évaluation ouest, janvier 2020. Secr. can. de consult. sci. du MPO, Rép. des Sci. 2020/014. (Erratum : février 2020)*


ⵊⵔⵉⵔⴰⵣⵓⵛ: **ⴰⵔⵍⵛⴻⵙⵜ:** X **ⴰⵊⵛⵔⵙⵜⵓⵔⵜ:** X


ⵏⵜⵏⵓⵖ: **ⴰⵔⵍⵛⴻⵙⵜ ⴰⵏ ⴰⵊⵛⵔⵙⵜⵓⵔⵜ ⴻⵔⴻⵙⵜⵛⵓⵛ ⴻⵔⴻⵙⵜⵛⵓⵛ ⵊⵔⵉⵔⴰⵣⵓⵛ ⵔⵜⵛⵜⵛ ⴻⵔⴻⵙⵜⵛⵓⵛ ⵊⵔⵉⵔⴰⵣⵓⵛ ⴰⵏ ⵜⵃⵔⵛⵜⵛⵜⵔⵓⵛ ⴰⵙⵃⵔⵜⵛⵜⵔⵓⵛ ⴰⵔⴰⵙⵓ ⴰⵏ ⴰⵛⵊⴰⵓⵛⵜⵏ ⵛⵔⵔⵜⵛⵜⵔⵓⵛ**

ⵔⵔⵜⵛⵜⵔⵓⵛ:

ⵊⵛⵔⵜⵛⵊⴰ – ⵃⵛⵔⵛⵏⵓ ⵜⵃⵔⵏⵜⵛⵊⴰⵓⵛⵜⵔⵓⵛ ⵔⵔⵜⵛⵜⵔⵓⵛ

ⵊⵛⵔⵔⵜⵛⵊⴰ – ⵏⵛⵏⵔⵛⵏⵓ ⵜⵃⵔⵏⵜⵛⵊⴰⵓⵛⵜⵔⵓⵛ ⵔⵔⵜⵛⵜⵔⵓⵛ





ⵜⵔⵜⵛⵜⵔⵓⵛ ⵔⵛⵊⵑⵑⴰⵛ (*Pandalus borealis*)



ⵜⵃⵔⵛⵔⵜⵛ ⵔⵛⵊⵑⵑⴰⵛ (*Pandalus montagui*)



ᐅᑭᐅᑕᖅᑐᒥ ᐃᓗᐊᑦ (*Boreogadus saida*)

ᓄᓇᓕᐊᔪᖅᓯᒪᓂᖓᑦ

ᐊᒪᖏᐅᑕᓪᓚᓴᒃᓯᑦ ᑲᓇᑕᒥ (DFO) ᐱᓕᓪᓚᖅᑎᖅᓯᒪᔪᑦ ᓄᓇᕗᒻᒥ ᐅᒪᔪᓕᓪᓚᓴᕐᔪᐊᒃᓯᑦ ᑲᑎᒪᔨᖓᓂᓂᓂ ᐊᒪᖏᐅᓂᖓ ᐅᒪᔪᓕᓪᓚᓴᓂᑦ (ᑲᑎᒪᔨᑦ) ᐱᒃᔪᑎᖅᑎᖅᑐᓂ ᐅᑭᐅᑕᖅᑐᒥ ᐅᓚᖅᓂ ᐊᖏᓪᓕᐅᑕᐅᓚᖏᑦᑐᑦᑕᖅᑐᓂ ᐅᑭᐅᑕᖅᑐᒥ ᑭᓯᐊᓂᓂ ᐊᒪᒪ ᖃᓄᖅᑕᓂᓂᓪᓚᖏᑦ, ᐱᒋᐊᖅᑲᖅᑐᓂ ᐅᖃᐊᖅᓴᖅᓂᓂ 2018. ᑐᓴᖓᐸᓪᓕᔾᔪᑎᓂᑦ ᖃᐅᔨᓴᖅᑎᓄᓂ ᐊᒪᒪ ᐊᖏᓪᓕᐅᑕᐅᔪᓂᑦ ᑐᑭᓯᐅᑦᑎᐅᓂᑦ ᐊᑐᐊᓱᓕᐅᔪᓂ ᐅᑭᐅᑕᖅᑐᒥ ᐅᓚᓴᓂᑦ (ᑕᑯᔭ ᐊᓕᔪᑕᖓ 1), ᓄᓇᒋᖓᓚᖅᑐᓂᑦ ᐊᐅᓕᑎᓂᓵᖅ (RM) ᖃᐅᔨᓴᖅᑐᑦ ᐱᔪᓱᓇᑎᑎᔪᒥᑦ ᖁᑯᐊᒥᓕᔪᓱᓂᖅᓴᒪᒥ ᐊᖏᓪᓕᑕᐅᓕᕐᓂᕐᑐᓂ ᐊᖏᓪᓕᑕᐅᓕᔪᓂ ᐱᖓᖅᑐᐊᐃᓂᖏᓪᓕᐅᓚᕐᓂᕐᑐᖅ ᐱᐅᓂᐅᓕᔪᓂ ᑕᒃᑯᓗᖓ ᐅᒋᐊᖅᑕᐅᓂ 2018/19 ᐊᒪᒪ 2019/20 ᐱᖓᓱᓇᖓᓂ ᐱᓱᖓᖓ (WAZ) (ᐃᓕᔪᑕᖓ 2). ᐃᓱᓇᕆᑦᔪᓕᐅᖅᑐᑦ, ᐱᓕᓪᓚᖅᑕᔪᖅ ᑐᓂᓯᓕᐅᖅᑐᖅ ᑐᐱᔪᑎᑎᑎᐅᑐᓂ ᐅᓚᖅᓂ ᐊᖏᓪᓕᑕᐅᓕᕐᓂᕐᑐᓂᑐᓂ ᐊᖏᓪᓕᑕᐅᓕᔪᓂ ᐊᐅᓕᑎᓂᓵᓂᒪᑦ ᑐᐱᔪᑎᑎᑎᐅᑐᓂ ᑎᑎᕋᕐᓂᖅ 2019 ᓄᖅᓚᓂᓕᓪᓚᖅᑐᓂ ᐅᖃᓕᓂᐅᔪᓂᑦ ᐊᐅᓕᑎᓂᓵᓂᒪᑦ ᑐᐱᔪᑎᑎᑎᐅᑐᓂ 2019 ᓄᖅᓚᓂᕐᑐᓄᓂ (ᐃᓕᔪᑕᖓ 3).

ᐱᓕᓪᓚᖅᑕᔪᖅ ᑲᔪᕐᑐᖅᓯᒪᔪᓂ ᖃᓄᓐᓄᓂ ᖃᐅᔨᓴᖅᑎᓂᓂ ᐅᖃᐅᔨᓴᕐᔪᐊᒃᓯᑦ ᒪᓕᒪᑕᑦᓴᕐᔪᐊᓂᓂ ᐃᖅᓯᓕᑕᐅᓕᕐᓂᕐᑐᓂᑐᓂ ᐃᖅᓯᓕᑕᐅᓕᔪᓂ ᐅᓚᖅᓂ ᓕᓕᓂᐊᕐᓂᖅ ᐅᖃᐅᔨᓴᕐᔪᐊᒃᓯᑦ ᐃᖅᓯᓕᑕᐅᓕᔪᓂ ᐊᐅᓕᑎᓂᓵᓂᒪᑦ ᑐᐱᔪᑎᑎᑎᐅᑐᓂ ᐱᑕᖃᓂᒃᑯᑦᑐᓂ (CSAS) ᐱᓴᓕᓕᐊᓂᖓ ᐱᓕᓪᓚᕐᓇᖅᑐᖅᑐᓂ ᐅᖃᐅᔨᓴᕐᔪᐊᒃᓯᑦ 2019 (ᑕᑯᔭ ᖃᐅᔨᓴᖅᑎᓂᓂ ᑐᐱᔪᑎᑎᑎᐅᑐᓂ).

ᐃᓕᓴᕐᔪᐊᔪᖅ, ᐱᓕᓪᓚᖅᑕᔪᖅ ᑲᔪᕐᑐᖅᓯᒪᔪᓂ ᓄᓇᒪᓂᕐᓂᓂᖃᖅᓴᓂᑦ ᐊᑯᓂᐅᓂᓂᖅᓴᒪᒥ ᐊᐅᓕᑎᓂᓵᓂᒪᑦ ᐅᓚᖅᓂ ᐃᖅᓯᓕᑕᐅᓕᕐᓂᕐᑐᓂᑐᓂ ᐃᖅᓯᓕᑕᐅᓕᔪᓂ ᐃᒪᐃᓐᓇᕆᑦ ᐅᒥᐊᕐᓂᕐᑐᓂ ᐃᖅᓯᓕᑕᐅᓕᔪᓂ ᐅᓚᖅᓂ ᐊᐅᓕᑎᓂᓵᓂᒪᑦ ᐊᒪᒪ ᓄᓇᒪᓂᖅ ᐃᖅᓯᓕᑕᐅᓕᔪᓂ ᐅᑭᐅᑕᖅᑐᒥ. ᓄᓇᒪᓂᖅᓴᒃᑯᓂᖅᓴᓂᓂ ᐃᖅᓯᓕᑕᐅᓕᔪᓂ ᐅᑭᐅᑕᖅᑐᒥ ᐊᒪᒪ ᓄᓇᒪᓂᖅ ᐃᖅᓯᓕᑕᐅᓕᔪᓂ ᐅᑭᐅᑕᖅᑐᒥ

ᐅᖃᓕᐃᓐᓇᕐᑎᒍᑦ ᑭᖑᑦᑕᑦᑕᐅᔪᓂᑦ ᐊᖏᕈᑎᔨᖅᑕᖓ ᐃᓐᓇᐅᔪᓕᑦ ᐅᑎᖃᓂᑦ ᐃᖅᓯᑕᑦᑕᐅᑎᐊᕐᓂᑐᓂ ᐃᖅᓯᑕᑦᑕᐅᔪᓂᑦ ᐃᓕᒍᑦᑕᐅᔪᓂᑦ SFA 1 (ᐃᓕᒍᑦᑕᖓ 2).

ᖃᐅᔨᓴᖅᑎᓂᑦ ᑐᑭᓯᐊᕐᓗᒃᑖᑦ

ᖃᐅᔨᓴᖅᑎᓂᑦ ᑐᑭᓯᐊᕐᓗᒃᑖᑎᑦ ᐅᑎᖃᓂᑦ ᑐᓂᔨᐅᓕᐅᖅᑐᑦ ᑲᑎᒪᔨᓄᑦ ᓇᓗᓇᐃᔨᐃᓂᕐᒥᑦ ᑐᑭᓯᐊᕐᓗᒃᑖᑎᓄᑦ ᑎᓯᐊᓕᒃ ᑲᑎᒪᔨᓄᑦ 1. ᓴᖅᑭᖅᑎᑕᐅᔪᓂᑦ ᐱᒋᐊᖅᑐᓂ ᐊᓯᐊᓂᒃ 2019 CSAS ᖃᒥᖅᑭᓂᕐᒥᑦ ᐱᓕᕆᓕᐊᓂᖅ ᐱᖅᓯᐅᑎᓪᓗᑎᑦ ᑎᑎᕋᖅᑕᐅᔪᓂ, ᖃᐅᔨᓴᖅᓂᕐᒧᑦ ᑎᑎᕋᖅᓯᒪᔪᓂ (ᓇᓗᓇᐃᔭᖅᓯᒪᔪᓂ ᐃᓕᒍᑦᑕᖓ 6) ᐊᒻᒪ ᖃᐅᔨᓴᖅᑎᓂ ᐅᖃᐅᔨᓴᐅᔪᓂᕐᒧᑦ ᐅᓂᒃᑲᓂ (ᓇᓗᓇᐃᔭᖅᓯᒪᔪᓂ ᐃᓕᒍᑦᑕᖓ 7), ᐊᓯᓇ ᐊᑐᐃᓐᓇᐅᑦᑕᐅᑦ.

ᐃᓱᒪᓕᓪᓕᒋᑦ ᒥᑭᓂᖅᓯᓇᓂᑦ ᑲᑎᑎᑦᑕᑦ ᑐᖃᓄᖅᑕᖅᑐᑦ (ᑐᖓᔪᓂ 0.1% ᐃᖃᓗᓂᑦ ᐊᖏᕿᔭᒪᓕᒡᓗᑦ ᑐᓴᓱᖁᔨᖅ ᖁᓱᓕᓂᑐᖅ ᑐᓴᖓᓂᖃᖅᖁᓂᑦ ᓂᓪᓚᕐᓇᖅᑐᓂᓄᑦ ᐱᑕᖃᓚᐊᖃᖏᓂᕐᒧᓄᑦ) ᐅᑭᐅᑕᖅᑐᒥ ᐅᓗᐃ ᐱᒋᐊᖅᓯᒪᓕᓪᓕᒋᑦ ᑲᑎᒪᔨᓄᑦ ᑲᑎᒪᔨᓄᑦ ᑭᓚᐅᑕᖅᑐᒥ ᐅᑭᐅᑕᖅᑐᒥ ᑭᖑᑦᑕᑦᑕᐅᔪᓂᑦ ᐱᖓᓱᖅᑐᒥᓂᑦ ᐃᓚᖓᑕᐅᓂᖅ, ᐃᓱᒪᑎᐃᓕᐊᓂᖅᖁᖅ ᐊᑦᑕᐃᐅᑕᖃᖏᓂᖅ ᑭᖑᑦᑕᓕᒃ ᑲᑎᒪᔨᓄᑦ ᐱᓕᕆᔨᒃᑲᓂᑦ ᐅᑎᖃᓂᑦ ᐊᒥᓱᓕᕆᔨᒃᑲᓂᑦ ᑭᖑᑦᑕᓕᒃ ᑭᖑᓂᖓᓄᑦ.

ᓇᓕᖏᐊᔪᓚᐅᑦ

1) ᓴᖅᑭᑎᑎᖅᑕᓂ ᐅᑭᐅᑕᖅᑐᒥ ᐅᑎᖃᓂᑦ ᐃᖃᓗᑦᑕᐅᑎᐊᕐᓂᑐᓂ ᐃᖃᓗᑦᑕᐅᔪᓂᑦ ᑭᓕᓕᖓᓂ ᐊᓂᖅᓴᐅᔪᒍᑦ 5% ᐅᖃᓕᐃᓐᓇᕐᑎᒍᑦ ᑭᖑᑦᑕᐃᑦ ᐅᖅᓯᓕᓵᓂᑦ 200kg ᐊᑕᕐᓂᖅᑐᑦ, ᑐᓴᖓᓂᖃᖅᑐᑦ ᐊᑐᖅᑕᐅᓚᔨᑦᑐᓂᑦ ᐱᖓᓱᓕᔨᖅᑐᓂᑦ ᑲᓕᓐᓂᐅᕐᑕᒪᔭᖅᑐᓂᑦ, ᐊᑕᕐᓂᖅᑐᑦ.

2) ᓴᖅᑭᑎᑎᖅᑕᓂ ᐅᑭᐅᑕᖅᑐᒥ ᐅᑎᖃᓂᑦ ᐃᖃᓗᑦᑕᐅᑎᐊᕐᓂᑐᓂ ᐃᖃᓗᑦᑕᐅᔪᓂᑦ ᑭᓕᓕᖓᓂ ᐊᓂᖅᓴᐅᔪᒍᑦ 10% ᐅᖃᓕᐃᓐᓇᕐᑎᒍᑦ ᑭᖑᑦᑕᐃᑦ ᐅᖅᓯᓕᓵᓂᑦ 200kg ᐊᑕᕐᓂᖅᑐᑦ, ᑐᓴᖓᓂᖃᖅᑐᑦ ᐊᑐᖅᑕᐅᓚᔨᑦᑐᓂᑦ ᐱᖓᓱᓕᔨᖅᑐᓂᑦ ᑲᓕᓐᓂᐅᕐᑕᒪᔭᖅᑐᓂᑦ, ᐊᑕᕐᓂᖅᑐᑦ.

ᑲᑎᒪᔨᑦ ᐃᓱᒪᓕᐅᕈᓇᖅᑐᑦ ᐊᒻᒪ/ᐅᕐᓂᓚᓵᓂᑦ ᐊᑐᓕᖏᓕᐅᕐᓕᓗᑎ, ᓇᓴᒪᑎᓪᓕᒍᔪ, ᐃᖃᓗᑦᑕᐅᑎᐊᑦᑐᓂ ᐃᖃᓗᑦᑕᐅᔪᓂᑦ ᑭᓕᓕᖓᓂ ᐊᑲᐅᔪᓂᕐᓂᖃᖅᖁᓂᑦ ᐊᑐᓂᖓᓂᓵᓂ, ᐊᒻᒪ /ᐅᕐᓂᓚᓵᓂᑦ ᐊᓕᓕᓂᕐᑕᒪᓂ ᐊᖏᓂᖓᓂᓵᓂᑦ ᑐᑭᓂᖓᓂᓵᓂᑦ ᐊᑐᓂᖓᓂᓵᓂᑦ ᑐᑭᓯᐊᕐᓗᒃᑖᑦ.

ᓇᓗᓇᐃᔨᖅᓯᒪᓂᖓᓂᑦ ᑐᓴᕐᕋᐅᑦ

ᐃᓱᒪᓕᓪᓕᒋᑦ ᖃᓄᐃᓂᑦ, ᐱᓕᕆᔨᐅᔪᖅᑭᑎᓂᑦ ᑲᑎᒪᔨᓄᑦ, ᐊᑐᓕᖅᑎᑕᐅᓂᖏᓐᓄᑦ ᐱᖓᓱᐃᓕᓴᕐᓂ 2020/21 ᐃᖃᓗᓕᕆᓂᐅᔪᖅ:

1) ᐃᓱᒪᓕᐅᕐᓂᖅ ᐅᑭᐅᑕᖅᑐᒥ ᐅᑎᖃᓄᑦ ᐃᖃᓗᑦᑕᐅᑎᐊᕐᓂᑐᓂ ᐃᖃᓗᑦᑕᐅᔪᓂᑦ ᐃᓗᐊᓂ (WAZ, ᐊᒻᒪ NU/NK-E) ᐊᒻᒪ ᐊᑐᓕᖏᓕᐅᕐᓕᒃᑐᑦ ᐅᑭᐅᑕᖅᑐᒥ ᐅᑎᖃᓄᑦ ᐃᖃᓗᑦᑕᐅᑎᐊᕐᓂᑐᓂ ᐃᖃᓗᑦᑕᐅᔪᓂᑦ ᑭᖑᑦᑕᓕᒃ ᐊᓕᓕᓂᕐᑕᒪᓂ ᐊᓕᓕᓂᕐᑕᒪᓂᖅᑐᑦ ᓄᓇᕗᒻᒥ.

ᑎᑎᕋᖅᑐᖅ: ᐃᒪᓕᕆᐅᑕᓪᓗᓴᐅᑉᑯᑦ ᐃᒪᓕᕆᔨᖅᑐᑎᖅᑐᓂᑦ ᐊᐅᓚᑦᑎᓂᕐᓄᑦ ᐊᒻᒪ ᑕᓪᓕᐅᕐᓕᕆᐅᑕᓪᓗᓴᐅᑉᑯᑦ ᑲᓇᑕᒥ

ᐅᑉᓗᖓ: ᓂᕆᐅᐊᓪ 7, 2020

ᐃᓕᔪᑦᖏᑦ

ᐃᓕᔪᑦᖅ 1 – ᐃᖅᓗᑦᑕᐅᒋᐊᖏᑦᑐᑦ ᐃᖅᓗᑦᑕᐅᖅᑕᖅᓯᒪᔾᔪᑎᖅᐃᑦ ᐅᑭᐅᑕᖅᑐᒥ ᔪᖓᓴᐸᓂᑦ ᐃᖅᓗᓚᕆᓂᖅᑎᑦ, ᐅᑭᐅᑕᖅᑐᒥ ᐅᓴᐃᑦ

ᐃᓕᔪᑦᖅ 2 – ᖃᓄᐃᓕᓂᕆᐊᓕᖏᑦ ᓪᓚᕆᓴᓯ ᐊᑲᖅᑭᐊᖅᑐᑦ ᐅᑭᐅᑕᖅᑐᒥ ᑭᖓᓴᐸᓄᑦ (2017-2019)

ᐃᓕᔪᑦᖅ 3 – ᑐᕆᕋᕆᐃᑦ (ᑐᑭᓯᐊᕈᑎᓄᑦ): ᐊᑐᓕᖅᑎᑕᐅᓂᖅᖓ ᐅᑭᐅᑕᖅᑐᒥ ᐅᓴᐃᑦ ᐃᖅᓗᑦᑕᐅᒋᐊᖏᑦᑐᑦ ᐃᖅᓗᑦᑕᐅᔾᔨᑦ ᐊᐅᓪᓕᑎᑎᐊᕈᑎᑦ 2019/20 ᐅᑭᐅᑕᖅᑐᒍᑦ ᐊᓪᒪᒃ ᖃᓄᖅᑕᓕᓂᓄᑦ ᔪᖓᓴᐸᓂᑦ ᐃᖅᓗᓚᕆᓂᖅᑎᑦ ᐱᖓᓱᓪᓗᓂ ᖃᐅᔨᓴᕐᕕᐅᔾᔨᑦ ᑭᓪᓕᖓ

ᐃᓕᔪᑦᖅ 4 – ᑲᑎᒪᑎᑦᑎᓂᕐᒧᑦ ᐃᓪᓕᐅᖃᑕᐅᔾᔨᑦ ᑎᑎᕋᖅᓯᒪᔾᔨᑦ (*ᔫᓂᓄᐊᖓᓚ 2020*)


ᐃᓕᔪᑦᖅ 5 – ᑲᑐᑎᔾᔨᑦ ᐃᖅᓗᓚᐱᓂᐅᔾᔨᑦ ᑭᐅᔾᔨᑦ ᐅᑭᐅᑕᖅᑐᒥ ᐅᓴᐃᑦ ᐃᖅᓗᑦᑕᐅᒋᐊᖏᑦᑐᑦ ᐃᖅᓗᑦᑕᐅᔾᔨᑦ ᐊᐅᓪᓕᑎᑎᐊᕈᑎᑦ ᐊᓪᒪᒃ ᒪᖅᑭᑎᑦᑎᓂᕆᑦ ᐊᑲᖅᑭᐊᖅᓯᒪᔾᔨᓂᑦ ᖃᓄᐃᓕᓂᕆᐊᓕᖏᑦ ᓪᓚᕆᓴᓯ (COL)

ᐃᓕᔪᑦᖅ 6 - ᓇᓗᓇᐃᔨᖅᓯᒪᓂᖏᑦ: ᖃᐅᔨᓴᖅᑕᐅᓂᖅᖓ ᐊᑦᑐᖅᑕᐅᔾᔨᓇᖅᑐᑦ ᐃᖅᓗᑦᑕᐅᒋᐊᖏᑦᑐᑦ ᐃᖅᓗᑦᑕᐅᔾᔨᑦ ᑐᔾᔭᖃᑦᑕᕐᓂᖏᑦ ᐅᑭᐅᑕᖅᑐᒥ ᐅᑎᖅᓄᑦ *(Boreogadus saida)* ᐱᑕᖃᖅᓂᖏᓐᓂᑦ ᐅᑭᐅᑕᖅᑐᒥ *(Pandalus borealis)* ᐊᓪᒪᒃ ᖃᐅᔨᖅᑕᓪᓕᑦ *(Pandalus montagui)* ᑭᖓᓴᐸᓂᑦ ᐃᖅᓗᓚᕆᓂᖅᑎᑦ ᑭᖓᓴᐸᓂᐊᖅᓱᐃᑦᑐᔾᔨᓂᑦ 1, 2, ᐊᓪᒪᒃ 3 (ᖃᐅᔨᓴᓂᖅᓯᒪᒍᑦ ᑎᑎᕋᖅᓯᒪᔾᔨᑦ 2019/073)


ᐃᓕᔪᑦᖅ 7 - ᓇᓗᓇᐃᔨᖅᓯᒪᓂᖏᑦ: ᖃᑭᖃᕐᓂᖅ ᐅᑭᐅᑕᖅᑐᑐ ᐅᑎᖅᓄᑦ *(Boreogadus saida)* ᐃᖅᓗᑦᑕᐅᒋᐊᖏᑦᑐᑦ ᐃᖅᓗᑦᑕᐅᔾᔨᑦ ᑲᓇᑕᒥ ᐅᑭᐅᑕᖅᑐᒥ ᑭᖓᓴᐸᓂᑦ ᐃᖅᓗᓚᕆᓂᖅᑎᑦ (ᖃᐅᔨᓴᖅᑎᑎᑦ ᐅᖃᖅᐳᓴᒋᐊᓂᒍᑦ ᐅᓂᒃᑲᑦ 2020/007)

ᐃᓕᔪᑦᖅ 8 – ᖃᐅᔨᓴᓂᒍᑦ ᑎᑎᕋᖅᓯᒪᔾᔨᑦ 2019/073: ᖃᐅᔨᓴᓂᖏᑦ ᐊᑦᑐᖅᑕᐅᔾᔨᓇᖅᑐᓂᑦ ᐃᖅᓗᑦᑕᐅᒋᐊᖏᑦᑐᑦ ᐃᖅᓗᑦᑕᐅᔾᔨᓂᑦ ᑐᔾᔭᖃᑦᑕᕐᓂᖏᑦ ᐅᑭᐅᑕᖅᑐᒥ ᐅᑎᖅᓄᑦ *(Boreogadus saida)* ᑭᖓᓴᐸᖏᓐᓂ ᐅᑭᐅᑕᖅᑐᒥ *(Pandalus borealis)* ᐊᓪᒪᒃ ᖃᐅᔨᖅᑕᓪᓕᑦ *(Pandalus montagui)* ᑭᖓᓴᐸᓂ ᐃᖅᓗᓚᕆᓂᖅᑎᑦ ᑭᖓᓴᐸᓂᐊᖅᓱᐃᑦᑐᔾᔨᓂᑦ 1, 2, ᐊᓪᒪᒃ 3

ᐃᓕᔪᑦᖅ 9 – ᖃᐅᔨᓴᖅᑎᓂᑦ ᐅᖃᐅᔨᔪᒋᐊᓂᒍᑦ ᐅᓂᒃᑲᑦ 2020/007: ᖃᑎᖅᑭᑐᓂᖅ ᐅᑭᐅᑕᖅᑐᒥ ᐅᓴᖏᓐᓂ *(Boreogadus saida)* ᐃᖅᓗᑦᑕᐅᒋᐊᖏᑦᑐᑦ ᐃᖅᓗᑦᑕᐅᔾᔨᑦ ᑲᓇᑕᒥ ᐅᑭᐅᑕᖅᑐᒥ ᔪᖓᓴᐸᓂᑦ ᐃᖅᓗᓚᕆᓂᖅᑎᑦ




ᐊᐅᓚᓵᒍᐊᓐᓄᑦ 1. ᐅᑭᐅᑕᖅᑐᒥ ᐅᓴᐃᑦ ᐃᖅᓗᓪᑕᐅᕈᑎᐊᖏᑦᑐᑦ ᐃᖅᓗᓪᑕᐅᕐᓂᑦ ᑭᖑᑦᑲᓂᐊᕐᕕᒃᓴᒥ 1 ᐊᖏᕐᒪᔪᓄᑦ 1979–2018.



ᐊᐅᓚᓵᒍᐊᓐᓄᑦ 2. ᐅᑭᐅᑕᖅᑐᒥ ᐅᓴᐃᑦ ᐃᖅᓗᓪᑕᐅᕈᑎᐊᖏᑦᑐᑦ ᐃᖅᓗᓪᑕᐅᕐᓂᑦ ᑭᖑᑦᑲᓂᐊᕐᕕᒃᓴᒥ 2 (ᑲᓇᐅᓇᖅ ᖃᐅᔨᓴᕕᐅᕐᓂᖅ ᑭᓚᖅᑐᖅ) ᐊᖏᕐᒪᔪᓄᑦ 1979–2018[1]

[1] ᑭᖑᑦᑲᓂᐊᕐᓂᑦ ᐃᖅᓗᓚᕆᓂᖏᑦᑕᑦ ᐊᐅᓪᓚᕆᑎᓂᖏᓪᒍᑦ ᐅᑎᖅᑕᖏᓂᖅ ᐊᓯᔾᓚᖅᑐᒃᓴᖅ ᐅᔾᓗᖅᓯᐅᑎᖓᔾᔪᓂ ᐊᖏᕐᒪᔪᒥᑦ ᑭᓇᐅᔭᕆᕆᓂᖏᓪᒍᑦ ᐊᖏᕐᒪᔪᒍᑦ 2003.



ᐊᐳᓂᖏᐅᐊᖅ 3. ᐅᑭᐅᑕᖅᑐᒥ ᐅᓗᐃᑦ ᐃᖅᓗᓪᑕᐅᑭᐊᖏᑦᑐᑦ ᐃᖅᓗᓪᑕᐅᓂᐸᓂ ᑭᖑᓪᓕᓂᖅᑐᑦ 3 (ᐱᖓᓂᓀᖓᓂ ᖅᐅᓵᓴᖅᐊᐅᓂᐸᒥᑦ ᑭᓚᓂᖏᓗ) ᐊᖏᕐᒍᒍᓂ 1979–2018[2].

ᑎᑎᕋᖅᓯᒪᔪᑦ 1. ᐅᑭᐅᑕᖅᑐᒥ ᐅᓗᖅᑕᐅᓯᒪᔪᑦ ᐅᑯᓀᓂ SFA 1, EAZ ᐊᒻᒪ WAZ ᐊᖏᕐᒍᒍᓂ 2018-19 ᑎᑭᑦᑐᒍ 2019-20 (ᐅᓀᐅᓚᖅᑎᓪᓗᒍ ᐄᓕᓂᐊᓀ 30, 2020). *ᑐᑭᓯᐱᑎᐊᕐᓇᑦ ᐃᓱᒪᒋᔭᐅᔪᑦ ᐱᑎᐊᓂᖏᓐᓄᐅᓪᓗᑎ.*

	2019-20 (ᐅᓀᐅᓚᖅᑎᓪᓗᒍ ᐄᓕᓂᐊᓀ 30, 2020)	**2018-19**
SFA 1	12.389	0.410
EAZ	12.103	12.935
WAZ	41.670	61.975

[2] ᑭᖑᓪᓕᓂᑦ ᐃᖅᓗᓚᑕᓂᕐᒥᑦ ᐊᐅᓪᓚᑎᓂᕐᒪᑦ ᐅᑎᖅᑕᕐᓂᖓ ᐊᓯᔾᓚᖅᑐᑎᖅᓂᐅᓂ ᐅᔾᓗᖅᓯᐅᑎᖅᑭᓂᓂ ᐊᕐᕌᒍᒥᑦ ᑭᓇᐅᔭᓚᑕᓂᕐᒪᑦ ᐊᕐᕌᒍᒪᑦ 2003.

| ᑭᖑᒍᑦᐸᓂᐊᖏᐃᐅᐅᔪᑦ | ᐊᖏᖅᑐᒍᖏᑐ ᖃᓄᐃᓕᓂᐊᕐᓂᑦ ᓯᐃᓴᖏᓪ (COL) ᐊᑐᓕᖅᑎᑕᐅᓂᖑᓪᓚᖅ | ᖃᓄᐃᓕᓂᓂᕆᔨᐊᓕᕐᓂᑦ ᓯᐃᓴᖏᓪ (ᐃᖃᓗᓕᑕᐅᔨᐊᖏᑦᑐᓂᑦ ᐃᖃᓗᓕᑕᐅᔪᓂᑦ ᒪᓕᖅᑎᑦᑐᓂᖓᓚᖅ ᐃᓕᒍᑕᖏᑦ)

 5.2
 "ᐃᓚᐃᓂᓂᔭᓂ ***ᑲᑎᑦᑐᑎᑦ ᐃ ᖃᓗᓕᑕᐅᔨᐊᖏᑦᑐᑦ ᐃᑲᓗᑕᐅᔪᑦ ᐊᑕᖏᑦᑐᑦ ᐃ ᖃᑲᒥᐅᑕᐃᑦ ᐅᒪᔪᓂᑦ ᐊᖏᔪᓂᑦᑐᑦ ᓇᓕᐊᓂᑐᐃᓇᖅ*** *ᐱᔾᐅᔪᓂᑦ ᐅᖓᓯᑕᐅᔭᔪᓂᑦ* ***ᐊᖏᓂᖅᓴᖓᓚᒃ 2.5 >ᓴᓂᑎᒃ ᐅᖏᐊᓂᖅᓯᑎᑐᒍᑎᑦ ᑭᖑᓂᖅ 100 kg,*** *ᓕᐃᓴᓂᓗᒥᑦ ᑎᒍᒥᐊᖅᑐᒍᑎ ᐃᖃᓗᓕᑕᐅᔨᐊᓂᑦᑐᑦ ᐅᖃᓕᓂᖅᓂᑦ ᐃᖃᓗᓕᑕᐅᔨᐊᓂᑦᑐᑦ ᐃᖃᓗᓕᑕᐅᔪᓂᑦ ᐃᖃᓗᓕᑕᐅᔨᐊᓂᑦᑐᑦ ᐃᖅᑲᐅᑕᑯᓂᑦ ᐃᖃᓗᓕᑕᐅᔨᐊᓂᑦᑐᑦ ᐃᖅᑲᐅᑕᑯᓂᑦ ᐃᖃᓗᓕᑕᐅᔪ

[ᑐᓴᖅᓴᐃᓪᓕᖅ (ᑐᑭᓯᐊᕐᕉᑎᓄᑦ): ᐊᑐᓕᖅᑎᑕᐅᓂᖏᑦ ᐅᑭᐅᑕᖅᑐᒥ ᐅᓴᐃᓪᓕᖅ ᐃᖏᓐᓂᑕᐅᕈᐊᖅᑎᑦᑐᑦ ᐃᖏᓐᓂᑕᐅᔪᑦ ᐊᐅᓚᑦᑎᓯᑎᐊᕐᕈᑎᑦ 2019/20 ᐅᑭᐅᑕᖅᑐᒍᑦ ᐊᒻᒪ ᖃᐅᖅᑕᓕᓐᓄᓪᓕᑦ ᔪᖓᒍᐸᓂᑦ ᐃᖏᓐᓂᓚᓂᖅᑎᑦ ᐱᖓᓐᓇᓂ ᖃᐅᔨᓴᖏᐅᔪᒥᑦ ᑭᓕᓚᖓ]

ᑲᓇᓂᐊᓗ 2020, ᐃᒪᖏᐅᑕᓕᓚᕆᔨᒋᑦ ᒪᖏᑭᐊᑎᑦᑐᑎᑦ ᑲᑎᒪᓕᐅᕐᖓᑦ ᐅᖏᓪᓕᐅᑎᕆᔨᒋᑦ ᑐᙵᕆᓐᓇᖅᑭᒪᓕᒃᓴᓂ ᐱᓕᕆᓂᐊᕐᓂᓂᑦ ᖃᓄᐃᓕᓴᐅᒪᔭᕐᓂᑦ ᓕᐃᓴᓇᓕ ᐱᓕᕆᐊᖃᖏᑦᓂᓂ ᐅᑭᐅᕐᖓᒍᑎ ᐅᖃᐅᑕᖓᒍᑎ ᐃᖃᓗᓕᐅᑕᐅᔪᒋᑦ ᐃᖃᓗᓕᑕᐅᔪᒋᑦ ᐅᑭᐅᕐᖓᒍᑎ. ᑎᑎᕋᖅᓯᒪᔪᑦ ᐃᓕᐅᕐᖃᐅᔪᓂᑦ ᐊᑐᐃᓐᓇᐅᔪᑦ ᐊᑦᑐᓂ:

ᐅᐸᙵᓯᒪ ᖃᓂᑕᒋᑦ ᐊᐃ ᕿᓚᓂ ᑲᑎᒪᑎᑎᓂᕐᒥᑦ ᐅᖃᓕᐅᑎᕆᔨᒋᑦ

ᐅᔪᕕᑦ ᓄᐊᑯᓵ - ᐃᒪᖏᐅᑕᓕᓚᕆᔨᒋᑦ ᖃᐅᔨᓴᖃᖅᑎᑦ
ᓯᐅᓗ ᐅᓱᓴᓵ - ᐃᒪᖏᐅᑕᓕᓚᕆᔨᒋᑦ ᐃᒪᖏᐅᑕᓂ ᐊᐅᓪᓕᑎᓂᕐᒧᑦ C&A
ᐊᓵᔨᓪ ᓴᖅ - ᐃᒪᖏᐅᑕᓕᓚᕆᔨᒋᑦ ᐃᒪᖏᐅᑕᓂ ᐊᐅᓪᓕᑎᓂᕐᒧᑦC&A
ᐃᓴᑎᓵ Hᐃᐊᓄᓗ - ᐃᒪᖏᐅᑕᓕᓚᕆᔨᒋᑦ ᐃᒪᖏᐅᑕᓂ ᐊᐅᓪᓕᑎᓂᕐᒧᑦ NL
ᑯᐊᑦᓂ ᑕᐅᐳ - ᐃᒪᖏᐅᑕᓕᓚᕆᔨᒋᑦ ᐃᒪᖏᐅᑕᓂ ᐊᐅᓪᓕᑎᓂᕐᒧᑦ NHQ
ᓪᓗ ᐃᐊᑦ - ᐃᒪᖏᐅᑕᓕᓚᕆᔨᒋᑦ ᐃᒪᖏᐅᑕᓂ ᐊᐅᓪᓕᑎᓂᕐᒧᑦ NHQ*
ᑎᐅᓗᔪ ᒪᕼᐅᓂ - ᐃᒪᖏᐅᑕᓕᓚᕆᔨᒋᑦ ᐃᒪᖏᐅᑕᓂ ᐊᐅᓪᓕᑎᓂᕐᒧᑦ NHQ

ᐊᑦᓱ ᓴᐃᔪᑦ – ᓄᓇᕗᓴᒥ ᐅᒪᔪᓕᓚᕆᔨᖏᔪᒋᑦ ᖃᑎᒪᔨᖏᑦᑕ
.ᐃᓴᓵ ᐊᖏᐊᑭᖅ - ᓄᓇᕗᓴᒥ ᐅᒪᔪᓕᓚᕆᔨᖏᔪᒋᑦ ᖃᑎᒪᔨᖏᑦᑕ*
ᑎᐊᓂᔪᑦ ᑎᓗ - ᓄᓇᕗᓴᒥ ᐅᒪᔪᓕᓚᕆᔨᖏᔪᒋᑦ ᖃᑎᒪᔨᖏᑦᑕ

ᕗᖅᓱᓄ ᑲᓵ–ᒧᖏᓵ - ᓄᓇᕕᔭ ᐃᒪᖏᐅᑕᓂᑦ ᐅᒪᔪᓂᑦ ᖃᑎᒪᔨᖏᑦᑕ *

ᐅᑐᔪ ᓴᖅᒪᓵ - ᖃᓇᑕᒥ ᖃᑐᐃᓐᖃᑎᒋᖏᑦᑕ ᓄᓕᒃᕿᓕᓚᐅᖅᑲᑦ
ᐊᓕᔭᑎᐅ ᐅᕆᐃᓕ – ᐅᓄᐅᑕᖅᑐᒥ ᖃᑐᐃᓐᖃᑎᒋᑦ
ᔨᐅᓗ ᕗᐊᖅᑦ - ᖁᓄᖅᑕᒧᓵᔭ ᑯᐊᐅᓱᓴᓵ
ᑯᓚᔪ ᕿᓭᓂᖃᓵ - ᖁᓄᖅᑕᒧᓵᒥ ᐃᖅᓗᓕᓚᕆᔨᒋᑦ ᖃᑐᐃᓐᖃᑎᒋᖏᑦᑕ
ᐳᕆᐃᔨᓵ ᐅᔭ - ᓄᓇᕗᓴᒥ ᐃᖅᓗᓕᓚᕆᔨᒋᑦ ᖃᑐᐃᓐᖃᑎᒋᖏᑦᑕ*
ᐳᕆᐃᔨᓵ ᒧᔭᓇᒪ - ᓂᐅᐸᐅᑕᓵᓂᑦ ᑎᒥᖓᒃ
ᐱᑕ ᑭᔪ - ᒪᑭᕕᔭᒃ ᑯᐊᐅᓱᓴᓵ

[ᑲᑐᑎᓴᑦ ᐃᖏᓗᓕᐱᓂᐅᓴᑦ ᑭᐅᓴᑦ ᐅᑭᐅᑕᖅᑐᑎ ᐅᑲᐱᐃᑦ ᐃᖏᓗᑦᑕᐅᕈᐊᖅᑎᑦᑐᑦ ᐃᖏᓗᑦᑕᐅᓴᑦ ᐊᐅᓪᓚᑎᓯᑎᐊᑭᑎᓱᑦ ᐊᐠᒪ ᒪᖅᑭᑎᑎᓂᕆᑦ ᐊᐧᖅᑭᑎᐊᖅᓯᒪᔨᓂᑦ ᖃᓄᐃᓕᓂᕆᐊᓕᖓᑦ ᓪᓚᐃᑦᓯᓗᑎᐊᓕᖓᑦ ᓚᐃᓴᓐᓯ (COL)]

ᖃᐅᔨᓴᕐᑐᑦ ᑎᑎᕋᖅᓯᒪᔭᖅᓴᖅ 2019/073

**ᓇᐃᓕᓐᓴᕐᑕᒃ: ᖃᒥᖅᑭᐅᐅᓂᖏᑦ ᑕᑯᓴᕈᑐᐃᓇᓇᓕᓪᓕᐊᓪᓕᑦ ᐸᐅᑐᐃᓂᖅᓂᖅᑐᑦ ᐱᓕᐅᓇᓯᓇᓕᑎᐅ ᑐᖏᑲᖅᑕᐅᔪᑦ ᐅᑭᐅᖅᑕᖅᑐᒃ ᐅᑲᓐᖏᑦ (ᓴᓇᐅᔨᑕᑦ, ᓴᐃᑐᒃ) ᑲᑎᕋᖅᓯᒪᓂᖏᑦ ᐱᓇᓯᑕᐅᑎᓪᓗᓂᑦ ᐅᑭᐅᖅᑕᖅᑐᒃ (ᕿᓪᑕᓪᓚᓵ, ᓴᐃᓇᓇᓚᓵ) ᐊᒻᒪᓗ ᑕᖅᓴᓪᓕᑦ (ᕿᓪᑕᓪᓚᓵ, ᒪᓕᓪᑕᓪᓚᓵ) ᑲᑎᒪᔨᖅ ᐃᓐᓇᖅᑐᓕᕐᓗᓂᑦ ᐅᓇᖅᓯᓪᓕᐊᓂᑦ ᐱᑭᖅᑐᓂᑦ ᐅᑭᐅᖅᑕᖅᑐᒃ (ᐃᓚᖅᓗ, ᑕᐃᑲᖅᓯᒪᓂᖅ 2017), ᑭᖓᔪᓂᓂᐊᓂᖅ ᐃᖅᑲᓇᐃᓕᓐᓂᖅᑐᑦ ᐅᒥᐊᔪᐊᖅᑭᓂᓂᖅ ᐊᐅᓚᑦᑎᐊᓂᖅ ᐃᓕᓴᕐᓂᖅᑐᑦ ᐃᓚᖅᑭᑦᑐᖅᓯᒪᓂᖅ ᐃᓕᓴᕐᓂᖅᑐᑦ ᐃᓕᓴᕐᓂᖅᑐᑦ ᐃᓕᓴᕐᓂᖅᑐᑦ

ᖃᐅᔨᓴᕐᑎᓄᑦᓯᑦ ᐅᖃᐅᓯᖅᓯᐊᕐᓚᒍᑦ 2020/007

ᓇᐃᓕᓕᐊᓗᓐᓂᕐᓴ: ᖃᒥᖅᑭᕆᐅᓂᖅᓴᖅ ᐅᑭᐅᖃᖅᑕᖅᑐᑦ ᐅᓚᖅᑎᑦ (ᓴᐊᓚᐅᑉᓯᑦ ᓴᐃᑎ) ᐱᓇᓱᑕᐅᓇᑎᒃ ᐃᖃᓱᒃᑕᓇᓄᑦ ᐅᖃᐅᖃᖅᑕᖅᑐᐊᓂ ᑭᖑᓱᒃᑐᑎᓱᒃᑐᑦᑎᓄᑦ ᐃᖃᓗᓯᓱᒃᑐᑎᑦᑎᓄᑦ

- ᑭᖑᓱᒃᓴᓱᒃᑎᑦ ᐃᖃᓗᓯᓱᓕᑦ ᑲᓇᑕᐅᑦ ᐅᑭᐅᖃᖅᑕᖅᑐᖓᓂ ᐊᑐᑎᖃᒡᓗᓕᖅᑐᑦ ᓱᓇᒃᑯᑦᑕᓚᕐᑎᑕ ᐃᓄᐃᑦ ᓄᓇᕗᒻᒥ ᓄᓇᖃᖓᓕᕐᓱᐊᑕᑕᐅᓕᓪᓚᒃᑎ ᑭᖑᓱᒃᐸᐃᑦ ᐃᖃᓗᓯᓱᑎᑎᓕᐅᑦᐸ ᐃᓂᖓᓂᑕ 1 (SFA 1) ᐊᓯᒪᓗ ᑕᒃᑯᐊᒃ ᑲᓇᓴᓂᓕᓪᓚ ᐱᖓᓱᐅᑉᖢᕆᔨᑎᑦᑕ ᑭᓚᖃᖅᑎᖓᑦ (EAZ/WAZ). ᑕᑯᐃᓴᓇᓚᓪᓚᔪ, ᐱᐅᔪᖅ ᐊᖑᑐᖃᖅᑎᑦᑕᑯᓘᑦ ᐃᓚᐃᑦᑐᐃᓐᔪᓚᓪᓚ ᑭᖑᓱᒃᐸᑕᐅᓱᓄᑦ ᐊᓯᖅᓯᐊᔪᑦ ᐃᓚᐃᑦᑐᔪᑦ ᑭᖑᓱᒃᐸᑕᐅᓱᓄᑦ ᐃᓚᐃᑦᑐᖓᓕᖅᑐᑦ (ᓴᓇᐅᑦᓴᑦ ᓴᐃᑎᑕ) ᐃᖃᓗᖃᓕᕆᔨᑦ ᐅᓪᓴᓇᐅᓕᕐᓚᔪᓂᑦ.
- ᐱᔨᔪᑎᑦᓚᓚᔪ ᐊᒥᓚᔪᐊᖅᑐᒥᑲ ᓇᐃᓴᐅᑎᑦᔪᑎᖅᓂᖅᓯᖓᓕᓄᑦ, ᖃᐅᔨᓴᖅᑐᑦ ᐅᖃᐅᔨᓴᖅᓯᖅᑐᑦ ᓴᖅᑭᔪᓂᓇᕐᓂᑎᑲ ᑭᓚᓕᑦᑕᐅᑦ ᐱᓚᒥᑎᑲ ᐊᐱᖅᔪᑎᑦᑎᓕᕐᓚᔪᓂᑦ ᑭᖑᓕᒍᑦᑕᐅᔪᓕᒪ ᐱᖓᓱᒃᑐᑦᑎᓄᑦ ᐊᓯᖅᓯᐊᔪᑦ ᐊᐱᖅᔪᑎᑦᑎᓕᕐᓚᔪᓂᑦ ᐊᐱᖅᔪᑎᑦᑎᓕᕐᓚᔪᓂᑦ (ᓴᓇᐅᑦᓴᑦ ᓴᐃᑎᑕ) ᐃᖃᓗᖃᖅᑎᑦᑎᓕᕐᓚᔪᓂᑦ ᐃᓚᐃᑦᑐᔪᑦ


ᓱᐅᓂᒃᑯᑦ ᓴᖅᑭᑎᑕᓂᕐᒥᒃ ᖃᐅᔨᓴᖅᑎᑦ ᐅᖃᐅᔨᔨᖅᓱᐊᕐᓂᒃᑐᓂᓕᕆᔨᐊᑎᓂᒃ ᓇᓪᓕᐊᓇᕐᓵᑎᓂᒃ ᐃᖃᓗᐊᓂᑦ ᐱᖃᓱᐅᕐᓴᖅᑎᑦ ᐃᖃᓗᓐᓄᑦ.

- ᑐᑭᖓᓪᓕᓇᕐᑐᖅ ᖃᐅᔨᓴᖃᓂᖅ ᐱᑕᖃᓇᐊᓕᒃ ᐅᑭᐅᖅᑕᖅᑐᑦ ᐅᑎᖅᑕ ᐅᒪᔨᔨᓱᖅᑐᓂᑦ, ᐊᐅᑕᓂᓇᓱᖅᑕ ᐅᒪᓂᐊᒍᑦ, ᐊᐅᓱᓂᖅᑐᓂᑦ, ᐊᓚᒡᑕᐅᖅ ᐅᒪᔨᓂᑦ ᐊᐅᑕᓪᓚᕐᓂᕐᕕᑕ ᑕᐅᑐᖓᐊᕐᑎᖅᑐᓂᑦ ᑐᑭᓯᓂᓪᓕᐊᓐᓂᒃᑯᑦ ᑲᓇᑕᒥ ᖃᑎᖅᑕᐅᑦᑕᐅᓚᖓᑕᑕ ᐅᒪᔨᓂᑦ ᓂᖃᖅᑐᖓᓂᒃᑯᑦ. ᑕᒪᓐᓇ ᖃᐅᔨᔨᖅᑎᓂᑦ ᓴᖅᑭᑦᓴᕆᔭᒃᑯᑦ ᓇᐃᔨᐅᑎᔨᐅᓂᖅᓴᖅᑯᑦ ᑲᑎᕐᓂᖅᑐᓂᑦ ᐊᐃᔨᐅᑎᓂᑦ ᓇᓕᐅᑦᑕᒃᓴᖅᑐᓂᑦ ᐱᐅᓂᐊᕐᓇᓕᒃ ᐃᓕᓴᕐᕕᐅᑎᓂᑦ ᐅᖃᐅᔨᓴᖅᑎᓂᑦ ᐱᓇᓱᐊᕐᓇᓕᒃ ᐃᖅᑲᓇᐅᑦᑕᐅᔪ


[ᖃᐅᔨᓴᓂᖅᓯᒪᔪᑦ ᑎᑎᕆᖅᓱᒪᔪᑦ 2019/073: ᖃᐅᔨᓴᓂᕐᒥᑦ ᐊᑦᑐᖅᑕᐅᕗᔪᓐᓇᖅᑐᓂᑦ ᐃᖃᓗᑦᑕᐅᕈᑎᐊᖏᑦᑐᑦ ᐃᖃᓗᑦᑕᐅᕗᔪᓂᑦ ᑐᑭᓯᖃᑦᑕᕐᓂᖏᑦ ᐅᑭᐅᑕᖅᑐᒥ ᐅᑎᖅᓄᑦ *(Boreogadus saida)* ᑭᖑᑦᑕᖏᓪᓂ ᐅᑭᐅᑕᖅᑐᒥ *(Pandalus borealis)* ᐊᓱᒪ ᖃᓄᖅᑕᓪᓕᑦ *(Pandalus montagui)* ᑭᖑᑦᑕᓂ ᐃᖃᓗᓕᒫᓂᕐᒥᑦ ᑭᖑᑦᑕᓂᐊᕐᕕᐅᔪᓂᑦ 1, 2, ᐊᓱᒪ 3 (ᖃᐅᔨᓴᓂᖅᓯᒪᔪᑦ ᑎᑎᕆᖅᓱᒪᔪᑦ 2019/073)]

ᐃᓕᔪᑕᖏᑎ 9

[ᖃᐅᔨᓴᖅᓂᕆᑦ ᐅᖃᐅᔨᔪᒋᐊᓂᖓᔪᑦ ᐅᓂᒃᑲᑦ 2020/007: ᑭᒥᖅᑐᓂᖅ ᐅᑭᐅᑕᖅᑐᒥ ᐅᑎ ᖃᒋᓂᓂ (Boreogadus saida) ᐃᖃᓗᑦᑕᐅᒋᐊᖏᑦᑐᑦ ᐃᖃᓗᑦᑕᐅᔪᑦ ᑲᓇᑕᒥ ᐅᑭᐅᑕᖅᑐᒥ ᑯᓘᓪᓚᓂ ᐃᖃᓗᓕᕆᓂᖓᔪᑦ]




ᐊᐅᓚᓵᒍᐊᓐᓄᑦ 1. ᐅᑭᐅᑕᖅᑐᒥ ᐅᓴᐃᑦ ᐃᖅᓗᓪᑕᐅᕈᑎᐊᖏᑦᑐᑦ ᐃᖅᓗᓪᑕᐅᕐᓂᑦ ᑭᖑᑦᑲᓂᐊᕐᕕᒃᓴᒥ 1 ᐊᖏᕐᒪᔪᓄᑦ 1979–2018.



ᐊᐅᓚᓵᒍᐊᓐᓄᑦ 2. ᐅᑭᐅᑕᖅᑐᒥ ᐅᓴᐃᑦ ᐃᖅᓗᓪᑕᐅᕈᑎᐊᖏᑦᑐᑦ ᐃᖅᓗᓪᑕᐅᕐᓂᑦ ᑭᖑᑦᑲᓂᐊᕐᕕᒃᓴᒥ 2 (ᑲᓇᐅᓇᖅ ᖃᐅᔨᓴᕕᐅᕐᓂᖅ ᑭᓚᖅᑐᖅ) ᐊᖏᕐᒪᔪᓄᑦ 1979–2018[1]

[1] ᑭᖑᑦᑲᓂᐊᕐᓂᑦ ᐃᖅᓗᓚᕆᓂᖏᑦᑕᑦ ᐊᐅᓪᓚᕆᑎᓂᖏᓪᒍᑦ ᐅᑎᖅᑕᖏᓂᖅ ᐊᓯᔾᓚᖅᑐᒃᓴᖅ ᐅᔾᓗᖅᓯᐅᑎᖓᔾᔪᓂ ᐊᖏᕐᒪᔪᒥᑦ ᑭᓇᐅᔭᕆᕆᓂᖏᓪᒍᑦ ᐊᖏᕐᒪᔪᒍᑦ 2003.



ᐊᐳᓂᖏᐅᐊᖅ 3. ᐅᑭᐅᑕᖅᑐᒥ ᐅᓗᐃᑦ ᐃᖅᓗᓪᑕᐅᑭᐊᖏᑦᑐᑦ ᐃᖅᓗᓪᑕᐅᓂᐸᓂ ᑭᖑᓪᓕᓂᖅᑐᑦ 3 (ᐱᖓᓂᓀᖓᓂ ᖅᐅᓵᓴᖅᐊᐅᓂᐸᒥᑦ ᑭᓚᓂᖏᓗ) ᐊᖏᕐᒍᒍᓂ 1979–2018[2].

ᑎᑎᕋᖅᓯᒪᔪᑦ 1. ᐅᑭᐅᑕᖅᑐᒥ ᐅᓗᖅᑕᐅᓯᒪᔪᑦ ᐅᑯᓀᓂ SFA 1, EAZ ᐊᒻᒪ WAZ ᐊᖏᕐᒍᒍᓂ 2018-19 ᑎᑭᑦᑐᒍ 2019-20 (ᐅᓀᐅᓚᖅᑎᓪᓗᒍ ᐄᓕᓂᐊᓀ 30, 2020). *ᑐᑭᓯᐱᑎᐊᕐᓇᑦ ᐃᓱᒪᒋᔭᐅᔪᑦ ᐱᑎᐊᓂᖏᓐᓄᐅᓪᓗᑎ.*

	2019-20 (ᐅᓀᐅᓚᖅᑎᓪᓗᒍ ᐄᓕᓂᐊᓀ 30, 2020)	**2018-19**
SFA 1	12.389	0.410
EAZ	12.103	12.935
WAZ	41.670	61.975

[2] ᑭᖑᓪᓕᓂᑦ ᐃᖅᓗᓚᑕᓂᕐᒥᑦ ᐊᐅᓪᓚᑎᓂᕐᒪᑦ ᐅᑎᖅᑕᕐᓂᖓ ᐊᓯᔾᓚᖅᑐᑎᖅᓂᐅᓂ ᐅᔾᓗᖅᓯᐅᑎᖅᑭᓂᓂ ᐊᕐᕌᒍᒥᑦ ᑭᓇᐅᔭᓚᑕᓂᕐᒪᑦ ᐊᕐᕌᒍᒪᑦ 2003.

| ᑭᖑᒍᑦᐸᓂᐊᖏᐃᐅᐅᔪᑦ | ᐊᖏᖅᑐᒍᖏᑐ ᖃᓄᐃᓕᓂᐊᕐᓂᑦ ᓯᐃᓴᖏᓪ (COL) ᐊᑐᓕᖅᑎᑕᐅᓂᖑᓪᓚᖅ | ᖃᓄᐃᓕᓂᓂᕆᔨᐊᓕᕐᓂᑦ ᓯᐃᓴᖏᓪ (ᐃᖃᓗᓕᑕᐅᔨᐊᖏᑦᑐᓂᑦ ᐃᖃᓗᓕᑕᐅᔪᓂᑦ ᒪᓕᖅᑎᑦᑐᓂᖓᓚᖅ ᐃᓕᒍᑕᖏᑦ)

 5.2
 "ᐃᓚᐃᓂᓂᔭᓂ ***ᑲᑎᑦᑐᑎᑦ ᐃ ᖃᓗᓕᑕᐅᔨᐊᖏᑦᑐᑦ ᐃᑲᓗᑕᐅᔪᑦ ᐊᑕᖏᑦᑐᑦ ᐃ ᖃᑲᒥᐅᑕᐃᑦ ᐅᒪᔪᓂᑦ ᐊᖏᔪᓂᑦᑐᑦ ᓇᓕᐊᓂᑐᐃᓇᖅ*** *ᐱᔾᐅᔪᓂᑦ ᐅᖓᓯᑕᐅᔭᔪᓂᑦ* ***ᐊᖏᓂᖅᓴᖓᓚᒃ 2.5 >ᓴᓂᑎᒃ ᐅᖏᐊᓂᖅᓯᑎᑐᒍᑎᑦ ᑭᖑᓂᖅ 100 kg,*** *ᓕᐃᓴᓂᓗᒥᑦ ᑎᒍᒥᐊᖅᑐᒍᑎ ᐃᖃᓗᓕᑕᐅᔨᐊᓂᑦᑐᑦ ᐅᖃᓕᓂᖅᓂᑦ ᐃᖃᓗᓕᑕᐅᔨᐊᓂᑦᑐᑦ ᐃᖃᓗᓕᑕᐅᔪᓂᑦ ᐃᖃᓗᓕᑕᐅᔨᐊᓂᑦᑐᑦ ᐃᖅᑲᐅᑕᑯᓂᑦ ᐃᖃᓗᓕᑕᐅᔨᐊᓂᑦᑐᑦ ᐃᖅᑲᐅᑕᑯᓂᑦ ᐃᖃᓗᓕᑕᐅᔪ

ⵊⵔⵉⵎⵓⵜⵏ ⵜⵉⵎⵓⵔⴰⵜⵏ
ⵜⵜⴰⵡⵉⵔⵉⵏ ⵏ ⵜⵎⵓⵔⴰⵜ ⵏ ⵜⵎⵓⵔⴰⵜ ⵏ ⵜⵎⵓⵔⴰⵜ ⵏ ⵜⵎⵓⵔⴰⵜ
ⵜⵉⵣⵔⵉ

ⵜⵓⵔⵉⵎ: X **ⵜⵉⵔⵉⵎⵜⵓ:** **ⵜⵣⵔⵜⵉⵔⵜⵏ:**


**ⵏⵜⵔⵓⵔⵜ ⵜⵣⵔⵜⵉⵔⵜⵓⵔⵜ ⵜⵉⵏⵜ ⵜⵜⵔⵉⵔⵜⵓⵔⵜ ⵜⵜⵔⵉⵔⵜⵓⵔⵜ ⵜⵜⵔⵉⵔⵜⵓⵔⵜ ⵜⵜⵔⵉⵔⵜⵓⵔⵜ ⵜⵜⵔⵉⵔⵜⵓⵔⵜ 2019/20 ⵜⵔⵜⵔⵜⵓⵣⵣ ⵔⵜⵣⵜⵓⵔⵜ ⵜⵔⵉⵔⵜⵓⵔⵜ ⵜⵔⵉⵔⵜⵓⵔⵜ ⵜⵔⵉⵔ


ᐃᖃᓗᖓᓂᒃ ᖃᓄᐃᓄᖅᑕᐅᓪᓗᑎᑎᒃ ᐅᒪᓄᓪᒪᓐᓕᒍᓐᓂᒃ ᐊᖏᕐᓄᖏᑦ ᓇᓱᒪᑦᑐᑎᒃ, ᐃᖃᓗᓪᑕᐅᖅᓯᖃᑎᖅᑕᖅᑐᑦ ᑭᖑᓪᓕᓴᕐᓄᓪᓕᐊᑭᑎᓂᒃ ᐃᖃᓗᓪᓚᓄᓕᕐᕆᑦ, ᓇᓗᓇᐃᖅᑭᔪᖅ ᐅᑎᐃᑦ ᐊᓱᒪ ᑭᖑᓪᓕᐊᑦ ᑕᓴᑭᖅᑐᑦ ᓇᔪᖅᑭᓭᑐᑦ. ᐅᑎᐃᑦ ᐱᓱᒪᓚᐅᔪᐸᑦ ᐱᑕᖃᖅᑐᑦ ᐃᒪᕆᒥ ᐊᐊᒋᑎᖅᓄᓪ , ᓂᕈᖃᖅᑎᑎᓪᓗᑎ ᐊᒥᑭᓱᓕᓴᒃᓂᒃ ᐅᒪᔪᓇᒃ ᑕᓚᐅᖅᑎᐅᑕᓇᒃ ᑎᓭᐊᓇᒃ, ᐃᒪᖅᑎᐅᑕᓇᒃ ᐅᒪᔪᓇᒃ, ᐊᓱᒪ ᐃᖃᓗᓴᓇᒃ.

ᑐᕈᑭᐊᕈᑎᑦ ᐅᑎᖅᓇᒃ ᓄᐊᑕᐅᔪᑦ ᐃᒪᖅᑎᑦᑐᑎᑦ ᖃᐅᔨᓴᖅᑎᓇᒃ ᐃᖃᓗᓪᓚᓂᖅᓯᒪᔪᑦ ᑎᑎᕋᖅᕕᓇᒃ, ᑭᑭᐊᓂᓪ ᐊᒥᑭᓱᓕᓴᒃᓂᒃ ᖃᓇᐃᓪᓚᓂᖅᓯᒪᓂᖅᓯᒪᓂᒃ ᖃᐅᔨᓴᖅᑕᐅᓪᓚᓵᒪᓕᖅᓯᒪᓂᑐᑦ ᖃᐅᔨᓴᖅᑎᓇᒃ, ᐊᓱᒪ ᑕᐃᒪᐃᓕᐅᔪᖅᓯᒪᓂᖅ ᖃᐅᔨᒪᓕᕆᐅᔪᓂᑐᑦ. ᐃᒪᖅᑎᐅᑕᓪᓚᕐᓂᖅᑐᑦ ᐱᓪᓚᓇᓪᓯᖅᑕᐅᖅᑐᑦ CSAS ᖁᒥᖅᑭᓂᒧᑦ ᐱᓭᓪᓕᐊᓂᖓ ᐊᓪᒪ 2019 ᐅᖃᐅᔨᑭᐊᖅᓗᑎ ᐅᑎᖅᓂᒃ ᐃᖃᓗᓪᑕᐅᖅᓱᖅᑕᖅᑐᓂᒃ ᑭᓪᓕᖅᑐᓄᒃ ᓄᓇᓕᖅᑐᓄᒃ WAZ, EAZ ᐊᓱᒪ SFA 1. ᐱᖃᑕᐅᓪᓕᐅᖅᑐᑦ ᑭᓱᖅᑐᐃᐅᑦ ᐃᒪᖅᑎᐅᑕᓪᓚᓵᒪᓕᖅᓯᒪᓂᖅ ᖃᐅᔨᓴᖅᑎᑦ ᐊᓱᒪ RM, ᐊᓱᒪ ᑲᑎᒪᐅᑦ.

ᒥᖏᐅᖏᖅᑕᐅᔪᑦ ᐱᑕᖃᖅᓂᓪᓚᔪᓇᖅᑕᖅᑭᑦ ᐅᑎᐃᑦ ᐊᒥᓱᓂᑦᑐᓄᒃ ᐊᐊᒋᑎᖅᓂ ᐱᖏᐅᓪᓕᐅᖅᑐᑦ ᐱᑕᖃᓚᐊᖃᖅᑎᑕᐅᔪᓂᒃ ᑎᑭᕐᑎᓪᓗᐊᑦᓗᑎ ᐱᖏᐅᔪᓂᑦ ᐊᓂᔪᓂᖅᖁᑦᑐᑦ ᑐᔪᓐᓇᖅᑐᓄᒃ ᐱᑕᖃᖅᑎᑦᑎᐊᑦᓗᑎ ᐱᖏᐅᔪᓂᑦ ᑐᔪᓐᓇᖅᑐᓄᒃ ᐊᐊᒋᑎᖅᓂ

ᑐᖃᓯᕐᑐᑦ – ᑲᓇᓐᓇᖅᓂ ᖃᐅᔨᓴᖅᑕᐅᔪ ᑭᓚᓐᖅ
ᑐᖃᓯᕗᕐᑐᑦ – ᐱᖅᓐᓇᖅᓂ ᖃᐅᔨᓴᖅᑕᐅᔪ ᑭᓚᓐᖅ





ᐊᓱᓴᓚᐊᖅ 1. ᐅᓪᓚᐃᑦ ᐃᖃᓗᑦᑕᐅᖏᓪᓗᒍ ᖃᑦᑕᖅᑐᑦ SFA 3 (ᐱᖏᓴᓐ ᖏᓂᓂ ᖃᐅᓵᓴᖅᑕᐅᔪᓄᑦ ᐅᓴᓕᓐᖏᓗᒍ) ᐅᑯᓂᖏᓐᓂᒃ 1979–2018[1].

[1] . ᑭᖓᓪᓕᐊᓂᕐᓗᑦ ᐊᐅᓚᑦᑎᓂᕐᓗᑦ ᐊᑐᖅᑕᐅᓂᖓᓐᓂ ᐊᓱᓵᓴᖅᑐᖅ ᐅᑉᓗᖅᓯᐅᑎᓂᑦ ᐊᖏᖁᔭᐅᓂ ᐊᖏᖁᔪᓗ ᑭᓇᐅᔭᓕᕆᓂᕐᓗᑦ ᐊᖏᖁᔪᒪᓕ ᐱᒋᐊᖏᓂᖅᑐᖅ 2003.a

9/26/2019

ᖃᐅᔨᒪᓕᐅᕆᓇᐊᓕᓗᒃ ᓕᐃᓴᓐᓂᐅ ᐅᑭᐅᑕᖅᑐᑎ ᑯᖓᔪᓕᓂᒃ ᐊᑲᑭᑎᐊᖅᑕᐅᔾᔪᑦ ᐃᒪᐃᓕᓚᓐᓱᓕᑎ:

5.2.2 ᐱᖅᓯᐅᑎᓐᓇᓂ ᐃᓕᓐᓴ 5.2 ᖃᐅᓪᓗᓂ, ᐃᖅᓱᓴᓯᑎᓯᓗᑦᑕ ᐃᒪᖏᓐᓂ ᐊᐅᓪᓚᑕᐅᔾᔪᓂᑦ ᓄᓇᕗᒻᒥ ᐱᖓᓐᓇᖓᓂ ᐊᐠᒪ/ᐅᕐᓂᓵᓐᓂᑦ ᓄᓇᕆᓇᖓᓂ ᑲᑎᑕᑐᒥᑦ ᐃᖅᓱᑦᑕᐅᖏᓐᖓᑦ ᐅᓴᐃᑦ ᐅᖓᑕᐅᑦᔨᔪᓂᑦ 5 ᐳᓴᓐᑎᓂᑦ ᐅᖃᐃᐃᓐᓂᖏᑎᒍᑦ ᐊᖓᔪᖅᑐᑦ 200kg, ᐊᑐᖅᑕᐅᑕᐅᔪᑦᑐᖅ ᑲᓪᓐᓂᐅᓪᓚᐅᖅᑐᓂᑦ ᐱᖓᓱᓂᔪᖅᑕᖅᑐᑎ, ᓕᐃᓴᓐᓂᒥᑦ ᑎᒍᒥᐊᖅᑐᖅ ᐅᖃᓕᓂᓂᑦ ᐅᒥᐊᖃᐊᒥᑦ ᐊᐅᓪᓚᑦᑎᑦᓂᐅᑎᒥᓕᓐᓄ ᓇᒻᓂᓐᑕᐅᑎᒥᓐᓄᑦ ᖃᐅᔨᓴᓐᓂᑦ (10) ᐃᒪᓐᓇᖅᑐᑦ ᒪᐃᔪᐃᑦ ᓇᒥᑐᐃᓐᓇᖅ ᐃᓂᒋᓐᓂᖓᓂ ᑲᑎᒪᔨᖅᑕᐃᓕᓐᓂᒃ ᐱᓇᓱᐊᖅᑕᐃᓕᓐᓂᓂ ᐃᖅᓱᓕᑕᐅᔾᔪᑎᖅᑐᓂᑦ ᐅᑎᕐᓂᑦ. ᓄᓕᐅᖅᑎᓐᓂᒥᑦ, ᑲᑎᑕᑐᒥᑦ ᐃᖅᓱᓕᑕᐅᔾᔪᑎᖅᑐᓂᑦ ᐅᑎᐃᑦ ᐅᖓᑕᐅᔾᔪᔨᒧᑎ ᐊᖏᓂᓄᖅᓴᒥ 5 ᐳᓴᓐᑎᒥ ᐅᖃᐃᐃᓐᓂᖏᑎᒍᑦ ᑯᖓᔪᖅᑐᑦᓂᐅᑎᒥᓐᓄᑦ, ᐅᑭᐊᖃᐊᓂᑦ 200 ᑭᓚᒧᑦᓱ. ᐊᑐᖅᑕᐅᑕᐅᔪᓕᐅᖅᑐᖅᑐᑦ 6–ᕿᖅᑎᖅᑐᑎᑦ ᑲᓪᓐᓂᐅᖏᓐᖓᑦᓭᓚᐅ, ᐅᒥᐊᖃᐊᖅ ᓄᖏᓐᓇᓗᐊᓕᓐᓂᒃ 10 ᐃᒪᓐᓇᖅ ᒪᐃᔪᓂᑦ ᓇᒥᑐᐃᓐᓇᖅ ᐃᓂᒋᓐᓂᖓᓂ ᐱᖓᓱᓂᔪᖅᑐᓂᑦ ᑲᓪᓐᓂᐅᑕᐅᓕᓐᓂᖅᑐᓂᑦ ᐊᓂᖓᓐᓂᑦ ᐃᖅᓱᓕᑕᐅᒪᓂᖃᑦᑐᑦ. ᓕᐃᓴᓐᓂᒥᑦ ᑎᒍᒥᐊᖅᑐᖅ ᐅᖃᐊᖃᐊᒥᑦ ᐊᐅᓪᓚᑕᐅᔾᔪᑎᖅᑐᓂᑦ ᑎᑎᕋᓕᐊᓕᓐᓂᑦ ᑎᑎᕋᖃᓚᒥ ᐱᓕᐅᑦᑕᐃᓐᓕᒪᓕᒃᑐᑦ ᐅᖓᑕᐅᑦᔨᔪᓂᑦ ᐅᑎᕐᓂᑦ ᐃᖅᓱᓕᑕᐅᖓᔾᔪᖅᑐᑦ. ᓕᐃᓴᓐᓂᒥᑦ ᑎᒍᒥᐊᖅᑐᖅ ᐅᖃᐊᖃᐊᒥᑦ ᐊᐅᓪᓚᑕᐅᑎᓂᖅᑐᖅ ᑎᑎᕋᓕᐊᖅᑐᖅᓴᖓᑦᑕ ᐅᑎᕐᓂᑦ ᐃᖅᓱᓕᑕᐅᖓᔾᔪᖅᑐᑦ (ᓴᓂᒧᑦ ᑐᑭᒧᑦ ᓄᓇᖓᒍᒥᑦ) ᐅᓴᐃᑦ ᐃᖅᓱᓕᑕᐅᖓᔾᔪᖅᑐᓂᑦ ᐊᖓᔪᓂᖓᑦ ᐃᖅᓱᓕᑕᐅᔾᔪᑦ ᐅᖃᐃᐃᓐᓂᖏᑎᒍᑦ ᑭᓚᒧᑦᓱ.

ᑕᑯᓇ ᐊᑲᑭᑎᐊᑭᑎᐅᔾᔪᖅ ᐱᑎᐊᖃᓂᑲᖅ ᐅᓪᓗᓚᖓᓂ ᑐᓂᐅᔾᔨᐅᔨᒥᑦ ᒧᑦᓯ 31, 2020.

ᑲᓇᓂᐊᓗ 2020, ᐃᒪᖏᐅᑕᓕᓚᕆᔨᒋᑦ ᒪᖏᑭᐊᑎᑦᑐᑎᑦ ᑲᑎᒪᓕᐅᕐᖓᑦ ᐅᖏᓪᓕᐅᑎᕆᔨᒋᑦ ᑐᙵᕆᓐᓇᖅᑭᒪᓕᒃᓴᓂ ᐱᓕᕆᓂᐊᕐᓂᓂᑦ ᖃᓄᐃᓕᓴᐅᒪᔭᕐᓂᑦ ᓕᐃᓴᓇᓕ ᐱᓕᕆᐊᖃᖏᑦᓂᓂ ᐅᑭᐅᕐᖓᒍᑎ ᐅᖃᐅᑕᖓᒍᑎ ᐃᖃᓗᓕᐅᑕᐅᔪᒋᑦ ᐃᖃᓗᓕᑕᐅᔪᒋᑦ ᐅᑭᐅᕐᖓᒍᑎ. ᑎᑎᕋᖅᓯᒪᔪᑦ ᐃᓕᐅᕐᖃᐅᔪᓂᑦ ᐊᑐᐃᓐᓇᐅᔪᑦ ᐊᑦᑐᓂ:

ᐅᐸᙵᓯᒪ ᖃᓂᑕᒋᑦ ᐊᐃ ᕿᓚᓂ ᑲᑎᒪᑎᑎᓂᕐᒥᑦ ᐅᖃᓕᐅᑎᕆᔨᒋᑦ

ᐅᔪᕕᑦ ᓄᐊᑯᓵ - ᐃᒪᖏᐅᑕᓕᓚᕆᔨᒋᑦ ᖃᐅᔨᓴᖃᖅᑎᑦ
ᓯᐅᓗ ᐅᓱᓴᓵ - ᐃᒪᖏᐅᑕᓕᓚᕆᔨᒋᑦ ᐃᒪᖏᐅᑕᓂ ᐊᐅᓪᓕᑎᓂᕐᒧᑦ C&A
ᐊᓵᔨᓪ ᓴᖅ - ᐃᒪᖏᐅᑕᓕᓚᕆᔨᒋᑦ ᐃᒪᖏᐅᑕᓂ ᐊᐅᓪᓕᑎᓂᕐᒧᑦC&A
ᐃᓴᑎᓵ Hᐃᐊᓄᓗ - ᐃᒪᖏᐅᑕᓕᓚᕆᔨᒋᑦ ᐃᒪᖏᐅᑕᓂ ᐊᐅᓪᓕᑎᓂᕐᒧᑦ NL
ᑯᐊᑦᓂ ᑕᐅᐳ - ᐃᒪᖏᐅᑕᓕᓚᕆᔨᒋᑦ ᐃᒪᖏᐅᑕᓂ ᐊᐅᓪᓕᑎᓂᕐᒧᑦ NHQ
ᓪᓗ ᐃᐊᑦ - ᐃᒪᖏᐅᑕᓕᓚᕆᔨᒋᑦ ᐃᒪᖏᐅᑕᓂ ᐊᐅᓪᓕᑎᓂᕐᒧᑦ NHQ*
ᑎᐅᓗᔪ ᒪᕼᐅᓂ - ᐃᒪᖏᐅᑕᓕᓚᕆᔨᒋᑦ ᐃᒪᖏᐅᑕᓂ ᐊᐅᓪᓕᑎᓂᕐᒧᑦ NHQ

ᐊᑦᓱ ᓴᐃᔪᑦ – ᓄᓇᕗᓴᒥ ᐅᒪᔪᓕᓚᕆᔨᖏᔪᒋᑦ ᖃᑎᒪᔨᖏᑦᑕ
.ᐃᓴᓵ ᐊᖏᐊᑭᖅ - ᓄᓇᕗᓴᒥ ᐅᒪᔪᓕᓚᕆᔨᖏᔪᒋᑦ ᖃᑎᒪᔨᖏᑦᑕ*
ᑎᐊᓂᔪᑦ ᑎᓗ - ᓄᓇᕗᓴᒥ ᐅᒪᔪᓕᓚᕆᔨᖏᔪᒋᑦ ᖃᑎᒪᔨᖏᑦᑕ

ᕗᖅᓱᓄ ᑲᓵ–ᒧᖏᓵ - ᓄᓇᕕᔭ ᐃᒪᖏᐅᑕᓂᑦ ᐅᒪᔪᓂᑦ ᖃᑎᒪᔨᖏᑦᑕ *

ᐅᑐᔪ ᓴᖅᒪᓵ - ᖃᓇᑕᒥ ᖃᑐᐃᓐᖃᑎᒋᖏᑦᑕ ᓄᓕᒃᕿᓕᓚᐅᖅᑲᑦ
ᐊᓕᔭᑎᐅ ᐅᕆᐃᓕ – ᐅᓄᐅᑕᖅᑐᒥ ᖃᑐᐃᓐᖃᑎᒋᑦ
ᔨᐅᓗ ᕗᐊᖅᑦ - ᖁᓄᖅᑕᒧᓵᔭ ᑯᐊᐅᓱᓴᓵ
ᑯᓚᔪ ᕿᓭᓂᖃᓵ - ᖁᓄᖅᑕᒧᓵᒥ ᐃᖅᓗᓕᓚᕆᔨᒋᑦ ᖃᑐᐃᓐᖃᑎᒋᖏᑦᑕ
ᐳᕆᐃᔨᓵ ᐅᔭ - ᓄᓇᕗᓴᒥ ᐃᖅᓗᓕᓚᕆᔨᒋᑦ ᖃᑐᐃᓐᖃᑎᒋᖏᑦᑕ*
ᐳᕆᐃᔨᓵ ᒧᔭᓇᒪ - ᓂᐅᐸᐅᑕᓵᓂᑦ ᑎᒥᖓᒃ
ᐱᑕ ᑭᔪ - ᒪᑭᕕᔭᒃ ᑯᐊᐅᓱᓴᓵ

ⵊⵔⵉⵎⵓⵜⵏ ⵜⵉⵎⵓⵔⴰⵜⵏ
ⵜⵜⴰⵡⵉⵔⵉⵏ ⵏ ⵜⵎⵓⵔⴰⵜ ⵏ ⵜⵎⵓⵔⴰⵜ ⵏ ⵜⵎⵓⵔⴰⵜ ⵏ ⵜⵎⵓⵔⴰⵜ
ⵜⵉⵣⵔⵉ

ⵜⵓⵔⵉⵎ: X **ⵜⵉⵔⵉⵎⵜⵓ:** **ⵜⵣⵔⵜⵉⵔⵜⵏ:**


**ⵏⵜⵔⵓⵔⵜ ⵜⵣⵔⵜⵉⵔⵜⵓⵔⵜ ⵜⵉⵏⵜ ⵜⵜⵔⵉⵔⵜⵓⵔⵜ ⵜⵜⵔⵉⵔⵜⵓⵔⵜ ⵜⵜⵔⵉⵔⵜⵓⵔⵜ ⵜⵜⵔⵉⵔⵜⵓⵔⵜ ⵜⵜⵔⵉⵔⵜⵓⵔⵜ 2019/20 ⵜⵔⵜⵔⵜⵓⵣⵣ ⵔⵜⵣⵜⵓⵔⵜ ⵜⵔⵉⵔⵜⵓⵔⵜ ⵜⵔⵉⵔⵜⵓⵔⵜ ⵜⵔⵉⵔ


ᐃᖃᓗᖓᓂᒃ ᖃᓄᐃᓄᖅᑕᐅᓪᓗᑎᑎᒃ ᐅᒪᓄᓪᒪᓐᓕᒍᓐᓂᒃ ᐊᖏᕐᓄᖏᑦ ᓇᓱᒪᑦᑐᑎᒃ, ᐃᖃᓗᓪᑕᐅᖅᓯᖃᑎᖅᑕᖅᑐᑦ ᑭᖑᓪᓕᓴᕐᓄᓪᓕᐊᑭᑎᓂᒃ ᐃᖃᓗᓪᓚᓄᓕᕐᕆᑦ, ᓇᓗᓇᐃᖅᑭᔪᖅ ᐅᑎᐃᑦ ᐊᓱᒪ ᑭᖑᓪᓕᐊᑦ ᑕᓴᑭᖅᑐᑦ ᓇᔪᖅᑭᓭᑐᑦ. ᐅᑎᐃᑦ ᐱᓱᒪᓚᐅᔪᐸᑦ ᐱᑕᖃᖅᑐᑦ ᐃᒪᕆᒥ ᐊᐊᒋᑎᖅᓄᓪ , ᓂᕈᖃᖅᑎᑎᓪᓗᑎ ᐊᒥᑭᓱᓕᓴᒃᓂᒃ ᐅᒪᔪᓇᒃ ᑕᓚᐅᖅᑎᐅᑕᓇᒃ ᑎᓭᐊᓇᒃ, ᐃᒪᖅᑎᐅᑕᓇᒃ ᐅᒪᔪᓇᒃ, ᐊᓱᒪ ᐃᖃᓗᓴᓇᒃ.

ᑐᕈᑭᐊᕈᑎᑦ ᐅᑎᖅᓇᒃ ᓄᐊᑕᐅᔪᑦ ᐃᒪᖅᑎᑦᑐᑎᑦ ᖃᐅᔨᓴᖅᑎᓇᒃ ᐃᖃᓗᓪᓚᓂᖅᓯᒪᔪᑦ ᑎᑎᕋᖅᕕᓇᒃ, ᑭᑭᐊᓂᓪ ᐊᒥᑭᓱᓕᓴᒃᓂᒃ ᖃᓇᐃᓪᓚᓂᖅᓯᒪᓂᖅᓯᒪᓂᒃ ᖃᐅᔨᓴᖅᑕᐅᓪᓚᓵᒪᓕᖅᓯᒪᓂᑐᑦ ᖃᐅᔨᓴᖅᑎᓇᒃ, ᐊᓱᒪ ᑕᐃᒪᐃᓕᐅᔪᖅᓯᒪᓂᖅ ᖃᐅᔨᒪᓕᕆᐅᔪᓂᑐᑦ. ᐃᒪᖅᑎᐅᑕᓪᓚᕐᓂᖅᑐᑦ ᐱᓪᓚᓇᓪᓯᖅᑕᐅᖅᑐᑦ CSAS ᖁᒥᖅᑭᓂᒧᑦ ᐱᓭᓪᓕᐊᓂᖓ ᐊᓪᒪ 2019 ᐅᖃᐅᔨᑭᐊᖅᓗᑎ ᐅᑎᖅᓂᒃ ᐃᖃᓗᓪᑕᐅᖅᓱᖅᑕᖅᑐᓂᒃ ᑭᓪᓕᖅᑐᓄᒃ ᓄᓇᓕᖅᑐᓄᒃ WAZ, EAZ ᐊᓱᒪ SFA 1. ᐱᖃᑕᐅᓪᓕᐅᖅᑐᑦ ᑭᓱᖅᑐᐃᐅᑦ ᐃᒪᖅᑎᐅᑕᓪᓚᓵᒪᓕᖅᓯᒪᓂᖅ ᖃᐅᔨᓴᖅᑎᑦ ᐊᓱᒪ RM, ᐊᓱᒪ ᑲᑎᒪᐅᑦ.

ᒥᖏᐅᖏᖅᑕᐅᔪᑦ ᐱᑕᖃᖅᓂᓪᓚᔪᓇᖅᑕᖅᑭᑦ ᐅᑎᐃᑦ ᐊᒥᓱᓂᑦᑐᓄᒃ ᐊᐊᒋᑎᖅᓂ ᐱᖏᐅᓪᓕᐅᖅᑐᑦ ᐱᑕᖃᓚᐊᖃᖅᑎᑕᐅᔪᓂᒃ ᑎᑭᕐᑎᓪᓗᐊᑦᓗᑎ ᐱᖏᐅᔪᓂᑦ ᐊᓂᔪᓂᖅᖁᑦᑐᑦ ᑐᔪᓐᓇᖅᑐᓄᒃ ᐱᑕᖃᖅᑎᑦᑎᐊᑦᓗᑎ ᐱᖏᐅᔪᓂᑦ ᑐᔪᓐᓇᖅᑐᓄᒃ ᐊᐊᒋᑎᖅᓂ

ᑐᖃᓯᕐᑐᑦ – ᑲᓇᓐᓇᖅᓂ ᖃᐅᔨᓴᖅᑕᐅᔪ ᑭᓚᓐᖅ
ᑐᖃᓯᕗᕐᑐᑦ – ᐱᖅᓐᓇᖅᓂ ᖃᐅᔨᓴᖅᑕᐅᔪ ᑭᓚᓐᖅ





ᐊᓱᓴᓚᐊᖅ 1. ᐅᓪᓚᐃᑦ ᐃᖃᓗᑦᑕᐅᖏᓪᓗᒍ ᖃᑦᑕᖅᑐᑦ SFA 3 (ᐱᖏᓴᓐ ᖏᓂᓂ ᖃᐅᓵᓴᖅᑕᐅᔪᓄᑦ ᐅᓴᓕᓐᖏᓗᒍ) ᐅᑯᓂᖏᓐᓂᒃ 1979–2018[1].

[1] . ᑭᖓᓪᓕᐊᓂᕐᓗᑦ ᐊᐅᓚᑦᑎᓂᕐᓗᑦ ᐊᑐᖅᑕᐅᓂᖓᓐᓂ ᐊᓱᓵᓴᖅᑐᖅ ᐅᑉᓗᖅᓯᐅᑎᓂᑦ ᐊᖏᖁᔭᐅᓂ ᐊᖏᖁᔪᓗ ᑭᓇᐅᔭᓕᕆᓂᕐᓗᑦ ᐊᖏᖁᔪᒪᓕ ᐱᒋᐊᖏᓂᖅᑐᖅ 2003.a

9/26/2019

ᖃᐅᔨᒪᓕᐅᕆᓇᐊᓕᓗᒃ ᓕᐃᓴᓐᓂᐅ ᐅᑭᐅᑕᖅᑐᑎ ᑯᖓᔪᓕᓂᒃ ᐊᑲᑭᑎᐊᖅᑕᐅᔾᔪᑦ ᐃᒪᐃᓕᓚᓐᓱᓕᑎ:

5.2.2 ᐱᖅᓯᐅᑎᓐᓇᓂ ᐃᓕᓐᓴ 5.2 ᖃᐅᓪᓗᓂ, ᐃᖅᓱᓴᓯᑎᓯᓗᑦᑕ ᐃᒪᖏᓐᓂ ᐊᐅᓪᓚᑕᐅᔾᔪᓂᑦ ᓄᓇᕗᒻᒥ ᐱᖓᓐᓇᖓᓂ ᐊᐠᒪ/ᐅᕐᓂᓵᓐᓂᑦ ᓄᓇᕆᓇᖓᓂ ᑲᑎᑕᑐᒥᑦ ᐃᖅᓱᑦᑕᐅᖏᓐᖓᑦ ᐅᓴᐃᑦ ᐅᖓᑕᐅᑦᔨᔪᓂᑦ 5 ᐳᓴᓐᑎᓂᑦ ᐅᖃᐃᐃᓐᓂᖏᑎᒍᑦ ᐊᖓᔪᖅᑐᑦ 200kg, ᐊᑐᖅᑕᐅᑕᐅᔪᑦᑐᖅ ᑲᓪᓐᓂᐅᓪᓚᐅᖅᑐᓂᑦ ᐱᖓᓱᓂᔪᖅᑕᖅᑐᑎ, ᓕᐃᓴᓐᓂᒥᑦ ᑎᒍᒥᐊᖅᑐᖅ ᐅᖃᓕᓂᓂᑦ ᐅᒥᐊᖃᐊᒥᑦ ᐊᐅᓪᓚᑦᑎᑦᓂᐅᑎᒥᓕᓐᓄ ᓇᒻᓂᓐᑕᐅᑎᒥᓐᓄᑦ ᖃᐅᔨᓴᓐᓂᑦ (10) ᐃᒪᓐᓇᖅᑐᑦ ᒪᐃᔪᐃᑦ ᓇᒥᑐᐃᓐᓇᖅ ᐃᓂᒋᓐᓂᖓᓂ ᑲᑎᒪᔨᖅᑕᐃᓕᓐᓂᒃ ᐱᓇᓱᐊᖅᑕᐃᓕᓐᓂᓂ ᐃᖅᓱᓕᑕᐅᔾᔪᑎᖅᑐᓂᑦ ᐅᑎᕐᓂᑦ. ᓄᓕᐅᖅᑎᓐᓂᒥᑦ, ᑲᑎᑕᑐᒥᑦ ᐃᖅᓱᓕᑕᐅᔾᔪᑎᖅᑐᓂᑦ ᐅᑎᐃᑦ ᐅᖓᑕᐅᔾᔪᔨᒧᑎ ᐊᖏᓂᓄᖅᓴᒥ 5 ᐳᓴᓐᑎᒥ ᐅᖃᐃᐃᓐᓂᖏᑎᒍᑦ ᑯᖓᔪᖅᑐᑦᓂᐅᑎᒥᓐᓄᑦ, ᐅᑭᐊᖃᐊᓂᑦ 200 ᑭᓚᒧᑦᓱ. ᐊᑐᖅᑕᐅᑕᐅᔪᓕᐅᖅᑐᖅᑐᑦ 6–ᕿᖅᑎᖅᑐᑎᑦ ᑲᓪᓐᓂᐅᖏᓐᖓᑦᓭᓚᐅ, ᐅᒥᐊᖃᐊᖅ ᓄᖏᓐᓇᓗᐊᓕᓐᓂᒃ 10 ᐃᒪᓐᓇᖅ ᒪᐃᔪᓂᑦ ᓇᒥᑐᐃᓐᓇᖅ ᐃᓂᒋᓐᓂᖓᓂ ᐱᖓᓱᓂᔪᖅᑐᓂᑦ ᑲᓪᓐᓂᐅᑕᐅᓕᓐᓂᖅᑐᓂᑦ ᐊᓂᖓᓐᓂᑦ ᐃᖅᓱᓕᑕᐅᒪᓂᖃᑦᑐᑦ. ᓕᐃᓴᓐᓂᒥᑦ ᑎᒍᒥᐊᖅᑐᖅ ᐅᖃᐊᖃᐊᒥᑦ ᐊᐅᓪᓚᑕᐅᔾᔪᑎᖅᑐᓂᑦ ᑎᑎᕋᓕᐊᓕᓐᓂᑦ ᑎᑎᕋᖃᓚᒥ ᐱᓕᐅᑦᑕᐃᓐᓕᒪᓕᒃᑐᑦ ᐅᖓᑕᐅᑦᔨᔪᓂᑦ ᐅᑎᕐᓂᑦ ᐃᖅᓱᓕᑕᐅᖓᔾᔪᖅᑐᑦ. ᓕᐃᓴᓐᓂᒥᑦ ᑎᒍᒥᐊᖅᑐᖅ ᐅᖃᐊᖃᐊᒥᑦ ᐊᐅᓪᓚᑕᐅᑎᓂᖅᑐᖅ ᑎᑎᕋᓕᐊᖅᑐᖅᓴᖓᑦᑕ ᐅᑎᕐᓂᑦ ᐃᖅᓱᓕᑕᐅᖓᔾᔪᖅᑐᑦ (ᓴᓂᒧᑦ ᑐᑭᒧᑦ ᓄᓇᖓᒍᒥᑦ) ᐅᓴᐃᑦ ᐃᖅᓱᓕᑕᐅᖓᔾᔪᖅᑐᓂᑦ ᐊᖓᔪᓂᖓᑦ ᐃᖅᓱᓕᑕᐅᔾᔪᑦ ᐅᖃᐃᐃᓐᓂᖏᑎᒍᑦ ᑭᓚᒧᑦᓱ.

ᑕᑯᓇ ᐊᑲᑭᑎᐊᑭᑎᐅᔾᔪᖅ ᐱᑎᐊᖃᓂᑲᖅ ᐅᓪᓗᓚᖓᓂ ᑐᓂᐅᔾᔨᐅᔨᒥᑦ ᒧᑦᓯ 31, 2020.

ᖃᐅᔨᓴᕐᑐᑦ ᑎᑎᕋᖅᓯᒪᔭᖅᓴᖅ 2019/073

**ᓇᐃᓕᓐᓴᕐᑕᒃ: ᖃᒥᖅᑭᐅᐅᓂᖏᑦ ᑕᑯᓴᕈᑐᐃᓇᓇᓕᓪᓕᐊᓪᓕᑦ ᐸᐅᑐᐃᓂᖅᓂᖅᑐᑦ ᐱᓕᐅᓇᓯᓇᓕᑎᐅ ᑐᖏᑲᖅᑕᐅᔪᑦ ᐅᑭᐅᖅᑕᖅᑐᒃ ᐅᑲᓐᖏᑦ (ᓴᓇᐅᔨᑕᑦ, ᓴᐃᑐᒃ) ᑲᑎᕋᖅᓯᒪᓂᖏᑦ ᐱᓇᓯᑕᐅᑎᓪᓗᓂᑦ ᐅᑭᐅᖅᑕᖅᑐᒃ (ᕿᓪᑕᓪᓚᓵ, ᓴᐃᓇᓇᓚᓵ) ᐊᒻᒪᓗ ᑕᖅᓴᓪᓕᑦ (ᕿᓪᑕᓪᓚᓵ, ᒪᓕᓪᑕᓪᓚᓵ) ᑲᑎᒪᔨᖅ ᐃᓐᓇᖅᑐᓕᕐᓗᓂᑦ ᐅᓇᖅᓯᓪᓕᐊᓂᑦ ᐱᑭᖅᑐᓂᑦ ᐅᑭᐅᖅᑕᖅᑐᒃ (ᐃᓚᖅᓗ, ᑕᐃᑲᖅᓯᒪᓂᖅ 2017), ᑭᖓᔪᓂᓂᐊᓂᖅ ᐃᖅᑲᓇᐃᓕᓐᓂᖅᑐᑦ ᐅᒥᐊᔪᐊᖅᑭᓂᓂᖅ ᐊᐅᓚᑦᑎᐊᓂᖅ ᐃᓕᓴᕐᓂᖅᑐᑦ ᐃᓚᖅᑭᑦᑐᖅᓯᒪᓂᖅ ᐃᓕᓴᕐᓂᖅᑐᑦ ᐃᓕᓴᕐᓂᖅᑐᑦ ᐃᓕᓴᕐᓂᖅᑐᑦ

ᖃᐅᔨᓴᕐᑎᓄᑦᓯᑦ ᐅᖃᐅᓯᖅᓯᐊᕐᓚᒍᑦ 2020/007

ᓇᐃᓕᓕᐊᓗᓐᓂᕐᓴ: ᖃᒥᖅᑭᕆᐅᓂᖅᓴᖅ ᐅᑭᐅᖃᖅᑕᖅᑐᑦ ᐅᓚᖅᑎᑦ (ᓴᐊᓚᐅᑉᓯᑦ ᓴᐃᑎ) ᐱᓇᓱᑕᐅᓇᑎᒃ ᐃᖃᓱᒃᑕᓇᓄᑦ ᐅᖃᐅᖃᖅᑕᖅᑐᐊᓂ ᑭᖑᓱᒃᑐᑎᓱᒃᑐᑦᑎᓄᑦ ᐃᖃᓗᓯᓱᒃᑐᑎᑦᑎᓄᑦ

- ᑭᖑᓱᒃᓴᓱᒃᑎᑦ ᐃᖃᓗᓯᓱᓕᑦ ᑲᓇᑕᐅᑦ ᐅᑭᐅᖃᖅᑕᖅᑐᖓᓂ ᐊᑐᑎᖃᒡᓗᓕᖅᑐᑦ ᓱᓇᒃᑯᑦᑕᓚᕐᑎᑕ ᐃᓄᐃᑦ ᓄᓇᕗᒻᒥ ᓄᓇᖃᖓᓕᕐᓱᐊᑕᑕᐅᓕᓪᓚᒃᑎ ᑭᖑᓱᒃᐸᐃᑦ ᐃᖃᓗᓯᓱᑎᑎᓕᐅᑦᐸ ᐃᓂᖓᓂᑕ 1 (SFA 1) ᐊᓯᒪᓗ ᑕᒃᑯᐊᒃ ᑲᓇᓴᓂᓕᓪᓚ ᐱᖓᓱᐅᑉᖢᕆᔨᑎᑦᑕ ᑭᓚᖃᖅᑎᖓᑦ (EAZ/WAZ). ᑕᑯᐃᓴᓇᓚᓪᓚᔪ, ᐱᐅᔪᖅ ᐊᖑᑐᖃᖅᑎᑦᑕᑯᓘᑦ ᐃᓚᐃᑦᑐᐃᓐᔪᓚᓪᓚ ᑭᖑᓱᒃᐸᑕᐅᓱᓄᑦ ᐊᓯᖅᓯᐊᔪᑦ ᐃᓚᐃᑦᑐᔪᑦ ᑭᖑᓱᒃᐸᑕᐅᓱᓄᑦ ᐃᓚᐃᑦᑐᖓᓕᖅᑐᑦ (ᓴᓇᐅᑦᓴᑦ ᓴᐃᑎᑕ) ᐃᖃᓗᖃᓕᕆᔨᑦ ᐅᓪᓴᓇᐅᓕᕐᓚᔪᓂᑦ.
- ᐱᔨᔪᑎᑦᓚᓚᔪ ᐊᒥᓚᔪᐊᖅᑐᒥᑲ ᓇᐃᓴᐅᑎᑦᔪᑎᖅᓂᖅᓯᖓᓕᓄᑦ, ᖃᐅᔨᓴᖅᑐᑦ ᐅᖃᐅᔨᓴᖅᓯᖅᑐᑦ ᓴᖅᑭᔪᓂᓇᕐᓂᑎᑲ ᑭᓚᓕᑦᑕᐅᑦ ᐱᓚᒥᑎᑲ ᐊᐱᖅᔪᑎᑦᑎᓕᕐᓚᔪᓂᑦ ᑭᖑᓕᒍᑦᑕᐅᔪᓕᒪ ᐱᖓᓱᒃᑐᑦᑎᓄᑦ ᐊᓯᖅᓯᐊᔪᑦ ᐊᐱᖅᔪᑎᑦᑎᓕᕐᓚᔪᓂᑦ ᐊᐱᖅᔪᑎᑦᑎᓕᕐᓚᔪᓂᑦ (ᓴᓇᐅᑦᓴᑦ ᓴᐃᑎᑕ) ᐃᖃᓗᖃᖅᑎᑦᑎᓕᕐᓚᔪᓂᑦ ᐃᓚᐃᑦᑐᔪᑦ

ᓱᐅᓂᒃᑯᑦ ᓴᖅᑭᑎᑕᓂᕐᒥᒃ ᖃᐅᔨᓴᖅᑎᑦ ᐅᖃᐅᔨᔨᖅᓱᐊᕐᓂᒃᑐᓂᓕᕆᔨᐊᑎᓂᒃ ᓇᓪᓕᐊᓇᕐᓵᑎᓂᒃ ᐃᖃᓗᐊᓂᑦ ᐱᖃᓱᐅᕐᓴᖅᑎᑦ ᐃᖃᓗᓐᓄᑦ.

- ᑐᑭᖓᓪᓕᓇᕐᑐᖅ ᖃᐅᔨᓴᖃᓂᖅ ᐱᑕᖃᓇᐊᓕᒃ ᐅᑭᐅᖅᑕᖅᑐᑦ ᐅᑎᖅᑕ ᐅᒪᔨᔨᓱᖅᑐᓂᑦ, ᐊᐅᑕᓂᓇᓱᖅᑕ ᐅᒪᓂᐊᒍᑦ, ᐊᐅᓱᓂᖅᑐᓂᑦ, ᐊᓚᒡᑕᐅᖅ ᐅᒪᔨᓂᑦ ᐊᐅᑕᓪᓚᕐᓂᕐᕕᑕ ᑕᐅᑐᖓᐊᕐᑎᖅᑐᓂᑦ ᑐᑭᓯᓂᓪᓕᐊᓐᓂᒃᑯᑦ ᑲᓇᑕᒥ ᖃᑎᖅᑕᐅᑦᑕᐅᓚᖓᑕᑕ ᐅᒪᔨᓂᑦ ᓂᖃᖅᑐᖓᓂᒃᑯᑦ. ᑕᒪᓐᓇ ᖃᐅᔨᔨᖅᑎᓂᑦ ᓴᖅᑭᑦᓴᕆᔭᒃᑯᑦ ᓇᐃᔨᐅᑎᔨᐅᓂᖅᓴᖅᑯᑦ ᑲᑎᕐᓂᖅᑐᓂᑦ ᐊᐃᔨᐅᑎᓂᑦ ᓇᓕᐅᑦᑕᒃᓴᖅᑐᓂᑦ ᐱᐅᓂᐊᕐᓇᓕᒃ ᐃᓕᓴᕐᕕᐅᑎᓂᑦ ᐅᖃᐅᔨᓴᖅᑎᓂᑦ ᐱᓇᓱᐊᕐᓇᓕᒃ ᐃᖅᑲᓇᐅᑦᑕᐅᔪ

Fisheries and Oceans Canada | Pêches et Océans Canada
Ecosystems and Oceans Science | Sciences des écosystèmes et des océans

Canadian Science Advisory Secretariat (CSAS)

Research Document 2019/073

Central and Arctic Region

Assessment of potential impacts of bycatch mortality on the Arctic Cod (*Boreogadus saida*) populations from the Northern (*Pandalus borealis*) and Striped (*Pandalus montagui*) Shrimp fisheries in Shrimp Fishing Areas 1, 2, and 3

Wojciech Walkusz, Sheila Atchison, Kevin Hedges, and David Deslauriers

Freshwater Institute
Fisheries and Oceans Canada
501 University Crescent
Winnipeg, Manitoba, R3T 2N6

January 2020

Canada

Foreword

This series documents the scientific basis for the evaluation of aquatic resources and ecosystems in Canada. As such, it addresses the issues of the day in the time frames required and the documents it contains are not intended as definitive statements on the subjects addressed but rather as progress reports on ongoing investigations.

Published by:

Fisheries and Oceans Canada
Canadian Science Advisory Secretariat
200 Kent Street
Ottawa ON K1A 0E6

http://www.dfo-mpo.gc.ca/csas-sccs/
csas-sccs@dfo-mpo.gc.ca

© Her Majesty the Queen in Right of Canada, 2020
ISSN 1919-5044

Correct citation for this publication:

Walkusz, W., Atchison, S., Hedges, K., and Deslauriers, D. 2020. Assessment of potential impacts of bycatch mortality on the Arctic Cod (*Boreogadus saida*) populations from the Northern (*Pandalus borealis*) and Striped (*Pandalus montagui*) Shrimp fisheries in Shrimp Fishing Areas 1, 2, and 3. DFO Can. Sci. Advis. Sec. Res. Doc. 2019/073. iv + 8 p.

Aussi disponible en français :

*Walkusz, W., Atchison, S., Hedges, K., et Deslauriers, D. 2020. Évaluation des impacts potentiel de mortalité par des prises accessoires sur les populations de morue arctique (*Boreogadus saida*) dans les pêches à la crevette nordique (*Pandalus borealis*) et crevette ésope (*Pandalus montagui*) dans les zones de pêche à la crevette 1, 2, et 3. Secr. can. de consult. sci. du MPO. Doc. de rech. 2019/073. iv + 9 p.*

TABLE OF CONTENTS

ABSTRACT iv
INTRODUCTION 1
REVIEW OF ARCTIC COD BIOLOGY AND ECOLOGY 2
DISTRIBUTIONAL PATTERNS AND BIOMASS INDICES 2
ECOSYSTEM REQUIREMENTS 3
ANALYSIS OF HISTORICAL BYCATCHES 4
UNCERTAINTIES OF ANALYSIS 6
Overall lack of knowledge of Arctic Cod biology in the studied area. 6
Insufficient data collection for the proper biomass index analysis. 6
Insufficient data on distribution and abundance of Arctic Cod predators. 6
Potential observer data bias. 7
DISCUSSION 7
REFERENCES CITED 7

ABSTRACT

Arctic Cod (*Boreogadus saida*) is considered a key forage species in Arctic marine ecosystems due to its pivotal role in the food web, serving as prey for numerous marine animals (seals, whales, birds, and fishes) and being an important consumer of secondary production (zooplankton). In recent years (i.e., since 2017), shrimp fishing vessels operating in Shrimp Fishing Areas 1 (Baffin Bay) and 3 (Hudson Strait), have occasionally reported large amounts (400–2300 kg per tow) of Arctic Cod bycatch. Fisheries and Oceans Canada (DFO) Resource Management approached DFO Science seeking advice on ecologically and biologically-responsible Arctic Cod removal levels in individual Shrimp Fishing Areas (SFAs) in the north. This research document reviews the Arctic Cod biology and ecology, considers distributional patterns, and analyzes historical bycatches in the Canadian Eastern Arctic shrimp fisheries. Also, an attempt was made to quantify Arctic Cod population size in the area of interest. Due to the limited data available for this analysis, the ecosystem requirements (i.e., consumption demands of Arctic Cod predators) were used as an indirect estimate of the population size. By contrasting the size of the estimated Arctic Cod population with the bycatch levels, the assumption was made that the impact of bycatch removal in each Shrimp Fishing Area on the Arctic Cod population is minimal. With a limited scope of the Arctic Cod biomass data and a high degree of uncertainty surrounding the biomass indices, ecologically and biologically-responsible Arctic Cod removal levels in individual Shrimp Fishing Areas (SFAs) cannot be established at present. Therefore, when amending the conditions of licence in response to the elevated Arctic Cod bycatch which would result in increased removals, caution is advised.

INTRODUCTION

Arctic Cod (*Boreogadus saida*; Figure 1) is considered a key (pivotal) forage species in Arctic marine ecosystems (Bradstreet et al. 1986). As a pelago-benthic dweller it occurs in both the pelagic realm and near the ocean floor (Geoffroy et al. 2016).



Figure 1. Arctic Cod (Boreogadus saida). Photo credit: Claude Nozères, DFO.

Larval and juvenile Arctic Cod concentrate mainly in the upper water column, taking advantage of the zooplankton (Copepoda) food base (Geoffroy et al. 2016). Upon reaching a certain age and length (age 1+, approx. 60 mm; Walkusz et al. 2013.) individuals descend to deeper depths and remain demersal for the rest of their life (Majewski et al. 2016). In benthic habitats, larger Arctic Cod feed on Amphipods and small fishes (Walkusz et al. 2013). Since younger and older individuals occur in both pelagic and benthic realms (Figure 2), it is difficult to study an entire Arctic Cod population as doing so would require the simultaneous use of pelagic and demersal fishing gear. Recently, hydroacoustic methods have proven useful for studying Arctic Cod populations; however, hydroacoustic signals still require ground-truthing to validate the remote observations.




Figure 2. Echogram recorded with SIMRAD EK-80 acoustic system (38 kHz) at a 353 m station in the Amundsen Gulf. The red rectangle (top) indicates a subsurface aggregation of young-of-the year fish, while the green rectangle (bottom) indicates a benthic aggregation of larger individuals. Note the vertical extension (thickness, 80 – 100 m) of each of the aggregations in contrast to the sampling capacity of fishing gear (few meters vertical opening). Courtesy of Andy Majewski (DFO, data unpublished).

Arctic Cod is extremely important in Arctic marine ecosystems as food for numerous marine animals (seals, whales, birds, and fishes) and as an important consumer of secondary production (zooplankton). Due to its key linking role in food webs, Arctic Cod is also of interest

to Indigenous Peoples who directly profit from healthy populations of marine resources, through subsistence harvests and/or commercial fishing. The intermediary position of Arctic Cod in Arctic marine food webs is reflected in the relatively large number of scientific projects devoted to understanding its distribution, feeding, and life cycle. Through these observations, Arctic Cod has been found in dense aggregations offshore (Majewski et al. 2016) and in dense nearshore schools (Welch et al. 1996). This patchy distribution pattern might increase the risk to Arctic Cod populations from commercial fishing activities when fishing locations overlap with areas of high Arctic Cod density.

Since 2017, shrimp fishing vessels operating in Shrimp Fishing Areas (SFA) 1 (Baffin Bay) and 3 (Hudson Strait), have occasionally reported large amounts (400–2300 kg per tow) of Arctic Cod bycatch. In Eastern Canada, Arctic Cod is listed as a groundfish species (Atlantic Fishery Regulations 1985), which means that elevated amounts of Arctic Cod bycatch can trigger a standard "move away" licence provision, as per the Conditions of Licence (DFO 2018), which states: "*In the event that the total incidental catch of all groundfish species in any set exceeds the greater of 2.5% or 100 kg total weight, the licence holder/operator must immediately change the vessel's fishing area by a minimum of 10 nautical miles from any coordinate during the last tow*". High occurrences of Arctic Cod in bycatch, while not common, pose a significant burden on the fishing industry and presently unknown ecological consequences. With limited time to harvest the allotted shrimp catch each year (because of a relatively short open-water period and long transit times from southern ports of origin), the fishing industry in the Eastern Arctic has asked Resource Management for increased flexibility for Arctic Cod bycatch, with move away provisions triggered based on amounts caught averaged over a number of consecutive tows.

As a result, DFO Resource Management approached DFO Science seeking advice on ecologically and biologically-responsible Arctic Cod removal levels in individual SFAs.

Given the absence of a comprehensive Arctic Cod stock assessment, the purpose of this report is to discuss ecologically and biologically tolerable levels for Arctic Cod removals in SFAs 1-3 and if it is possible to create a "cap'' system for the total annual bycatch level.

REVIEW OF ARCTIC COD BIOLOGY AND ECOLOGY

Arctic Cod is a relatively small fish (maximum size: 400 mm, mean: 140 mm; Matley et al. 2013) with a maximum life span of seven years. It sometimes co-occurs with Polar Cod (*Arctogadus glacialis*), which is similar in appearance, and the two species can be mistakenly identified for each other. Arctic Cod spawns in winter (Hop et al. 1995) and the larvae hatch between April and June depending on geographical location (Bouchard and Fortier 2011). Larval Arctic Cod feed initially on smaller prey items (e.g., Copepoda nauplii, Rotifers), advancing to larger prey items as they grow (e.g., *Calanus* copepods; Walkusz et al. 2011). Juvenile fish (approx. 35 mm in length) continue feeding on Copepods. After reaching the size of descent (approx. 50 mm; Falk-Petersen et al. 1986, Majewski et al. 2016) Arctic Cod adopt a demersal life and in addition to *Calanus* copepods start feeding on Amphipods and fishes, including cannibalism. Arctic Cod is considered a schooling fish (Hop et al. 1997), however, "aggregating fish" seems a more appropriate term considering the dense aggregations of different size/age individuals and non-unidirectional movement observed by Geoffroy et al. (2011, 2016).

DISTRIBUTIONAL PATTERNS AND BIOMASS INDICES

Arctic Cod is widely distributed in the Arctic. It is often dispersed in the water column (Figure 2) and is therefore hard to properly quantify using any one type of direct sampling gear (either demersal or pelagic). Walkusz et al. (2019) calculated Arctic Cod biomass indices for SFAs 1, 2,

and 3 based on biomass estimates from demersal scientific surveys only (2005–2017). These biomass estimates (maximum 138 t, 47,000 t, and 30,000 t in SFA 1, 2, and 3, respectively) were certainly underestimated as Arctic Cod can essentially occupy the entire water column and its distribution is extremely patchy. Arctic Cod has been observed in large nearshore schools in Barrow Strait (Welch et al. 1996), with aggregations being estimated to have a biomass of 20,000 t. Also, Welch et al. (1992) claimed that nearshore Arctic Cod aggregations along the Canadian Arctic Archipelago are common and were estimated to be approximately 75,000 t. Using acoustic methods that integrate biomass in the water column and near the bottom, Benoit et al. (2008) recorded large aggregations of Arctic Cod in Franklin Bay, with average biomass being 11.2 kg m^{-2} (equal to 11,200 t km^{-2} assuming continuous distribution) from February to April, with the maximum reaching 55 kg m^{-2} (equal to 55,000 t km^{-2} assuming continuous distribution). These high densities were assumed to be localized. Using similar methods, Geoffroy et al. (2011) detected a large winter aggregation in the Amundsen Gulf. They calculated that in February the biomass peaked at 0.732 kg m^{-2} (equal to 732 t km^{-2} assuming continuous distribution); however, the aggregation was most likely confined to the northern portion of the Amundsen Gulf.

ECOSYSTEM REQUIREMENTS

Using predator populations to estimate food base requirements has been used in marine research. For example Stenson (2013) estimated the energy requirements of individual prey populations required to sustain Harp Seal population. Walkusz et al. (2013) showed that for the Canadian Beaufort Shelf (area of 63,000 km^2), biomass calculated from demersal trawl surveys only accounts for a small fraction of all Arctic Cod biomass needed to meet the bio-energetic requirements of the ecosystem (i.e., feeding by seals, whales, birds, and fishes). Similar conclusions were presented by Welch et al. (1992) who claimed that biomass estimates from ‘dispersed’ fish produce significantly underestimated values. Walkusz et al. (2013) estimated that using the conservative assumption of approximately 20,000 beluga whales that consume a total of 440 t of fish per day and 500,000 ringed/bearded seals that consume a total of 2,500 t of fish per day require a total of 3,000 t of Arctic cod daily to sustain their predators' needs in the Beaufort Sea. It means that 500,000–1,000,000 t of fish are consumed over the course of the year (whales and birds are absent from the area after freeze-up). This amount, while high, would not be exceptional, considering Welch et al. (1992) estimated that 148,000 t of Arctic Cod is consumed annually in Lancaster Sound (area of 98,000 km^2).

Based on recent estimates for fish, bird, seal, and whale populations in Baffin Bay/Davis Strait (Table 1), Arctic Cod biomass requirement for ecosystem maintenance is similar (or higher) than the Beaufort Sea (i.e., > 500,000 t Arctic Cod annually, assuming 10% of all Northwest Atlantic marine mammal populations utilizing the area of interest). While this approach most likely puts the estimate of Arctic Cod biomass required to sustain their predators on the conservative side, it provides a general estimate as to the order of magnitude of the Arctic Cod biomass held in the ecosystem.

Table 1. Species and abundances of known Arctic Cod predators present in the Baffin Bay/Davis Strait area (source: Treble 2017, NIRB 2018).

Species	Abundance
Fishes (t)	
Greenland Halibut	200,000
Birds (individuals)	
Thick-billed Murre	860,000
Little Auk (dovekie)	30,000,000
Northern Fulmar	Not quantified but sizeable
Seals (individuals)	
Ringed seal	1,200,000
Bearded seal	Not quantified, but considered very abundant
Harp seal	7,400,000 in the Northeast Atlantic
Whales (individuals)	
Beluga whale	20,000
Narwhal	45,000

ANALYSIS OF HISTORICAL BYCATCHES

Arctic Cod, which is similar in size to shrimp, is a common bycatch species in the Eastern Canadian Arctic shrimp fishery. While the amount of Arctic Cod bycatch in a single tow is usually low, large catches are occasionally reported. Arctic Cod catches have no economic value and are discarded at sea.

Historical records for Arctic Cod bycatch in SFA 1 ranging between zero in years with no fishing to 143 t in 1993 (Figure 3; Walkusz et al. 2019). In SFA 2, the peak Arctic Cod bycatch was 31 t in 2004/05 (Figure 4).




Figure 3. Arctic Cod bycatch in SFA 1 for 1979–2018 (Walkusz et al. 2019).




Figure 4. Arctic Cod bycatch in SFA 2 for 1979-2018. Shrimp fishery management cycle changed from calendar year to fiscal year starting in 2003.

In SFA 3, incidental catches of Arctic Cod were low until recent years, when 47 t was caught in 2018 (Figure 5). The size of Arctic Cod catches is heavily related to aggregating processes, therefore incidental catches will reflect fluctuations in these processes.




Figure 5. Arctic Cod bycatch in SFA3 for 1979–2018. Shrimp fishery management cycle changed from calendar year to fiscal year starting in 2003.

UNCERTAINTIES OF ANALYSIS

Overall lack of knowledge of Arctic Cod biology in the studied area.

While general biology of Arctic Cod has been studied extensively in various Arctic locations, its biology is poorly understood in the geographic area considered in this review. That limitation applies to the timing of descent of juvenile Arctic Cod from pelagic to benthic habitats (i.e., when they become susceptible to mortality due to the shrimp fishery). Multi-seasonal sampling along with length analyses of individuals caught would provide insight into the ontogenetic vertical migrations of Arctic Cod.

Insufficient data collection for the proper biomass index analysis.

In this analysis, the biomass index was based solely on the benthic catches; however, Arctic Cod is considered a pelago-benthic dweller. The lack of information on the pelagic portion of the population can lead to significant underestimates of the biomass. Application of acoustic methods along with pelagic trawl sampling would address this knowledge gap.

Insufficient data on distribution and abundance of Arctic Cod predators.

Analysis of predation pressure on Arctic Cod was done using published abundances of predators with the assumption that a portion of the predator population (i.e., 10%) would be present in the area of interest. Using this assumption, the biomass of Arctic Cod required to sustain its predators was estimated. While this generalization provides some insight into Arctic

Cod abundance, it is not a direct measure of fish biomass and the uncertainty associated with this approach can be significant.

Potential observer data bias

All offshore vessels operating in the area of interest are obliged by the Conditions of Licence, to have an At Sea Observer (ASO) onboard. The data collected by the ASO is meant to be fishery independent; however, it can be expected that there are times when particular tows are not monitored due to the limitations of having only one ASO onboard, which may create bias in the information reported.

DISCUSSION

Considering the relatively small total mortality (less than 0.1% of the stock annually as proposed above based on predators' requirements) of Arctic Cod stemming from the Eastern Canadian Arctic shrimp fishery, it can be assumed that the impact of shrimp trawling on the Arctic Cod population is minimal. It should also be noted that the Arctic Cod bycatch is not retained by fishing vessels and gets discarded, thus, the biomass is not removed from the ecosystem.

With a limited scope of the Arctic Cod biomass data and a high degree of uncertainty surrounding the biomass indices, ecologically and biologically-responsible Arctic Cod removal levels in individual Shrimp Fishing Areas (SFAs) cannot be established at present. Therefore, when amending the conditions of licence in response to elevated Arctic Cod bycatch which would result in increased removals, caution is advised.

In order to answer the question of the ecologically and biologically-responsible Arctic Cod removal levels in individual SFAs, the following would have to take place:

- Study of Arctic Cod biology/ecology including migratory patterns, aggregating events, and vertical distribution patterns (population split between pelagic and benthic habitats); this could be achieved by a dedicated survey that would combine acoustic survey with pelagic and benthic trawl sampling to ground truth the acoustic data; and,
- Study of Arctic Cod role in the ecosystem, including its role as a consumer (of zooplankton) and food for higher trophic levels (fish, seals ,whales, birds); this could be achieved by studies of Arctic Cod stomach contents and biomarkers (stable isotopes and fatty acids), and through a better assessment of the marine mammals and bird populations in the area.

REFERENCES CITED

Atlantic Fishery Regulations, 1985. Canada Gazette Part II, 120 (1), SOR 86-21.

Benoit, D., Simard, Y., and Fortier, L. 2008. Hydroacoustic detection of large winter aggregations of Arctic cod (*Boreogadus saida*) at depth in ice-covered Franklin Bay (Beaufort Sea). J. Geophys. Res. 113: C06S90. doi:10.1029/2007JC004276

Bouchard, C., and Fortier, L. 2011. Circum-arctic comparison of the hatching season of polar cod *Boreogadus saida*: a test of the freshwater winter refuge hypothesis. Prog. Oceanogr. 90: 105–116.

Bradstreet, M.S.W., Finley, K.J., Sekerak, A.D., Griffiths, W.B., Evans, C.R., Fabijan, M.F., and Stallard, H.E. 1986. Aspects of the biology of arctic cod (*Boreogadus saida*) and its importance in arctic marine food chains. Can. Tech. Rep. Fish. Aquat. Sci. 1491: 193 pp.

DFO. 2018. Integrated Fisheries Management Plan (IFMP): Northern shrimp and striped shrimp – Shrimp fishing areas 0, 1, 4-7, the Eastern and Western Assessment Zones and North Atlantic Fisheries Organization (NAFO) Division 3M. Fisheries and Oceans Canada, Ottawa, ON. 84 p.

Falk-Petersen, I.B., Frivoll, V., Gulliksen, B., and Haug, T. 1986. Occurrence and size/age relations of polar cod, *Boreogadus saida* (Lepechin), in Spitsbergen coastal waters. Sarsia 71: 235–245.

Geoffroy, M., Majewski, A., LeBlanc, M., Gauthier, S., Walkusz, W., Reist, J.D., and Fortier, L. 2016. Vertical segregation of age-0 and age-1+ polar cod (*Boreogadus saida*) over the annual cycle in the Canadian Beaufort Sea. Polar Biol. 39: 1023–1037.

Geoffroy, M., Robert, D., Darnis, G., and Fortier, L. 2011. The aggregation of polar cod (*Boreogadus saida*) in the deep Atlantic layer of ice-covered Amundsen Gulf (Beaufort Sea) in winter. Polar Biol. 34: 1959–1971.

Hop, H., Graham, M. and Wordeau, V.L., 1995. Spawning energetics of Arctic cod (*Boreogadus saida*) in relation to seasonal development of the ovary and plasma sex steroid levels. Can. J. Fish. Aquat. Sci. 52: 541–550.

Hop, H., Welch, H., and Crawford, R., 1997. Population structure and feeding ecology of Arctic cod schools in the Canadian High Arctic. American Fisheries Society Symposium 19: 68–80.

Majewski, A.R., Walkusz, W., Lynn, B.R., Atchison, S., Eert, J., and Reist, J.D., 2016. Distribution and diet of demersal Arctic Cod, *Boreogadus saida*, in relation to habitat characteristics in the Canadian Beaufort Sea. Polar Biol. 39: 1087–1098.

Matley, J.K., Fisk, A.T., and Dick, T.A. 2013. The foraging ecology of Arctic cod (*Boreogadus saida*) during open water (July–August) in Allen Bay, Arctic Canada. Mar. Biol. 160: 2993–3004.

NIRB (Nunavut Impact Review Board). 2018. Strategic Environmental Assessment for Baffin Bay and Davis Strait: Environmental Setting and Review of Potential Effects of Oil and Gas Activities. Nunami Stantec. 584 pp.

Stenson, G.B. 2013. Estimating consumption of prey by Harp Seals, (*Pagophilus groenlandicus*) in NAFO Divisions 2J3KL. DFO Can. Sci. Advis. Sec. Res. Doc. 2012/156. iii + 26 p.

Treble, M. 2017. Report on Greenland halibut caught during the 2016 trawl survey in Divisions 0A and 0B. NAFO SCR Doc. 17-028: 30 p.

Walkusz, W., Hedges, K., Atchison, S., and Deslauriers, D. 2019. Arctic Cod (*Boreogadus saida*) Bycatch in Shrimp Fishing Areas 1-3: 1979-2018. Can. Tech. Rep. Fish. Aquat. Sci. 3340: v + 16 p.

Walkusz, W., Majewski, A.R., and Reist, J.D. 2013. Distribution and diet of the bottom dwelling Arctic cod in the Canadian Beaufort Sea. J. Mar. Syst. 127: 65–75.

Walkusz, W., Paulic, J.E., Williams, W.J., Kwasniewski, S., and Papst, M.H. 2011. Distribution and diet of larval and juvenile Arctic cod (*Boreogadus saida*) in the shallow Canadian Beaufort Sea. J. Mar. Syst. 84: 78–84.

Welch, H.E., Bergmann, M.A., Siferd, T.D., Martin, K.A., Curtis, M.F., Crawford, R.E., Conover R.J., and Hop, H. 1992. Energy flow through the marine ecosystem of the Lancaster Sound Region, Arctic Canada. Arctic 45: 343–357.

Welch, H.E., Crawford, R.E., and Hop, H. 1996. Occurrence of Arctic Cod *(Boreogadus saida)* schools and their vulnerability to predation in the Canadian High Arctic. Arctic 46: 331–339.

Fisheries and Oceans Canada | Pêches et Océans Canada
Ecosystems and Oceans Science | Sciences des écosystèmes et des océans

Central and Arctic Region

Canadian Science Advisory Secretariat
Science Advisory Report 2020/007

REVIEW OF ARCTIC COD (*BOREOGADUS SAIDA*) BYCATCH IN CANADIAN ARCTIC SHRIMP FISHERIES



*Arctic Cod (*Boreogadus saida*).*




Figure 1. Echogram recorded with SIMRAD EK-80 acoustic system (38 kHz) at a 353 m station in the Amundsen Gulf. The red rectangle (top) indicates a subsurface aggregation of young-of-the-year fish, while the green rectangle (bottom) indicates a benthic aggregation of larger individuals. Note the vertical extension (thickness, 80–100 m) of each of the aggregations in contrast to the sampling capacity of fishing gear (few meters vertical opening). Courtesy of Andy Majewski (DFO, data unpublished).

Context:

*Incidental bycatch and discard of non-targeted species occur in many fisheries. An objective of an Ecosystem Approach to Fisheries Management is to control incidental mortality of non-targeted species. Arctic Cod (*Boreogadus saida*) is a ubiquitous species occurring in a wide range of habitats in the Arctic. Given this widespread distribution and tendency to form large aggregations, it is a common bycatch species in various northern fisheries including the Northern and Striped Shrimp fisheries in the Eastern Canadian Arctic.*

Fisheries and Oceans Canada (DFO) Resource Management requested that DFO Science provide advice on a sustainable overall bycatch limit of Arctic Cod (in tonnes) for Shrimp Fishing Area 1 and the Eastern and Western Assessment Zones (EAZ and WAZ) as well as establishing suitable references to indicate when that mortality is unacceptable. In providing the requested advice, DFO Science was asked to consider the current "move away" provision on commercial licences.

*This Science Advisory Report is from the July 3–4, 2019 meeting on Advice on Arctic Cod (*Boreogadus saida*) Bycatch Limits in Northern Shrimp Fisheries in the Canadian Arctic. Additional publications from this meeting will be posted on the Fisheries and Oceans Canada (DFO) Science Advisory Schedule as they become available.*

February 2020

Canada

SUMMARY

- The shrimp fishery in the Canadian Arctic is an important resource for Inuit of Nunavut and Nunavik and the commercial shrimp industry in Shrimp Fishing Area 1 (SFA 1) and the Eastern and Western Assessment Zones (EAZ/WAZ). In general, a positive correlation between the total shrimp catch and Arctic Cod (*Boreogadus saida*) bycatch has been observed.
- Due to insufficient data, Science Advice could not be provided in regards to a “cap” for Arctic Cod removal and specifications of the “move away” provision.
- Due to the scarcity of data and the patchy spatial and temporal distribution of Arctic Cod, there is high uncertainty in the biomass index estimates obtained from the surveys that are not designed to target Arctic Cod. This uncertainty leads to low confidence in determining the impacts of Arctic Cod removal by the shrimp fishery. If the biomass index estimates are considered accurate, then the percentage of Arctic Cod removed from the total biomass is likely very low but given the high uncertainty, precaution is advised.
- The absence of a dedicated survey and the limited data on life history of Arctic Cod preclude the use of the Guidance Related to Bycatch and Discards in Canadian Commercial Fisheries (DFO 2012) to determine a sustainable catch limit.
- Canada is committed domestically and internationally to implementing conservation and management measures to address incidental bycatch, including Arctic Cod in the Northern and Striped Shrimp fisheries in the Eastern Canadian Arctic.
- Arctic Cod is a small bodied, short-lived forage fish with an atypical life history for gadoids as it occupies the pelagic zone for a portion of its life beyond the larval and juvenile stages. Despite these characteristics, Arctic Cod is specified as groundfish in the Atlantic Fishery Regulations and therefore “move away” provisions in the offshore Northern shrimp Conditions of Licence apply.
- Arctic Cod transfers energy from lower to higher trophic levels and thus is considered a pivotal species in the Arctic marine ecosystem, providing food for numerous species of seabirds, marine mammals, and fishes. A substantial total biomass of Arctic Cod is required for ecosystem maintenance.
- Arctic Cod have a circumpolar distribution and are considered seasonal migrants that use different habitats throughout the year (e.g., summer aggregations, under-ice habitat use).
- Continued reporting of both target and bycatch species from the At-Sea Observer program and vessel logbooks will be necessary to provide future Science Advice regarding any species caught in this fishery.
- Dedicated research is needed on Arctic Cod biology, ecology, population dynamics, and ecosystem modelling to better understand their pivotal role in the food web. This would also allow for greater certainty in biomass index estimates to improve Science Advice on sustainable bycatch removals by the shrimp fishery.

INTRODUCTION

The shrimp fishery in the Eastern Arctic operates in Shrimp Fishing Areas 1–3 (SFA 1–3; Figure 2). SFA 1 is located in Northwest Atlantic Fisheries Organization (NAFO) Division 0A within Baffin Bay and SFA 2 falls within Davis Strait. For the purpose of the shrimp stock assessment process, SFA 2 is analogous to the EAZ (Eastern Assessment Zone) (DFO 2017). SFA 3 is located in Hudson Strait and is analogous to the WAZ (Western Assessment Zone). The fishery in SFA 1 operates on a calendar year basis (i.e., January 1 to December 31). Historically, SFAs 2 and 3 were also managed on a calendar year basis, however, a new policy was adopted in 2003 that implemented a fiscal year management cycle (i.e., April 1 to March 31).




Figure 2. Location of Shrimp Fishing Areas (SFAs) used until 2013 and Western and Eastern Assessment Zones used since 2013.

Since 2017, shrimp fishing vessels operating in Shrimp Fishing Areas 1 (Baffin Bay) and 3 (Hudson Strait), have occasionally reported large amounts (400–2,300 kg per tow) of Arctic Cod bycatch. In Eastern Canada, Arctic Cod is listed as a groundfish species (Atlantic Fishery Regulations 1985), which means that elevated amounts of Arctic Cod bycatch can trigger a standard "move away" provision, as per the Northern and Striped Shrimp Conditions of Licence, which states: "*In the event that the total incidental catch of all groundfish species in any set exceeds the greater of 2.5% or 100 kg total weight, the licence holder/operator must immediately change the vessel's fishing area by a minimum of 10 nautical miles from any coordinate during the last tow*".

High occurrences of Arctic Cod bycatch, while not common, pose a significant burden on the fishing industry and have unknown ecological consequences. With limited time to harvest the allotted shrimp catch each year due to the relatively short open-water period and long transit

times from southern ports of origin, the fishing industry in the Eastern Arctic has asked DFO Resource Management for increased flexibility in Arctic Cod bycatch, with “move away” provisions triggered by the amount of Arctic Cod bycatch averaged over a number of consecutive tows rather than triggered by the amount of bycatch in one tow.

As a result, DFO Resource Management approached DFO Science seeking Science Advice on ecologically and biologically-responsible Arctic Cod removal levels in individual Shrimp Fishing Areas (SFAs). Given the absence of a comprehensive Arctic Cod stock assessment, the purpose of this process is to discuss ecologically and biologically acceptable levels for Arctic Cod removals in SFA 1, EAZ, and WAZ and whether it is possible to identify a “cap’’ for the total annual bycatch level of Arctic Cod for the three areas.

Arctic Cod Biology

Arctic Cod is a relatively small fish (mean size: 140 mm, maximum size: 400 mm; Matley et al. 2013) with a maximum life span of seven years. Arctic Cod spawns in winter (Hop et al. 1995) and the larvae hatch between April and June depending on geographical location (Bouchard and Fortier 2011). Larval Arctic Cod feed initially on smaller prey items (e.g., *Copepoda nauplii*, Rotifers) and prey on larger items as they grow (e.g., *Calanus copepods*; Walkusz et al. 2011). Juvenile fish (~35 mm in length) feed on copepods. After reaching the size of descent (~50 mm; Falk-Petersen et al. 1986, Majewski et al. 2016), Arctic Cod adopts a demersal life and, in addition to *Calanus* copepods, starts feeding on amphipods and fishes, including cannibalism. Once associated with the bottom, Arctic Cod becomes part of the demersal community, along with Northern and Striped Shrimp, and is incidentally caught in the shrimp fishery.

ASSESSMENT AND ANALYSIS

Arctic Cod Historical Catches

All vessels fishing for shrimp in the Eastern Canadian Arctic are required, as a condition of their licence, to have an At-Sea Observer (ASO) present on board at all times (i.e., 100% observer coverage). Data on bycatch (e.g., species caught, total weight caught for each species, measurements from individual animals) are collected by dedicated personnel (ASO) during shrimp fishing trips. Data collected by ASO are independent from the fishery records (logbooks) kept by the vessel’s captain and include information about both the target and bycatch species. Based on the data collected by ASO, Walkusz et al. (2020a) found a positive correlation (with the exception of SFA 2 where no correlation was established) between the amount of shrimp caught and the amount of Arctic Cod bycatch.

The fishery does not retain the incidentally caught Arctic Cod, thus, contrary to other fisheries in which bycatch is landed, the shrimp fishery returns the biomass to the sea. However, since the commercially executed fishing tows are 2–3 h in duration, the survival of released Arctic Cod is expected to be low.

Arctic Cod Population Biomass Estimates

Arctic Cod is considered a schooling fish (Hop et al. 1997), however, “aggregating fish” seems a more appropriate term considering the dense aggregations of different size/age individuals and non-unidirectional movement observed by Geoffroy et al. (2011, 2016; Figure 1). Arctic Cod is often dispersed throughout the water column and is therefore hard to properly quantify using any one type of directed sampling gear (either demersal or pelagic).

A biomass index of bottom-dwelling Arctic Cod was estimated using data from scientific surveys. Available data were combined from surveys targeting Northern and Striped Shrimp (200–600 m depth range) and groundfish, in particular Greenland Halibut (400–1,500 m depth range). The groundfish surveys conducted by DFO in Div. 0A (SFA 1) and Div. 0B (SFA 2) used an Alfredo III trawl with 20 mm cod-end mesh. Groundfish surveys are not conducted in SFA 3. Initial stock assessment (2006–2013) surveys for Northern and Striped Shrimp in SFA 3 were conducted by DFO with a Cosmos trawl with 12.8 mm cod end mesh. Since 2006 in SFA 2 and 2014 in SFA 3, surveys were designed by DFO and conducted by the Northern Shrimp Research Foundation (NSRF) using a Campelen 1800 trawl with 12.8 mm cod end mesh. All of the aforementioned surveys are strictly demersal and lack information on pelagic zone.

The results of these demersal surveys were used to quantify Arctic Cod biomass in SFAs 1–3, which resulted in an estimated biomass index that does not exceed 70,000 t in all SFAs combined (Walkusz et al. 2020b; Figures 3–5). This estimate likely represents a greatly underestimated index because of the scarcity of data, the patchiness of Arctic Cod occurrence, and the fact that only the demersal component of the ecosystem was sampled during the surveys.




Figure 3. Biomass index of bottom-dwelling Arctic Cod from the DFO Groundfish Survey in Div. 0A (SFA 1) for 2005–2017. Bars indicate standard deviation of the Arctic Cod biomass index. Note that the scale of Arctic Cod Biomass differs between SFAs (i.e., Figures 3–5).




Figure 4. Biomass index of bottom-dwelling Arctic Cod from the Northern Shrimp Research Foundation (NSRF) survey in SFA 2 for 2005–2017. Bars indicate standard deviation of the Arctic Cod biomass index. Note that the scale of Arctic Cod Biomass differs between SFAs (i.e., Figures 3–5).




Figure 5. Biomass index of bottom-dwelling Arctic Cod in SFA 3 from the NSRF survey for 2005–2017. Bars indicate standard deviation of the Arctic cod biomass index. Note that the scale of Arctic Cod Biomass differs between SFAs (i.e., Figures 3–5).

This underestimation most likely stems from the fact that Arctic Cod is a pelago-benthic fish and an unknown portion of its population may reside in the water column during the survey. Since Arctic Cod tend to aggregate, the surveys may not detect the aggregations because a given survey is a ''snap-shot'' of fish biomass at a specific time as opposed to an active tracking approach. Finally, Arctic Cod migrations are poorly understood in the Canadian Eastern Arctic, which can lead to surveys missing large cohorts of fish at certain times of the year.

By using recent estimates for fish, bird, seal, and whale populations in Baffin Bay/Davis Strait, Walkusz et al. (2020b) found that the Arctic Cod biomass requirement for ecosystem maintenance is > 500,000 t Arctic Cod annually. While this approach most likely produced a conservative estimate of the Arctic Cod biomass required to sustain its predators, it provides a general estimate of the order of magnitude of the Arctic Cod biomass required by the ecosystem.

Sources of Uncertainty

Overall lack of knowledge of Arctic Cod biology in the areas of interest.

While the general biology of Arctic Cod has been studied extensively in various Arctic locations, its biology is poorly understood in the geographic areas considered in this review. The lack of information on the timing of descent of juvenile Arctic Cod from pelagic to benthic habitats (i.e., when they become susceptible to fishing mortality in the shrimp fishery) represents therefore a significant knowledge gap. Multi-seasonal sampling along with length analyses of individuals caught would provide an insight into the ontogenetic timing of vertical migrations of Arctic Cod.

Insufficient data collection for biomass index analysis.

In this analysis, the biomass index was based solely on the benthic catches; however, Arctic Cod is considered a pelago-benthic species. Consequently, the lack of information on the pelagic portion of the population likely leads to significant underestimates of the Arctic Cod biomass. Application of acoustic methods (e.g., echosounder) along with pelagic trawl sampling would address this knowledge gap.

Insufficient data on distribution and abundance of Arctic Cod predators.

Analysis of predation pressure on Arctic Cod was conducted using published abundances of predators with the assumption that a portion of the predator population (i.e., 10%) would be present in the areas of interest. Using this assumption, the biomass of Arctic Cod required to sustain its predators was estimated. While this generalization provides some insight into Arctic Cod abundance, it is not a direct measure of fish biomass and the uncertainty associated with this approach is significant.

Potential observer data bias.

All offshore vessels operating in the areas of interest are obliged by the Conditions of Licence to have an At-Sea Observer (ASO) onboard. The data collected by the ASO is meant to be fishery independent; however, it can be expected that particular tows are not monitored due to the limitations of having only one ASO onboard, which may create an underestimation bias in the information reported.

CONCLUSIONS AND ADVICE

Considering the relatively small total fishing mortality of Arctic Cod stemming from the bycatch of the Northern shrimp fishery in the Eastern Canadian Arctic, i.e., less than 0.1% of the estimated Arctic Cod stock annually as estimated using predators' requirements, it can be assumed that the impact of the shrimp fishery on the Arctic Cod population is minimal in each of the three areas.

Consequently, bycatch mortality is unlikely to result in negative ecosystem impacts (i.e., predator requirements). However, given the high uncertainty in the Arctic Cod biomass index estimates, precaution is advised. Given the scarcity of data, Science cannot currently provide an ecologically and biologically acceptable Arctic Cod bycatch removal limit for the areas of interest.

Considering the methods used in shrimp fishery (e.g., length of the tow, characteristics of the fishing gear, processing of the catch), the probability of survival for incidentally caught Arctic Cod is low.

A dedicated sampling program to learn more about the species' biology and an acoustic survey to study the spatial and temporal distribution of Arctic Cod is strongly recommended to better inform future management decisions.

LIST OF MEETING PARTICIPANTS

All participants at this science peer review meeting are expected to participate as objective and knowledgeable individuals on the subject matter under review; not advocates or representatives of any interest group.

Name	Organization/Affiliation
Eva Enders (Chair)	DFO – Science, Central and Arctic Region
Joclyn Paulic	DFO – Science, Central and Arctic Region
Shelia Atchison	DFO – Science, Central and Arctic Region
Sheri Friesen	DFO – Resource Management, Central and Arctic Region
Hannah Murphy	DFO – Science, Newfoundland and Labrador Region
David Deslauriers	DFO – Science, Central and Arctic Region
Susan Thompson	DFO – Science, National Capital Region
Robert Moshenko	Nunavik Marine Region Wildlife Board
Amber Giles	Nunavut Wildlife Management Board
Leigh Edgar	DFO – Resource Management, National Capital Region
Ross Tallman	DFO – Science, Central and Arctic Region
Wojciech Walkusz	DFO – Science, Central and Arctic Region

SOURCES OF INFORMATION

This Science Advisory Report is from the July 3–4, 2019 meeting on Advice on Arctic Cod (*Boreogadus saida*) Bycatch Limits in Northern Shrimp Fisheries in the Canadian Arctic. Additional publications from this meeting will be posted on the Fisheries and Oceans Canada (DFO) Science Advisory Schedule as they become available.

Bouchard, C., and Fortier, L. 2011. Circum-arctic comparison of the hatching season of polar cod *Boreogadus saida*: a test of the freshwater winter refuge hypothesis. Prog. Oceanogr. 90: 105–116.

DFO. 2012. Guidance related to bycatch and discards in Canadian commercial fisheries. DFO Sci. Advis. Sec. Sci. Advis. Rep. 2012/022.

DFO. 2017. Assessment of Northern Shrimp, *Pandalus borealis*, and Striped Shrimp, *Pandalus montagui*, in the Eastern and Western Assessment Zones, February 2017. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2017/010. (Erratum: April 2017)

Falk-Petersen, I.B., Frivoll, V., Gulliksen, B., and Haug, T. 1986. Occurrence and size/age relations of Polar Cod, *Boreogadus saida* (Lepechin), in Spitsbergen coastal waters. Sarsia. 71: 235–245.

Geoffroy, M., Robert, D., Darnis, G., and Fortier, L. 2011. The aggregation of Polar Cod (*Boreogadus saida*) in the deep Atlantic layer of ice-covered Amundsen Gulf (Beaufort Sea) in winter. Polar Biol. 34: 1959–1971.

Geoffroy, M., Majewski, A., LeBlanc, M., Gauthier, S., Walkusz, W., Reist, J.D., and Fortier, L. 2016. Vertical segregation of age-0 and age-1+ Polar Cod (*Boreogadus saida*) over the annual cycle in the Canadian Beaufort Sea. Polar Biol. 39(6): 1023–1037.

Hop, H., Graham, M. and Wordeau, V.L. 1995. Spawning energetics of Arctic Cod (*Boreogadus saida*) in relation to seasonal development of the ovary and plasma sex steroid levels. Can. J. Fish. Aquat. Sci. 52: 541–550.

Hop, H., Welch, H., and Crawford, R. 1997. Population structure and feeding ecology of Arctic cod schools in the Canadian High Arctic. Am. Fish. Soc. Symp. 19: 68–80.

Majewski, A.R., Walkusz, W., Lynn, B.R., Atchison, S., Eert, J., and Reist, J.D. 2016. Distribution and diet of demersal Arctic Cod, *Boreogadus saida*, in relation to habitat characteristics in the Canadian Beaufort Sea. Polar Biol. 39: 1087–1098.

Matley, J.K., Fisk, A.T., and Dick, T.A. 2013. The foraging ecology of Arctic Cod (*Boreogadus saida*) during open water (July–August) in Allen Bay, Arctic Canada. Mar. Biol. 160: 2993–3004.

Walkusz, W., Paulic, J.E., Williams, W.J., Kwasniewski, S., and Papst, M.H. 2011. Distribution and diet of larval and juvenile Arctic Cod (*Boreogadus saida*) in the shallow Canadian Beaufort Sea. J. Mar. Syst. 84(3–4): 78–84.

Walkusz, W., Hedges, K., Atchison, S., and Deslauriers. D. 2020a. Arctic Cod (*Boreogadus saida*) bycatch in Shrimp Fishing Areas 1-3: 1979–2018. Can. Tech. Rep. Fish. Aquat. Sci. 3340: v + 16 p.

Walkusz, W., Atchison, S., Hedges, K., and Deslauriers, D. 2020b. Assessment of potential impacts of bycatch mortality on the Arctic Cod (*Boreogadus saida*) populations from the Northern (*Pandalus borealis*) and Striped (*Pandalus montagui*) Shrimp fisheries in Shrimp Fishing Areas 1, 2, and 3. DFO Can. Sci. Advis. Sec. Res. Doc. 2019/073. iv + 8 p.

THIS REPORT IS AVAILABLE FROM THE:

Center for Science Advice (CSA)
Central and Arctic Region
Fisheries and Oceans Canada
501 University Crescent
Winnipeg, Manitoba R3T 2N6

Telephone: 204-983-5131
E-Mail: xcna-csa-cas@dfo-mpo.gc.ca
Internet address: www.dfo-mpo.gc.ca/csas-sccs/

ISSN 1919-5087
© Her Majesty the Queen in Right of Canada, 2020



Correct Citation for this Publication:

DFO. 2020. Review of Arctic Cod (*Boreogadus saida*) bycatch in Canadian Arctic Shrimp Fisheries. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2020/007.

Aussi disponible en français :


MPO. 2020. Examen des prises accessoires de morue arctique (Boreogadus saida) dans les pêches à la crevette de l'Arctique canadien. Secr. can. de consult. sci. du MPO, Avis sci. 2020/007.

⊃ơϟᐅᐳᐟᑦ ᐅᐅᖏᒐ
ᓄᓇᕗᒻᒥ ᐅᒪᐟᓕᓂᓱᐟᐟᐊᐢᓄᑦ ᑲᑎᒪᔨᖏᑦ ᐊᖏᒪ ᓄᓇᕆᐢ ᐃᒪᖏᓂ ᐅᒪᐟᓂᑦ ᑲᑎᒪᔨᑦ
ᐅᔪᖏᒐ

ᑐᑭᓯᐊᕈᑎᑦ: **ᐃᓱᒪᓕᐅᕐᓂᖅ:** X **ᐊᑐᓕᕐᓄᐅᐳᐟᖅ:**

ᐱᕐᐟᑎᖏᒐ: **ᐋᖅᑕᐅᓂᖏᑦ ᐃᖅᓗᑦᑕᐅᑭᐊᖏᑦᑐᑦ ᐃᖅᓗᑦᑕᐅᐳᐟᑦ ᓇᓗᓇᐃᖅᑕᐅᓂᖏᑦ ᐅᑯᓄᖏᒐ**
***Pandalus borealis* ᓄᓇᕗᒻᒥ ᐊᖏᒪ ᓄᓇᕆᐢ ᑲᓇᑕᖏᒐ**

ᓄᓇᖏᔪᐊᖅ:
ᑐᖏᐟᖅᑐᐃᑦ – ᑲᓇᑕᖏᒐᓂ ᖃᐅᔨᓴᖅᑕᐅᓂᕐᒧᑦ ᑭᓚᓂᖏᒐ
ᑐᖏᐟᕈᐟᑦᑐᐃᑦ – ᐱᖏᓕᓇᖏᒐᓂ ᖃᐅᔨᓴᖅᑕᐅᓂᕐᒧᑦ ᑭᓚᓂᖏᒐ





ᐅᑭᐅᑦᑐᑎ ᑭᖏᐟᐸᐃᑦ (*Pandalus borealis*)

ᓇᓗᓇᐃᔨᖅᓯᒪᓂᖅᓯᑦ

ᒪᖁᑉᐳᒃ ᑭᒧᐸᐊᐸᑦ (*Pandalus montagui* ᐊᖏᒪ *Pandalus borealis*) ᓴᖅᑭᖅᑕᖅᑐᑦ ᐅᑭᐅᑕᖅᑐᒥ ᑭᒧᐸᐊᓂ ᐃᓕᓴᕐᓂᖅᑎᑦ ᑭᒧᐸᐊᓂᐊᖅᑕᖅᑐᑦ ᖃᑭᖅᑕᒍᓕᐸᓪᓚ ᓄᓇᖓᒃ ᐃᑭᕐᓴᖓᓂ ᐊᖏᒪ ᑲᓇᓂᖓᓂ ᓯᓚᐊᓗᒃ, ᐱᖅᓯᐅᑉᐸᔨᖅ ᐃᓕᖓᓂᓂ ᓄᓇᒪᒍ ᓇᔨᖅᑕᐅᔨᒋᑦ (NSA) ᐊᖏᒪ ᓄᓇᕆᐃᒃ ᐃᒪᖓ ᐊᕆᑦᑐᖅᓯᒪᔨᖅ (NMR). ᑲᑎᑦᑐᒋᑦ ᐃᖅᓗᑦᑕᐅᔨᓇᖅᑐᑦ ᐊᑐᓂ ᐊᐳᑦᓯᖓᑦᑐᓄᑦ ᐊᖅᑭᖅᑭᓯᒪᔨᑦ ᒪᖁᑉᐸᓂᖅ ᖃᐅᔨᓴᖅᓯᒪᓂᖅ ᖃᐅᔨᓴᖅᓯᒪᔨᑦ ᑭᐅᔨᓴᖅᓯᒪᓂᖅ, ᐱᖓᓂᓇᖓᓂ ᖃᐅᔨᓴᖅᓯᒪᔨᑦ ᑭᐅᔨᓴᖅᓯᒪᓂᖅ (WAZ) ᐊᖏᒪ ᑲᓇᓇᖓᓂ ᖃᐅᔨᓴᖅᓯᒪᔨᑦ ᑭᐅᔨᖓ (EAZ) (ᑕᓄᔪᒃ ᓄᓇᓕᒫᖅ).

ᑭᐅᔨᖓ ᐊᐃᑦᑎᑎᔨᖅ ᓄᓇᕗᑦ / ᓄᓇᐃᒃ ᑲᓇᓇᖓ (NU/NK E), ᐊᖏᒪ ᖃᑭᖅᑕᒍᓕᐸᓪᓚ ᐃᓕᓴᕐᓂᖅᑎᑦ ᐱᖓᓇᓇᖓ (DSW) ᐊᐅᓪᑕᐅᔨᒋᑦ ᐅᓂᒃ EAZ ᐊᐅᓪᓂᕐᑦ ᓇᓗᓇᐃᑦᑎᐊᖅᓯᑦ NSA ᐊᖏᒪ NMR ᐃᒪᖓᑦ ᐃᒪᐃᓇᖅᒋᑦ ᖃᑭᖅᑕᒍᓕᐸᓪᓚ ᐃᓕᓴᕐᓂᖅᑎᑦ (ᑕᓄᔪᒃ ᓄᓇᕗᑦ). ᓄᓇᕗᑦ ᐊᖏᒪ ᓄᓇᐃᒃ ᑭᖓᐳᓪᓕ ᐃᓪᓕᐅᕙᒃᑲᑕᐅᔨᓪᓕ ᑕᒪᑲᓇᓂ ᐊᓂᕆᑭᖓᓇᓂ ᑭᓪᓕᖓᓂ. ᐱᔭᔪᑎᑦᓯᓚᑦ ᐃᖅᓯᑦᑕᐅᔨᑦ ᓄᓇᕗᓪᓕᑦ ᐊᖏᒪ ᓄᓇᐃᓇᐃᓪᓕᐅᕙᒃᖃᖅᑕᐅᔨᑦ ᑎᑎᕋᖅᑕᐅᔨᓕᔪᑦ ᐊᐅᓪᑕᐅᑎᐃᓕᐅᖅᑕᖅᑐᑦ ᒪᖁᑉᐸᓂᖅ ᐱᐅᔪᓐᓇᖅᑐᑦ ᐊᖏᒪ ᐃᓄᐃᑦᑐᓂᖅᓯᒪᔨᑦ ᐃᖅᓯᑦᑕᐅᓕᕆᔨᒋᑦᑐᓂᖅᓯᒪᔨᑦ ᐃᖅᓯᑦᑕᐅᔨᓴᖅᓯᒪᓂᖅᓯᑦ ᐊᐅᓪᑕᐅᑎᐃᓕᐅᖅᑕᖅᑐᑦ ᒪᖁᑉᐸᓂᖅ ᐱᐅᔪᓐᓇᖅᑐᑦ ᐃᖅᓯᑦᑕᐅᔨᓴᖅᓯᒪᓂᖅᓯᑦ ᐊᐅᓪᑕᐅᑎᐃᓕᐅᖅᑕ

<u>ᓇᓗᓇᐃᔨᖅᓯᒪᓂᖅᓄᑦ ᑐᓴᕋᐅᓪᓗᑦ</u>

ᐱᓕᕆᔨᐅᔾᔨᖅ ᖃᓄᖅᑐᖅ ᑲᑎᒪᔨᓂᑦ:

1) ᐃᓱᒪᓕᐅᓂᖏᑦ ᐊᓯᔾᔨᖅᑕᐅᓗᓂ ᓇᓗᓇᐃᖅᑕᐅᓂᖅᓯᓐᓄᑦ Borealis ᐃᖅᓗᑦᑕᐅᑎᐊᖅᑎᑦᑐᓂᑦ ᐃᖅᓗᑦᑕᐅᔾᔨᓂᑦ ᑐᕆᖅᑕᐅᔾᔨᓄᑦ ᐊᓯᔾᔨᖅᑎᑦ ᐅᕙᓂ NU/NK E.

ᑎᑎᕋᖅᑐᖅ: ᐃᖅᓗᓕᕆᓂᕐᒧᑦ ᐊᐅᓚᑦᑎᓂᕐᒧᑦ, ᐃᒪᖏᑎᐅᑕᓕᕆᔨᒃᑯᑦ ᑲᓇᑕᒥ

ᐅᑉᓗᖓ: ᕿᑎᐳᐊᓗ 7, 2020

<u>ᐃᓚᒋᓕᐅᑎᓯᒪᔪᑦ</u>

ᐃᓚᒍᑦᖓ 1- ᑐᓴᖅᑎᑦᑎᓂᕐᒧᑦ ᓇᓗᓇᐃᔨᖅᓯᒪᓂᖅᓄᑦ (ᑲᑎᓄᐊᓗ 2020)

ᑐᓴᖅᑎᓂᒃᓯᒪᓂ ᓇᓗᓇᐃᔨᖅᓱᒪᓂᕐᓂᑦ (ᑲᓇᓂᐅᕐᓇ 2020)

ᐱᓕᕆᔨᐃᓐᓇᖅ ᐊᐅᓚᓪᓕᑎᓕᒪᓕᐅᖅᑐᖅ ᖃᕋᓴᐅᓴᖅᓯᑦ ᐃᖃᓗᓕᕆᓚᖃᖅᑕᖅᑐᓂᑦ ᐃᑲᔪᕐᓂᖏᓐᓂᓂᑦ ᐊᓱᐦᐸᓗᑎᑎ ᓇᓗᓇᐃᖅᑕᐅᓂᖏᑦ ᐃᖃᓗᓪᓕᐅᕐᓂᖏᑦᑐᓂᑦ ᐃᖃᓗᓪᓕᐅᔪᓂᑦ ᑐᕽᖅᑕᐅᔪᓄᑦ ᐊᐱᕆᖏᑦ ᐅᓂᒃᓂ NU/NK E.

ᐊᖏᕐᖅᑐᐃᓐᓇᐅᓕᐅᖅᑐᑦ ᐃᑲᔪᓱᖅᑐᑎᑎ ᑕᕐᓱᓕᒥᖓ ᐊᓱᐦᐸᓗᑎᑎ, ᑭᐅᐹᔪᕆᑦ ᐱᓴᐅᓪᓗᑎᑎ ᐅᑯᓇᖓᑦ ᑎᒍᒥᐊᖅᑎᐅᖃᕆᐊᓂᑦ:

- ᐱᑕ ᑭᐹ – ᒪᑭᕕᓴ ᑯᐊᐅᓱᓴᓐ
- ᐅᕋᐃᔨᓐ ᐅᓐᖅ, ᖃᒍᐅᑭᐅᑕᖅᑐᒥᑦ ᑲᑐᔪᖃᑎᒋᔭᖏᑦ (ᑭᓴᑐᖅᑐᒥᑦ ᖁᑭᖅᑕᒪᓗᓴ ᑯᐊᐅᓱᓴᓐ ᐊᓴᒪ ᖁᑭᖅᑕᒪᓗᒥᒥ ᐃᖃᓗᓕᕆᓚᖅᑎᓪᓗᑎᑦ ᑲᑐᔪᖃᑎᒋᔭᖏᑦ)

ⵊⵓⵖⵔⵜ ⵔⵓⵣⵍ
ⵙⵙⵣⵏⵔ ⵔⵍⵢⵛⵒⵒⵢⵄⵄⵏⵔ ⵃⵎⵍⵒⵛⵔ
ⵔⵊⵍⵍ

ⵊⵔⵢⵔⵄⵔⵚⵎⵔ: **ⵄⵢⵍⵛⵔⵔⵓⵏ:** X **ⵄⵊⵛⵏⵢⵏⵔⵔⵏ:** X

ⵑⵒⵢⵎⵏⵍ: **ⵃⵎⵔⵊⵔⵔ ⵄⵚⵒⵊⵔⵛⵔⵔⵔⵒⵛⵚⵊ ⵏⵢⵏⵣⵢⵒⵛⵛⵔ ⵔⵔⵔⵛⵚⵊⵔ (*Pandalus borealis*) 2020 ⵢⵚⵊⵏⵣⵛⵄⵛⵏⵔ ⵕⵚⵊⵏⵣⵛⵄⵏⵔⵔⵔ 0**



ⵔⵕⵔⵛⵚⵊⵔ ⵢⵚⵊⵏⵣⵚ (*Pandalus borealis*)

ⵒⵒⵄⵚⵒⵒⵛⵛ

ⵄⵚⵒⵒⵛⵛⵔⵔⵓⵛⵔⵔⵔ ⵔⵕⵔⵛⵚⵊⵔ ⵕⵚⵊⵏⵣⵒⵔ (*Pandalus borealis*) ⵑⵛⵚⵚⵊⵚ ⵕⵚⵊⵏⵣⵛⵄⵏⵔⵔⵔⵔⵛ (SFA) 0, ⵚⵕⵕⵚⵛⵒⵒⵏ ⵏⵢⵢⵛⵊⵄⵒⵒ ⵄⵕⵕⵏⵚⵚⵛⵒⵛⵊⵚ. SFA 0 ⵄⵒⵛⵕⵏⵚⵚⵛⵛⵔⵔⵔⵏⵛⵏⵚ ⵏⵏⵍ ⵄⵒⵛⵚ ⵄⵒⵒⵏⵢⵛⵚⵊ ⵒⵒⵄ ⵄⵛⵕⵏⵚⵚⵛⵛ ⵒⵒⵄⵔ ⵄⵛⵕⵏⵚⵚⵛⵛ (NSA) (ⵛⵢⵊⵖ ⵒⵒⵄⵚⵊⵏⵚ ⵑⵚⵕⵢⵔⵔⵏⵢⵒⵚ 1).

ⵑⵒⵢⵎⵏⵛⵒⵒⵊ ⵃⵊⵒⵛⵛⵔⵔⵓⵏⵚⵚⵚ, ⵄⵢⵒⵒⵛⵔⵔⵚⵛⵔⵔⵔ ⵏⵍⵍ ⵏⵊⵛⵒⵢⵛⵔⵔⵔ ⵃⵎⵊⵛⵛⵔ ⵄⵚⵒⵒⵛⵛⵔⵔⵔⵏⵛⵏⵚ (TAC) ⵏⵍⵍ ⵕⵚⵊⵏⵣⵛⵛⵔⵔⵊⵛⵛ ⵚⵢⵢⵛⵛⵔⵔⵔⵛⵛ Borealis ⵔⵛⵛⵔ SFA 0 ⵊⵒⵢⵚⵚⵊⵢⵛ ⵒⵒⵄⵔ ⵔⵊⵍⵢⵛⵔⵔⵓⵏⵢⵏⵣⵛⵄⵢⵛ ⵃⵎⵍⵒⵛⵛⵔⵔⵔ (NWMB). ⵒⵒⵄⵔⵍⵍ ⵔⵊⵍⵢⵛⵔⵔⵓⵏⵢⵏⵣⵛⵄⵢⵛ ⵃⵎⵍⵒⵛⵛⵔⵔⵔ ⵏⵊⵛⵛⵔⵔⵔⵛⵛ ⵏⵊⵛⵛⵔⵔⵔⵛⵛ ⵏⵊⵛⵛⵔⵔⵔⵛⵛ ⵄⵚⵒⵒ

ᖃᐅᔨᓴᖅᑎᓂᑦ ᑐᑭᓯᐊᕐᓗᒍᑦ

ᐱᒋᖏᓐᓂᕐᓗᓂ ᖃᐅᔨᓴᖅᑕᐅᓂᓂᑦ ᐅᑭᐅᑕᖅᑐᑐᒥ ᑭᖑᓪᓕᓂ ᖃᐅᔨᓴᖅᑕᐅᓯᒪᓂᓕᑦ ᐅᓂᒃ SFA 0 ᐃᖃᓗᓕᕆᔨᖃᑎᒋᔭᐅᓂᖅᓱᓕᓂ SFA 0 ᑭᖑᓪᓕᖅᑕᐅᔪᖅ 2010 ᑐᓴᓱᓂᖃᖅᑐᓂᑦ 2006 ᐊᒻᒪ 2008 (ᑲᓇᑕᒥ ᐅᖃᐅᓯᐅᕐᓂᑦ ᐅᓂᒃᑲᑦ 2010/024) (ᓇᓗᓇᐃᔨᖅᓯᒪᓂᖅᓂᑦ ᐃᓕᔪᑕᓴᓂ 2).

ᑕᒃᑯᐊ ᖃᐅᔨᓴᖅᓂᐅᔪᓂ ᓇᓗᓇᐃᖅᓯᓯᒪᔪᓂ ᑲᑎᓂᑐᑎᓂ ᐊᒥᓱᓂᖏᓂ ᓇᒃᓂᓂᖏᓂ ᐊᓂᑎᓂᑐᓂ. ᐃᖃᓗᓂᑕᐅᔪᓇᖅᑐᓂ ᑲᑎᓂᑐᑎᓂ ᐊᒥᓱᓂᖏᓂ ᓇᒃᓂᓂᖏᓂ ᐅᓇᐅᓕᐅᖅᑐᖅ 750 t (2006) ᐊᒻᒪ 1,100 t (2008) ᐊᖏᓇᐃᓂ ᓯᕿᐃᔨᖅᑐᑎᓂ ᐅᔨᓇᓂ 580 t (2006) ᐊᒻᒪ 800 t (2008). ᐊᑭᓐᑐᖃᐅᑎᔪᖅ TAC ᐅᔨᓇᓂ 500 t, ᐱᔨᐅᓇᐃᔭᑎᑐ, ᓴᖅᑭᑎᓂᑎᐸᑲᔨᖅᑐᖅ ᐊᑐᓂᕆᓂ ᖃᔪᓂᖏᓂ 40% (2008) ᑎᑭᓂᑐᒍ 70% (2006) ᑐᓇᓴᓂᖃᖅᑐᓂ ᖃᐅᔨᔨᐅᔪᓂ ᑲᑎᓂᑐᑎᓂ ᓇᒃᓂᓂᖏᓂ ᑕᐃᓵᓯᒪᓂᐅᔪᒥᓂ.

ᒪᖏᑭᐃᓇᐅᓵᑐᑎ ᖃᐅᔨᓴᖅᓂᐅᔪᓱᒃ, ᑲᑎᓂᑐᑎᓂ ᐊᓯᓵᓂᖏᓂ ᐊᓯᔨᖃᑕᓂᕆᓂᖏᓂ ᓇᓗᓇᐃᖅᑕᐅᔪᓇᖅᖏᓂᑐᑎᓂ ᐊᒻᒪ ᐃᓇᓇᐅᔪᒥᓂ ᑲᑎᓂᑐᑎᓂ ᐊᑎᓂᑐᑎᓂ ᖃᔪᓂᖏᓂ ᐊᒻᒪ ᐱᓯᒪᔨᐅᓂᖏᓂᓄᑦ ᓄᓇ ᓇᓗᓇᖅᑐᑎ. ᓇᒃᓂᓂᖏᓂ ᐃᓇᓇᐅᔪᒥᓂ ᐃᖃᓗᓐᓄᓯᕐᓴᐃᐅᖏᓂᑐᖅ ᐊᒻᒪ ᑐᓴᖅᓴᐃᓂᓕᑦᑐᑎᓂ ᖃᐅᔨᓴᖅᐅᓯᒪᔪᓂ ᑲᑎᓂᑐᑎᓂ ᐊᑎᓂᑐᑎᓂ, ᓯᐅᓂᓴᓕᒥ ᐱᔪᒪᔨᐅᔪᓇᖅᑐᑎ ᐃᖃᓗᓐᓂᕆᓂᔭᒍ ᑭᓇᓂᖅᑐᑎ. ᐃᖃᓗᓐᓂᕆᖃᖅᑐᑎ ᐱᔪᒪᓂᖃᖅᓯᖅᑕ ᑕᒫᓂ, ᖃᐅᔨᓴᖅᑎᓂᑦ ᖃᐅᔨᓴᖅᑕᐅᓂᖅᓱᓕᓂ ᑐᓴᕐᓯᖏᓐᓂᖅᑐᑎ. <ᕐᓇᐅᑎᑕᖃᖅᖏᓂᑐᑎᓂ ᓯᐅᓂᓴᓕᒥ ᖃᐅᔨᓴᖅᓂᐅᔭᕐᓂᓂ ᒧᓇᕐᓂᑦ.

ᐊᑐᓕᖁᔨᐅᔪᖅ

ᐊᑐᓕᖁᔨᐅᔪᖅ ᑕᒃᑯᐊ TAC ᐅᐅᖓ SFA 0 ᐊᖏᕐᕕᒍᑦ ᐊᓯᐊᓄᖅᑕᐅᓗᓂᓂ 500 t 2020–ᒍᑦ ᐊᓯᔨᐅᖓᓕᓗᑎᒃ ᐱᔪᓐᓇᐅᑎᒃ.

ᓇᓗᓇᐃᔨᖅᓯᒪᓂᖏᓂ ᑐᓴᓯᖅᐅᓂᑦ

ᐱᓕᕆᓂᐅᔪᓂᖅ ᑐᓴᓯᕐᓴᖅᑐᖅ:

1) ᐃᓯᒪᓕᐅᕈᑎᓂᑦ ᐃᖃᓗᓕᑕᐅᔪᓇᖅᑐᓂᑦ ᖁᔭᓕᓯᔪᓂᖏᓂ ᐃᓗᐊᓂ NSA ᐊᒻᒪ ᐊᑐᓕᖁᔨᐅᔪᓕᑦ ᐅᔨᓄᖓ TAC ᐅᒍᖓ SFA 0.


ᑎᑎᕋᖅᑐᖅ: ᐃᖃᓗᓕᕆᓂᖅᓴᒍᑦ ᐊᐅᓚᓕᑎᓂᖅᓴᒍᑦ, ᐃᒪᖏᐅᑕᓕᕆᓂᔨᕐᔪᑦ ᑲᓇᑕᒥ

ᐅᕙᓗᖓ: ᐃᓕᓴᕐᕕᐊᓗ 7, 2020

ᐃᓚᔪᑕᖏᓂᑦ

ᐃᓚᔪᑕᖓ 1 - ᓄᓇᖓᔪᐊᖅ ᐃᖅᑲᐅᐅᑕᓄᑦ ᐃᖃᓗᓐᓄᑦ ᐊᒻᒪ ᑭᖑᓪᓕᓂᑦ ᐊᐅᓚᓕᑎᓂᑦᑐᑎᓂ ᐊᑦᑦᓚᑎᑲ ᑲᓇᑕᒥ (ᐱᖃᓕᐅᑎᓪᓗᓂ ᑭᖑᓪᓕᖅᑐᓂᑦᖃᐃᓕᐅᔪᖅ 0)

ᐃᓚᔪᑕᖓ 2 - ᓇᓗᓇᐃᔨᖅᓯᒪᓂᖏᓂ: ᖃᐅᔨᓴᖅᑕᐅᓂᖏᓂ ᐅᑭᐅᑕᖅᑐᑐᒥ ᑭᖑᓪᓕᖅᑐᐃᓂ (*Pandalus borealis*) ᐅᐅᖓ SFA 0, 2, 3 ᐊᒻᒪ ᖃᐅᔨᖅᑕᓪᓕᓂ ᑭᖑᓪᓕᖅᑐᐃᓂ (*Pandalus montagui*) ᐅᐅᖓ SFA 2, 3 ᐊᒻᒪ 4 ᐱᖓᓐᓇᖓᓂ 63°W (ᖃᐅᔨᓴᖅᑎᓂᑦ ᐅᖃᐅᓯᐅᑕᑭᓐᑦ ᐅᓂᒃᑲᑦ 2010/024)


Fisheries and Oceans Canada
Pêches et Océans Canada
Canada
GROUNDFISH AND SHRIMP ADMINISTRATIVE AREAS IN ATLANTIC CANADA
February 18, 2014
Produced by Legislation and Regulatory Affairs, Fisheries and Oceans Canada
Cartography by René Duplain, Integrated Oceans Management Branch, Fisheries and Oceans Canada
0 150 300 600 900 1,200
Kilometers
0 75 150 300 450 600
Nautical Miles
Projection: Miller Cylindrical
Legend
Administrative Areas
Shrimp Management Areas
Shrimp Sub-Management Unit
Northwest Atlantic Fisheries Organization (NAFO) Divisions
Inuit Settlement Areas and Marine Regions
Areas of Equal Use and Occupancy
Nunavut Settlement Area
Nunavik Marine Region
Labrador Settlement Area
Other
Communities
Canada Landmass
Landmass (Other)
Description: Solid green lines indicate where shrimp management areas differ from NAFO Divisions. Shrimp Fishing Area 4 through 7 are located West of the EEZ. Shrimp Fishing Area 6 strides NAFO Divisions 2J and 3K and Area 7 overlaps with NAFO Division 3L. Dashed green line indicates shrimp sub-management units.
This is not an authoritative map. It has been prepared for illustrative purposes only.
Devon Island
Qaanaaq
Pond Inlet
Baffin Bay
Baffin Island
Clyde River
Foxe Basin
NU
Qikiqtarjuaq
Pangnirtung
Iqaluit
Kimmirut
Salluit
Quaqtaq
Puvirnituq
Kangirsuk
Aupaluk
Inukjuak
Kuujjuaq
QC
Ungava Bay
NU/NKW
Davis Strait
GREENLAND
Uummannaq
Ilulissat
Aasiaat
Sisimiut
Maniitsoq
Nuuk
Paamiut
Qaqortoq
Nain
Makkovik
NL
Cartwright
Happy Valley - Goose Bay
St. Anthony
Gaspe
St. John's
0
0A
1A
1
1B
1C
1D
1E
1F
0B
2G
4
5
2H
2J
6
3K
4S
4R
3Pn
4T
4Vn
3Ps
7
3L
3M
3O
3N
4W
4Vs
4X
5Zw
5Ze
60°30'W
65°00'W
70°W
DSW
DSE
NAFO
80°W
76°W
72°W
68°W
64°W
60°W
56°W
52°W
48°W
78°N
74°N
70°N
66°N
62°N
58°N
54°N
50°N
46°N
42°N

ᖃᐅᔨᓴᖅᑎᓂᑦ ᐅᖃᐅᔨᔪᒥᐊᕐᒋᑦ ᐅᓂᒃᑲᑦ 2010/024

ᓇᓗᓇᐃᔾᖅᓯᒪᓂᖏᑦ: ᖃᐅᔨᔪᖅᑕᐅᓂᖏᑦ ᐅᑭᐅᑕᖅᑐᒥ ᑭᖑᔪᑦᐸᐃᑦ (*Pandalus borealis*) ᐅᕝᓂ SFA 0, 2, 3 ᐊᒻᒪ ᖃᐅᔨᖅᑕᓕᑦ ᑭᖑᔪᑦᐸᐃᑦ (*Pandalus montagui*) ᐅᕝᓂ SFA 2, 3 ᐊᒻᒪ 4 ᐱᖓᓱᓇᖓᓂ 63°W

SFA 0 – *ᐅᑭᐅᑕᖅᑐᒥ ᑭᖑᔪᑦᐸᐃᑦ (Pandalus borealis)*

ᐃᖃᓗᓯᓚᐅᓂᖅ

- ᐃᖃᓗᓯᓚᐅᓯᒪᖓᑦᑐᑦ ᐊᖏᔪᕿᑦᑐᖅᓯᓯᒪᔪᓂᑦ.

ᑲᑎᑦᑐᒥᑦ ᐊᒥᓱᓂᖓᑦ ᓇᒡᓂᖓᑦ

- ᑭᓱᓂᖓᑦ ᑐᑭᓯᒥᐊᑭᑎᑦ ᑐᓂᖓᓯᖃᖅᑐᑦ ᒪᖏᑉᐹᓂᓂᐅ ᖃᐅᔨᔪᖏᓂᐅᔪᓂᓂᐅ 2006 ᐊᒻᒪ 2008.
- ᐃᖃᓗᑦᑕᐅᔪᓇᖅᑐᑦ ᑲᑎᑦᑐᒥᑦ ᐊᒥᓱᓂᖓᑦ ᓇᒡᓂᖓᑦ ᐅᓇᐅᓪᓗᐅᖅᑐᖓᓄᑦ 750 t (2006) ᐊᒻᒪ 1,100 t (2008).
- ᐊᖏᓇᐃᑦ ᓯᖏᐃᔾᖅᑐᑦ ᐊᒥᓱᓂᖓᑦ ᑲᑎᑦᑐᒥᑦ ᐊᒥᓱᓂᖓᑦ ᓇᒡᓂᖓᑦ ᐅᓇᐅᓪᓗᐅᖅᑐᖓᓄᑦ 580 t (2006) ᐊᒻᒪ 800 t (2008).

ᓄᑦᑎᓂᑦ ᓇᓂᓱᓂᖅ

- ᓄᑦᑎᓂᑦ ᓇᓂᓱᓂᖅ ᓇᓗᓇᖅᑐᖓᓄᑦ

ᑐᖏᖏᑦᑕᓂᖓᑦ

- ᐊᑭᑦᑐᕿᐅᑎᔪᒥᑦ TAC 500 t ᓴᖅᑭᑎᑦᑎᔪᓇᖅᑐᖓᓄᑦ ᐊᑐᖅᑕᐅᓂᖓᑦᓄᓇᑦ ᖏᖏᐃᔾᖅᑐᑦ 40%-70% ᑐᓴᕐᓂᖅᖃᖅᑐᑦ ᖃᐅᔨᓴᐅᓯᒪᔪᓂᑦ ᐊᑦᓴᑦᖓᑦᑐᑐᑎᓂᑦ ᐅᒪᔪᓂᑦ. ᐊᑦᑎᓂᖅᓴᒥᑦ TAC ᐊᑐᓕᖏᐅᔪᖅ.

ᒪᓇᐅᔪᒥᑦ ᑕᑯᓇᖅᑕᐅᔪᑦ ᐊᒻᒪ ᓯᐅᓂᓴᒥ ᓴᖅᑭᑕᓇᑐᐃᓇᐊᓇᓕᓕᑦ

- ᓯᐅᓂᓴᒥᑦ ᓴᖅᑭᑕᓇᑐᐃᓇᐊᓇᓕᓕᑦ ᐃᖃᓗᓯᓚᐅᓂᖏᑦ ᑭᓕᓚᖃᖅᑎᖅᑕᐅᔪᓂᑦ.
- ᐸᖏᓇᐅᑎᑕᖃᖅᑎᑕᖅᑐᑦ ᓯᐅᓂᓴᒥᑦ ᖃᐅᔨᔪᖏᓂᐅᔪᓂᑦ ᑕᒪᒪᓂᑦ.

ᑐᓂᔨᐅᔪᓂ ᐅᑭᖏᑦ
ᓄᓇᕗᒻᒥ ᐅᒪᔪᓕᕆᓂᖅᓴᖅᑐᑎᓂ ᑲᑎᒪᔨᖏᑦᑕ
ᐃᓕᓴ 2020

ᐅᔨᖏᑦ

ᑐᑭᓯᐅᑦᑕᕐᓂᑦ: X **ᐃᓱᒪᓕᐅᕙᓂᖅ**:

ᐱᔪᓀᖏᑦ: ᐃᖃᓗᓕᑐᑎᐊᖅ ᐃᖃᓗᐃᓕ

ᓕᐊᑦ ᕼᐃᐅᒪᔭ – ᖃᐅᔨᓴᕙᓂᖅ
ᑕᐃᓗ ᖃᓄᐅ ᐊᒻᒪ ᑕᐃᓕᓄ ᑐᓴᓱᐅ – ᓄᓇᒋᖏᓐᓄᑦᑐᓂ ᐊᐅᓕᕆᓂᕙᓂᖅ

ᐃᖃᓗᓕᒫᓂᕐᒧᑦ ᐊᐅᓕᕆᓂᓂᕐᒧᑦ ᑐᑭᓯᐅᑦᑕᕐᓂᑦ:

- ᑭᓇᐅᖏᓕᕆᔨᖅᓂᕐᒧᑦ ᐃᖃᓗᓕᒫᓂᕐᒧᑦ ᓴᖅᑭᓕᐅᕙᓪᑐᑦ 5 ᓇᔨᖅᑕᐅᔨᓂᑦ [ᐃᖃᓗᓕᑐᒃ, ᕿᓕᕐᔨᐊᖅ (Surrey), ᕼᐊᓕᒃᓘᒃ (Thirty-Mile), ᖃᐃd, ᐊᒻᒪ ᓄᓴᓕᖅ (Byron Bay) ᑐᖅᑭᑦ]. ᒪᓕᓕᑐᒍ ᐊᐅᓕᕆᓂᕐᒧᑦ ᐊᑐᖅᑕᐅᔨᑦ ᓇᓗᓇᐃᖃᖅᓭᒪᔨᑦ ᐃᓕᓕᐅᑎᓭᒪᔨᓂ ᐃᖃᓗᓕᒫᓂᕐᒧᑦ ᐊᐅᓕᕆᓂᕐᒧᑦ ᕿᓇᐅᑦ (ᐊᖏᑭᖅᑕᐅᔨᖅ 2014), ᖃᐅᔨᑕᒪᑦ ᑐᓴᖅᑎᑭᖅᑕᕐᓂᖅ ᐊᒻᒪ ᑯᑕᖏᓐ ᐊᐅᓕᑕᐅᓂᖏᓐ ᐱᓕᕆᐊᖏᔨᑦ ᖁᑎᖅᓭᐅᑦ ᓂᖁᑦ ᐃᖃᓗᓐᓄᐊᖅᑐᖅᑎᑯᓚᒍ, ᐃᖃᓗᓕᒫᓘᒻᒥ ᖃᐅᔨᓴᕙᓂᖅ ᐱᓕᕆᓂᕐᒧᑦ ᓄᓇᕗᑦᑎᔨᑦ ᐃᖃᓗᓕᒫᓂᕐᒧᑦ ᑐᐊᖏᓐᓄᑦᑐᓂ ᖃᐅᔨᓴᕙᓂᖅ ᑐᑭᓯᐅᑦᑕᕐᓂᑦ; ᐊᒻᒪ ᐃᖃᓗᓕᑕᒪᓕᕆᓂᑦ ᐃᖃᓗᓐᖏᑦᑕᕆᔨᓪᓚᑦ ᐃᖃᓗᓐᖏᑦᑕᐅᔨᓂ ᐊᑐᐃ ᐱᓇᓕᔨᖏᓪᓗᑎᑦ, ᐃᖃᓗᓐᖏᑦᑕᐅᑎᖏᓐᓄᑦᑐᓂ ᐃᖃᓗᓐᖏᑦᑕᐅᔨᓂ ᐊᒻᒪ ᐃᑎᑕᐅᔨᓂ ᑐᑭᓯᐅᑦᑕᕐᑎᓐᓄᑦᑐᓂ ᐊᑐᖅᑕᖅᑐᑎᓂ ᑎᑎᕋᖏᓐᓄᑦᑐᓂ.

2019 ᑭᓇᐅᖏᓕᕆᔨᖅᓂᕐᒧᓂᒃᑯᓂ ᐃᖃᓗᓐᖏᑦᑕᐅᔨᓂᑦ ᑭᓪᓕᓕᓂᐊᕐᓇᖅᑕᐅᓕᕆᔨᓂ

ᐃᖃᓗᓐᖏᑦᑕᕆᓂᖅ ᓇᒧᓐᓂᖏᓐ (ᐊᑎᐅᔪᓚᑦᑐᖅ)	ᑭᓇᐅᖏᓕᕆᔨᖅᓂᕐᒧᑦ ᑯᑕᖏᓐ ᑭᓇᐅᔭᖓ, ᖃᐅᓂᒃᖏᓐᓄᓇ ᐅᖃᓕᐃᓇᓐᓂᖏᓐ	2019 ᑭᓇᐅᖏᓕᕆᔨᖅᓂᕐᒧᑦ ᐃᖃᓗᓐᖏᑦᑕᐅᔨᓂ ᑭᓇᐅᔭᖓ, ᖃᐅᓂᒃᖏᓐᓄᓇ ᐅᖃᓕᐃᓇᓐᓂᖏᓐ	2019 % ᓄᓇᖃᖅᑕᐅᔨᓂ ᓴᓂᐊᓂ ᑯᑕᖏᓐ
ᐃᖃᓗᓇᒃ ᑯᖏᓐ (ᖁᐃᓕᓚᑕᓇ)	20,000	16,699	83
ᕼᐊᓗᓇᒃ (Thirty-Mile)	5,000	5,061	101
ᖃᐃd	17,000	12,481	73
ᕿᓕᕐᔨᐊᖅ (Surrey ᑯᖏᓐ)	9,100	8,884	98
ᓄᓴᓕᖅ ᑯᖏᓐ (Byron Bay	9,100 (*5,000)	5,061	101
ᑲᑎᓕᒪᖃᑦᑕᒥᑦ	**56,100**	**48,186**	**86**

*ᑐᑭᖃᖅᑕᐅᔨᖅ ᑯᑕᖏᓐ

- ᓄᓴᓕᖅ ᑯᖏᓐ ᒪᓇᐅᔨᔪᑎᑦ ᐃᖃᓗᓐᖏᑦᑕᕆᓂᖅ ᐊᑕᑎᓕᓕᑎᐊᖅᓭᒪᔨᓂᑦ, ᑐᑭᖃᖅᑕᐅᔨᓂᑦ, ᑭᓇᐅᖏᓕᕆᔨᖅᓂᕐᒧᑦ ᕿᓂᖅᓭᓕᐅᕙᓂᖅ 5,000 ᐅᓗᑭᓚᒃ. ᑕᓇᓐᓇ ᒪᓕᓇᓂᖅᑲᖅᑐᑦ ᐱᐅᖅᓭᐊᓂᕐᒧᑦ-ᑐᓇᓚᔪᓂ ᐊᐅᓕᕆᓂᕐᒧᑦ ᐊᑐᖅᑕᐅᔨᓂ ᑐᓴᓱᖏᓚᔪᓂ ᐃᖃᓗᓐᖏᑦᑕᕆᐊᖅ ᐃᖃᓗᓐᖏᑦᑕᐅᔨᑦ ᐃᓕᓕᐅᑎᓭᒪᔨᓂ

ᐃᖃᓗᓪᓚᕆᓂᖅ ᐊᐅᓪᓚᑎᓂᖅ <ᖃᓄᐅᑦ ᐱᓪᓚᓴᑦ, ᐊᒻᒪ ᐃᑲᔪᖃᕐᓂᖅᓯᒪᔪᓂ ᓄᓇᕗᒻᒥ ᐅᒪᓴᓪᖁᔭᐅᑉᑯᓂ ᑲᑎᒪᔨᖏᓐᓄᑦ ᓱᓂ 6 2017 ᐃᓱᒪᓪᓕᕐᓂᑎᓪᓗᑎᑦ ᑎᑎᕐᓂᖅᓱᒪᔪᓂ ᐃᓄᐃᑦ ᑲᑎᒪᓂᑎᓪᓗᒍ ᐱᔪᓐᓇᖅᑐᖅᖃᖅᑐᑦ ᐃᓕᓴᕐᓂᕐᕕᐊᓂᓗᒍ ᑯᑕᖏᑦ ᓴᓐᓴᓄ ᐃᓕᓴᕐᓂᐃᓕᓴᕐᓂ. ᐃᖃᓗᓯᑕᐅᕙᓪᓕᐊᓂᖅ ᖃᐅᔨᓴᖃᓂᖅ ᐊᒻᒪ ᑐᓴᖅᑎᑎᓂᓪᓗᑎᑦ ᐃᓂᓕᓴᐅᓐᖃᖅ, ᐃᑲᔪᖃᖅᑕᐅᔪᖅ ᐱᓪᓚᓴᔪᓂ ᑲᔪᖏᖅᖃᓕᓂᖏᓪᓕ ᑲᑐᑎᓂᖅᑐᑦ ᐱᐅᓂᖅᑐᑦ ᐊᒻᒪ ᐱᐅᓴᒥᑦ ᐊᐅᑦᑕᐅᕙᓪᓕᐊᓂᖅ ᐊᐅᑦᑕᐅᕙᓪᓕᐊᓂᓗᒍ.

- ᖃᑉᑎᖏᑦ ᓂᖅᐳᑦ ᑎᑎᕐᓯᒪᓕᒥᑦ ᐊᒥᑎᓪᓚᕆᑎᓂᕐᓯᓚᔪᓂ ᑐᖏᖅᑕᖏᓐᓄᑦ ᐃᓕᓴᖅᑕᐅᕙᓪᓕᐊᓂᓗᒍ ᐃᑲᔪᐃᑦ ᐊᒻᒪ ᓴᐃᑯ ᐱᔨᔭᑎᑎᓪᓗᒍ ᖃᑯᖓᓕᕐᓂᖅᑕᐅᓂᖏᓐᓇ ᐃᓕᓴᕐᓂᕐᕕᔪᓐᓇᖅᑐᑦ ᓴᓐᓴᓄ ᐃᓕᓴᓐᓂ, ᓱᖃᓪᓚᖅᑐᓂᖅ ᐅᖓᓇᑕᐅᕙᓪᓕᐊᓂᖅ ᐃᓕᓴᕐᓂᕐᕕᔪᓐᓇᖅᑐᑦ ᐃᓕᓴᖅᑕᐅᕙᓪᓕᐊᓂᓗᒍ ᐃᓕᓴᕐᓂᕐᕕᔪᓐᓇᖅᑐᑦ ᐅᖓᓇᑕᐅᕙᓪᓕᐊᓂᓗᒍ.

- ᐃᓕᓴᖅᑐᑎᓂᕐᒪᐊᒥ IFMP (ᐃᓚᒋᑕᐅᑎᓯᓚᕐᓂ ᐃᓕᓴᖅᑕᐅᓂᓪᓗᑎᑦ ᐊᐅᑦᑕᐅᓂᓪᓗᑎᑦ <ᖃᓄᐅᑦ) ᐊᑐᑕᖅᑎᑕᐅᒋᐅᑕᖅᑐᖅ 2014 ᑭᓇᐅᔨᑕᐅᕙᓯᓚᖅᑐᓄᑦ ᐃᓕᓴᖅᑕᐅᕙᔪᓂᑦ ᐃᓕᓴᖅᑐᓂᑦ, ᓱᑲᒋᑕᖅᐸᖓᔪᖅ ᑑᓴᔨᓄᖓ ᐃᓕᓴᖅᑐᓄᑦ ᑲᓇᑕᒥ. ᐃᓚᒋᑕᐅᑎᓯᓚᕐᓂᑦ ᐃᓕᓴᖅᑕᐅᓂᓪᓗᑎᑦ ᐊᐅᑦᑕᐅᓂᓪᓗᑎᑦ <ᖃᓄᐅᑦ ᖃᑎᒌᖃᐅᔨᔪᓐᓇᖅᑐᑦ ᐊᒻᒪ ᐅᒋᔭᒋᔪᖓᔨᓕᕐᓂᖅᑐᑦ ᐃᓕᓴᖅᑕᐅᓂᓪᓗᑎᑦ ᐊᐅᑦᑕᐅᓂᓪᓗᑎᑦ ᐊᒻᒪ ᐱᓪᓚᕆᐊᖃᓪᓚᕆᐊᖅᑐᓂᑦ ᐊᒻᒪ ᓄᑦᓯ ᑐᑭᓯᐅᑎᓂᑦ ᐃᓕᓴᖅᑕᐅᓂᓪᓗᑎᑦ ᐊᐅᑦᑕᐅᓂᓪᓗᑎᑦ <ᖃᓄᐅᑦ ᐱᓪᓚᕆᔨᓂ ᖃᑎᒌᖃᐅᔨᔪᓐᓇᖅᑐᑦ ᐃᓚᒋᑕᐅᑎᓯᓚᕐᓂᑦ ᐊᐅᑦᑕᐅᓂᓪᓗᑎᑦ <ᖃᓄᐅᑎᐅᑕᐅᕙᓪᓕᐊᓂᖅᑐᑦ, ᐊᒻᒪ ᑐᓂᓴᐅᓂᐊᕐᓂᑦ ᓄᓇᕗᒻᒥ ᐅᒪᓴᓪᖁᔭᐅᑉᑯᓂ ᖃᑎᒌᖃᐅᔨᔪᓐᓇᖅᑐᑦ ᐊᐅᑦᑕᐅᓂᓪᓗᑎᑦ ᐊᐅᑦᑕᐅᓂᓪᓗᑎᑦ <ᖃᓄᐅᑎᐅᑕᐅᕙᓪᓕᐊᓂᖅᑐᑦ, ᐊᒻᒪ ᑐᓂᓴᐅᓂᐊᕐᓂᑦ ᓄᓇᕗᒻᒥ ᐅᒪᓴᓪᖁᔭᐅᑉᑯᓂ ᖃᑎᒌᖃᐅᔨᔪᓐᓇᖅᑐᑦ ᐊᒻᒪ ᐊᖏᕐᓂᖅᑕᐅᕙᓪᓕᐊᓂᖅ 2020 ᑭᓇᐅᔨᑕᐅᓕᕐᓂᓪᓗᑎᑦ ᐃᓕᓴᖅᑕᐅᓂᓪᓗᑎᑦ.

ᖃᐅᔨᓴᐃᓂᖅ:

- ᐃᓕᓴᖅᑐᑎᓂᕐᒪᐊᒥ ᑭᓇᐅᔨᑕᐅᓕᕐᓂᓪᓗᑎᑦ ᐃᓕᓴᖅᑕᐅᓚᐅᖅ ᖃᐅᔨᓴᓂᓪᓗᑎᑦ ᐱᑕᓚᐊᖅ (ᐃᓕᓴᖅᑕᐅᓚᓪᓗᑎᑦ ᑐᓴᖏᓚᕆᖅᖃᖅᑐᑦ ᖃᐅᔨᓴᓂᓪᓗᒥᑦ) ᑲᔪᓯᓂᖅᑎᐊᖅᓂᕐᒥᓚᐅᖅᓯᔪᖅ ᐱᑕᖅᖃᖅᑎᑎᓪᓗᒍ ᐃᒪᓕᕐᒥᑕᕐᓚᓪᓕᐊᖅᑯᓂ ᑲᑎᒪᓂᖅᑕᐅᓚᐅᖅ 200 ᐃᓕᖅᑐᓂᓂ ᐊᑐᓂ ᑭᓇᐅᔨᑕᐅᓕᕐᓂᓪᓗᑎᑦ ᐃᓕᖅᑐᖓᔪᐅᔪᓂᓂ ᐃᒪᖏᓐᓂᓂ.

- ᐃᒪᖏᐅᑕᕐᓚᓪᓕᐊᖅᑯᓂ ᑲᓇᑕᒥ (DFO) ᑲᔪᓯᓂᖅᑎᑎᐊᓕᐅᖅᑐᖅᑕᑦ ᐃᓕᓴᖅᑕᐅᓚᓂᓪᓗᑎᑦ ᐃᒪᒥᖅᑯᓂ ᖃᐅᔨᓴᓂᖅᑎᑦ 200 ᐃᓕᖅᑐᓂᓂ ᓴᓐᓴᓄ ᐃᓕᓴᖅᑐᓂᓂ ᐊᑐᓂ ᑭᓇᐅᔨᑕᐅᓕᕐᓂᓪᓗᑎᑦ ᐃᓕᓴᖅᑐᖓᔪᐅᔪᓂᓂ ᐃᒪᓐᓇᓂᓂ. ᖃᐅᔨᓴᖃᖅᑐᐅᔪᓂ ᑐᑭᓯᐅᑎᓂᑦ ᐱᓪᓚᕆᐊᖃᓪᓚᕆᐊᖅᑐᑦ ᓄᓇᕗᒻᒥ ᐊᐅᑦᑕᐅᓂᓪᓗᑎᑦ ᐃᒪᒥᖅᑯᓂ ᐱᔨᔭᑎᑎᓪᓗᒍ ᐊᐅᑦᑕᐅᓂᓪᓗᑎᑦ ᐃᓕᓴᖅᑐᓂᓂ ᐊᑐᓂ ᑭᓇᐅᔨᑕᐅᓕᕐᓂᓪᓗᑎᑦ ᐃᓕᓴᖅᑐᓂᓂ ᐊᑐᓂ ᐊᐅᑦᑕᐅᓂᓪᓗᑎᑦ ᐃᓕᓴᖅᑐᓂᓂ

- 30 ᐃᖃᓗᐃᑦ ᓇᓗᓇᐃᔭᓕᔪᖅᑕᐅᓕᐅᖅᑐᑦ ᓂᐊᓕᓐ ᑯᓘᓂᒃ 2019.

ᓄᓇᓕᒻᒥ ᐱᖃᑕᐅᑎᑕᑎᓂᖅ (ᖃᐅᔨᓴᖅᓂᖅ/ᓄᓇᒋᓐᓇᖅᑐᓂᑦ ᐊᐅᓚᑕᑎᓂᖅ):

- ᐊᒥᓱᓚᓕᐃᑦ ᐳᓚᕆᔅᓂᐅᔪᑦ/ᑲᑎᒪᓂᐅᔪᑦ ᐃᖃᓗᑕᑐᑎᐊᖅ ᐅᒪᔪᕆᓂᐊᑎᒃᑯᑦ ᑲᑎᒪᔨᖏᓕᓗ ᐊᐅᔨᐅᑎᓕᓗᒍ (ᔨᑦᓂ-ᐊᖏᑭᓱ 2019).
- ᐱᔨᑦᖃᓱᒪᔪᑦ ᓇᓗᓇᐃᔨᖃᓱᒪᔪᑦ ᐊᖏᒪ ᖃᑭᒥᖅᑭᐅᓂᖏᑦ ᖃᑲᐅᔨᕆᖃᑎᒌᓂ ᖃᑲᐅᔨᕆᓂᕐᒧᑦ ᐱᓕᓚᐊᑦ ᑐᓄᖅᑐᐅᓂᖅᑐᑦ ᖃᖅᑎᒥᐅᓂᖅ ᑲᑎᒪᔨᐅᑎᓂᒃᑯᑦ ᐅᒪᔪᓕᓚᔨᐅᑦᑐᑦ ᑲᑎᒪᔨᐅᑎᓂᒃᑯᑦ ᐊᐅᓚᑕᑎᓂᖅ ᐱᓕᓚᔨᐅᑎᓂᒃᑯᑦ ᑲᑎᒪᓕᕆᔨᐊᑎᒃᓴᓂᑦ (ᓱᑎᐱᓚ 2019).
- ᐱᔨᑦᖃᓱᒪᔪᑦ ᓇᓗᓇᐃᔨᖃᓱᒪᔪᑦ ᐊᖏᒪ ᖃᑭᒥᖅᑭᐅᓂᖏᑦ ᖃᑲᐅᔨᕆᖃᑎᒌᓂ ᖃᑲᐅᔨᕆᓂᕐᒧᑦ ᐱᓕᓚᔨᐅᑎᓂᒃᑯᑦ ᐊᖏᒪ ᐃᖃᓗᓕᓚᓂᕐᒧᑦ ᐊᐅᓚᑕᑎᓂᕐᒧᑦ ᐱᓕᓚᔨᐅᑎᓂᖅ ᑐᓄᖅᑐᐅᓂᖅᑐᑦ ᐃᖃᓗᑕᑐᑎᐊᖅ ᐅᒪᔪᓕᓚᔨᐅᑦᑐᑦ ᐃᖃᓗᓕᓚᓂᕐᒧᑦ ᐊᐅᓚᑕᑎᓂᒃᑯᑦ ᐃᖃᓗᓇᓕᕆᔨᐊᑎᒃᓴᓂᑦ ᑲᑎᒪᓕᕆᔨᐊᑎᒃᑯᑦ ᐅᒪᔪᓕᓚᔨᐅᑦᑐᑦ (ᐅᓐᓄᓚ 2020).
- ᐃᖃᓗᓇᓱᓐᓇᐅᑕᐅᖅᑎᓐᓇᒍ ᐱᖃᑕᐅᑎᑎᑐᓂᕐᒧᑦ ᑲᑎᒪᕆᑎᖃᓂᖅ ᐃᖃᓗᑕᑐᑎᐊᖅ ᐅᒪᔪᓕᓚᔨᐅᑦᑐᑦ ᑲᑎᒪᔪᓐᓇᖅᑐᓂ, ᖃᑭᑎᕆᑦᓂᕐᒧᑦ ᓂᖅᑭᑦ, ᐊᖏᒪ ᑭᓇᐅᔨᑕᐅᑎᔪᓂᖅᓂᖅᑐᑦ ᐃᖃᓗᓇᓱᓂᑲᑦ ᐊᖏᒪ ᐃᒪᓕᕆᑕᑕᓚᔨᐅᑦᑐᑦ ᑲᑎᒪᔪᓐᓇᖅᑐᓂ

⊃σ⁴⊳⊲ᐟᑦ ⊳◈ᖏᓚ
⌀⌀◈⁴Γ ⊳L⊲ᐟ⊂ᓱᔪᖏ⊲⁰ᑯᐟ ⁰ᑎLᔪᖏᑦ
ᒧᖏ 2020

⊳ᒧᖏᓚ

⊃Pᔪ⊂⊲P⌂ᖏ: X **ᐃᔪL⊂⊳ᖏσᖏ⁰:**

ᐱ⊲ᐟᑎᖏᓚ: ᐃLᖏΓ⊳⊂⊂ᓱᔪ⁰ᑯᐟ ⁰⌀ᓭΓ – ⊃Pᔪ⊂⊲P⌂ᖏ

⊃Pᔪ⊂⊲P⌂ᖏ:

ᐃᖏ⁰⌀ᓱᐃ⊲ᖏ⌂ᖏ:

- ⊲ᑯ⁰ᑲ <ᐃᔪ⁰ ᐃ⊂Γ⊂⊳⊲⊂⊂⊳ᖏ⊃ᖏ ᐃLᖏΓ⊳⊂⊂⊂ᓱᔪ⁰ᑯᐟ⁰⊙ᐟ (DFO) ⊳P⊳⊂ᖏ⊃Γ ⊲⊳⊂σᖏᑎ⁰⊙ᐟ ⊙⌂ᓴᐱᓱΓ 2019 ⊲ᖏᑎᔪᖏᑲᖏᒧᑦ⊃σ ᐃᖏ⊃ᓱᑎσᖏᒧᑦ ⊲⊳⊂ᔪᑎσᖏᒧᑦ ⊳Lᔪσᖏᑦ ᖏ⊳⊳ᔪᖏᖏᑎ.

ᐃLᖏ⊳⊂ᐃᐟᑦ ⊳Lᔪᐃᐃᑦ:

1) ⊃ᑲᑢᓪᖏ:

- ᑦ⁰⊙⊲ᓱΓ 2020, DFO ᖏ⊳⊳ᔪᖏ⌂ᑦ ⊳ᑎᖏᑎᖏ⊂⊳⊂⊃ᖏ⊃ᑦ ⌀⌀◈⁴Γ ⊳L⊲ᔪ⊂ᓱᔪ⁰ᑯᐟ⁰⊙ᐟ ⁰ᑎLᔪᖏᑦ⁰σᖏᑦ (NWMB), ⌀⌀◈ᑦ ⊃⁰ᖏᑲᐃ⁰ ᑎΓᖏᑢᑦ (NTI), ᖇPPᖏᑲᓪᔪΓ ⊳Lᔪᔪᑦᓱᔪ⁰ᑯᐟ ⁰ᑎLᔪᖏᑦ (QWB), ⊲ᓭL ⊲ᔪ⊃ᖏᑎ⊳⊲ᔪᑦ ⊳Lᔪᔪσᐃᑎ⁰ᑯᐟ ⁰⊃ᑦᔪᖏᑎ⌂ᑦ⁰ᑦ (HTOs) ◈ᐃσᔪᐃᑢᓭ ⊳ᖏ⊳⊳ᔪᖏᖏᑎ⊃ᑎ ᔪᓫᔪᒪσᖏᑲ ⊃ᔪᔪᑲᖏᒧᓱᖏ 2020 ⁰⌀⁰ᓫᖏᑲᔭ ᖇPPᖏᑲᓪᔭ⁰, Admiralty Inlet, and Eclipse Sound ⊃ᑲᑢᓪσᑦ ᖏᖏᑢᑎᔪᒪ⁰ᑯᐟ ᖏ⊳⊳ᔪᖏσᑎᒥᑦ.
- ⊃ᑲᑢᓪᑎ⊳⊲ᔪσᑦ σ⊿ᖏᑢᑲᑦ ⊳ᑎᖏᑎ⊂⊳⊲ᔪ ⊲ᓭL ⊃ᑲᑢᓪᑎ⊳⊲ᔪ ⊲ᒥᔪᐟσᖏᑦ⊂ ⊃⊙ᐟᑦ⊃σᖏᑦ ⊃⊳⊲ᑦᑎ⊲ᑲᓪᑦ ᐃᒪᔪᖏᑦσᑎᒧᑦ⊃ᑦ ⁰⌀⊃ᔪᖏᑎ⌂ᑦ⁰ᑦ⁰⊙ᐟ ᐃLᖏΓ⊳⊂⊂ᓱᔪ⁰ᑯᐟ⁰⊙ᐟ ᖇPᓪᔭᑢᑢᖏᔪᐟ⁰⊿⊳⊂⊳ᖏᑎᑦᔭᑢᑦ, ᒪᖏ 31, 2020. ⊃ᑲᑢᓪᑎ⊳⊲ᔪ ⊃Pᔪ⊂⊲P⌂ᑎᖏᑦ Harvest data will be reconciled ᔪᐅσ⊲ᒧᑦ 2020 ᖇPᓪᔭᑢᑢᖏᑎ⊳⊲ᔪᒧᑦσ⊿ᖏᑢᑲᑦ ᐱᔪ⊳⊲ᔪ ⊲ᓭL ⊲ᖏᑦᒧᒧᒪΓ ⊲ᔪ⊿ᓴᑲᖏᑦ ⊲ᑦᖇᑦᑦ⊳σ⊲ᖏᑲᑦ.

2) ⊲ᐃ⊿ᑦ:

- Γσᔭ⊂ᒧᑦ ᐃᔪL⊂⊳ᖏσᖏᑢ σᓱ⊳⊂ᔪ⊳⊲ᔪᖏ ⊿ᖏσ⊲ᓱ 14, 2020 ⌀⌀◈⁴Γ ⊳L⊲ᔪ⊂ᓱᔪ⁰ᑯᐟ⁰⊙ᐟ ⁰ᑎLᔪᖏᑦ ⊲ᖏᑦᖏ⊂⊳σᖏᑢ⊙ᐟ 76 ⊲ᐃ⊿ᑦ ⊲ᐃᖏᖏᑢᔪ⊃ᐃ⁰ᓫσᖏᒧᑦ 2020-ᒧᑦ.
- ᐃLᖏ⊳⊂⊂ᓱᔪ⁰ᑯᐟ ⊃σᔪᔪᔪᑢ⊂⊃ᑦ ⊃ᔪᔪ⊳⊲ᔪΓᑲ ᐃᔪL⊂⊳σᖏᒧᑦ⁰ᑦ ⌀⌀◈⁴Γ ⊳L⊲ᔪ⊂ᓱᔪ⁰ᑯᐟ⁰⊙ᐟ ⁰ᑎLᔪᖏᑦ⁰⊙ᐟ⁰ᑲ 2020–Γᑦ ᔪᖏᑲᑎᖏᑢᑎᐃ⊃ ⁰ᑎᖏ⊃⊂ᑦ ⊳L⊲ᔪᖏ⊂⊳⊲ᔪ⁰ᓫᖏ⊃ᑦ (TAH) ᖏᔪᖏᔪ⁰σᑢᖏ⊃ᑦ ⊲ᐃᐃᖏᔪᒧᑦ⁰⊙ᐟ ⊲ᐃᐃᖏᔪᒧᑦ ⊲ᒥᔪᐟσᖏᑦ ⊲ᒥᔪᐟ⊂⊳⊲ᔪσᑦ ⌀⌀◈⁴Γ.
- ⌀⌀◈⁴Γ ⊲ᐃᐃᖏᔪᐟ⊂ᓱᔪ⁰ᑯᐟ ⁰ᑎLᔪσᑢᖏᔪ⁰⊙ᐟ ᔪᖏᓫᑦ⊳⊲ᔪᔪᖏᑲᑦ ᒪᔪᑦ 13, 2020 ᐃᖏᑦᓱ⁰σ ᖏᖏ⊳⊲ᔪᖏᑦᓱ⊃ᑎ ⊲ᐃ–ᖏᔪᑦ⊳⊲ᔪσᑦ ⊃ᔪᖏᑎᖏᑢᑎᖏᑦ⊂σᒧᑦ⊃ᑦ ⊲ᓭL ⌀⌀◈⁴Γᒧᑦ ⊲ᒥᔪᐟ⊂⊳⊲ᔪᑦ ᖏ⊳⊳ᔪᖏσᒧᑦ⊃ᑦ ᐱᔪᓫᑦᒪ⊲ᑦ.
- ᐱ⊂ᓫᖏᑎᖏᑢᑎᖏ⊃ᑎ ᖏσᖏᑲᑎᑦ ⊳L⊲ᔪσᐃᑎ⁰ᑯᐟ ⁰⊃ᑦᔪᖏᑎ⌂ᑦ⁰σ, ⌀⌀◈⁴Γᒧᑦ ⊲ᒥᔪᐟ⊂⊳⊲ᔪᑦ ⊳L⊲ᔪᖏ⊂⊳⊲ᔪσᑦ ᖏ⊳⊳ᔪᖏσᒧᑦ⊃ᑦ ᐱᔪᓫᑦᒪ⊲ᑦ ⊲ᐃᐃᖏᔪᒧᑦ ᖏσᖏᑲᑎᑦ σᓫᑦ⊳⊲ᔪᔪᖏᑲᑦ ᑲ⊿ᔪσ⊲ᒧᑦ⊃ᑦ 2020–Γᑦ.
- ᐱ⊂ᓫᖏᑎᖏᑢᑎᖏ⊃ᑎ ⊲ᐃᐃᖏᔪᐟ ⊳L⊲ᔪσᐃᑎ⁰ᑯᐟ ⁰⊃ᑦᔪᖏᑎ⌂ᑦ⁰σ, ⌀⌀◈⁴Γᒧᑦ ⊲ᒥᔪᐟ⊂⊳⊲ᔪᑦ ⊳L⊲ᔪᖏ⊂⊳⊲ᔪσᑦ ᖏ⊳⊳ᔪᖏσᒧᑦ⊃ᑦ ᐱᔪᓫᑦᒪ⊲ᑦ ⊲ᐃᐃᖏᔪᒧᑦ ᔪᑦᑦσ⊃ᑦ ᐱᒥ⊲ᖏᑎ⊂⊳σ⊲ᖏᑲᑦ⊃ᖏ 2020–Γᑦ.
- ᐃLᖏ⊳⊂⊂ᓱᔪ⁰ᑯᐟ ᐱ⊂⊳⊲ᔪ⊃ᑦ ⊲ᒥᔪᒪ⊂⁰σᑦ ⊃ᔪᖏᑎᖏᒧᑎσᑦ 2019 ⊲ᐃᖏᔪᒪ⁰ᑲ⊃ᐃ⁰ᓫσᒧᑦ⊃ᑦ ⊲ᒥᔪᐟ ᖏ⊳⊳ᔪᒥ⊲ᖏᔪᖏᖏᑢᑎᖏ⊃ᑦ ⊲ᒥᑦᑦ⊂⊳ᔪᔪσᑦ ᑦ⁰ᑯᐟ⊲ ᐱ⊃ᑦ ᐱᑦᖏᑢᔪ⊲ᖏᖏᑎᑦ⊳⊲ᔪ⊃ᑦ ᑦᑦσ ⊂⊿ᔪᖏᑲᔪᒪ⁰ᑯᐟ ᖏ⊳⊳ᔪᒥ⊲ᖏᔪᖏᖏᑢᑎᖏ⊃ᑦ ᖏᖏᑢᑎᖏᑦ⊂σᒧᑦ⊃ᑦ ᖏᑦᑦ⊂σᒧᑦ⊃ᑦ ᐃᒪᔪᑦᒪ⊲ᑦ ⊃ᔪᖏᑎᖏᑢᑎᖏᑦ⊂σᒧᑦ⊃ᑦ ᐱᔪᓫᑦᒪ⊲ᑦ ⊲ᒥᔪᐟ⊂⊳⊲ᔪᑦ ⊳L⊲ᔪᖏ⊂⊳⊲ᔪσᑦ ᖏ⊳⊳ᔪᖏσᒧᑦ⊃ᑦ ᐱᔪᓫᑦᒪ⊲ᑦ ⊲ᐃᐃᖏᔪᒧᑦ ᔪᑦᑦσ⊃ᑦ ᐱᒥ⊲ᖏᑎ⊂⊳σ⊲ᖏᑲᑦ⊃ᖏ 2020–Γᑦ.
- ᐃLᖏ⊳⊂⊂ᓱᔪ⁰ᑯᐟ ᐱ⊂⊳⊲ᔪ⊃ᑦ ⊲ᒥᔪᒪ⊂⁰σᑦ ⊃ᔪᖏᑎᖏᒧᑎσᑦ 2019 ⊲ᐃᖏᔪᒪ⁰ᑲ⊃ᐃ⁰ᓫσᒧᑦ⊃ᑦ ⊲ᒥᔪᐟ ᖏ⊳⊳ᔪᒪ⊲ᖏᔪᖏᖏᑢᑎᖏ⊃ᑦ ⊲ᒥᑦᑦ⊂⊳ᔪᔪσᑦ ᑦ⁰ᑯᐟ⊲ ᐱ⊃ᑦ ᐱᑦᖏᑢᔪ⊲ᖏᖏᑎᑦ⊳⊲ᔪ⊃ᑦ ⊲ᒥᑦᑦ⊂σᒧᑦ⊃ᑦ ᐱᔪᓫᑦᒪ⊲ᑦ ⊃ᔪᖏᑎᖏᑢᑎᖏᑦ⊂σᒧᑦ⊃ᑦ ⊲ᒥᔪᐟ⊂⊳⊲ᔪᑦ ⊳L⊲ᔪᖏ⊂⊳⊲ᔪσᑦ ᖏ⊳⊳ᔪᖏσᒧᑦ⊃ᑦ ᐱᔪᓫᑦᒪ⊲ᑦ ⊲ᐃᐃᖏᔪᒧᑦ ᔪᑦᑦσ⊃ᑦ ᐱᒥ⊲ᖏᑎ⊂⊳σ⊲ᖏᑲᑦ⊃ᖏ 2020–Γᑦ.
- ᐃLᖏ⊳⊂⊂ᓱᔪ⁰ᑯᐟ ᐱ⊂⊳⊲ᔪ⊃ᑦ ⊲ᒥᔪᒪ⊂⁰σᑦ ⊃ᔪᖏᑎᖏᒧᑎσᑦ 2019 ⊲ᐃᖏᔪᒪ⁰ᑲ⊃ᐃ⁰ᓫσᒧᑦ⊃ᑦ ⊲ᒥᔪᐟ ᖏ⊳⊳ᔪᒪ⊲ᖏᔪᖏᖏᑢᑎᖏ⊃ᑦ ⊲ᒥᑦᑦ⊂⊳ᔪᔪσᑦ ᑦ⁰ᑯᐟ⊲ ᐱ⊃ᑦ ᐱᑦᖏᑢᔪ⊲ᖏᖏᑎᑦ⊳⊲ᔪ⊃ᑦ ⊲ᒥᑦᑦ⊂σᒧᑦ⊃ᑦ ᐱᔪᓫᑦᒪ⊲ᑦ ⊃ᔪᖏᑎᖏᑢᑎᖏᑦ⊂σᒧᑦ⊃ᑦ ⊲ᒥᔪᐟ⊂⊳⊲ᔪᑦ ⊳L⊲ᔪᖏ⊂⊳⊲ᔪσᑦ ᖏ⊳⊳ᔪᖏσᒧᑦ⊃ᑦ ᐱᔪᓫᑦᒪ⊲ᑦ ⊲ᐃᐃᖏᔪᒧᑦ ᔪᑦᑦσ⊃ᑦ ᐱᒥ⊲ᖏᑎ⊂⊳σ⊲ᖏᑲᑦ⊃ᖏ 2020–Γᑦ.
- ᐃLᖏ⊳⊂⊂ᓱᔪ⁰ᑯᐟ ᐱ⊂⊳⊲ᔪ⊃ᑦ ⊲ᒥᔪᒪ⊂⁰σᑦ ⊃ᔪᖏᑎᖏᒧᑎσᑦ 2019 ⊲ᐃᖏᔪᒪ⁰ᑲ⊃ᐃ⁰ᓫσᒧᑦ⊃ᑦ ⊲ᒥᔪᐟ ᖏ⊳⊳ᔪᒪ⊲ᖏᔪᖏᖏᑢᑎᖏ⊃ᑦ ⊲ᒥᑦᑦ⊂⊳ᔪᔪσᑦ ᑦ⁰ᑯᐟ⊲ ᐱ⊃ᑦ ᐱᑦᖏᑢᔪ⊲ᖏᖏᑎᑦ⊳⊲ᔪ⊃ᑦ ⊲ᒥᑦᑦ⊂σᒧᑦ⊃ᑦ ᐱᔪᓫᑦᒪ⊲ᑦ ⊃ᔪᖏᑎᖏᑢᑎᖏᑦ⊂σᒧᑦ⊃ᑦ ⊲ᒥᔪᐟ⊂⊳⊲ᔪᑦ ⊳L⊲ᔪᖏ⊂⊳⊲ᔪσᑦ ᖏ⊳⊳ᔪᖏσᒧᑦ⊃ᑦ ᐱᔪᓫᑦᒪ⊲ᑦ ⊲ᐃᐃᖏᔪᒧᑦ ᔪᑦᑦσ⊃ᑦ ᐱᒥ⊲ᖏᑎ⊂⊳σ⊲ᖏᑲᑦ⊃ᖏ 2020–Γᑦ.
- ᐃLᖏ⊳⊂⊂ᓱᔪ⁰ᑯᐟ ᐱ⊂⊳⊲ᔪ⊃ᑦ ⊲ᒥᔪᒪ⊂⁰σᑦ ⊃ᔪᖏᑎᖏᒧᑎσᑦ 2019 ⊲ᐃᖏᔪᒪ⁰ᑲ⊃ᐃ⁰ᓫσᒧᑦ⊃ᑦ ⊲ᒥᔪᐟ ᖏ⊳⊳ᔪᒪ⊲ᖏᔪᖏᖏᑢᑎᖏ⊃ᑦ ⊲ᒥᑦᑦ⊂⊳ᔪᔪσᑦ ᑦ⁰ᑯᐟ⊲ ᐱ⊃ᑦ ᐱᑦᖏᑢᔪ⊲ᖏᖏᑎᑦ⊳⊲ᔪ⊃ᑦ ⊲ᒥᑦᑦ⊂σᒧᑦ⊃ᑦ ᐱᔪᓫᑦᒪ⊲ᑦ ⊃ᔪᖏᑎᖏᑢᑎᖏᑦ⊂σᒧᑦ⊃ᑦ ⊲ᒥᔪᐟ⊂⊳⊲ᔪᑦ ⊳L⊲ᔪᖏ⊂⊳⊲ᔪσᑦ ᖏ⊳⊳ᔪᖏσᒧᑦ⊃ᑦ ᐱᔪᓫᑦᒪ⊲ᑦ ⊲ᐃᐃᖏᔪᒧᑦ ᔪᑦᑦσ⊃ᑦ ᐱᒥ⊲ᖏᑎ⊂⊳σ⊲ᖏᑲᑦ⊃ᖏ 2020–Γᑦ.
- ᐃLᖏ⊳⊂⊂ᓱᔪ⁰ᑯᐟ ᐱ⊂⊳⊲ᔪ⊃ᑦ ⊲ᒥᔪᒪ⊂⁰σᑦ ⊃ᔪᖏᑎᖏᒧᑎσᑦ 2019 ⊲ᐃᖏᔪᒪ⁰ᑲ⊃ᐃ⁰ᓫσᒧᑦ⊃ᑦ ⊲ᒥᔪᐟ ᖏ⊳⊳ᔪᒪ⊲ᖏᔪᖏᖏᑢᑎᖏ⊃ᑦ ⊲ᒥᑦᑦ⊂⊳ᔪᔪσᑦ ᑦ⁰ᑯᐟ⊲ ᐱ⊃ᑦ ᐱᑦᖏᑢᔪ⊲ᖏᖏᑎᑦ⊳⊲ᔪ⊃ᑦ ⊲ᒥᑦᑦ⊂σᒧᑦ⊃ᑦ ᐱᔪᓫᑦᒪ⊲ᑦ ⊃ᔪᖏᑎᖏᑢᑎᖏᑦ⊂σᒧᑦ⊃ᑦ ⊲ᒥᔪᐟ⊂⊳⊲ᔪᑦ ⊳L⊲ᔪᖏ⊂⊳⊲ᔪσᑦ ᖏ⊳⊳ᔪᖏσᒧᑦ⊃ᑦ ᐱᔪᓫᑦᒪ⊲ᑦ ⊲ᐃᐃᖏᔪᒧᑦ ᔪᑦᑦσ⊃ᑦ ᐱᒥ⊲ᖏᑎ⊂⊳σ⊲ᖏᑲᑦ⊃ᖏ 2020–Γᑦ.

3) ᑦ⁰σᑦ⊃ᑲᑦ ᐃPᖏᑢᑎᑕ ᖇPᓪᔭᑢᑢᖏᔪᐟ⊂⊳⊲ᔪ⊃ᑦ:

- ᖏᑎᔪᑎᖏᑦ 41 ᖇPᓪᔭᑢᑢᖏᑎ⊳⊲ᔪᑦ ⊃ᔪᖏᑎᖏᑢᑎᐃᒧᑎᑕ⊳⊲ᔪ⊃ᑦ ᑦ⁰σᑲᑎᑦ ⊲ᐅᑦᖏᑢᑢᖏᑦ⊂⊳⊲ᔪ⊃ᑦ ᑦ⁰σᑲᑎᑦ ⊲ᒥᔪᐟ⊂⊳⊲ᔪᑦ ⊳L⊲ᔪᖏ⊂⊳⊲ᔪσᑦ ᖏ⊳⊳ᔪᖏσᒧᑦ⊃ᑦ ᐱᔪᓫᑦᒪ⊲ᑦ ⊲ᒥᔪᐟ⊂⊳⊲ᔪᑦ ⊳L⊲ᔪᖏ⊂⊳⊲ᔪσᑦ ᖏ⊳⊳ᔪᖏσᒧᑦ⊃ᑦ ᐱᔪᓫᑦᒪ⊲ᑦ ⊲ᐃᐃᖏᔪᒧᑦ ᔪᑦᑦσ⊃ᑦ ᐱᒥ⊲ᖏᑎ⊂⊳σ⊲ᖏᑲᑦ⊃ᖏ 2019.
- ᒪᐃᖏᑎᔪᑎᖏᑦ 14, 2019, ᖏᑎᒪσᖏᑦ⊂⊳⊲ᔪ⊃ᑦ ᑦ⁰σᑦ⊃ᑲᑦ ᖇPᖏᑢᑎᖏᑦ⊂⊳⊲ᔪ⊃ᑦ⊂σᑦ ᑦ⁰σᑦ⊃ᑲᑦ ᖇPᖏᑢᑎᖏᑦ⊂⊳⊲ᔪ⊃ᑦ⊂σᑦ ᑦ⁰σᑦ⊃ᑲᑦ ᐱᔪᓫᑦᒪ⊲ᑦ ⊳L⊲ᔪᖏ⊂⊳⊲ᔪσᑦ ᖏ⊳⊳ᔪᖏσᒧᑦ⊃ᑦ

ᑲᑐᔪᖅᑎᑎᕐᓂᑦ, ᓄᓇᕗᑦ ᑐᓴᖅᓱᓚᐅᖅ ᑎᒥᖅᑎᑦ, ᓄᓇᕗᓴᒥ ᐅᒪᔨᓕᕆᔨᕐᕕᐊᖅᑯᑦ ᑲᑎᒪᔨᖅᓂᑦ, ᐃᒪᖃᐅᑕᓕᕆᔨᖅᑯᑦ ᐊᒥᒪ ᖃᐅᔨᖅᑕᐃᓗᒥ ᐅᒪᔨᓕᕆᔨᖅᑯᑦ ᑲᑎᒪᔨᖅᓂᑦ. ᐃᒪᖃᐅᑕᓕᕆᔨᖅᑯᑦ ᑐᓴᕐᕋᓕᐅᖅᑐᑦ <ᓴᓂᖅᑐᖅ ᐃᐱᖅᓯᓂ ᕿᑭᖅᑕᒃᓯᒪᔪᓂ ᐱᑎᐊᖅᑲᑎᑎᑦᓂᑦ ᐊᐅᑕᕐᓇᖅᑲᑎᑎᓂᖅᓂᑦ ᐱᓕᕆᖅᑲᑎᑎᑦᓂᑦ ᐊᐅᑕᕐᓇᖅᑲᑎᑎᑦᓂᑦ ᐱᑎᐊ

2) <ᓐᓂᖅᑐᖅ:

- 22,426 kg ᐃᖃᓗᐃᑦ ᑐᙵᖅᑎᑎᒍᑕᐅᓕᐅᖅᑐᑦ ᐃᖃᓗᓯᑕᐅᔪᓂ 2019 ᐊᐅᓚᑦᑯᓂ ᐃᖃᓗᓕᕆᓂᖅᓯᒪᓂ, 20,607 kg ᖃᐅᔨᓴᖅᓯᐅᔪᓐᓇᖅᑐᑦ ᐃᓕᖅᑎᐅᓂ ᐊᒻᒪ 1,819 kg ᑭᓇᐅᔭᕆᐅᕆᔨᓐᓂᐅᑯᓂ ᐃᓕᖅᑎᐅᓂ.
- <ᓐᓂᖅᑐᖅ ᐅᓪᓚᔪᓂᐊᑎᑯᓂ ᑲᑐᔨᓴᖅᑎᑦᑎᖅᑐᑦ ᐅᖃᖅᑐᔨᒋᐊᕈᐅᖅᑐᑦ ᐃᓕᖅᒥᐅᕈᓐᓇᖅᑐᓂ ᑐᓂᔪᖅᑕᖅᓂᐊᕈᓐᓇᖅᑐᑎᑎᑐᒃ ᓕᐃᓴᓘᔪᓂ 2019-20 ᐅᑭᐅᐅᖅᑯᓂ ᐃᖃᓗᓕᕆᓂᖅᓯᒪᓂ ᐱᑎᐊᖅᔪᒪᓪᓗᓂ ᑲᓐᓂᖅ y 21, 2020.

<u>ᐊᑯᑭᑐ ᖃᓕᓴᕐᓂᑦ (ᓇᓪᓕᐊᔪᓐ):</u>

1) <ᓐᓂᖅᑐᖅ ᐃᑭᖓᓂᓂ ᖃᓕᓴᕐᓇᓂ ᐃᖃᓗᑐᓇᓂᖅ:

- <ᓐᓂᖅᑐᖅ ᐅᓪᓚᔪᓂᐊᑎᑯᓂ ᑲᑐᔨᖃᑎᒌᓕᐅᖅᑐᑦ ᑭᓇᐅᔭᕆᐅᕆᔨᓐᓂᐅᓂᔪᓂ ᓕᐃᓴᓘᔪᓂ 2020 ᐅᑭᐅᐅᖅᑯᓂ ᖃᓕᓴᕐᓇᓂ ᐃᖃᓗᑐᓇᓂᖅᑎᑦ ᑐᓂᔭᐅᓗᓂ ᑲᓐᓂᖅ 21, 2020.
- ᐃᓕᖅᑎᐅᑕᕐᓇᓕᐅᖅᑯᓂ ᑲᑎᖃᑎᖃᓕᐅᖅᑐᑦ ᐱᔪᒪᓂᖃᖅᑐᖅᓂ <ᓐᓂᖅᑐᐃᑎ ᖃᓕᓴᕐᓇᓂᐅᐊᖅᑎᓂ ᑲᓐᓂᖅ 28, 2020 ᐅᖃᖅᔨᓴᖃᖅᑐᑎ ᑎᑎᕋᖅᓂᐊᖅ ᐊᒻᒪ ᓕᐃᓴᓘᔪᓂ ᐊᒻᒪ ᑭᐅᓗᑐ ᐃᖃᓗᓯᑎᑦ ᐊᐱᖅᑯᑎᕐᓂᓐᓂ.

2) 2019 ᐃᓚᐃᓐᓇᕆᑦ ᐃᖃᓗᑐᓇᓂᖅ:

- ᑲᑎᑦᑐᑦ 9,551 mt ᐃᖃᓗᑦᑕᐅᓕᐅᖅᑐᑦ ᐅᐊᓐᓇᖅᓯᒪᓂ ᐱᖅᓯᓐᓇᖅᓯᓂ ᐊᑦᓭᓐᑎᒃ ᐃᖃᓗᑐᓇᓂᕐᒧᑦ ᑲᑐᔨᖃᑎᒌᖅᓯᒪ (NAFO) Division 0A.
- ᑲᑎᑦᑐᑦ 4,944 mt ᐃᖃᓗᑦᑕᐅᓕᐅᖅᑐᑦ NAFO ᐊᖏᔪᑐᖅᓯᒪᓂᖅᓯᒪ 0B ᓄᓇᕗᒻᒥ ᓕᐃᓴᖅᓯᓂᓪᓗᓂ ᑎᒍᒥᐊᖅᑎᓄᑦ.

3) ᑲᓇᓐᓇᖅᓯᓂ ᐅᑭᐅᑕᖅᑐᒥ ᐃᖅᑲᒥᐅᑕᓂ ᐃᖃᓗᓐᓂᑦ ᐅᖃᖅᐅᔨᑦᑕᐊᖅᑎᑦ ᑲᑎᒪᔨᕿᓐᓂᑦ:

- ᐊᖏᕐᕈᒍᒪᓕᑦ ᑲᑎᒪᓂᖅᓯᒪ ᑲᓇᓐᓇᖅᓯᒪᓂ ᐅᑭᐅᑕᖅᑐᒥ ᐃᖅᑲᒥᐅᑕᓂ ᐃᖃᓗᓐᓂᑦ ᐅᖃᖅᐅᔨᑦᑕᐊᖅᑎᑦ ᑲᑎᒪᔨᕿᓐᓂᑦ (EAGSAC) ᐃᖃᓗᓐᓂᓄᓕᓐᓂᖅᑐᑦ ᐃᓕᓴᕐᓂᐊᕐᓇᖅ 25-26, 2020.

4) ᐃᓕᖏᑦ ᐅᒥᐊᖅᑐᖅᑎᐅᓂᖏᓪᓗᑦ ᑲᑎᒪᔨᑦ ᓇᓗᓇᐃᔭᑯᓂᖅᓯᒪᓂᖏᓪᓗᑦ:

- ᑎᓯᐱᓕᐅᑎᓪᓗᒍ 2019, SA0 ᑲᑎᕐᓂᓐᓂ ᐃᖃᓗᑕᓕᓂᖏᓪᓗᑦ ᐱᓕᐅᖅᑐᑦ ᐃᓕᖏᑦ ᐅᒥᐊᖅᑐᖅᑎᐅᓂᖏᓪᓗᑦ ᑲᑎᒪᔨᑦ (MSC) ᒪᑭᒪᑎᑎᓂᖏᓪᓗᑦ ᓇᓗᓇᐃᔭᑯᓂᑐᖃᓂᖅ.

<u>ᐅᑭᐅᑕᖅᑐᒥ ᑭᖑᓪᓕᐃᓕᐅᑦ:</u>

1) ᓄᓇᕗᒻᒥᑦ ᐃᖃᓗᑕᓕᓂᐅᔨᕐᓇᖅ ᑯᓚᑦᓪᓕᐅᑦ ᐃᖃᓗᔨᓯᓂᐅᐊᔨᓂᐅᑦ ᐊᑦᑐᐊᓂᒃᓴᑦ ᓄᓇᕗᒻᒥᑦ (2019/20):

- ᑲᑎᑦᑐᑦ 2.1 mt ᐃᖃᓗᑦᑕᐅᓕᐅᖅᑐᑦ ᖃᑭᑭᖅᑕᔭᓪᓕ ᐊᑯᑭᑦᑐᐃᓐᓗ ᓄᓇᖑᒃ ᐃᑭᕆᖑᓂ ᑲᓇᓐᓇᖅᓯᒪᓂ.
- ᑲᑎᑦᑐᑦ 1,619 mt ᐃᖃᓗᑦᑕᐅᓕᐅᖅᑐᑦ ᖃᑭᑭᖅᑕᔭᓪᓕ ᐊᑯᑭᑦᑐᐃᓐᓗ ᓄᓇᖑᒃ ᐃᑭᕆᖑᓂ ᐱᖅᓯᓐᓇᖅᑖᓂ.
- ᐃᖃᓗᔨᓯᑦᑐᕆᐅᕈᓐᓇᑦᑐᖅ ᑭᖑᓪᓕᐃᓕᐅᓂᐊᕆᔨᒪᔨᓕᕐᓂᑦ 1.
- ᑲᑎᑦᑐᑦ 34 mt ᐃᖃᓗᑦᑕᐅᓕᐅᖅᑐᑦ ᑲᑐᔪᓐᓇᖅᓯᒪᓂ ᖃᐅᔨᓴᖅᑕᐅᔪᓕᑦ ᑭᓪᓕᖓᓪᓗ.
- ᑲᑎᑦᑐᑦ 4,556 mt ᐃᖃᓗᑦᑕᐅᓕᐅᖅᑐᑦ ᐱᖅᓯᓐᓇᖅᓯᒪᓂᖃᐅᔨᓴᖅᑕᐅᔪᓕᑦ ᑭᓪᓕᖓᓪᓗᒍ.

2) ᐅᑭᐅᑕᖅᑐᒥ ᑭᖑᓪᓕᐃᓕᐅ ᐅᖃᖅᐅᔨᑦᑕᐊᖅᑎᑦ ᑲᑎᒪᔨᕿᓐᓂᑦ:

- ᐊᖏᕐᕈᒍᒪᓕᑦ ᑲᑎᒪᓂᖅᓯᒪ ᐅᑭᐅᑕᖅᑐᒥ ᑭᖑᓪᓕᐃᓕᐅ ᐅᖃᖅᐅᔨᑦᑕᐊᖅᑎᑦ ᑲᑎᒪᔨᕿᓐᓂᑦ (NSAC) ᑲᑎᒪᓕᓂᖅᑐᑦ ᐄᐊᓕᓪᓚᕆᒃ ᒥᓕᐅᖅᑐᖅ ᐊᐅᓚᑦᑯᓂ ᒪᓕᐅᖅᑐᖅ 4-5, 2020.

ᑎᑎᕋᖅᑐᖅ: ᐃᖃᓗᐃᑦ ᐊᑦᓕᒪᔨᓐᓇ – ᐃᓕᖏᑦᐅᑕᕐᓇᓕᐅᖅᑯᓂ ᑲᓇᑕᒥ

ᐅᓪᓗᓐᓇ: ᓇᖕᒪᐊᓇ 7, 2020

ᐃᖏᓕᕐᓂᖅ
ᐊᐅᓚᕐᑎᓯᐅᖃᕐᓂᖅᓯᒪᔪᑦ
ᑲᑎᒪᔨᕐᓂᑦ

IRNIURVIIT AREA
CO-MANAGEMENT
COMMITTEE

7 ᐃᑉᐊᓗ 2020

ᓄᓇᑉᓕᐹᑦ ᐅᒪᔨᓚᓗᐃᔨᐊᕐᒪᑦ ᑲᑎᒪᔨᖏᑦ
ᐱᖓᓱᐊᕐᓂᐅ ᖃᐅᔨᓂᐅ ᓇᑎᖏᑦ, ᐊᑦᓚᐃᒪᑦ ᐃᒃᓗᖃᒪᖓᓱᓂᐅ
ᑎᑎᕋᖅᓂᐊᖏᐃᓐ 1379
ᐃᖅᑲᓗᐃᑦ, ᓄᓇᑉᑦ X0A 0H0
ᕿᓚᑕᐅᔭᒪᑦ ᑎᑎᕋᖅᑕᐃᖓᓱᓄᑦ: tsataa@nwmb.com

ᓄᓇᑉᓕᐹᑦ ᐅᒪᔨᓚᓗᐃᔨᐊᕐᒪᑦ ᑲᑎᒪᓂᖅᓴᓇᓱᖓᓱᓂᐅᒃ ᑲᑎᒪᓂᖓᓱᓂ, ᓇᔭᐅᑎᓚᒃ RM 001-2020: ᐱᖓᓚᓚᒃᑭᒥᒃ ᑎᑎᕋᕐᓯᒪᓚᖅᑐᑦ ᐊᐅᓚᕐᑎᓂᕐᒪᑦ ᐸᖏᕐᓂᖅᑐᓂᐅᒃ ᐊᓪᒪᓗ ᐃᒃᑯᑐᐃᕐᒥᒃ ᑎᑎᕋᖅᑕᐃᖓᓱᓂᐅ, ᐅᒪᔨᓚᓗᐃᔨᐊᕐᒪᑦᓄᓇᓕᖅᑐᓂ ᐊᖓᕐᓂᖅᑐᓂᐅ

ᖃᒃᖏᓴᐅᖅᑐᕐᒥ (ᖃᒃᑯᓗᓇᑎᑐᑦ East Bay-Γ) ᐊᓪᒪᓗ ᐃᒃᑯᑐᐊᑦ (HΔᐅᓇ ᒋᐸᓴᓓ) ᑎᓐᓕᕐᖏᓴᐃᒃ ᓄᓇᑉᑦ ᓴᑭᑉᖏᓴᐅᖅᑐᓂ ᓴᓕᕐᓂᐅᒃᑐᓗᐊᓂᐅ ᐱᓇᓱᖏᑎᑎᐅᓯᐅᖏᓴᓕᓵᑦ ᑕᒫᓂ 1959-ᖓᒍᑉᑐᓗᒍ. ᒪᓕᑐᖏᑎᑎᐅᓂᐅᓕᕐᓂᓄᑦ ᑕᐃᑯᓇᓓᓄᑦ ᐊᐅᓚᕐᑎᓂᕐᒪᑦ ᐊᖓᕐᓂᑎᑎᒍᔪᑦ, *ᐃᓄᐃᑦ ᐊᑭᖏᑎᑎᐅᓂᐅᓕᕐᓂᓄᑦ ᐃᑲᔪᓴᖏᑕᒃᑭᕐᓂᓄᑦ ᐊᖓᕐᓂᖅᑐᓂᐅ ᐊᓪᒪᓗ ᐃᓄᐃᑦ ᓄᓇᖓᓄᑦ (ᐃᓄᐃᑦ ᐃᑲᔪᓴᖏᑕᒃᑭᕐᓂᓄᑦ ᐊᖓᕐᓂᖅᑐᓂᐅ) ᓴᑉᑕᓕᕐᓂᒥ ᓇᖓᑎᑕᐅᖏᓴᐅᑎᑎᐅᓯᐅᖏᓴᓕᓵᑦ 2006-ᖓᒍᑉᑐᓗᒍ>, ᐊᓪᒪᓗ ᐊᖓᕐᓂᑲᑎᒋᔪᒍᑕᐅᓂᐊᖏᑲᓂᓚᐊᓕᓕᐅᖏᓴᐅᓴᐅᑎᑦ ᑕᒫᓂ 2016-Γ. ᓄᓇᑉᑦ ᑐᓐᖓᕆᐃᖏᑦ. ᐱᖓᓱᓴᓕᓗ ᐊᐃᖏᑐᖏᓴᐅᓴᐅᑎᑦ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖃᑎᒋᑉᑐᓂ ᐊᓪᒪᓗ ᑲᓇᑕᒥ ᐊᐅᓚᕐᑎᓂᕐᒪᑦ ᐃᓄᐃᑦ ᐊᖓᕐᓂᖅᑐᓂᐅ (ᐊᐅᓚᕐᑎᓂᕐᒪᑦᓄᓇᓕᖅᑐᓂ ᓴᓕᕐᓂᐅᒃᑐᓗᐊᓂᐅ ᐊᓪᒪᓗ ᐃᓄᐃᑦ ᐃᑲᔪᓴᖏᑕᒃᑭᕐᓂᓄᑦ ᐊᖓᕐᓂᖅᑐᓂᐅ) ᓴᓇᓕᕐᓂᒥ ᓴᖅᑭᑐᓗᐊᓂᐅ (2006-ᖓᒍᑉᑐᓗᒍ>, ᐊᓪᒪᓗ ᐊᖓᕐᓂᑲᑎᒋᔪᒍᑕᐅᓂᐊᖏᑲᓂᓚᐊᓕᓕᐅᖏᓴᐅᓴᐅᑎᑦ ᑕᒫᓂ 2016-Γ. ᓄᓇᑉᑦ ᑐᓐᖓᕆᐃᖏᑦ. ᐱᖓᓱᓴᓕᓗ ᐊᐃᖏᑐᖏᓴᐅᓴᐅᑎᑦ ᐃᓄᐃᑦ ᐊᖓᕐᓂᖅᑐᓂᐅ) ᓴᓇᓕᕐᓂᒥ ᐊᐅᓚᕐᑎᓂᕐᒪᑦ ᐃᓄᐃᑦ ᐊᖓᕐᓂᖅᑐᓂᐅ ᐊᓪᒪᓗ ᐃᓄᐃᑦ ᓄᓇᖓᓄᑦ (ᐃᓄᐃᑦ ᐃᑲᔪᓴᖏᑕᒃᑭᕐᓂᓄᑦ ᐊᖓᕐᓂᖅᑐᓂᐅ ᐊᓪᒪᓗ ᐃᓄᐃᑦ ᐃᑲᔪᓴᖏᑕᒃᑭᕐᓂᓄ



ᐅᒪᔨᓪᓚᔪᕐᐊᖅᑯᑦ Δᓗᒪᓪᐅᔪᖅᑎᐅᕈᓐᓂᒋᑦ ᓴᐳᵒσᐊᖅᑕᐅᕐᐃᓴᓂσᖅᑕᐅᓗᒪᔨᓄᑦ ᑐᖏᓐᒐᔨσᖅ ᐊᐅᓪᓗᑎσᓕᒧᑦ <ᕐᓇᐅᑎᓪᓚᐊᓚᐅᓗᒪᕈᑐᐃσᖅ (*Δᓄσᐃᑦ Δᑲᔨᓗᐊᓴᖏᵒᓄᑦ ᐊᖏᑭᑎᑦ Δᓗᓪᓂᒋᑦ Lᓪᖅᑕᐅᑦᓗᑎᖅ 3.6.1*).

ᑎᑎᕆᖅᑐᖅ,,



ᓄᐊ ᑲᑦᓪᖅ
Δᖅᓯᖇᐅᑕᖅ, Δᓕσᐅᔪᕐᐃᖅᑯᑦ ᐊᐅᓪᓗᑎᓚᐅᔪᖅᑲᑎᑎᒋσᓕᒧᑦ ᑲᑎᒪᓂᕆᓪᑦᓯᖓᵒᓄᑦ

ᓇᖅᓯᐅᑎᓴᐅᓯᒪᓗᑎᖅ Lᑯσᓂᑲ:

1. ᑎᑎᕆᖅᓂᑦ “ᕆᖅᕈᓴᐅᖅᑐᑎᒥ ᑎᓂᒥᐊᕆᖅᕐᓇᓂᒥᖅ ᐊᐅᓪᓗᑎσᓕᒧᑦ <ᕐᓇᐅᑲᑦ: ᑭᖓᔪᓪᓕᒥ ᑎᑎᕆᖅᓯᒪᓂᖅᓂᑦ, ᖃᓄᐊᓗ 2020”
2. ᑎᑎᕆᖅᓂᑦ “Δᑲᑐᐊᒥ ᑎᓂᒥᐊᕆᖅᕐᓇᓂᒥᖅ ᐊᐅᓪᓗᑎσᓕᒧᑦ <ᕐᓇᐅᑎᑦ: ᑭᖓᔪᓪᓕᒥ ᑎᑎᕆᖅᓯᒪᓂᖅᓂᑦ, ᖃᓄᐊᓗ 2020”
3. ᐅσᓂᑲᓪᓕᐊᑦ “ᕆᖅᕈᓴᐅᖅᑐᑎᒥ (East Bay-Γ) ᐊᓘᒪᓗ Δᑲᑐᐊᒥ (HΔᐅᓗ ᑎᐸᵒᔪ) ᑎᓂᒥᐊᕆᖅᕐᓇᓂᒥᖅᓂᑦ - ᑭᖓᔪᓪᓕᖅᐸᖓᓂ ᑎᑎᕆᖅᓯᒪᓂᖅᓂᑦ ᐊᐅᓪᓗᑎσᓕᒧᑦ <ᕐᓇᐅᑎᖅ ᓄᓇᕗᒻᒥ ᐅᒪᔨᓪᓚᔪᕐᐊᖅᑯᑦ ᑲᑎᒪᓂᖅᓂᒋᓚᐅᔪᖅᑕᐅᓗᒪᕈᑐᐃσᖅ”
4. ᑎᑎᕆᖅᓂᑦ “2016-Γᑲ 2023-Lᑦ ΔᓄΔᑦ ᐊᖅᑐᖅᑕᐅᓯᒪᓂᖅᓂᒋᓄᑦ Δᑲᔨᓗᐊᖅᑕᒋᑎᑎᒋᓚᖅᓂᒋᓄᑦᓗ ᐊᖏᑭᑎᑦ ᑲᓇᑕᒥ ᐅᒪᔨᓪᓚᔪᕐᐊᖅᑯᑦ ᑎᑎᕆᖅᓯᒪᓂᖅᓂᒋᓄᑦ ᐊᓘᒪᓗ ᑎᓂᒥᐊᕆᖅᕐᓇᓂᒥᖅᓂᑦ ᓄᓇᕗᒻᒥ ΔᓄΔᑦ ᓄᓇᕗᑦᓱᒪᔨᖏᵒᓄᓕᕆᓂᒋᓄᑦ”
5. ᑎᑎᕆᖅᓯᐅᑎᓂᑦ “ᑎᓂᒥᐊᕆᖅᕐᓇᓂᒥᖅ ᐊᐅᓪᓗᑎσᓕᒧᑦ <ᕐᓇᐅᑎσᖅ ᖃᑭᕐᕕᐊᕐσᓕᒧᑦ ᐅᕆᖅᐅᓯᖅᓴᕆᖅᕐᓇᐅσᓕᒫᓂᑦ ᐊᓘᒪᓗ ᑐᖃᓯᒪᔨᓄᑦ ᑎᑎᕆᖅᓯᒪᓂᖅᓂᑦ”

ᐊᔨᔪᖏᵒσᖅ ᑐᔪᕆᖅᑕᐅᓗᑎᖅ:

ᓄᓇᕗᒻᒥ ᑲᑎᒪᓂᖅᓂᑦ ᓴᓇᔨᓴᕐᓂᖅ ᐅᒪᔨᓪᓚᔪᕐᐊᖅᑯᑦ ᐊᐅᓪᓗᑎσᓕᒧᑦ <ᕐᓇᐅᑎᓂᑦ: ᑭᖓᔪᓪᓕᒥ ᑎᑎᕆᖅᓯᒪᓂᖅᓂᑦ, ᖃᓄᐊᓗ 2020”

ᑲᑎᒪᓂᖅᓂᑦ

ᑲᑎᒪᓂᖅᓂᑦ

Environment and Climate Change Canada | Environnement et Changement climatique Canada



2020

ᖃᒧᑕᕐᓂᖅ ᑎᒥᐅᔪᓂᖅ ᐱᓕᕆᔨᒃᑯᑦ ᐊᕐᓇᐅᑎᓪ

[ᐅᓄᐊᓪ 2020]

ᐃᓕᑕᒪᖃᐅᔪᓂ:

İᓴᓇᐅᓕᕐᒪᑐᒃᓴᖅ ᓯᖅᓂᐊᒍᓕᓇ ᑲᑎᒪᔨᐅᓕᐅᕐᒪᑐᒃᓴᖅᑐᓕ, ᑲᑎᒪᔨᐅᓕᕐᒪᑐᒃᓴᖅᑐᓕ ᐃᔅᓂᐅᕐᖓᓂᖅᓯᓕᕐᓂᓴᖅᓯᓕᕐᓂᓴᖅᑐᑦ ᐊᐅᓕᕐᑎᑦᐅᕐᒪᑐᒃᓴᖅᑐᑦ ᐊᐅᓕᕐᑎᑦᐅᕐᒪᑐᒃᓴᖅᑐᑦ (ᐃᔅᓂᐅᕐᖓᓂᖅᓯᓕᕐᓂᓴᖅᑐᑦ ᐊᐅᓕᕐᑎᑦᐅᕐᒪᑐᒃᓴᖅᑎᑎᑦ) ᓴᖅᑭᑎᑕᓕᐅᕐᒪᑐᒃᓴᖅᓴᓐᓂ ᐅᓂᐊᓂᖅᓴᓐᓂ ᐊᐅᓕᕐᑎᑦᐅᕐᒪᑐᒃᓴᖅᑐᑦ ᐊᐅᓕᕐᑎᑦᐅᕐᒪᑐᒃᓴᖅᑐᑦ ᑲᑎᒪᔨᐅᓕᕐᒪᑐᒃᓴᖅᑐᑦ ᐊᐅᓕᕐᑎᑦᐅ

Environment and Climate Change Canada
Public inquiries centre
Fontaine Building 12th floor
200 Sacré-Coeur Blvd
Gatineau, QC. K1A 0H3
Telephone: 819-938-3860
Toll-free: 1-800-668-6767 (in Canada only)
Email: ec.enviroinfo.ec@canada.ca

ᖃᒃᓱᓕᔪᖅ ᐊᐃᔨᔪᐊᑎᓱᒃ: © ᐊᕿᑎᓕᓗᓂᔨᕐᕕᐊᒃᑯᓪ ᓯᓚᐅᔪᓐᓇ ᐊᓯᐅᓴᖅᓯᓕᓕᐊᔪᖅᓱᒃ ᑲᓇᒥᒥ, ᖃᐅᔨ-ᑲᑎᕐᓯᐊ ᑐᑲᐊᖅ
© ᑯᐃᐧᑎᑦᓴᖅ, ᐱᓕᕆᓇᐅᑎᖃᖅᑎᓱᒃᓴᓪᓗ ᑲᓇᒥᒥ, ᑭᓱᖅᑐᖅᑕᐅᔪᓐᓇᓂᐅ ᒥᔪᑦᑕᖅᓴᓪᓗ ᐊᕿᑎᓕᓗᓂᔨᕐᕕᐊᒃᑯᓪ ᓯᓚᐅᔪᓐᓇ ᐊᓯᐅᓴᖅᓯᓕᓕᐊᔪᖅᓱᒃᓴᓪᓗᑦ, 2020

Aussi disponible en français
ᐊᑐᐃᓐᓇᐅᑎᒋᔭᐅᑦ ᐅᐃᓴᖅᑎᑐᑦ

ᐱᓴᓯᑎᑭᓪᓗᑭᓂ ᐊᕐᖓᑎᓕᓚᓴᕋᐊᒃᓯᓂ ᓱᓕᐅᖅᓯᓪᓕᐊᓂᖅᑐᓄᓚᓂ ᑲᓇᑕᐅᖅ ᐊᐅᓚᑦ ᐃᓕᓴᕐᕕᖏᓐᓂᒃ ᐊᑐᖅᑐᕐᓂᖅᑐᓂ ᐊᓂᔪᓂᖅᐸᑦᓕᐊᓂᖅᓄᓚᓂ ᓄᓇᕗᑦ ᐅᖃᓕᒫᓂ ᐊᑐᖅᑐᕐᓂᖅᑐᓂ ᐃᓚᖓᑕᐅᓂᖅ ᐊᑐᖅᑐᕐᓂᖅᑐᓂ ᐊᑐᖅᑐᕐᓂᖅᑐᓂ ᐊᑐᖅᑐᕐᓂᖅᑐᓂ

ᖃᓄᖅ ᐊᖏᑎᑎᓯᓂ ᐊᓯᑎᓕᓪᓚᕆᔨᓂ ᓱᓕᐅᔨᓗ ᐊᓱᔨᕐᓴᖅᓯᓕᐅᖅᓯᓚᐅᓂ ᑲᓇᑕᒥ ᓴᓇᔨᖅᑐᑦᓕᐅᑎᓂᒃᑯᐱᒪᖏᖅᑐᓂ?

ᐃᓕᓇᐅᔨᖅ ᓴᓇᔨᖅᑐᑦᓕᐅᑎᓂᒃᑯᐱᒪᕐᒪᖔᑦ ᐱᓕᓪᓕᖏᐅᔨᓂ 55-ᖅᑎᑯᓕᖅᑐᑦ ᑲᓇᑕᓕᒫᒥ ᐅᒪᔨᖅᑎᖏᐅᑎᑕᐅᔨᓂᐅ ᐊᒻᒪᓗ 92-ᖅᔪᓕᑎᑯ ᑎᖅᒥᐊᖅᑎᖏᐅᔨᓂ, ᐊᖅᑎᓂᖅᑲᖅᑎᑎᓕᓗᑎᑯ ᓄᓇᖅᑎᓂ ᐅᖅᓚᒌᓂ 14 ᒥᓕᔪᓂᒃ ᐄᐃᐊᐳᒌᓂᒃ ᐅᐸᑐᕋᐅᑎᓕᓪᓂᒃ ᑲᓇᑕᒥ ᓇᓂᑐᐃᓐᓇᖅ.

ᓱᓇᐅᖅᓚᒌ ᐊᐅᓕᕆᑎᔨᐸᑎᐃᓂ ᐸᖁᓇᐅᑦᑐᓂ?

ᐊᐅᓕᕆᑎᔭᒪᓂᕐᒧᑦ ᐸᖁᓇᐅᑦᑐᓂ ᐱᑕᖁᖏᐃᐅᖅᑐᑦ ᐱᓕᓚᑐᑕᐅᔨᓴᖅᓂᕐᒧᑦ, ᐊᐅᓕᕆᑎᐅᔨᐅᔨᓄᑦ ᐃᓱᒪᓕᐅᕈᑕᐅᓴᓚᕆᓕᓕᒫᓄᑦ. ᑐᑦᖅᓚᐅᖅ ᐊᑐᖅᑕᐅᓂᕐᒧᑦ ᐊᖅᑎᓂᖅᑲᖅᑎᑎᓕᓗᑎᑯ ᓄᓇᖅᑎᓂ ᐱᑕᖁᖏᐃᐅᖅᑐᑦᓕᐅᑎᓂᒃᑯᐱᒪᖏᖅᑐᓂ ᐊᐅᓕᕆᑎᐅᔨᐅᔨᓄᑦ ᐃᓱᒪᓕᐅᕈᑕᐅᓴᓚᕆᓕᓕᒫᓄᑦ ᐱᓕᓪᓕᖏᐅᔨᓂ ᐊᓯᔨᓴᖅᑐᑦᓕᐅᑎᓂᒃᑯᐱᒪᖏᖅᑐᓂ ᐊᒻᒪᓗ ᐃᖅᑭᐅᒪᓕᖅᑎᓯᓚᕆᓂ ᐊᐅᓕᕆᑎᐅᔨᐅᔨᓄᑦ ᐱᓕᓪᓕᖏᐅᔨᓂ ᐊᓯᔨᓴᖅᑐᑦᓕᐅᑎᓂᒃᑯᐱᒪᖏᖅᑐᓂ. ᐊᖅᑎᓂᖅᑲᖅᑎᑎᓕᓗᑎᑯ ᐅᑎᕐᓂᖅᑐᑦ ᐊᐅᓕᕆᑎᐅᔨᐅᔨᓄᑦ ᐊᓯᔨᓴᖅᑐᑦᓕᐅᑎᓂᒃᑯᐱᒪᖏᖅᑐᓂ ᐊᐅᓕᕆᑎᐅᔨᐅᔨᓄᑦ ᐱᓕᓪᓕᖏᐅᔨᓂ

ᖃᓄᐃᔨᑎᖏᑕ ᓴᓇᓂᕆᐊᖅᑕᐅᓯᒪᔪᓂ ᓄᓇᐃᓂ ᐊᐅᓪᓚᑕᐅᕙᒃᑐᑐᑦ?

ᐊᐅᓪᓚᕈᑎᐅᔪᓂᑦ ᐊᑐᖅᑕᐅᕙᒃᑐᑐᑦ ᒪᓯᓄᖏᓪ ᓯᔅᓗ ᑐᕐᖏᓂᕐᒧᑦ, ᓇᐅᔨᑎᖅᑐᐃᒃᓪᓚᐊᖅᑕᕈᑦᑕᐃᓐᓇᕐᓂᕐᒧᑦ, ᐋᖅᑭᐅᒪᔨᑎᔨᑎᐊᕐᒪᓂᕐᒧᑦ, ᐱᐅᓯᕈᐊᖅᑎᔨᑎᔪᓐᓇᖅᓂᕐᒧᑦ ᐅᒪᔪᖅᑕᐅᕙᒃᖏᓂᖅᓄᑦ, ᓇᔪᖅᑕᐅᕙᒃᓂᖅᑭᓐᓂᖅᓄᑦ, ᐃᓕᓴᐅᔨᓯᑐᓗ ᖃᐅᔨᓴᖅᑕᐅᖅᑕᖏᓂᖅᓄᑦ, ᒪᓕᑐᖏᓂᖅᓄᑦ ᒪᓕᖅᑕᐅᒃᖃᐃᐅᓚᐊᖅᖏᓂᖅᓄᑦ ᐊᓱᒪᓗ ᐱᖏᔨᑎᖅᑭᓐᓂᖅᓄᑦ ᐃᓚᓗᖏᔨᑎᖅᑐᓗ ᐋᖅᑭᐅᒪᔨᐊᓚᐊᖅᖏᓂᖅᓄᓗᒃ. ᖃᐅᔨᓴᖅᑕᐅᖅᑕᖏᓂᕐᒪᕐᕕᖅ ᐱᓚᒪᓚᐊᑐᑎᒃᓴᖅᑕᕐᕕᖅ ᐅᓇᑕᒥ ᐱᓕᕐᓂᖅᓄᑦ ᐃᓚᓗᖏᔨᑎᖅᑭᓐᓂᖅᓄᑦ ᐃᓕᓴᐅᔪᖅᑕᐅᖅᑕᖏᓂᖅᓄᑦ ᐅᖃᓕᒫᓂᖅᓄᑦ ᐊᑐᖅᑕᐅᕙᒃᖏᓂᖅᓄᑦ ᐃᓕᓴᐅᔪᖅᑕᐅᖅᑕᖏ

ᖃᒃᑲᓇᐅᑲᐃᔭᕐᑎ ᑎᓕᐊᖏᖃᕐᓂᒃ

ᖃᒃᑲᓇᐅᑲᐃᔭᕐᑎ ᑎᓕᐊᖏᖃᕐᓂᒃ (ᖃᒃᑲᓇᐅᑲᐃᔭᕐᑎᑲᖅ) ᐃᓕᓴᒃ ᑕᐅᔪᓂ ᑲᓇᓐᓇᖅᑕᓕᓴᓂ ᓴᖅᓴᕐᑐᐃᔨᓂ, 35 ᑭᓯᒥᑕᓂᒃ ᐅᖏᓕᓕᓴᓂᖓᒥ, ᑭᔪᖅᓴᐅᓂ ᐊᕕᒃᑐᓂᓕᓴᓕᓂᓕᓂ, ᓄᓇᕗᓕᒥ. ᖃᒃᑲᓇᐅᑲᐃᔭᕐᑎ ᑎᓕᐊᖏᖃᕐᓂᕕᐅᓂᖅ ᐊᑕᑭᕗᖅ, ᐃᒪᓐᔪᑦ ᑕᓇᐅᕗᔪᓂᓕᓂ 50 ᑭᓯᒥᑕᓴᓐᔪᑦ ᑕᑭᓂᓴᓐᔪᑦ, ᐊᓚᒐᓂ ᑕᒪᐃᓐᓂᒃᑲᒃ ᐊᑕᑭᓪᓗ ᑕᐅᕗᕐᓂᓕ

ᐊᓂᑎᔨᑎᑕᐅᓂᖅᑭᓐᓄᑦ ᐱᖓᓐᓇᖅᑕᓕᓴᓂᖓᔪᓄᑦ ᓄᓇᔪᑦ ᑎᑭᐅᕐᔪᓄᑦ, ᑐᓂᒋᑎᐅᑦ ᓄᓚᕗᓕᓂᖅᑐᑦ (ᖃᒃᓪᓗᓇᑎᑐᑦ Boothia Fox Shield-ᒥᒃ ᑕᐃᔨᐅᕐᔪᓂ ᐅᔨᕋᑦᑲᓪᓗᓇᖅᑭᓐᓄᑦ), ᓄᓇᒥᐅᑕᓕᐊᔨᓐᓄᑦ ᐅᒃᑯᓂᖅᓴᕐᓂᖅ ᖃᒃᑯᕐᓂᖅᓴᖅ ᐅᔨᕋᖅᓂᖅ. ᐃᒪᐃᔨᓪᓗᐊᓂᔪᖅᑭᑐᓂᓪᓗ ᑕᒪᓐᓇ, ᖃᒃᑲᐅᑉᖃᒃᑲᐃᕐᓂᕆᐊᖑᓂᖅᓱᒃ, ᐱᑕᕋᐃᐅᓂᒪᔪᖅ ᑭᓕᓇᖅᓂᖅ ᐊᑕᑭᓪᓗᑕᐅᓂᖅᑭᓐᓄᓂᖅ ᓄᓇᒥ, ᐊᒥᓂᒍᓕᐊᓂᖅ ᑎᑭᐅᓂᑕᓕᐅᑉᐸᓂᓕᓂ ᐃᑲᑕᐃᓕᓂ ᑕᓂᒍᓐᓇ. ᐃᒪᐃᔨᓪᓗᐊᓂᔪᖅᑭᑐᓂᓕᓂ ᑲᑎᒪᔨᖃᕐᓂᖅᑭᓐᓄᑦ ᑕᐃᔨᐅᕐᔪᓂ ᑭᓚᐅᖅᑭᑦ ᐊᑕᑭᓪᓗᑕᐅᓂᖅᑭᓐᓄᓂᖅ ᓄᓇᒥ, ᐊᒥᓂᒍᓕᐊᓂᖅ ᑲᑎᒪᔨᖃᕐᓂᖅᑭᓐᓄᓂᖅ ᑕᐃᔨᐅᕐᔪᓂ ᑲᑎᒪᔨᖃᕐᓂᖅᑭᓐᓄᓂᖅ ᐱᖓᓐᓇᖅᑕᓕᓴᓂᖅᑭᓐᓄᓂᖅ ᐊᑕᑭᓪᓗᑕᐅᓂᖅᑭᓐᓄᓂᖅ ᓄᓇᒥ, ᐊᒥᓂᒍᓕᐊᓂᖅ ᑲᑎᒪᔨᖃᕐᓂᖅᑭᓐᓄᓂᖅ ᐱᖓᓐᓇᖅᑕᓕᓴᓂᖅᑭᓐᓄᓂᖅ ᑕᐃᔨᐅᕐᔪᓂ

ᐱᓕᒪᓐᓄᓂᖅ ᖃᖅᑳᓕᓄᖅᑐᑦ ᑎᖏᑕᐊᖅᖃᕐᓗᖅ

ᐃᓐᓇᓴᖅ ᑎᖏᑕᓯᓂᑦ ᖃᓂᒪᓂᑕᓂᓕᑎᓂᒃ ᐊᔪᓪᓚᓄᖅᑕᓂᒃᓴᓪᒪᓂᓂᓄᓂᑦ ᓄᓇᖏᑦ ᑕᓂᖅᓂ ᖃᓄᖏᔭᖏᑐᓪᓗ ᑐᑭᓚᓂᑭᑕᓂᓕᓂᖅᑐᑦ ᓇᒪᓂᑦᑕᖅᑕᓂᓄᖏᑦ, ᑎᓪᓗᖅᖃᐊᖅᓯᓂᓄᖏᓐᓂᓪᓗ ᐊᒥᓯᓂᖅᖃᖅᓂᖅ ᑕᐃᓂ 234,000-ᓂᒃ ᑲᖓᖅᓂᒃ (ᖃᓄᖏᔭᖏᑐᓵᓕᑐᒃ ᑕᐃᓂ ~2% ᐳᓴᓐᑎᓂᒃ ᒥᑭᐃᑎᐅᓂᖅᓴᐃᑦ ᑲᖓᐃᑦ ᓇᓂᑐᐃᓐᓇᕐᑎᐊᖅ ᐱᑕᖅᓯᖓᓂᖏᓂᒃ ᑲᓇᑕᐅᓪᓗ ᓄᓇᖅᑐᓯᐊᓵᖅᑎᑕᕐᒥᔪᓂᒃ). ᓇᓪᐅᑦᖅᑕᐅᓯᒪᒋᐅᑦ ᑕᐃᓵᓂᖅᓴᓂᖅ 5,000 ᐅᓪᓗᖅᓯᖓᓂᓂᑦ ᓂᕆᓪᓚᐃᑦ Atlantic brant (3% ᐳᓴᓐᖏᓵᓵᕐᑐᒃ ᐱᑕᖅᖃᓪᓚᐅᖅᑐᒃ ᑭᓯᐊᓂ ᐃᓐᓇᐅᓪᓚᖅᑐᖅ ᖃᐅᔨᓴᒪᐅᔪᓵᓵᖅᑐᑦ ᖃᓄᐅᑎᑦ ᐱᑕᖅᓯᖓᓪᓚᖕᖏᓐᓄᑦ. ᐊᑕᐅᓯᓂ ᑕᐃᒪ ᒥᑎᖅᑕᖅᓄᖅᖢᓂᑕᐅᐊᖓᔪᖅ ᓄᓇᑕᐅᑦ ᐅᑭᐅᖅᑕᖅᑐᑦᓴᓂᑦ ᖃᓄᐅᑎᑦ ᐅᑭᐅᖅᑕᖅᑐᑦᓴᓂᑦ, ᖃᑭᑭᖅᑕᕐᒥᑦ. ᐊᑕᐅᓯᓂ ᑕᐃᒪ ᒥᑎᖅᑕᖅᓄᖅᖢᓂᑕᐅᐊᖓᔪᖅ ᓄᓇᑕᐅᑦ ᐅᑭᐅᖅᑕᖅᑐᑦᓴᓂᑦ, ᐊᑐᓂᖅᑐᑦ ᐃᓐᓇᐅᓪᓚᖅᑐᖅ ᐱᑕᖅᓯᖓᓪᓚᖕᖏᓐᓄ

ᑕᒪᑲᑯᓇᖅᓯᒪᓕ. ᖃᓄᖅᓱᐃᓐᓇᕐᓂᖅ ᑎᑎᒋᔭᖅᑎᓯᐅᑦᑕᕐᓄᑦ ᑎᑭᑦᑕᐃᔨᓂᕐᓄᑦ ᐅᑭᐅᖅᑯᑦ, ᓴᓪᓕᒋᐅᓪᓗ ᖃᐅᔨᑎᑦ ᑕᐅᑐᖅᑯᓯᒪᓕᕐᓂᖅ ᐊᓯᓚᑖᓂᖅ ᐊᑐᐊᓕᖅᑐᑦ ᐊᐅᓚᑦᑎᓯᐅᑎᓂᖅ ᐊᑭᓐᓇᐅᑉ ᐊᑭᓐᓇᐅᑉ ᑕᐅᑐᖅᓯᒪ.

ᐊᐅᓚᑦᑎᓴᐅᖅᑎᑦᑕᐅᓂᕐᓂ ᐊᓯᒪᓗ ᐊᖏᕈᑕᐅᔨᓴᓚᐃᓪᓕᑦ ᑲᒪᒋᔭᐅᓯᖏᓐᓄᑦᓂᕐᓂ

ᒪᓕᖅᑕᐅᔨᓴᐊᕐᓂᖅᓯᖏᓐᓄᑦᓂᕐᓂ ᓄᓇᕗᓂᒥ ᐊᖏᕈᑐᐃᑦ, ᐃᓄᐃᑦ ᐊᐅᑐᑕᐅᓕᒪᓂᖏᓐᓄᑦᓂᕐᓂ ᐃᑲᔪᓕᐊᖅᑕᕈᑎᐅᓴᖏᓐᓄᑦᓂᓪᓗ ᐊᖏᕈᑐᑎᑦ ᑲᑎᒪᔨᖅᑎᑦᑎᓂᖅᓂᕐᓂ ᐊᓯᒪᓗ ᑎᑎᒋᔭᖅᑎᓯᐅᑦᑕᕐᓄᑦ ᐊᓯᔾᔨᕈᓘᔪᐊᕐᓂᖅᓯᒪᓕᕐᒥ, (ᐃᓄᐃᑦ ᐃᑲᔪᓕᐊᖅᓴᖏᓐᓄᑦᓂᕐᓂ ᐊᖏᕈᑐᐃᑦ) ᓕᓚᐅᑦᒥ ᐱᓱᓴᖅᑕᐅᔨᓴᖃᓕᓴᖅᑐᑦ 2006-ᖓᑎᓪᓗᒍ, ᐊᑐᖅᑕᐅᓕᓕᖅᑐᓂᖅ 7-ᓂᓪᓗ ᐊᖅᑭᒍᓂᑦ, ᐊᖏᕐᓂᖅᑎᑦᑎᐊᖅᖃᓂᕐᔪᓕᓕᓚᓪᓕᑎᑐᖅ 7-ᑲᓂᓕᐅᓕᐊᓕᒥᔨᓂᓪᓗ ᐊᖅᑭᒍᓂᑦ. ᒪᓕᓚᐃᑦ ᐊᐃᓴᑐᖅᓯᒪᓂᖏᓐᓄᑦ 3-ᖓᓚᓵᑐᑐᑦ (ᐱᔨᔪᑎᖅᓂᖅᑐᑦ ᐊᐅᓕᕐᑎᓴᐅᓂᖅᑎᑦᑕᐅᔨᔪᓂᑦ) ᑕᒪᑲᑯᓇᖅᑎᑦ ᐃᓄᐃᑦ ᐃᑲᔪᓕᐊᖅᓴᖏᓐᓄᑦᓂᕐᓂ ᐊᖏᕈᑐᐃᑦ ᑎᑎᖅᑲᐃ ᐅᖅᓂᖅᓯᒪᔨᖅᓂᖅᐳᑦ ᒪᓐᓇᑐᓂᖅᓂᕐᓂᓪᓗ ᐃᓕᓴᕐᓂᖅᓂᕐᓂᓪᓗ:

a) ᐊᑦᑭᐅᒪᕕᑎᐊᖅᑐᓂᖅ ᐊᐅᓕᕐᑎᓴᐅᑎᓂᖅᑎᑦᑎᓂᖅ ᓄᓇᕗᓂᒥ ᐅᒪᔨᕐᓂᖅᓯᐃᓐᔨᓂᖅ ᑎᑎᒋᔭᖅᑎᓯᐅᓕᖅᓴᓂᐅᑦ; ᑭᒃᑯᐊ ᐃᓄᐅᑯᑎᑎᖅᓂᕐᓂ ᐊᖏᓪᓗ ᐅᒪᔨᕐᓂᕆᔨᒃᑯᑦ, ᒪᓕᖅᑕᐅᔨᓂᐊᑦᑎᓂᖅ ᓄᓇᕗᓂᒥ ᓄᓇᕗᑦᑎᓂᖅᓂᕐᓂ ᐊᖏᓪᓗ ᐱᓕᕆᔨᖅᑐᓂᖅᓂᕐᓂ ᐊᖏᓪᓗ ᐱᓕᕆᔨᖅᑐᓂᖅᓂᕐᓂ ᑕᐃᒃᑯᓐᓇᖅᑐᑦ ᑎᑎᖅᑲᐃᓐᓇᖅᑐᓂᖅ ᑕᐃᒃᑯᓐᓇᖅᑐᑦ ᐃᓄᐃᑦ, ᑭᒃᑯᐊ 9-ᒥ, 5-ᒥᓗ;
b) ᐃᓱᒪᓕᐅᕙᓪᓕᐅᕙᑎᓂᖅᑎᑦᑎᓂᖅ ᓄᓇᕗᓂᒥ ᐅᒪᔨᕐᓂᖅᓯᐃᓐᓇᖅᓴᖏᓐᓄᑦᓂᓪᓗ ᐱᓗᒪᓐᓇᐅᔨᓴᐅᑐᑦ ᑐᑭᐅᓱᒪᓕᐅᕙᑎᐊᖅᑎᓯᐅᑦᑕᕐᓂᓪᓗᑎᑦᑎᓂᖅ ᐊᑐᐊᓐᓇᓪᓗᑎᑦᑎᓂᖅ ᐃᓄᐃᑦ ᖃᐅᔨᓴᖅᑐᓂᖅᓂᕐᓂ ᐊᖏᓪᓗ ᐱᓗᒪᓐᓇᐅᔨᓴᐅᑐᑦ ᑎᑎᖅᑲᐃᓐᓇᖅᑐᓂᖅᓂᕐᓂ ᑕᐃᓚᐅᔨᓴᐅᑐᑦ ᐊᖏᓪᓗ
c) ᓄᓇᓕᓂᕐ ᐃᓄᐃᑦ ᐃᓕᓐᓄᐃᓕᖅᑕᐅᓂᖅᑎᑦᑎᓂᖅ <ᕐᓇᐃᓕᓐᓂᖅᓂᕐᒧᑦ ᐊᐅᓕᕐᑎᓴᐅᑎᓂᖅᑕᐅᓂᕐᒧᑦᓴᓪᓗ ᓄᓇᕗᓂᒥ ᐅᒪᔨᕐᓂᖅᓯᐃᓐᓇᖅᓴᖏᓐᓄᑦ ᐊᖏᓪᓗ ᑎᑎᖅᑲᐃᓐᓇᖅᓴᖏᓐᓄᑦ.

ᑲᓇᑕᒥ ᐅᒪᔨᕐᓂᕆᔨᒃᑯᑦ ᐊᐅᓕᕐᑎᓂᐅᑉᑐᓂ ᖃᓄᖅᓱᐃᓐᓇᕐᓂᖅᑎᑦᑎᓂᖅᓂᕐᓂᑲ ᐃᓕᓐᓂᑦᑐᓂᕐᓂ ᐃᓅᓂᐅᑦᓯᓂᖅᓂᕐᓂ ᐊᖏᓪᓗᐊᓕᖅᑐᓂᖅᓂᕐᓂᓪᓗ ᐊᐅᓕᕐᑎᓴᐅᑎᓂᖅᑕᐅᓂᕐᒧᑦᓴᓪᓗ ᑲᑎᒪᔨᖅᑎᑦᑎᓂᖅᓂᕐᓂ (ᐃᓅᓂᐅᑦᓯᓂᖅᓂᕐᓂ ᐊᐅᓕᕐᑎᓴᐅᑎᓂᖅᑕᐅᓂᕐᒧᑦᓴᓪᓗ ᑲᑎᒪᔨᕐᕕᒃᓴᓂᖅ) ᓴᓪᓕᒋᐅᓪᓗ ᐊᖏᓪᓗᐊᓕᖅᑐᓂᖅᓂᕐᓂ ᐃᓅᓂᐅᑦᓯᓂᖅᓂᕐᓂ

ᒪᓪᒃᑕᐅᓪᓗᓂᒃ ᐃᓄᐃᑦ ᐃᑲᔪᓯᐊᕐᓴᖅᓄᓄᑦ ᐊᖏᕈᑎᒋᑦ, ᐃᓕᓴᕐᕕᖏᓂᑦ ᐊᐅᓪᓚᕆᐅᕐᕕᖃᑦᑕᐅᔪᓂ ᑎᑎᕋᓚᐅᕐᕕᖓᑦᑐᑦ ᐊᐅᓚᕆᓴᕐᓗᒍ <ᕐᓇᐅᑎᓴᒃ ᖃᕐᕕᓴᐅᕐᑐᐃᓐᓇᓴᒃ, ᐃᓄᐃᑦ ᑐᐱᔪᓴᐊᕐᕕᕐᕕᖃᑦᑕᕐᔪᑎᒃ, ᐊᓚᒪᓗ ᑭᖅᑦᓚᕐᒥᐅᑦ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖃᑎᒋᖅᑎᑦ, ᑐᵒᖏᓚᕐᑲᖏᓐᓂᓗ ᐊᓚᒪᓗ ᓄᓇᓚᖅᓴᒃ ᐃᓄᖅᓴᒃ ᐃᓚᐅᔪᓚᔪᓴᒃ ᓴᑦᓚᓐᓴ. ᐊᐅᓚᕆᓂᔪᕐᕕᖃᑦᑕᐅᔪᓂ ᖃᑎᒌᓐᓇᕐᔪᓚᕆᓴᕐᕕᖃᑦᑕᐅᓴᖅᓴᑦ ᐱᓕᕐᓂᖅᑕᐅᔪᓚᕆᓴᕐᕕᖃᑦᑕᐅᓴᖅᓴᒃ ᐊᐅᓚᕆᓂᔪᓐᓇᖅᑐᑦ <ᕐᓇᐅᑎᓂᒃ ᑕᐃᖁᓄᓐᓚᒃ ᓄᓇᓕᖅᑐᑦ ᐅᓪᓗᕈᓘᑦᑐᒥᐅᑕᐅᓂᖅᑐᑦ (ᐅᑐᔾᔨᑎᒃ ᐃᓄᐃᑦ *ᐃᖁᔪᓴᐊᕐᓴᖅᓄᓄᑦ ᐊᖏᕈᑎᒋᑦ 3.6.2*). ᒪᓕᓴᓗᑎᒃ ᐃᓱᒪᓕᐅᕐᕕᖃᑎᐅᔪᓐᓇᖅᑐᓂᖅᓴᒃ ᐱᔪᓐᓇᕐᓂᕐᑎᑦ ᐊᑕᒍᑦ ᓄᓇᕗᓂᑦ ᓄᓇᕈᑕᐅᑎᔪᑦ ᐊᖏᕈᑎᒋᖅᓄᑦ, ᐅᒃᖢᔾᔨᖅᑕᐅᓂᑦ ᑭᓴᕐᑕᐅᔪᖅ ᐊᖏᕈᑎᖃᖅᑎᕐᕕᑦ, ᑕᒪᐃᓐᓇᖅ, ᐃᓕᓴᕐᕕᖏᓂᑦ ᐅᑎᖅᑎᑕᐅᔪᓂ ᐊᐅᓚᕆᓴᕐᕕᖃᑦᑕᐅᔪᓂ ᐅᑎᖅᑎᓂᒃᓴᖅᑐᑦ, ᐃᓱᒪᒃᓴᖅᓱᐅᕈᑕᐅᓚᐅᖃᖅᑐᓴᖅᑐᓂᒃ, ᐃᓱᒪᒃᓴᖅᓱᐅᕈᑎᐊᖅᑐᓴᖅᑐᓂᖅᓴᖅᓱᐅᓗᒍ ᐊᐅᓚᕆᓂᔪᓐᓇᖅᑐᑦ <ᕐᓇᐅᓱᐊᑦ ᑐᓴᕐᐅᖅᑐᓴᖅᑐᓂᖅᓴᖅᓱᐅᓗᖏᑦ ᑭᖓᔪᓪᓗᕆᑦ ᐃᓱᒪᓕᐅᕈᑕᐅᑎᔪᑦ ᐅᓪᓗᕈᓘᑦᑐᒥᐅᑕᐅᓂᖅᑐᑦ ᐃᓕᓴᕐᓂᕐᕕᖃᑦᑕᐅᓴᖅᓴᑦ ᓄᓇᕗᓂᑦ. ᒥᓂᑕᐅᔪᖅ ᐊᖏᕈᑎᖃᖅᖢᑦ ᐊᐅᓚᕆᓴᕐᕕᖃᑦᑕᐅᔪᓂ <ᕐᓇᐅᑎᐅᓴᖃᑦᑕᐅᔪᓂᖅᓴᒃ ᒥᓂᑕᐅᔪᖅ ᑲᑎᒪᔨᖅᑎᑦ ᐃᓕᓴᕐᓂᕐᕕᖃᑦᑕᐅᔪᓂ ᐊᖏᕈᑎᖃᖅᑎᕐᕕᑦ ᑕᒪᐃᓐᓇᖅ ᓴᓇᔨᓴᒃᑯᑦ ᐃᓄᐃᑦ ᐊᐅᓚᕆᓴᕐᕕᖃᑦᑕᐅᔪᓂ ᐊᐅᓚᕆᓂᔪᓐᓇᖅᑐᑦ <ᕐᓇᐅᑎᓂᒃ ᐃᓄᐃᑦ ᐊᐅᓚᕆᓴᕐᕕᖃᑦᑕᐅᔪᓂ ᐊᐅᓚᕆᓂᔪᓐᓇᖅᑐᑦ ᐃᓕᓴᕐᓂᕐᕕᖃᑦᑕᐅᓴᖅᓴᑦ ᓄᓇᕗᓂᑦ ᐊᖏᕈᑎᖃᖅᑎᕐᕕᑦ ᐃᓕᓴᕐᓂᕐᕕᖃᑦᑕᐅᓴᖅᓴᑦ ᓄᓇᕗᓂᑦ ᐊᐅᓚᕆᓴᕐᕕᖃᑦᑕᐅᔪᓂ ᐊᐅᓚᕆᓂᔪᓐᓇᖅᑐᑦ ᐊᐅᓚᕆᓴᕐᕕᖃᑦᑕᐅᔪᓂ ᐊᐅᓚᕆᓂᔪᓐᓇᖅᑐᑦ ᐃᓕᓴᕐᓂᕐᕕᖃᑦᑕᐅᓴᖅᓴᑦ ᓄᓇᕗᓂᑦ ᐊᖏᕈᑎᖃᖅᑎᕐᕕᑦ ᐃᓕᓴᕐᓂᕐᕕᖃᑦᑕᐅᓴᖅᓴᑦ ᓄᓇᕗᓂ

ᒪᐅᐱᓚᐃᓪ ᖃᐅᓄᐃᓕᓐᓄᓂᓐᓂᕐᓂ

ᑎᑎᕎᖅᑕᐅᓂᖅᓂᕐᓂ ᐊᐃᓴᐅᑐᓕᔪᓂᒃ ᑕᑯᑎᕎᑎᓵᔪᓱᓪᓕᓪ xi
ᑎᑎᕎᖅᓯᒪᓂᖅᓂᕐᓂ ᑕᑯᔾᔨᖅᓴᐃᓪᓕᓪ xi
ᑎᑎᕎᖅᓯᒪᓂᖅᓂᕐᓂ ᓇᐃᓚᓕᒋᐊᕎᖅᑕᐅᓯᒪᔪᓂᕐ ᑎᑎᕎᖅᐃᓪᓕᓪ ᖃᓄᖅᑲᓇᑕᑎᑐᑦ xii
1 ᐅᕎᖅᐅᓯᐅᑦᑎᐊᕎᓂᖅᓂᕐᓂ ᔪᐳᵒᓂᐊᖅᑕᐅᓯᒪᔪᓂᕐ ᓄᓇᐃᓪᓕᓪ 1
1.1 ᐊᐃᓴᐅᑐᖅᓯᒪᓂᖅᓄᑕ ᖃᓄᐃᓕᓐᓄᓂᓐᓄᖅ 9
1.2 ᖃᓄᐃᖅ ᐱᓕᕐᓂᐊᓕᐅᕎᓪᓗᓐᓂᓪ 10
1.3 ᓄᓇᕎᔨᑎᑎᑦᓴᐅᓂᖅᓂᕐᓂ ᐊᒻᒪᓗ ᐊᑐᖅᑕᐅᔪᵒᓇᕎᓂᖅᓂᕐᓂ 11
1.4 ᐃᓕᓴᕎᔨᑎᑎᑦᓴᐅᔪᓂᕐ ᐱᕎᔨᑦᓴᓪᓗ 13
2 ᐊᕐᕈᑎᖅᓂᕐ ᐊᕐᕈᑎᑎᐅᑕᕎᖅᓂᖅᓂᕐᓂᓪᓗ 15
2.1 ᓄᓇᒥᐅᑕᐃᓪᓗ ᑕᓚᐅᕎᒥᓗ ᐃᒪᕎᒥᐅᑕᐃᓪᓂ ᐅᒪᔪᖅᑲᓵᓪᓂ 15
2.2 ᑎᖅᒥᐊᓪᓂ 17
2.2.1 ᑲᖓᔾᐃᓪᓂ 17
2.2.2 ᒥᑦᓴᵒ 21
2.2.3 ᓯᓚᕆᖅᐸᓯᖅᒥᐅᑕᐃᓪᓂ ᓯᓚᕆᕎᓚᐊᑐᓚᐃᓪᓂ 22
2.2.4 ᐊᓯᖅᑭᓂ ᐃᒪᐅᓂᕐᒋᓯᓂ ᑎᖅᒥᐊᓪᓂ 22
2.3 ᐊᓯᖅᓂ ᐅᒪᔪᐃᓪᓂ 23
2.3.1 ᓄᓇᒥᐅᑕᐃᓪᓂ ᐅᒪᔪᐃᓪᓂ ᓂᕐᔪᑦᓴᓪᓂ ᐊᐱᒪᵒᑎᕎᑎᓯᓯᓂ ᐱᐊᑎᕐᒥᓂᵒᖅ 23
2.3.2 ᑕᓚᐅᕐᒥᐅᑕᐃᓪᓂ ᐅᒪᔪᐃᓪᓂ 24
2.3.3 ᑕᓚᐅᕐᖓᑭᵒᓂᕐᒥ ᑕᓯᕐᒥᐅᑕᐃᓪᓂ ᑕᓚᐅᕐᒥᐅᑕᐃᓪᓂᓪ ᐃᕐᖢᓗᑎᓴᐃᓪᓂ 24
2.4 ᐱᐅᕐᖢᑎᖅᓂᖅᓄᒃ ᓄᓇᓪ 25
2.5 ᐅᒪᔪᐃᓪᓂ ᐊᒥᓱᖅᑭᓪᓚᐊᓕᕐᖢᑎᖅᑐᓂ 25
2.6 ᑕᐃᓂᕎᒥᐅᑕᑎᖅᐅᓇᖅᑭᕐᖢᑎᖅᑐᓂ ᐅᒪᔪᐃᓪᓂ/ᕎᔨᐸᕎᕿᐃᓪᓚᵒᓂᓪ, ᐱᐅᕐᖢᑎᓪᓚᵒᓂᓪ ᐱᑕᕎᑎᓂᖅᓂᕐᓂ 29
3 ᐃᓕᓴᕐᔨᓯᑐᑎᕎᒥᔪᑦ ᐊᑐᓗᓚᕆᐅᕐᖓᓂᖅᓂᕐᓂ ᓄᓇᒥᐅᑕᕐᑕᐃ 30
3.1 ᐃᓕᓴᕐᔨᓯᑐᑎᕎᒥᔪᑦ ᓄᓇᒥ ᐱᔨᔨᑎᐃᓚᓂᓪ ᐊᑐᖅᑕᐅᕐᓭᒪᐅᑎᑦᓴᓂᖅᓂᕐᓂ ᑐᑭᖅᑎᓂᖅᓂᕐᒪᓂᓪᓗ ᖃᐅᔨᓴᖅᑕᐅᓯᒪᔪᓂᕐ 30
3.2 ᑕᐃᒪᓐᓇᓂᵒ ᐃᓄᐃᑦ ᐊᑐᖅᑎᑕᑎᓂᖅᓂᕐᓂ ᓄᓇᒥᵒ 31
3.3 ᐊᓐᓇᐅᓕᓴᐅᑐᖅ ᐃᓄᐃᑦ ᐊᑐᖅᑎᑕᑎᓂᖅᓂᕐᓂ ᓄᓇᒥᵒ 36
4 ᑐᔅᓚᓚᖅᐅᔪᓂ ᐊᒻᒪᓗ ᐱᖅᓵᖅᑕᐅᔪᓯᒪᔪᓂᕐ 38
4.1 ᑕᐅᑐᓚᓚᖅᐅᔪᓂ 38
4.2 ᑐᔅᓚᓚᖅᐅᔪᓂ ᐊᒻᒪᓗ ᐱᖅᓵᖅᑕᐅᔪᓯᒪᔪᓂᕐ 38
5 ᐊᐅᓪᓕᕐᓂᓂᕐᒥᔪ ᐃᓱᒪᓕᖅᑲᓱᐅᕐᑐᑎᓕᖅᑲᓇᐃᓕᓐᓂᕐ 42
6 ᐊᐅᓪᓕᕐᓂᑎᐅᑐᓂᕎᒧᑦ ᖃᓄᐃᓕᓐᓄᑕᐅᕐᑐᑎᓕᖅᑲᓇᐃᓕᓐᓂᕐ 49
6.1 ᐱᐅᓱᑐᖅᑎᒧᑦ ᓄᓇᒥᐅᑕᓂᔪᒃ ᐊᐅᓪᓕᕐᓂᓂᖅᓄᒃ 49
6.1.1 ᐱᐅᓱᓚᕆᒧᑦ ᐊᒻᒪᓗ ᐃᓄᕎᔨᑎᖅᓄᓪᓕᓪ ᐊᑐᖅᑕᐅᑦᖃᐃᓕᵒᓇᕎᓂᖅᓂᕐᓂ 49
6.1.2 ᐅᒪᔪᓕᖅᑲᓵᓪᓂ ᐱᓕᒪᓚᐅᔪᓂ ᐃᓄᖅᓄᓪᓕᓪ 50
6.1.3 ᐃᓄᐃᑦ ᑕᐅᔭᓂ ᑕᐃᔪᓱᐅᑐᖅᓂᕐᓂ ᓄᓇᒧᑦ ᐃᒪᓕᓪᓕᓪ 51
6.2 ᐅᒪᔪᓕᓚᓂᑎᓗᒃ ᐊᒻᒪᓗ ᐅᒪᔪᓕᖅᑲᓵᐅᔭᓂᖅᓄᓂᵒᓂᓗᒃ ᐊᐅᓪᓕᕐᓂᓂᖅᓄᒃ 51

6.2.1 ᐊᓯᔨᖅᑕᐅᓂᖏᑦ ᐃᓕᑕᒪᔭᐅᓯᒪᔪᖅ ᖃᓄᐃᔅᑐᓂᖏᓐᓄᓂ ᓴᓇᓂᐊᖅᓯᒪᕐᓂᐊᐅᓂᖅ.....51
6.2.2 ᐊᒥᓱᓗᐊᓕᕆᓂᖏᑦ ᑲᖏᐃᔅ.....52
6.2.3 ᒥᑎᖅᑲᓵᑦ.....53
6.2.4 ᐅᒪᔪᐃᔅ ᐊᒥᓱᖏᑦᑐᓗᐊᓕᕆᓂᖏᑦ.....53
6.2.5 ᑕᒧᓂᒥᐅᑕᑐᖅᐅᓕᖏᑦᑕᑦ ᐅᒪᔪᐃᔅ/ᖃᐅᔨᐱᕐᐃᔅᓗᓐᓂᑦ ᐱᑕᖅᑐᑦᓗᓐᓂᑦ ᐱᓕᕆᖅᓂᖏᑦ..53
6.2.6 ᐊᖏᐃᓇᓱᐊᖅᓂᖅ.....54
6.2.7 ᑎᖏᒥᓱᑦ ᐃᖏᕐᕕᒃᓂᖏᑦ ᑕᐅᖓᓂ.....54
6.2.8 ᑕᓚᐅᖏᒥ ᐅᒥᐊᖅᑐᕆᓂᖅ/ᐅᒥᐊᕐᔪᐊᖅᑲᕆᓂᖅ.....55
6.3 ᓇᐅᕐᓂᖅᑐᐃᓴᖅᑐᕆᐃᐅᕗᑦᑕᓂᖏᑦ ᖃᐅᔨᓴᕆᐃᐅᖓᒃᖏᓪᓗ.....56
6.4 ᐃᓄᖃᑎᑎᑭᐅᔪᓂᒃ ᖃᐅᔨᓚᑎᕐᓂᑎᓂᖅᓴᐅᓚᐊᖅᑲᕆᓂᖅ ᐊᓪᓚᓗ ᖃᐅᓚᓴᐅᓴᓚᐊᓕᕆᓂᖓᓂᒃ ᐊᐅᓕᕐᓂᐊᖓᓂᖅ.....57
6.4.1 ᓄᓇᓕᑦ ᐅᓴᐅᔨᔪᓂᖅᓴᐅᔪᓐᓇᖅᑐᓂᖏᑦ ᑐᑭᔨᐅᒪᔭᓚᐊᖅᑲᓂᖅᑐᓕᓪᓗ.....57
6.4.2 ᐃᓄᖃᑎᑎᑭᐅᔪᑦ ᐅᓴᐅᔨᔪᓂᖅᓴᐅᑕᐅᔪᓐᓇᖅᑐᖏᑦ.....58
6.4.3 ᐃᓕᓐᓂᐊᖅᑐᓄᑦ ᑐᕆᖓᒃᓱᑦ ᐱᓕᓚᐊᓴᓕᖅᑕᑦ.....59
6.4.4 ᓂᐅᕈᐊᖅᓂᒃ >ᓕᕆᕐᑐᓕᓚᓂᕆᒃ ᐱᓕᓚᐊᖅᑲᓂᖅᓴᖅ.....60
6.4.5 ᓄᓇᒥᑕᖅᑐᖅᓯᒪᔭᖓᓂᖏᑦ ᑕᕽᒪᕐᖓᓂᒃ.....60
6.4.6 ᐃᓄᖓᓄᑦ ᐱᕐᔪᑎᑎᑭᐅᔪᑦ ᓄᓇᐃᑦ.....61
6.4.7 ᒪᓕᓚᓴᐅᔪᓂᒃ ᒪᓕᓴᒥᐅᑲᐃᓴᖅᑕᕐᓂᖓᓂᖏᑦ.....62
6.4.8 ᐊᐃᔅᑐᖅᑐᓯᒪᔭᖓᓂᒃᓴᓂ ᖃᐅᓄᐃᓕᓕᐅᕈᐅᔭᖅᑐᑦ ᐱᑎᐊᖅᑎᑎᑕᐅᓯᒪᓕᓴᖅᑐᕕᓕᓗ.....63
7 ᐊᖓᑭᐅᕈᑕᐅᓯᒪᔾᔪᑎᒋᓕᑦ, ᐊᔪᕐᓇᖅᑎᑎᑕᐅᔪᑦ ᐱᓕᓚᐊᐃᐃᐅᔪᓐᓇᕐᓂᕐᔪᑦ ᐊᓪᓚᓗ ᐊᔪᖅᑎᑎᑕᐅᔪᑦ ᑎᑭᓯᒪᐃᐅᔪᓐᓇᕐᓂᖓᓂᒃᓄᓂᑦ.....64
7.1 ᐊᐅᓕᕐᑎᔨᓄᑦ ᐊᖓᑭᖅᑕᐅᓯᒪᔾᔪᑎᒋᓕᑦ.....65
7.1.1 ᐊᖓᑭᖅᑕᐅᖅᑲᑕᕐᓂᖏᑦ ᐱᔪᓇᐅᔨᑦᑐᑦ.....65
7.1.2 ᓄᓇᕗᓪᓕᒥ ᐃᓄᐃᑦ ᐱᓕᓚᔪᓇᕐᓂᖏᑦ ᐱᔪᓇᐅᔨᑎᖅᖓᓂᒃᓴᑎᑐᒃ.....66
7.1.3 ᓄᓇᕗᓪᓕᒥᐅᓂ ᐃᓄᐃᑦ ᐱᓕᓚᐊᓚᔪᓇᖅᑕᕐᓂᖏᑦ ᑭᓯᐊᓂ ᐱᔪᓇᐅᔨᑎᖅᖓᓂᒃᑎᑐᒃ.....67
7.1.4 ᐃᓄᓕᓚᐅᖓᓂᑕᐃᓂᑦ ᐱᓕᓚᐊᖅᑲᓇᓇᕐᓂᖏᑦ ᐱᔪᓇᐅᔨᑎᖅᖓᓂᒃᓴᑎᑐᒃ.....67
7.1.5 ᐃᓄᓕᓚᐅᖓᓂᑕᑦ ᐱᓕᓚᔪᓇᕐᓂᖏᑦ ᐊᖓᑭᖅᑕᐅᔪᓇᖅᑕᑦ ᑭᓯᐊᓂ ᐱᔪᓇᐅᔨᑎᖅᓴᓗᑎᑐᒃ 67
7.1.6 ᐱᓕᓚᔪᓇᕐᓂᖅ ᐊᖓᑭᖅᑕᐅᓯᒪᔪᑦ ᐱᔪᓇᐅᔨᑎᖅᓴᓗᑎᑐᒃ ᐃᓄᖓᓄᑦ ᐱᖁᔨᓚᐅᔪᑦ ᓄᓇᓂ 68
7.1.7 ᑲᓇᑕᒥ ᐱᔪᓇᐅᔨᑎᓂᒃ ᐱᓇᓯᐊᖅᑎᑕᑭᓇᕐᓂᕐᔪᑦ ᐱᐅᓯᓚᐅᔨᖅᑕᑦ ᐊᑐᖅᑕᐅᔾᔨᓚᐊᓕᓕᓂ..68
7.2 ᐊᓯᖏᑦ ᑲᓇᑕᐅᑉ ᒐᕐᒪᑲᑯᖏᓇᓄᑦ ᐊᓪᓚᓗ ᓄᓇᕗᓪᓕᒥ ᓄᓇᕗᒪᖏᓇᓄᑦ ᐊᖓᑭᐅᑕᐅᕐᖅᑐᑦ ᐊᓪᓚᓗ ᐱᔪᓇᐅᔨᑎᐅᔨᖅᑐᕐᔪᑦ.....69
8 ᓄᓇᖓᓂᑦ ᑎᖏᒥᐊᖅᑎᖅᓴᐅᑉ ᖃᐅᔨᓄᑕᐅᔪᖅᓴᐅᕐᓂᖓᓂᒃᓄᓂᑦ ᑕᐃᔨᐅᓂᖓᓂᑦ ᐃᓕᑕᒪᔭᐅᓯᒪᓂᖏᓐᓄᓗ.....71
9 ᐋᓐᓂᖅᑕᐅᑦᑕᐃᓕᒪᓕᑎᑦᑎᓂᖅ, ᐋᓐᓂᐊᖅᑐᓗᒪᓂᖅ ᐊᓪᓚᓗ ᐅᓗᒪᐊᓇᖅᑐᓗᒪᓂᖅ.....77
9.1 ᓄᖕᖓᓇᓗᔨᖅᑐᖅ ᐋᓐᓂᐊᓕᕆᑕᐅᔪᖅᓴᐅᕐᓂᖅ ᒥᑎᓄᓄᑦ.....78
9.2 ᓇᓄᐃᑦ ᑐᖅᑯᓕᐅᔪᖓᓂᖏᑦ.....78
10 ᐊᑐᓕᖅᑎᑦᑎᖕᖏᓕᒪᓂᐊᖅᓂᖅ.....79
11 ᐸᖁᓇᐅᑦᑐ ᐊᑐᓕᖅᑎᑕᐅᓂᖏᑦ.....81
11.1 ᐊᐅᓕᕐᑎᓂᖓᓄᑦ ᐸᖁᓇᐅᑦᑐ ᐋᖁᑭᐊᖅᑕᐅᔪᓇᖅᑕᐅᓂᖏᑦ.....84
11.2 ᐊᐅᓕᕐᑎᔨᐅᔪᓇᖅᑕᐅᓂᖓᓄᑦ ᑎᓕᔭᐅᓯᒪᔪᑦ.....84
12 ᐱᓕᓚᖅᑎᑎᑦᑎᐊᑭᓇᖅᑕᐅᓂᖅ.....85
12.1 ᐃᓄᐃᑦ ᐊᓪᓚᓗ ᑲᖏᒪᓄᑦ ᐱᓕᓚᖅᑎᑎᑦᔨᓇᖅᑕᐅᓂᖏᑦ.....85

12.2 ᓄᓇᕗᑦ ᓴᖅᓯᒪᔨᑦ 85
13 ᐅᖃᓕᒫᓂᐊᕆᔭᓯᒪᔭᖏᑦᑐᓂᒃ ᑐᑭᓯᓂᐊᖃᖅᐳᑎᓯᒪᔪᓕᒃᑕ 87
ᑎᑎᕋᖅᐃᑦ ᐃᓕᓴᖅᓂᖅᓯᕗᑦ 92
ᑎᑎᕋᖅᐃᑦ ᐃᓕᓴᖅᓂᖅᓯᕗᑦ A: ᒪᓕᒐᕋᖅᑎᒍᑦ ᐅᓂᒃᑲᖅᑕᐅᓕᐊᖃᓂᖅᓯᕗᑦ ᓄᓇᐃᑦ ᓇᓵᖅᓂᖅᓯᕗᑦ 93

ᑎᑎᕆᖅᑕᐅᓂᖏᑦ ᐊᐃᔪᐃᑐᒍᓂᒃ ᑕᑯᑎᑕᑎᓴᓯᕐᓂᑦ

ᐊᐃᔪᐃᖅᑐᒍᓂᒃ ᑕᑯᑎᑕᑎᓴᓯᑦ 1: ᖃᖅᓴᐅᖅᑰᖏᑦ ᑎᖏᑕᐊᖃᖅᓯᖏᑐᒃ ᓇᐃᓇᖅᑐᒍᓂᒃ ᖃᐅᔨᒪᑎᑕᑎᑎᕈᐊᑭᕆᑦ.....3

ᐊᐃᔪᐃᖅᑐᒍᓂᒃ ᑕᑯᑎᑕᑎᓴᓯᑦ 2: ᐊᐃᔪᐃᖅᑐᒍᓂᖓᓚᑕ ᖃᓄᐃᑕᖏᓐᓄᓂᖏᑦ ᐊᓕᒪᓗ ᐅᒪᔨᖃᖅᓂᓂᖏᑦ ᐊᓂᓗᐊᓂ ᓇᓗᓇᐃᓄᑦᑯᑎᑕᐅᓂᒍᓂᖏᑦ ᑎᖏᑕᐊᖃᖅᓯᓇᖏᑦ.....10

ᐊᐃᔪᐃᖅᑐᒍᓂᒃ ᑕᑯᑎᑕᑎᓴᓯᑦ 3: ᐃᓗᓗᓂᑯᑎᐅᔨᔪᑦ ᐱᖅᑯᑲᓖᓗ ᑎᖏᑕᐊᖃᖅᓯᓇᖏᑦ.....13

ᐊᐃᔪᐃᖅᑐᒍᓂᒃ ᑕᑯᑎᑕᑎᓴᓯᑦ 4: ᐅᒪᔨᐃᑦ ᐊᒥᓱᖏᓗᐊᓕᕐᓂᖓᓂᖏᑦ ᑎᖏᑕᐊᖃᖅᓯᓇᖏᑦ.....26

ᐊᐃᔪᐃᖅᑐᒍᓂᒃ ᑕᑯᑎᑕᑎᓴᓯᑦ 5: ᐃᓄᐃᑦ ᐃᓕᓴᐅᔪᑐᖃᖅᓂᖓᓂᒃ ᑕᐃᔭᓱᑦ ᓄᓇᓂᒃ ᐃᒪᐃᓕᓐᓂᑦ ᑎᖏᑕᐊᖃᖅᓯᓇᖏᑦ (ᐋᖅᑭᐊᖃᖅᑕᐅᑎᐊᖅᓂᖅᑐᑦ ᐃᓄᐃᑦ ᐱᐅᔪᑐᖃᖅᓂᒃ <<ᑎᑎᕐᓂᑦ ᑎᑎᖃᖅᑎᑦᓂᓂᖅ)).....33

ᐊᐃᔪᐃᖅᑐᒍᓂᒃ ᑕᑯᑎᑕᑎᓴᓯᑦ 6: ᐊᐅᓪᓚᑎᓂᔭᒍᑦ ᑐᖁᑎᓕᕆᐅᔨᔪᑦ ᐊᓕᒪᓗ ᐱᓴᖅᑕᐅᔨᔫᑦ.....39

ᐊᐃᔪᐃᖅᑐᒍᓂᒃ ᑕᑯᑎᑕᑎᓴᓯᑦ 7: ᓇᐃᓇᖅᑕᐅᔪᒪᔨᑦ ᐊᐅᓪᓚᑎᓂᖓᓂᖅ ᐃᔪᒪᓴᖅᐅᔪᕈᑎᖅᓴᐃᓪᓗ ᐊᓕᒪᓗ ᖃᓄᐃᓪᐅᔪᕈᑎᖅᓴᖓᓂᖅ.....46

ᐊᐃᔪᐃᖅᑐᒍᓂᒃ ᑕᑯᑎᑕᑎᓴᓯᑦ 8: ᒪᓚᑎᓴᐅᔨᔪᑦ ᖃᓄᐃᑕᖏᓐᓄᓴᑎᐊᖃᖅᓯᖓᓂᖅᓄᑦ ᑎᑯᔨᐊᖅᑕᐅᔨᐅᓇᖅᓯᐊᕈᓂ ᖃᖅᓴᐅᖅᑰᖅᓵᖅᑕᐅᓇᖅᑐᒥᒃ ᐅᒪᔨᖃᖅᓯᐃᐅᓂᖅᓗᑦ.....72

ᐊᐃᔪᐃᖅᑐᒍᓂᒃ ᑕᑯᑎᑕᑎᓴᓯᑦ 9: ᓇᐃᓇᖅᑕᐅᔪᒪᔨᑦ ᖃᐅᔨᓴᖅᑎᐅᓪᓚᐅᖅᑐᑦ ᓇᐅᑦᑎᖅᑐᐃᐅᔪᖅᑐᖃᑦᑕᖅᑭᒪᔨᓪᓗ ᑎᖏᑕᐊᖃᖅᓯᓇᖏᑦ.....74

ᐊᐃᔪᐃᖅᑐᒍᓂᒃ ᑕᑯᑎᑕᑎᓴᓯᑦ 10: ᑕᓪᓕᒪᓂᑦ ᐊᖏᖅᓯᒪᓂᑦ ᐊᐅᓪᓚᑎᓂᖓᓗᓕᑦ <ᖁᓇᐅᑎᓴᓂᒃ ᐊᑐᓕᖅᑎᑦᑎᓭᓕᒫᓂᖅ..82

ᑎᑎᕆᖅᑭᒪᓂᖓᓂᑦ ᑕᑯᔭᓴᕿᐃᐅᑦ

ᑕᑯᔭᓴᖅ 1: ᐃᓂᓪᓗ ᑎᖏᑕᐊᖃᖅᓯᐅᓪᓕ ᐊᓕᒪᓗ ᐃᓄᐃᑦ ᐅᖃᐅᔨᑐᖃᑦᑕᒃᑯᒍᑦ ᓄᓇᓕᑦ ᑕᐃᔭᐅᓂᖓᓂᑦ (ᐱᔭᐅᔨᒪᔨᑦ ᐃᓄᐃᑦ ᐱᐅᔨᑐᖃᖅᓂᖅᑐᑦ <<ᑎᑎᕐᓂᑦ ᑎᑎᖃᖅᑎᑦᓂᓂᖅ). 8

ᑕᑯᔭᓴᖅ 2: ᐃᓄᐃᑦ ᓄᓇᑯᓂᑎᔭᖓᓂᑦ ᐃᓗᐊᓂ ᑎᖏᑕᐊᖃᖅᓯᐅᓪᓕ. 12

ᑕᑯᔭᓴᖅ 3: ᓇᔨᓴᓚᐅᓪᓗᓗᐊᑕᖅᑕᖅᓂᖓᓂᖅᓂᖅ ᐊᐃᔪᐃᖅᑕᐅᔨᔪᒪᓂᖓᓂᑦ ᑎᖏᑕᐊᖃᖅᓯᐃᐅᓪᓕ (ᒪᓕᒃᑕᐅᔨᒪᓪᓗᑎᒃ ᐱᓕᕆᓚᐅᖅᑐᑦ F᐀ᓴᑦᑎᓵ ᐊᓕᒪᓗ ᐃᓗᓕᓇ-Fontaine and Mallory, 2011). 16

ᑕᑯᔭᓴᖅ 4: ᐃᓄᐃᑦ ᖃᐅᔨᒪᔪᑐᖃᖅᓂᖓᓂᑦ ᓇᓂᔨᕿᔪᓂᖓᓂᒃᓄᑦ ᑲᕿᐃᑦ ᑎᖏᑕᐊᖃᖅᓯᐃᐅᓂᖏᑦ (ᒪᓕᒃᑐᑦ ᑲᑕᑦ, ᐱᓕᕆᖃᑎᖅᓂᓪᓗ-Carter et al. 2018-ᒥ ᖃᐅᔨᓴᖅᓂᓂᓇᓂᖓᓂᒃᓂᖅ). ᐱᑕᖃᖅᓯᓇᐅᓂᖓᓂᑦ ᓴᖅᑭᐅᒪᔨᔫᑦ ᑕᒪᐃᓐᓂᖅᑐᑦ ᐃᓄᐃᑦ ᖃᐅᔨᒪᔪᑐᖃᖅᓂᖓᓂᒃ ᒪᓕᒃᑕᓴᑎᑐᒃ 21-ᓂᒃ ᐃᓄᖅᓂᒃ ᐃᓕᖃᓕᐅᖅᓯᒪᑕ.....19

ᑕᑯᔭᓴᖅ 5: ᑲᖅᓯᐃᑦ ᐅᓪᓗᖃᖅᓯᖓᓂᑕ ᓇᓂᖅᓂᖓᓂᑦ ᑐᑭᓯᐅᖅᑕᐅᓗᑎᒃ 2014-ᒥ ᓴᓪᓕᐅᑦ ᑭᑭᖅᑕᓗᐊᓂᒃ ᖁᔨᓕᐅᑎᓪᓗᑎᒃ ᖃᐅᔨᒪᔨᖅᑕᐅᓂᖓᓂᖅᓂ ᐊᑲᓕᐅᖅᑕᐅᓕᐅᖅᑐᑐᓂᒃ.....20

ᑕᑯᔭᓴᖅ 6: ᑕᐃᒪᖓᓂᒃ ᐃᓄᐃᑦ ᐊᑐᖃᑕᑎᓂᖓᓂᑦ ᓄᓇᒥᒃ ᓴᓪᓕᐅᑦ ᑭᑭᖅᑕᓗᐊᓂᒃ. 35

ᑕᑯᔭᓴᖅ 7: ᑕᐃᒪᖓᓂᒃ ᐊᓈᓇᐅᓕᖅᑐᖅᓓᓗ ᐃᓅᔨᔪᑎᑦᓴᖓᓂᒃ ᐊᖏᓇᓱᐊᖓᓂᑦ ᓴᓪᓕᖅᑎᐅᑦ ᑭᑭᖅᑕᓗᐊᓂ. 37

ᑕᑯᔭᓴᖅ 8: ᑎᖏᑕᐊᖃᖅᓯᒪᔪ ᐱᔨᓇᓕᐅᑎᑕᕋᓱᐊᖅᓂᖅᓕᑦ ᐅᐊᑐᑕᑦᑕᐅᑦ ᖃᓄᐃᑕᐅᑕᑕᐅᔪᓂᖓᓂᑦ. 70

ᑎᑎᕎᖅᓯᒪᓂᖅᓄᑦ ᓇᐃᓕᓕᑎᐊᖅᑕᐅᓯᒪᔪᑦ ᑎᑎᖅᑐᐃᑦ ᖃᖅᑎᓗᓀᑎᑐᑦ

ACMC	**ᐊᕐᓇᐅᑐᖅᓯᒪᓂᖅᓄᓐᓄᑦ ᐊᐅᓚᕋᑎᓕᐅᖅᑕᐅᑐᓂᔪᑦ ᑲᑎᒪᓕᕎᓪᓚᑦ**
COSEWIC	ᑲᓇᑕᒥ ᑲᑎᒪᓕᕎᓪᓚᖅᓄᑦ ᖃᖅᓄᐃᑦᓚᓂᖅᓂᖅᓄᑦ ᐊᒥᓱᔪᖅᓂᖅᑐᓪᓗᕈᓐᓇᕐᑕᐅᑐᓂᖅᓄᑦ ᐅᒪᔪᐃᑦ
CWA	*ᑲᓇᑕᒥ ᐅᐊᓪᔪᑕᓇᓂᖅᓪᓗᑦ ᒪᑕᑎᐃᑦ*
CWS	ᑲᓇᑕᒥ ᐅᒪᔪᑕᓇᓕᓐ ᐱᓕᕆᑎᕎᐃᓕᓂᖅᑐᑦ
IIBA	*ᐃᓄᐃᑦ ᐊᖅᑐᖅᑕᐅᓱᒪᓂᖅᓄᑦ ᐃᑲᔪᓱᐊᖅᑕᓪᓕᐅᑎᖅᓴᖅᑐᓄᑦ ᐊᖅᑎᕐᕈᑦ ᑲᓇᑕᓪᓕᒥ ᐅᒪᒃᓕᓂᖅᑕᐅᓱᒪᔪᓄᑦ ᓄᓇᕗᒻᒥ ᓄᓇᑦᓴᐃᐅᓱᒪᔪᓂᑦ ᓄᓇᖏᑎᖅᑐᑦ (ᐃᓄᐃᑦ ᐊᖅᑐᖅᑕᐅᓱᒪᓂᖅᓄᑦ ᐊᖅᑎᕐᕈᑦ)*
CIRNAC	ᑲᓇᑕᐅᑉ-ᓄᓇᖅᑲᖅᑐᑐᖅᐃᓕᓐᓂ ᐱᑕᓚᖅᑎᑎᑦᓴᖅᑐᑦ ᐅᑭᐅᖅᑕᖅᑐᒥᓂ ᐱᓕᑦᑕᔪᑕᓚᔪᑦ ᑲᓇᑕᒥ
DFO	ᐃᖅᑲᓂᕐᓚᕆᔪᐊᖅᑯᑦ ᑕᓚᐅᖅᒥᐅᑕᑕᓚᓕᑦᓂ ᑲᓇᑕᒥ
ECCC	ᐊᕐᓂᑎᑕᓚᕆᔪᐊᖅᑯᑦ ᓱᓕᐅᑦᓂ ᐊᓱᓕᖅᑕᑦᑕᖅᓚᖅᓴᖅᓄᑦ ᑲᓇᑕᒥ
GN	ᓄᓇᕗᑦ ᑐᕐᒪᐃᑦ
HTO	ᐊᖏᓇᓱᖅᑏᑦ/ᒪᖅᑲᐃᑏᑦ ᒥᑭᒧᐊᖅᔅᐊᖅᑎᑦᓂᓂ ᑲᑐᓕᖅᑲᑎᑎᕐᖅᑐᑦ
IHT	ᐃᓄᐃᑦ ᐱᐅᓱᑐᖅᖏᓪᓗᒃ ᑭᑭᑎᑎᓕᑦ
IOL	ᐃᖏᓄᖏᑦ ᐱᖁᓂᑎᕐᔪᐅᓪᓚᑦ ᓄᓇᐃᑦ
IQ	ᐃᓄᐃᑦ ᖃᑲᐅᓕᕐᔪᑐᖅᑐᑦ
KivIA	ᑭᓚᑦᓂᕐᒥ ᐃᓄᐃᑦ ᑲᑐᓕᖅᑲᑎᑎᑦ
MBCA	*ᑎᖏᒥᐊᖁᓗᒃ ᐅᑎᖅᑭᒃᑐᑐᓗᒃ ᑲᑎᓴᒪᐅᑕᐅᓗᑎᒃ ᒪᓕᑕᑎᓕᐊᖏᓕᓪᓚᑦ*
MBS	ᑎᖏᒥᐊᖅᑲᓇᐃᑦ
MBSR	*ᑎᖏᒥᐊᖅᑲᓇᖏᓄᑦ ᒪᓕᑕᑦᑕᑐᐃᓪᓚᑦ*
NIRB	ᓄᓇᕗᓴᒥ ᐊᓚᑎᑦ ᐊᒃᑐᖅᑕᐅᔨᓐᓇᖁᓕᖅᑐᓂᖅᓄᑦ ᑲᑎᒪᓕᐅᑦ
NA	*ᐊᖏᕈᑕᐅᓱᒪᔪᑦ ᐃᓄᖏᓄᑦ ᓄᓇᑦᓴᑲᑕᐅᓱᒪᔪᓂᑦ ᓄᓇᕗᓴᒥ ᑯᐃᓐᒧᑦᑕᓗ ᐱᔨᓐᓇᐅᑎᖅᑲᓂᖅᓄᑦ ᑲᓇᑕᒥ, ᑕᐃᔨᐅᓚᖏᓪᓕᔭᖅ ᓄᓇᕗᓴᒥ ᐊᖏᕈᑎᑎᓂᒃ*
NPC	ᓄᓇᕗᓴᒥ ᓄᓇᐃᑦ ᐊᑐᑎᒃᑲᖏᓐᓂᒃ ᑭᖅᓇᐃᓕᑦ ᑲᒥᓱᓇᐃᑦ
NTI	ᓄᓇᕗᑦ ᑐᓐᖏᓚᕕᒃᑯᑦ ᑎᒥᖅᑯᑎᖏᓕᑦ
NuPPAA	*ᓄᓇᕗᓴᒥ ᑭᖅᓇᐃᓂᔪᑦ ᐱᑕᓚᐊᒃᑲᓂᔪᓐᓇ ᖃᐅᓕᑲᓐᓇᖅᓂᖅᓂᔪᑦ ᒪᑕᑎᐃᑦ*
NWA	ᑲᓇᑕᒥᒻᒥ ᐅᒪᔪᑕᓇᖅᑲᖅᓯᐅᔨᖅ
NWMB	ᓄᓇᕗᓴᒥ ᐅᒪᔪᑕᓇᓕᓐᓂᔪᐊᖅᑯᑦ ᑲᑎᒪᓕᖅᑎᑦ
SARA	*ᐅᒪᔪᐃᑦ ᐊᒥᓱᓐᓂᑕᓂᓂᖅᓄᓐᓄᑦ ᒪᑕᑎᐃᑦ*

1 ᐅᖃᐅᓯᐅᑎᐊᖃᓂᒃᑕᒥᐅᖃᖅᑕᐅᓯᒪᔪᓂ ᓄᓇᕗᑦ

ᑕᒪᓐᓇ ᖃᐅᖅᓴᐅᖅᑐᑎᒥ ᑎᓂᒥᐊᖃᐅᖃᕐᓗᒃ ᐱᓐᔭᑎᖅᑕᐅᓕᐅᖅᓯᒪᔪᖅ 1959-ᖁᑎᓪᓗᔪ, ᑲᓇᑕᓕᒫᒥ ᑎᓂᒥᐊᓪ ᓂᕆᑭᓴᐅᓕᒃᑐᑦ ᑎᑭᓪᓚᕆᐊᑦ ᑲᑎᒪᖃᕐᑕᐅᓂᓂᓐᓄᑦ ᒪᓕᓕᒫᓂ, 1994-ᒥ (MBCA-ᒪᓕᑕᒪᓕᐊᖅᑎᔪᑦ) ᓴᓐᓂᐊᖅᑕᐅᑭᐊᖅᓯᑎᒃ ᐊᒥᓯᑦᖅᓯᐊᓚᐅᐅᔪᒪᓪᓗᑎᒃ ᑎᑭᑦᔪᑕᒫᒃ ᑎᓂᒥᐊᓪ. ᑎᓂᒥᐊᖃᖅᑐᕐᓗᒃ ᑕᒫᓂ ᑭᑦᓯᒃᑐᓗᓂᒃ 1,124 km^2 ᑭᓪᒥᑕᓂᒃ ᐊᓂᑎᓂᓕᒃ, ᓄᓇᖏᑐ ᐊᓚᒪᓗ ᑕᓚᐅᖓᓂᖏᑐ, ᐃᒪᖏᑐ, ᓴᓪᓕᐅᐸ ᖁᑭᖅᑖᓗᐊᓕᓕᒋᖅ, ᓄᓇᕗᒻᒥ (ᑕᑯᐅᓴᓴᖅ 1). ᑲᓇᖏᓇᖅᓴᓯᖏᑐ ᑎᒥᐊᖃᖅᑐᐃᐅᐸ ᑭᓪᓕᖃᖅᑐᐃᖏᑐᓂ ᑕᒫᖅᓯᒪᓂ ᐊᒡᖏᒪᑕ ᖃᐅᖅᓴᐅᖅᑐᑎᒃᓴᐸ ᑕᓚᐅᖏᓪᓂ ᐊᓚᒪᓗ ᐊᒥᑦᑐᐸ ᑕᓚᐅᖏᓪᓂ ᑭᓪᓕᖅᑎᑦ ᑎᑭᒥᐊᖃᖅᑐᐃᖏᑐᓂᐸ ᐅᑎᖓᓚᐊᖅᓯᒪᔪᖅ 60 ᑭᓯᒥᑕᓂᒃ ᓄᓇᖅᐱᐅᓂᔪᑦ, ᐃᓕᒋᕐᐅᓗᑎᒃ ᖃᐅᖅᑲᖅᖃᖅᑐᓂᖅᑐᓂᒃ ᑭᓕᖅᐃᓐᓇᖅᓂᒃ ᐱᕐᖅᐃᐅᓯᒪᓂᖅᑐᓂᒃ ᓄᓇᕗᑦ ᐊᓚᒪᓗ ᖁᔨᓂᖅᓴᓕᓂᖅᑐᓂᒃ ᓯᓚᖅᑲᓇᓯᐅᓂᒥᒃ ᐊᖓᔪᓚᐊᓂᑦᑐᓂ ᖃᐅᖅᑲᐅᖅᑐᐸ ᑲᑎᒪᖃᕐᑕᐅᓂᓂᓐᓄᑦ ᐅᑎᖓᓚᐊᖅᑐᐃᖏᑐᐸ ᑲᑎᒪᖃᕐᑕᐅᓂᓂᓐᓄᑦ ᑲᓇᐃᔪᐅᓚᐅᖅ, ᐅᖃᐅᓯᐅᑎᐊᐠᖅᑐᓗᓂᒃ ᑕᒪᓐᓇ ᑎᑎᖃᖅᑕᐃᓐ ᐃᓕᓴᐅᓕᐅᓂᒃᑐᓂᒃ 1-ᒥ. ᑲᓇᔪᐅᖖᓕᓚᐊᓕᖅᑎᓪᓗᑎᓂ ᐊᐅᓚᕆᑎᔨᐊᑎᖅᑕᓂᒃ ᑎᓂᒥᐊᖃᖅᑐᐃᖏᑐᐸ ᑭᓪᓕᖅᑎᑦ ᑕᓚᐅᖏᓪᓂ ᐊᓚᒪᓗ ᐃᓕᓴᐅᓕᐅᓂᒃᑐᓂᒃ ᑎᓂᒥᐊᖃᖅᑐᐃᖏᑐᐸ ᑭᓪᓕᖅᑎᑦ ᑕᓚᐅᖏᓪᓂ ᐊᒡᖏᒪᑕ ᖃᐅᖅᓴᐅᖅᑐᑎᒃᓴᐸ ᑕᓚᐅᖏᓪᓂ ᐊᓚᒪᓗ ᐃᓕᓴᐅᓕᐅᓂᒃᑐᓂᒃ ᑎᓂᒥᐊᖃᖅᑐᐃᖏᑐᐸ ᑭᓪᓕᖅᑎᑦ ᑕᓚᐅᖏᓪᓂ.

ᑎᓂᒥᐊᖃᖅᑐᐃᖏᑐ ᓄᓕᒥᐊᖃᖅᑐᐃᐅᓚᐅᖅᓴᖅ ᐊᐅᓵᑎᓕᑐᑎᐅᓚᐅᖅᓄᑦ (ᑐᑭᓕᐅᖅᑕᐅᓕᐅᓯᒪᔪᓂᒃ ᐅᖓᓚᑕᓂᒃᑐᓂᒃ 1% ᓴᓕᓂᑎᓂᒃ ᒥᑭᓂᖅᓴᐅᓕᔪᓂᒃ ᑲᖖᓚᖅ, ᓂᕆᓕᖅᓇᖓᒃ Atlantic brant-ᓂᒃ ᐊᓚᒪᓗ ᒥᑎᓂᒃ (ᐅᖓᓚᑦᓂᕆᕐᑕᐅᓕᐅᓗᑎᒃ ᐊᒥᓯᓂᖓᓕ ᑎᑎᕆᑕᐅᑕᖅᑲᓯᒪᔪᓂ 90-ᖓᕐᓂᕐᑐᓂ ᑎᓂᒥᐊᓕ, ᑕᐃᒃᑯᐊᓗ 9-ᖁᔨᓂ ᐅᒪᔨᐃᓕ ᐊᒥᓯᓂᖓᑐᓗᓕᖅᑐᓂ (ᒪᓕᑎᕐᔨᐊᖃᖅᑎᔪᓂ ᐅᒪᔨᐃᓕ ᐊᒥᓯᓂᖓᑐᓗᓕᖅᓂᕐᓂᖅᑐᓂᓂ ᒪᓕᑎᖃᖅᑎᔪᓂ ᐃᓕᓴᐅᓂᒃᑐᓂ COSEWIC-ᑯᓂ ᑲᑎᒪᔨᖅᑐᓂᓂ-ᑲᓇᑕᒥ ᑲᑎᒪᔨᕆᓵᖅᑐᓂᓂ ᖃᓄᖅᑲᐃᓕᖅᓯᒪᓂᖅᑐᓂᓂ ᐊᒥᓯᔨᓐᓂᖅᓴᕐᓕᐊᓂᖃᖅᑐᓂᒃᑐᓂᒃ ᐊᒥᓯᔨᖅᑐᓂᓂ ᐅᒪᔨᐃᓕ). ᖃᐅᖅᑲᐅᖅᑐᑎᒥ ᑎᓂᒥᐊᖃᖅᑐᐃᖏᑐᕐᑕᐅᖅ ᐊᑐᖅᑕᐅᓕᖅᑭᒥᔪᖅ ᐊᖃᖅᑐᓕᓚᐅᖅ ᐅᖓᓚᑕᓂᒃᑐᓂᒃ ᐃᓕᓴᐅᓂᒃᑐᓂᒃ, 13-ᖁᔨᓄᓂ ᐅᒪᔨᓄᓂ ᐱᐊᖃᖓᓕᖅᑐᓂᒃ ᐊᐃᒪᓕᑎᑎᕆᓚᕐᓗᒃᑐᓂᒃ ᐊᒥᓯᔨᓐᓂᖅᓴᕐᓕᐊᓂᒃ ᐃᓕᓴᐅᓂᒃᑐᓂᒃ ᐊᒥᓯᔨᓐᓂᖅᓴᕐᓕᐊᓂᒃ.

ᑕᐃᒪᖏᓂᒃ 2000-ᒋᖏ, ᖃᐅᐅᓴᓓᖅᑕᐅᓯᒪᔪᓕᒪᓐᐊᔭᓕᖅᑐᓓᖅ ᐃᒪᒃᐅᓂᖓᒋᐅᓯᒪᔪᓂ ᑎᓂᒥᐊᓪ, ᓇᐅᕗᐃᓐᓕᔭᓗ, ᒥᑏᓕᔭᓗ ᐊᒥᓯᔨᓕᔭᓗ ᐊᓯᖏᑎᓴᓂ ᓯᓚᕆᐊᖃᓴᕆᔨᓕᐃᓂ ᖃᐅᖅᑲᐅᖅᑐᑎᒃᓴᐸ ᑕᐅᓂᖅᑐᐅᑕᓄᑦ ᐊᒥᓯᔨᓐᓂᖅᓴᕐᓕᐊᓂᒃ ᐊᓯᖏᑎᓴᓂ ᐊᒥᓯᔨᖅᑐᓂᓂ ᐱᕆᔭᐃᓐᓂᖅᓴᕐᓕᐊᓂᒃ ᐊᒥᓯᔨᖅᑐᓂᓂ ᐃᓕᓴᐅᓂᒃᑐᓂᒃ ᐊᓯᖏᑎᓴᓂ ᐊᒥᓯᔨᖅᑐᓂᓂ ᐃᓕᓴᐅᓂᒃᑐᓂᒃ ᑎᓂᒥᐊᖃᖅᑐᐃᖏᑐᕐᑕᐅᖅ ᐊᒥᓯᔨᓐᓂᖅᓴᕐᓕᐊᓂᒃ ᐃᓕᓴᐅᓂᒃᑐᓂᒃ ᐊᒥᓯᔨᓐᓂᖅᓴᕐᓕᐊᓂᒃ.

ᑕᐃᒪᖏᓂᒃ ᑎᓂᒥᐊᖃᖅᑐᐃᖏᑐᕐᑕᐅᖅ ᐃᒪᒃᐅᓂᖓᒋᐅᓯᒪᔪᓂ ᑎᓂᒥᐊᓪ, ᓇᐅᕗᐃᓐᓕᔭᓗ, ᒥᑏᓕᔭᓗ ᐊᒥᓯᔨᓕᔭᓗ ᐊᓯᖏᑎᓴᓂ ᐊᒥᓯᔨᖅᑐᓂᓂ ᐱᕆᔭᐃᓐᓂᖅᓴᕐᓕᐊᓂᒃ ᐊᓯᖏᑎᓴᓂ ᐊᒥᓯᔨᖅᑐᓂᓂ ᐃᓕᓴᐅᓂᒃᑐᓂᒃ ᐊᓯᖏᑎᓴᓂ ᐊᒥᓯᔨᖅᑐᓂᓂ ᐃᓕᓴᐅᓂᒃᑐᓂᒃ ᐊᒥᓯᔨ

ᑎᖃᐊᖅᑲᓇᖅᓄᑦ ᒪᓕᒐᓪᓗᑕᓴᐃᑦ, ᑕᐅᖓᓂᑐᓐᓇᖅᓯᑎᑦᓗ ᑭᓯᐊᓂ ᐱᔪᖅᓂᐅᐅᑎᓂᖅ ᑐᓂᔅᐅᓯᒪᔪᑎᖅ ᐊᖓᑎᓕᓴᕆᐊᖅᑯᓐᓄᑦ ᓯᓚᐅᑦ, ᐅᒪᔪᓕᓴᕐᓂᖅᓄᑦᓂᖅ. ᒪᑯᓂᖅᑐ ᐊᔪᖅᑎᑕᐅᑎᑉᑐᑦ ᐊᑐᐊᖅᑲᓇᐃᖅᑲᓪᓚᑦ 1) ᑎᖃᐊᖅᓯᐅᑐᓇᖅᓄᖅ, 2) ᐸᖅᓚᕆᐃᔪᓐᓇᖅᓄᖅ, ᓯᕆᖅᑎᓇᔪᓐᓇᖅᓂᒃᑯᑦ ᑎᖃᐊᑦ ᒪᓐᖅᑎᓐᓂᖅ, 3) ᐱᓯᒪᔪᓐᓇᖅᓄᖅ ᐅᒪᔪᓂᖅ ᑎᖃᖅᓄᖅ, ᑎᒥᖅᑎᓂᒃᑯᑦ, ᐊᒥᖅᑎᓂᖅ, ᐅᑦᓗᖅᑎᓂᖅ, ᒪᓐᖅᑎᓂᖅ ᑎᖃᐊᑦ, 4) ᖃᓄᐅᓯᕆᐅᓐᓇᖅᓄᖅ ᐊᓯᖅᑎᓂᖅᓗᑯᑦ ᐊᖏᓇᓯᑐᑦᓯᐅᑎᖅ ᐊᑐᐅᓐᓇᖅᓄᖅ.

ᐊᕐᓚᑲᖏᑦᑐᕐᓂᒃ ᑕᐅᑎᑕᑎᓴᒃ 1: ᖃᖏᖏᓴᐅᖏᑲᒍᕐᓂ ᑎᓐᓕᐊᖏᖃᕐᓚᓐᓂᒃ ᓇᐃᓇᖏᑦᓂᓕᕐᓂᒃ ᖃᐅᔨᓴᒪᑎᑎᑕᓕᕐᑐᑦ


ᓴᓪᓕᐊᖏᑕᐅᓂᒪᖕᓚᓐᓄᓂ ᑎᒃᓂᐊᖏᑕᐅᓂᒪᓐᓇᑦ	ᑎᓐᓕᐊᖏᖃᕐᓚᒃ ᓴᓪᓕᐊᖏᑕᐅᓂᒪᓚᒃ
ᒪᑕᑦᓂᒃ ᒪᑕᔪᑕᐅᓪᓕᑦ ᑎᒃᓂᐊᖏᑕᐅᓂᒪᓐᓇᓚᓂᒃ ᓴᓪᓕᐊᖏᑕᐅᓂᒪᖕᓚᒃᓂᐊᕐᓂᑦ	ᐃᓂᖏᖃᕐᓚᐅᓪᓕᐊᖏᓴᓪᑦ ᐃᓂᓯᒪᓴᐅᓂᐊᖏᓴᓪᑦ ᓴᖅᑭᑎᑎᕐᖏᐅᓪᓕᐊᖏᓴᓪᑦ ᓴᓪᓕᐊᖏᑕᐅᓂᒪᖕᓚᒃ ᑎᓐᓕᐊᖏᖃᕐᓚᐅᓂᖏᓴᓪᑦ ᒪᓂᓐᓇᑦ ᒪᑕᓕᔪᑕᐅᑭᑐᒃ ᐊᑕᐅᓂᓯᓚᐅᔪᑎᑎᒃ ᐊᑕᐅᓂᓯᓚᐅᖏᓴᓪᑦ ᐊᒥᓂᓪᓕᖏᖃᕐᓚᐅᓂᖏᓴᓪᑦ

ᐅᑭᐅᖅᑕᖅᑐᖅᓯᓗᐅᓂᖏᑦ ᐊᐃᒃᑐᖅᓗᒥᓂᖅ	
ᐊᐃᒃᑐᖅᓗᒥᓂᕐᓂᑐᑦ	ᑭᓐᓕᖅ
ᐃᓕᐅᕈᓴᐅᓂᖏᑦ ᖃᐅᔨᑐᑎᑦᑐᑦ ᐊᒃᑐᖅᑕᐅᓗᒪᔪᑦ ᓄᓇᓕᓐᓂ	ᓴᓐᓕᖅ
ᓄᓇᖏᐊᑎᒍ ᓴᓂᒧᑦ ᐊᖏᓪᓗ ᑐᑭᓴᐅᑎᓂᖅ ᓇᓱᐅᓂᖅᓯᒪᓂᖅ	64°00’ N/82° 00’ W
ᐊᖓᕈᓂᖅ	1,124 km2,ᑭᐸᓴᓵᑐᓂᒃ ᑭᓕᒋᑕᓂᒃ, ᐃᒪᖃᖅᓯᑎᒃ 286 km2 ᑭᐸᓴᓵᑐᓂᒃ ᑭᓕᒋᑕᓂᒃ ᑕᓵᐅᖏᒥ ᐃᒪᔪᖃᖅᖁᐅᓂᖓᓂᓂ
ᐳᖅᑐᓂᖃᖅᑎᑎᓂᖅ (m ᒋᑕᒃᔪᑦ)	ᑕᓵᐅᐸ ᓇᑎᐊᓂᒃ ᑕᒪᐅᖅ ᖃᔪᓗᑦ ᐃᒪᖅᑎᑕ ᐳᖅᑐᓂᖅᐸᓵᓴᖅᓄᓇᓂᓂ ᑎᑭᐅᒪᓂᖅ ᖃᖅᑐᕈᓴᖅ 122 m ᒋᑕᑕᖅᓂᑦ
ᐊᖏᖅᔪᒍᒥ ᓴᖏᑭᑎᑕᐅᓯᒪᔪᑦ ᒪᑕᓚᑕᐊᑦ ᐅᖏᖅᑕᒻᒪᔪᖅᑎᒍᑦ ᓚᓯᔪᑦᔪᓕ-Gazette-ᔪᑦ)	1959
ᖃᐅᔨᐃᓕᓂᖅᓄᓇᓕᓂᑦ ᓴᐳᓂᖅᑕᐅᓯᒪᔪᑦ ᑎᑭᑦᑎᐊᖅᑐᒍᑦ ᓴᐳᓂᖅᑕᐅᓯᒪᔪᑦ ᑎᑭᑦᑎᐊᖅᑐᒍᑦ	ᑕᐃᒪᖅᓄᓂᒃ: ᓴᖏᑭᑎᑕᐅᓕᓕᐅᖅᐳᖅ ᓴᐳᓂᓂᐊᖅᑕᐅᖏᔪᓕᑕᑦ ᐃᓕᖓᐃᐅᓕᖓᓂᓄᓇᓂᓂ, ᐅᓕᓗᖏᖅᓯᐊᖅᑕᐅᖏᔪᓕᑕᑦ ᓄᓇᓕᓐᓂ ᑲᖏᖅᓂᓂᓄᓇᓂ Atlantic brant-ᓂᒃ (3% ᐳᓴᓂᑎᕐᓂᕐᑐᓂᒃ ᓄᓇᕋᐊᓕᒫᒥ) ᐃᓂᓇᐅᓕᖅᑐᖅᓄᖅ: ᓄᓇᓕᓐ ᐱᑕᖅᖏᓐᐅᕋᖅ 2% ᐳᓴᓂᑎᕐᓂᓵᓕᖅᑐᓂᒃ ᑲᓇᑕᐅᐸ ᓄᓵᓗᐊᑕ ᖃᑎᖅᑐᓴᐊᓵᓂᖓᓴᖅᑕᕐᑕᖅᑐᓂᒃ ᒥᑭᓂᖅᑎᐅᓂᖅᑕᐅᕋᓂᒃ ᑲᖏᖅᑕᐅᓂᖓᓂᓂ, 1% ᐳᓴᓂᑎᕐᓂᓵᓕᖅᓂᓗ ᓂᕈᓂᖅᓇᓂᓂ Atlantic brant-ᓄᓂ) (ᑕᐃᒪᐃᓇᓴᑎᐅᑐᖅᓄᐃᓇᖅᑐᓂᒃ) ᐊᒻᒪᓗ 5% ᐳᓴᓂᑎᕐᓂᓵᓕᖅᑐᓂᒃ ᒥᑎᖅᓂᒃ ᐅᑭᐅᖅᑕᖅᑐᒥ ᐱᑕᖅᓴᖓᓂᕐᓂᓵᓂᒃ
ᓴᐳᓂᖅᑕᐅᓯᒪᔪᓕᕐᓂᓵᓄᓂ ᐃᓗᐃᐃᕐᖅᑕᐅᓯᒪᓕᒡᔪᐊᑎᓂᕐᓂ	ᐊᐃᒃᑐᖅᓴᓕᕐᓗᒃ A-ᒥᒃ, ᖃᔪᕎᓂᑐᖅᑐᒥᒃ ᐱᑕᖅᑲᓚᐊᖅᑭᒪᓚᓂᕐᓂᓄᓇᓂᓂ, ᐃᓕᑕᓚᐅᐅᓪᓚᑎᓐᓂᕐ ᐅᒪᔪᓂᖅᓂᕐᓂᓵᓄᓂ ᓀᔪᖃᐃᓴᖅᓂᕐᓂᓵᓂᕐᓂᓚᓂᖅᓂᕐᓂᓄᓇᓂᓂᕐᓗᓂᓐᓂᓐᓂᕐ ᓴᐳᓂᖅᑕᐅᓯᒪᓕᒡᓗᑎᒃ
ᓄᓇᖅᔪᒍᒥ ᓄᓇᕐᓴᔪᒍᑦ ᑕᐱᓵᓂᕐᑦ ᓴᐳᓂᖅᔪᖅᓂᖅᓯᒪᑦ ᐱᐅᔨᑐᖅᓴᖅᑭᑐᐃᓚᐅᖅ IUCN-ᔪᑦ ᐃᓕᑕᓚᐅᐅᑎᑕᓐᓄᑦ	ᐊᐃᒃᑐᖅᓴᓕᕐᓗᒃ 1b (ᐅᒪᔪᖅᑭᐃᐅᖅᑎᑕᑐᓴᖅᓂᕐᓂᓄᓇᓂᓂ)

ᑲᖏᒪᐃᑦ ᒥᓂᑕᐃᑦ ᑲᑎᒪᔨᖅᑐᓄᑦ ᒪᓕᒐᓕᐊᖓᔪᒧᑦ ᓱᓴᐅᑉ	P.C. 1959-629 (SOR/74-514)
ᓱᓕᓇᐃᒃᑯᑦᑎᒋᑦ ᑲᓇᑕᐅᕐᓂ ᓄᓇᕗᑎᖅᑐᓂᒃ ᐱᖓᓐᓄᑦᑐᓂᒃ ᓱᓴᐅᑉ	ᐊᑐᖅᑎᑦᑐᖅ
ᐃᓄᖅᓄᑦ ᐱᖓᓐᓄᑦᑐᕐᓂᑦ ᓄᓇᕗᑦ (ᓄᓇᕗᑦ ᐊᐦᐃᓄᖅᑐᑦ ᐃᓕᓴᕐᓂᖅ ᐊᑐᖅᑐᓂᒃ)	ᐊᐦᐃᑐᖅᓯᒪᔨᑦ ᓱᓕᓯᒪᓂᖅ ᓄᓇᕗᑦ CH-13 ᖃᖅᓴᓂᑦ ᓄᓇᕗᑦ	ᓄᑎᓪᓗᑎᑦ ᐊᖏᑐᖏᑦ (km2) 1,734 km2 ᑭᓪᓕᔪᖅᑐᑦ ᑭᓇᒥᑕᐃᑦ
ᐊᓯᖅᓱᓂᖏᓐᓂᖅᑕᐅᔪᖅ ᑎᑎᖅᑕᐅᓯᒪᓂᖏᓐᓂᒃ	• ᖃᖅᑯᓯᐅᖅᑐᒍᑎᒋᔭᐅᓂᒃ ᑕᒪᐅᖅᓴᓂ ᐱᓕᒪᐅᑦᓂᖅ ᑎᖏᒥᐊᖅᑯᖅᓯᐅᓕᖅᓯᖅ (NU023) • ᐱᓕᒪᐅᑦᓂᖏᓐᓂᒃ ᑎᖏᒥᐊᖅᑯᖅᓴᕕᓂ ᓄᓇᕗᓴᒥ (ᓄᓇᖏᓴ ᐊᓕᐊᐅᑎᓪᓗᒃ 16) • ᐱᓕᒪᐅᑦᓂᖏᓐᓂᒃ ᓄᓇᕗᑦ ᑎᖏᒥᐊᖅᓄᑦ (ᓄᓇᕗᓴᒥ ᓄᓇᖏᓴ ᐊᓕᐊᐅᑎᓪᓗᒃ 44) • ᐱᓕᒪᐅᑦᓂᖏᓐᓂᒃ ᑎᖏᒥᐊᖅᑯᖅᓴᕕᓂ (ᓄᓇᕗᓴᒥ ᓄᓇᖏᓴ ᐊᓕᐊᐅᑎᓪᓗᒃ 24)
ᐊᓂᒃᑲᓂᖏᓐᓂᓂᐅᑎᑦᑕᐅᔭᖅ ᑎᒃᑯᐊᖅᑕᐅᓂᒋᓕᖏᓐᓂᒃ	• ᐊᖏᔪᕐᓂᖅ ᐱᓕᒪᐅᔨᓂᒃ ᑎᖏᒥᐊᖅᑯᖅᓴᕕᓂᓐᓇᖅᑐᖏᓐᓄᓕᓂ ᒪᓕᑐᕐᓂᒃ ᒪᓕᓕᑎᐊᖅᑯᓂ ᓄᓇᕗᓴᐋᖅᑎ: ᐃᒪᐅᓂᖅᑭᓯᓂ ᑎᖏᒥᐊᖅᑯᓂᖏᓐᓄᓕᓂ; ᓄᓇᕗᓴᐋᖅᑎ ᐱᓕᒪᐅᑦᓂᖅᑐᓂᖅᓯᑦ: ᑎᑭᓯᒪᓐᕕᖅᑯᕐᑕᐃᓇᓯᓂᖏᓐᓄᓕᓂᓗ ᐱᓕᒪᐅᑦᓂᖅᓯᑦ ᐅᒪᔨᓄᑦ • 5% ᐳᖃᐅᖏᑦᓂᓕᒪᒃᑕ ᐅᑭᐅᖅᑕᕐᑐᖅᑲᓯᔪᖅᑎᐅᑦ ᒥᑎᑦ ᓄᓕᐊᖅᑲᖅᑐᑦ ᑕᒪᓐᓂ • 2% ᐳᖃᐅᖏᑦᓂᓕᒪᒃᑕ ᑲᓇᑕᐅᕐᓂ ᖃᓄᖅᑲᓯᐊᓂᓯᓯᒪᓂᖏᓐᓂᒃ ᑲᖓᐃᑦ ᑕᒪᓐᓂ ᓄᓕᐊᖅᑲᕐᓕᒪᒃ • 3% ᐳᖃᐅᖏᑦᓂᓂᑦ ᓂᕐᑕᖅᓇᐃᑦ Atlantic brant ᑕᒪᓐᓂ ᓄᓕᐊᖅᑲᓕᐅᖅᑐᑦ • ᐱᓕᒪᐅᔨᖅᓴ ᐊᐅᓴᖅᑯᑦ ᖃᓄᖅᓱᓕᐃᑦ ᐃᓕᖅᑕᖅᑲᓯᓚᑦ, ᓄᓕᐊᖅᑲᓴᓯᑎᓕᓗ, ᓂᕐᓄᖅᓴᖅᓯᐅᖅᑲᖅᑲᓯᑎᓕᓗ ᑕᒪᓐᓂ • ᐱᓕᒪᐅᔨᖅᓴ ᓇᓄᕐᓄᑦ ᐃᓗᓕᖅᑲᖅᑎᐅᓂᖅᑲᓯᐅᓂᖅᓴᖅ/ᓯᑎᓂᖅᑲᖅᑎᐅᓂᖅᓴᖅ ᐊᐅᓴᖅᑯᑦᓕᓗ ᓂᕐᓄᖅᓴᖅᓯᐅᖅᑲᖅᑎᐅᓂᖅᓴᕐᓂᒃ ᓇᓄᕐᓄᑦ • ᐱᓕᒪᐊᓕᔭᕐᓂᖅ ᓇᑦᑐᑕᖅᑲᖅᑕᕐᑎᐅᑕᓄᑦ ᑐᖅᑕᓄᑦ ᓄᕐᓇᖅᑎᐅᓂᖅᓴᕐᓂᒃ ᐊᐅᓴᖅᑯᑦᓕᓗ ᓂᕐᓄᖅᓴᖅᓯᐅᖅᑲᖅᑎᖅᓴᕐᓂᒃ • ᐱᓕᒪᐊᓕᔭᕐᓂᖅ ᐊᐃᖅᓴᖅᑲᓯᔪᖅᓱᓂ ᓂᕐᓄᖅᓴᖅᓯᐅᖅᑲᖅᑎᓂᖅᓂᖅᓄᓕᓂ ᑕᒪᓐᓂ	
ᑕᒪᓂᕐᒥᐅᑕᑐᖅᑲᐅᓂᖅᑕᓄᑦ ᐅᒪᔨᓄᑦ/ᖃᐅᔨᐱᕐᓄᑦ ᐱᑕᖅᑲᖅᓯᐅᓕᖅᓴᓂ ᖃᐅᖅᑲᖅᑕᐅᓂᖅᑕᖅᑲᓯᐅᓕᖅᓴᓂ ᓯᓚᓕ	ᖃᐅᐅᔨᒪᐅᔨᖅᑲᖅᑐᖅᑕᖅᑐᖅ ᓯᓚ
ᐅᒪᔨᓄᑦ ᐊᒥᓯᖏᓐᓄᓪᓕᐊᓕᕐᓂᖏᓐᓂᒃ	ᐃᓕᑕᒪᐅᐅᔨᓕᖏᓐᓂᒃ ᑲᓇᑕᒥ ᐅᒪᔨᓄᑦ ᐊᒥᓯᖏᓐᓄᓪᓕᐊᓕᕐᓂᖏᓐᓂᒃ ᓄᓇᕗᓴᐋᖅᑐᓂᒃ ᓄᖓᒍᓐᓇᖅᑐᖅ:

	ᓇᐅᔨᓴᖅ ᓗᓪᓕᐊᓪᓕᐊᓂᖅ (ᓴᐅᕐᕋᓪᓕᒃ) <u>ᐅᓗᓚᐊᓇᖅᑐᒋᓕᖅᑐᖅ:</u> ᖃᐅᔨᒪᓂᖅ Bank swallow ᑐᖓᔪᖅᑐᕐᑲᓐᓇᖅᑐᑦ ᖃᐅᔨᒪᓂᕐ Barn swallow ᐊᑦᑯᐊᖅ/ᐊᑦᑯᐊᒃᓴᓗᒃ ᓇᐅᔨᓴᓐ ᑕᖅᓴᑐᖅᓯᖅᓴᖅ ᐊᑦᑯᐊᖅ/ᐊᕐᑯᐊᒃᓴᓗᒃ <u>ᐊᐳᐅᐅᖏᑕᐃᒥᒃ ᐃᓕᓴᒪᓗᑕᐅᔪᓂ:</u> ᑐᑭᒧᐊᒃᑐᓂᒃ ᑎᒥᖏᓚᔪᑦ ᑕᖅᓴᓕᓂᒃ ᐊᑦᑯᐊᖅ/ᐊᑦᑯᐊᒃᓴᓗᒃ ᑭᖑᓚᐃᐊᕐᓂᒃ/ᑲᔪᑦᖅ ᓇᓂᕐᖅ ᐊᐅᓪᓛᓕᓂᒃ ᓴᕐᕋᖅ ᒥᑭᵒᓂᖅᓴᖅ ᐅᒃᐱᒃ ᖃᖅᖁᖓ ᖃᐅᐅᔨᓴᖅᑕᐅᓕᒪᔪᓂ COSEWIC-ᑯᵒᓂᑦ ᑲᓇᑕᒥ ᑲᑎᒪᔨᕕᒋᖖᵒᓂᑦ ᖃᓄᓂᐃᓕᓐᖓᓂᖖᑭᵒᓂᑦ ᐊᒥᓗᐊᑭᵒᓂᓪᓚᕐᓕᐊᓂᕕᖅᑕᐅᓕᒪᔪᓂᑦ ᐅᒪᔪᐃᑦ <u>ᐅᓗᓚᐊᓇᖅᑐᒋᓕᖅᑐᖅᑐᑦ:</u> ᓇᓚᔨᖅᑐᖅᑕᐃᓕᖅᑎᐅᑦ ᑐᑦᑐᐃᑦ ᓂᐱᒃᓴ ᖃᑯᖅᓗᖅᑐᖅ <u>ᐊᐳᐅᐅᖏᑕᐃᒥᒃ ᐃᓕᓴᒪᓗᑕᐅᔪᓂ:</u> ᐊᐃᐃᓴᑦ ᖃᑭᓇᓕᓴᐃᑦ ᐊᖏᓴᑦ ᖃᑯᔪᓂᖅ Harris’ sparrow ᑐᐃᓕᓐᑦ ᖃᑭᓇᓕᓴᐃᑦ ᓇᖏᑲᑦ ᑕᓪᐅᐳᑦ ᓇᑎᐊᓂ ᖃᑲᓕᓕᓐᑦ ᑲᑭᑦᓕᐅᑎᖃᖅᑐᕐᑐᖅᑐᑦ ᖃᓴᖅᑐᑦ – ᐃᖃᓴᓕᔪᐊᖓᓂᖅᑕᐅᐴᖖᒥᔪᑦ, ᖃᑲᑦᓕᓇᓗᑎᑐᑦ ᑕᐃᔨᐅᔪᑦ -Thorny Skate–ᒥᓴ
ᐊᐅᓕᖖᑎᓕᐅᓂᖏᓐᓄᑦ ᑎᒥᖏᔪᑎᑭᔨᐃᑦ	ᐊᐴᑎᓕᓪᓚᔨᕕᐊᒃᑯᑦ ᓱᓕᐅᑦᓕᓗ ᐊᓱᓕᖅᑲᑦᓕᓕᓪᓚᔨᕕᓕᓪᓚᖖᑦ ᑲᓇᑕᒥ (ᑲᓇᑕᒥ ᐅᒪᔪᓕᓪᓚᓄᑦ ᐱᓕᖖᑎᑎᐃᓕᖖᑦ) ᐃᓕᑭᑦᓕᓗᓂᑦᑐᑦ ᐃᖏᓂᐅᔨᓴᖅᑕᐅᓕᒪᔪᓂᑦ ᐊᐃᒃᑐᖅᓱᓕᖖᑭᵒᓂᑦ ᐊᐅᓕᖖᑎᓕᐅᓂᖏᓐᓄᑦᑕᐅᓕᒪᔪᓂᑦ ᑲᑎᒪᔨᕕᓕᖖᑦ

ᐃᓄᖅᓄᑦ ᑎᑭᓯᒪᔪᖅᓴᖅᑎᓕᕐᓂᑦ ᐊᒻᒪᓗ ᐊᑐᖅᑕᐅᔪᖅᓴᖅᑎᓕᕐᓂᑦ	ᐃᓄᐃᑦ ᓄᓇᕗᒻᒥᐅᑦ ᐊᑭᖏᖅᖏᕐᕉᒥᒃ ᐊᒻᒪᓗ ᐃᓱᒪᖏᖅᓱᓱᖅᑎᒃ ᑎᑭᓯᒪᔪᖅᓴᖅᑐᑦ ᑕᒪᐅᖅᑲ ᓄᓇᒧᑦ ᐊᖏᖅᓴᖅᑎᓂᖅᓯᒪᔪᑦ, ᑎᑭᓯᒪᔪᖅᓴᖅᑕᐃᑎᒃ ᐃᒪᖏᓄᑦ ᐊᒻᒪᓗ ᑕᕆᐅᖅᑲᓄᑦ ᐃᓗᐊᓂᕐᕉᓂᒃ ᑎᖏᒥᐊᖏᖅᖃᕐᐳᑦ (ᒪᓕᒃᑕᐅᑦᓗᑎᒃ ᐃᓗᓕᖅᑎᑦ 5-ᖅᑲᓂᕐᕉᑦ ᐃᓄᐃᑦ ᐃᑲᔪᓯᐊᕆᖅᑎᓕᓄᑦ ᐊᖅᑎᕈᑎᖅᑎᑎᑦ) ᐊᒻᒪᓗ ᒪᓕᒃᑕᐅᔾᔪᑕᐊᖏᖅᕕᓗᑎᒃ ᐃᓗᓕᖅᑎᑦ 5.7.18-ᒋᑦᑐᑦ ᓄᓇᕗᒻᒥ ᐊᖅᑎᕈᑎᓂ). ᐱᔪᓐᓇᐅᑎᖅᕕᐊᖏᖅᕈᓐᓇᖅᑎᑦ ᐃᓄᐃᑦ ᕿᓇᐅᔾᔨᑕᐅᕈᑎᕆᔭᓂᒃ ᐱᑕᕕᓂᐊᕈᑎᒃ. ᐃᓄᒋᑕᕕᐅᓴᖅᑎᑐᑦ ᐊᑐᕈᓐᓇᖅᑐᑦ ᑎᖏᒥᐊᖏᖅᖏᖅᓱᖅᓂ ᐃᖅᑭᐊᖅᐳᐃᔾᔨᐅᑕᐅᓂᖅᓗᑦ ᐊᔪᖅᑎᓄᒋᓗ, ᐅᓪᓚᔪᓂᒃ ᕈᔪᐊᓂ ᐱᔪᓐᓇᐅᑎᖅᕈᑎᒃ, ᒪᑕᒃᑕᐅᒋᓗᑎᒃ ᑎᖏᒥᐊᖅᓂᒃ ᐅᑎᖅᑮᑐᑐᓂᒃ ᖃᑎᒌᒪᐅᑕᐅᓗᑎᒃ ᒪᑕᑐᐊᖓᔪᓂᖅᑐᑦ.	



ᑕᓂᕐᓴᓂᖅ 1: ᐃᓕᖅᑐ ᑎᖏᑕᐅᖅᑲᐃᔨᑦ ᐊᒻᒪᓗ ᐃᓄᐃᑦ ᐅᖅᑲᐅᓯᑐᖅᑏᑦ ᓄᓇᐃᑦ ᑕᐃᔨᐅᓕᖅᑎᓕᒃ (ᐱᔨᐅᓯᒪᔪᑦ ᐃᓄᐃᑦ ᐱᐅᓯᑐᖅᑲᓕᖅᑎᓕᒃ ᐊᑎᖅᑲᓇᑎᖅᑐᓂᖅᓴᓕᒃ).

1.1 ᐊᕕᒃᑐᖅᓯᒪᓂᖅᓯᑕ ᖃᓄᐃᑦᑕᓂᖅᓂᖅ

ᖃᐅᖅᓴᐅᖅᑐᑦ ᑎᖅᒥᐊᖅᖃᕕᖅ (ᓄᓇᖕᓴᐊᑎᔪᑦ ᑎᑎᖅᑲᓕᖅ 64°00’N, 82°00’W) ᐃᓂᓕᒃ ᓴᑦᑕᐅᑦ ᕿᑭᖅᑕᒋᐊᑕ ᓂᒋᖅᓯᓯᐊᑕ ᑲᓇᐃᖅᑐᓂᖅᓯᓂ, ᐊᖏᕐᓯᐊᒍᒃ ᕿᑭᖅᑕᑕᐅᖅᓯᒪᑦ ᐅᐊᖏᓇᖅᓯᓯᖏᓂ ᑕᒪᓐᓇ ᕿᐊᑦᓴᖏᐊᑦ. ᓴᑦᑕᐅᑦ ᕿᑭᖅᑕᒋᐊᐊ ᐃᓕᑎᕆᐅᐅᖅ ᑭᐅᑦᑕᖅᑎᐅᓄᑦ, ᓄᓇᕗᒻᒥ, ᑲᓇᑕᒥ. ᓴᑦᑕᖅ, ᑕᒧᓂ ᐅᖏᓯᖏᓂᑕᒃ 30 km ᑭᓕᒥᑕᓂᒃ ᑎᖅᒥᐊᖅᑲᓇᖅᑎᒃ, ᓄᓇᓕᑎᕆᐅᔨᑉᐅᖅ ᕿᑭᖅᑕᓂᖅᓯᒪ, ᐃᓄᖅᑲᖅᓯᓂᓗ ᑕᒧᓂᖅ 891-ᖅᑎᑦᑐᑐᓂᒃ (ᐋᓴᐃᔨᑦ ᑲᓇᑕᒥ, 2016 ᐃᓄᐃᑦ ᐊᒥᓯᓂᖅᑎᑦ).

ᑕᒪᖏᓇ ᑎᖅᒥᐊᖅᑲᓇᕐᓂᒃ ᐃᓕᓴᕆᔨᖅᑐᖅ ᖃᐅᖅᓴᐅᖅᑑᑦ ᑕᒪᐅᖏᓂᓂᒃ ᐃᒪᖅᒥᒃ, ᑲᖏᖅᓴᖅᑕᐅᓂᖅᓯᓂ 50 km ᑭᓕᒥᑕᓂᒃ ᑕᑭᓂᓂᖅᒥᒃ, ᐊᓗᒪᓗ ᑕᒪᐃᓐᓂᑦᓴᑎᒃ ᓄᓇᖅᑎᑦ ᐊᑎᑎᑉᑕᐅᓂᖅᑎᑦ ᖃᐅᖅᑲᒃᑲᖅᖏᓂᐅᐅᔪᑦ ᑎᑭᐅᒪᔪᑦ ᑐᓴᐅᒪᔪᑦ ᑐᓂᒋᖅᓯᓂᓄᑦ ᕿᑭᖅᑕᐅᓂᓄᑦ, ᐱᖏᖅᓇᖅᑕᕐᓯᐊᓂ ᑭᓯᓕᖅᑲᓇᕐᓂᒃ ᑎᖅᒥᐊᖅᑲᓇᐅᒃ ᐊᑕᑉᑐᑦ ᕿᑭᖅᑕᐅᓂᓄᑦ ᐃᓕᓴᕆᔨᖅᑐᖅᓯᓂᓄᑦ ᐊᓪᓚᖅ ᑕᒪᐅᖏᓂ ᐃᓗᐃᓐᓇᕐᓂᓄᑦ ᐊᑎᒍᓐᓃᑦ ᑕᒪᐅᖏᓂᓄᑦ ᐃᒪᓕᒧᑦ (Foxe Channel-ᒧᑦ).

ᓴᑦᑕᐅᑦ ᑲᓇᖅᑐᖅᑕᐊ, ᑎᖅᒥᐊᖅᑲᓇᖅᓱᒃ, ᓴᓇᘐᐅᓕᖅᓯᒪᓕᕐᓂᕆᓂᖅ ᐊᖅᑎᓕᕐᑲᒃ ᐊᖅᒪᓚᓗ ᐊᑲᑦᑕᖅᑕᐅᓕᖅᓯᒪᓗᐊᕐᓂᑦ. ᐃᓅᓴᐸᑎᑲᓂᒃ ᐊᖏᓇᓕᖅᑲᘐᖅᓂᖅᓱᒃ ᑲᔪᓯᓇᖅᓴᐅᓂᒃ ᑕᒥᓂᑦ ᐱᒻᒪᓇᑕᐅᔭᓂᖅᓱᒃᓄᑦ ᐃᓕᓴᕆᔨᖅᑐᖅᓯᓂᓄᑦ, ᐃᓅᓴᖅᑎᑭᖅᑐᓂᓄᑦ ᐊᖅᒪᓚᓗ ᐃᓅᓴᐸᑎᑲᓂᒃᑲᖅᖏᓂᓂᓄᑦ ᓴᑦᑕᐅᑦ (ᐅᓴᖅᑲᓕᖅᓯᓂᑕᑲᒃᑲᖅᑐᓂᑦ ᑎᑎᕋᖅᑕᐅᓂᖅᑐᖅ 3.2 ᐊᖅᒪᓚᓗ 3.3-ᒥ). ᐃᓅᓇᑦ ᓅᑦᐸᑐᑦ ᕿᑭᖅᑕᕐᓱᖅᓂ ᐱᓚᐊᖅᑐᑦ ᓴᑦᑕᕐᒥᓂ ᐅᑭᐅᖅᓂ ᐊᖅᒪᓚᓗ ᐊᐅᓴᖅᓂ ᕿᐊᑦᑕᖅᓂ ᐅᒥᐊᖅᓂᑦᓗ. ᐅᖏᓯᖏᓂᖅ ᓄᓇᖅᓄᑦ ᓄᓇᖅᓂᑲᑎᑕᓇᖅᑎᓕᖅᑕᐊᖅᑐᓂᖅᓄᓄᑦ ᓴᓇᔨᐅᖅᓴᓴᓴᖅᑲᖅᓴᖅᑎᑦᑐᓂ

ᐊᕕᐅᑐᖅᓯᒪᔨᓂᖅ ᑕᔪᑎᑕᑎᓴᔨᑦ 2: ᐊᕕᐅᑐᖅᓯᒪᓂᖏᓪᑕ ᖃᓄᐃᑦᓂᖏᓪᓂᖅᑎᑦ ᐊᒻᒪᓗ ᐅᒪᔨᖃᖅᑎᓂᖅᑎᑦ ᐊᓐᓇᓗᐊᓂ ᓄᓇᓕᒫᖅ ᐅᓗᓀᐃᐅᔪᑎᖅᑕᐅᓯᒪᓂᖅᑎᑦ ᑎᓐᓕᐊᖃᖅᓴᖅᓕ

ᐊᕕᐅᑐᖅᓯᒪᓂᖏᓪᑕ ᖃᓄᐃᑦᓂᖏᓪᓂᖅᑎᑦ	ᑲᓇᑕᐅᐸ ᓄᓐᓗᓗᐊᓂᒃ ᑕᐃᔨᓯᒪᓂᑐᖅᓯᒪᒐᒃ ᐅᓴᕋᖅᓯᐸᓪᓕᐊᓕᒫᖅᓂᖅᑎᑦ
ᓄᓇᖏᑕ ᑲᓇᑕᒥ ᐊᕕᐅᑐᖅᓯᒪᓂᖏᓕ	ᕼᐊᓪᓚᕆᓴᐸ ᐊᕕᐅᑐᖅᓯᒪᓂᖏᓕᓂᑦ ᐃᓕᑎᓴᐅᔪᖅ ᑯᐃᔨᐊᑦᑐᓗᐸ, ᒪᓂᑐᐸᒥ ᐊᕕᐅᑐᖅᓯᒪᓂᖏᓕᓂ
ᐊᖓᑎᒥᐅᑕᓂᑦ ᖃᓄᐃᙳᑐᖅᓂᖅᑎᑦ	2 ᒪᖅᑭᒃ ᐅᐊᖏᓇᖏᓂ ᐅᑭᐅᖅᑕᖅᑐᑐᓂᐅᐸ 3 ᐱᖏᓕᓪᑦ ᓂᒋᐊᖅᓂ ᐅᑭᐅᖅᑕᖅᑐᑐᓂᐅᐸ
ᓄᓇᖅᑦ ᐅᒪᔨᖃᖅᓂᖅᑭᓄᑦᓗ ᐊᔪᐃᖅᑭᓂᖏᓪᓂᖅᑎᑦ	2.5 ᐃᖅᑕᕿᐅᐸ-ᐃᓗᓕᐅᑦᓗ ᑕᓇᐅᕋᓗᐊ 3.2 ᑭᖕᓕᓕᐅᐸ ᐊᑎᑎᐅᑐᑎᐅᓂᖅᑎᑦ ᓄᓄᓇᐃᑦ
ᓄᓇᖅᑦ ᐅᒪᔨᖃᖅᓂᖅᑭᓄᑦᓗ ᐊᔪᐃᖅᑭᓂᖏᓪᓂᖅᑎᑦ	2.5.30 ᐅᖃᓄᔪᓕᕆᐸ ᐅᔪᕽᑦᓭᖓᔪᓕᐊᓗᐊ 3.2.46 ᓕᑦᓕᐅᐸ ᖃᑭᖅᑕᒍᐊ ᓇᐸᖅᑐᑐᖅᓯᒪᓂᖅ
ᓄᓇᖅᑦ ᐅᒪᔨᖃᖅᓂᖅᑭᓄᑦᓗ ᐊᖅᑐᖅᑕᐅᓗᐊᖅᑕᐃᓕᒪᔨᑦ (ᓄᓇᐃᑦ ᓄᓇᒥᐅᑕᖅᑦᓗ)	2.5.30.130 ᖃᖅᖅᓕᐅᖅᑐᑐᖅ ᑲᖏᑎᖅᑐᖓ ᖃᑭᑭᖅᑕᐃᑦᓗ 3.2.46.184 ᐊᖅᐸᖏᔨᐊᐸ ᑐᓂᖏᒥᐅᑦᓗ ᐊᔪᒥᖓᓕᒥ ᑕᓇᐅᖓᓕᑕ ᐃᓕᖓᓕ (Fisher Strait) ᖃᑭᖅᑕᐃᑦᓗ
ᑕᓇᐅᖏᒥᐅᑕᐃᑦ ᐅᒪᔨᖃᖅᕼᐃᓕᓴᐅᓂᖅᑎᑦ	ᐊᓱᓕᕆᓂᑦᑐᑎᓂᒃ ᕼᐊᓪᓚᕆᓴᐱᖓᓂᖅᑎᑦ

1.2 ᖃᓄᖅ ᐱᖓᑦᓂᐊᓕᐅᖏᓐᓇᑦ

ᐃᓄᐃᑦ ᖃᐅᐅᑎᑦᑎᓕᐅᖅᓯᒪᔨᑦ ᑲᖓᒍᓕᐊᑐᓂᖓᓂᒃ ᖃᖅᖅᓕᐅᖅᑐᑐᖅᓗ ᑕᐃᔨᓯᒥᖓ ᐊᓕᓐ ᑐᓗᒥᒃ-Alan Loughrey ᑲᓇᑕᒥ ᐅᒪᔨᓕᓚᓱᓂᒃ 1952-ᖓᑎᓪᓗᐃ. ᐅᒪᔨᓕᓚᓱᑦ ᖃᐅᐅᔨᕼᐊᓕᓕᓕᐅᖅᑐᑦ ᖃᖅᖏᓐᓄᓕᒃᑯᑦ ᓯᓇᑕᖃᖅᓂᖓᓂᓂᒃ ᑕᒧᓂ 1952 (A. Loughrey) ᐊᒻᒪᓗ 1955 (F. G. Cooch), ᑭᖓᔪᓕᐊᔪᑦ ᒪᖅᑭᖕᓂᒃ ᐊᖅᑎᔪᖖᓂᒃ ᓄᓇᒋᑦᓯᓗᑎᒃ ᖃᖅᐅᐅᔨᕼᐊᓕᐅᖅᑐᑦ (T.W. Barry). 1957-ᒥ ᑐᓂᓯᔨᖃᓕᐅᖅᑐᖅ ᑐᕈᓯᕆᐅᑎᓂᒃ ᓴᓪᓂᓕᐊᖅᑕᐅᖅᑎᑦᓗᐃᑦ ᐅᑦᓗᖅᑭᕆᐊᓗᓂᖖᓐᓇ ᐃᕐᖕᕆᐊᕐᕆᓂᖖᓐᓇᓂᒃ ᕿᑭᔪᑎᐅᓂᓕᓐᓂᖅᑐᖅᑎᑦ ᐃᓕᓴᕐᓂᖅᑐᑦ ᐊᒻᒪᓗ ᓂᑦᑦᔨᓇᓕᑦ, Atlantic brant ᐅᓴᖏᓂᐊᒃᓴᖅᓯᐅᑦᑭᐊᕐᕆᓂᖓᓂᖅᑐᒃ ᓴᓪᓂᓕᐊᖅᑕᐅᖅᑎᑦᓗᐃᑦᓗᓂᑦ ᐊᒻᒪᓗ ᓴᓕᑦᑎᐅᐅᖅᑐᓂᒃ ᑎᑭᓴᖅᑐᓂᒃ (Cooch & Barry, 1957). ᓱᖅᓕᒪᐅᓕᐅᖅᑐᑦ ᑕᐃᔨᓯᒪᓂ ᐃᓕᑕᖅᓯᓕᓕᐅᖅᑐᑦ ᑕᐃᓴᖖᓂᒃ “ᑲᐅᖅᓴᐅᖅᑕᐅᒥᒃ”-“Kouksauktow” ᖃᖅᑎᓗᓀᑐᑦ ᐃᓄᖖᓄᑦ ᑕᐃᓴᐅᓂᑕᖅᑕᐅᑎᓗᓂ ᑕᐃᒪᓐᓇᒃ, ᖃᖅᖅᓴᐅᖅᑐᑐᖅᓂᖅᑎᓄᑦ ᑕᐅᖖᓐᓇᓗ ᑕᓯᐅᓴᕐᔪᐊᖓᖖᓐᓇ. (Cooch & Barry, 1957). ᒪᐃ 21, 1959-ᒥ, ᖃᖅᖅᓴᐅᖅᑐᑐᖅ ᑎᑎᕈᐊᖅᑭᓴᒃ ᐱᖓᖓᑎᐅᖅᑕᐅᑎᓕᐅᖅᑐᖅ ᒥᓵᑕᐅᔪᑦ ᑲᑎᒪᓴᖏᓐᓄᓄᑦ, ᑲᓇᑕᒥ ᒪᑕᓚᑕᐅᖅᓯᓗᒃ, (P.C. 1959-629).

ᑎᑎᕆᖃᐅᔾᑲᓕᐅᖅᑐᑦ ᐊᐅᓪᓕᐅᓂᖅᓴᓄᑦ <ᕆᓇᐅᑦᑐᑦ 1986-ᖓᑎᓪᓗᑐ, ᓯᐅᓂᐊᒍᑦ ᓄᓇᕗᒻᒥ ᓄᓇᒋᓪᓕᐅᖏᓂᖅᓴᓂᖅ ᓯᓗ ᐊᒻᒪᓗ ᐊᑎᓕᐅᖅᑕᐅᓪᓕᐅᖏᓂᖅᓴᖏᓂ ᓯᓗ ᐃᓄᐃᑦ ᐃᑲᔪᓯᐊᓴᖏᓂᓄᑦ ᐊᖏᕈᑎᑦ (Stephenson & McCormick, 1986).


1.3 ᓄᓇᕗᑎᑎᕐᕆᐅᓂᖅᓴᓂᑦ ᐊᒻᒪᓗ ᐊᑐᖅᑕᐅᔪᓐᓇᕐᓂᖅᓴᓂᑦ

ᑕᒪᓐᓇ ᓄᓇᖓ ᐊᖏᕐᓂᖅᓴᐅᔪᖅ ᑕᒪᐃᓐᓂᖅᓴᖅ ᐃᓄᓐᓄᑦ ᐱᕐᓂᑎᑎᕐᕆᐅᔪᕆᒪᓕᑦ ᓄᓇᒥ. ᐊᑕᐅᓯᖅ ᐃᖏᒥᔨᖓᓴᖅ ᓄᓇᐅᓂᖓ ᑕᐃᓂ ᓇᓗᓇᐃᑲᓯᑕᓗᒃ ᑎᖏᒥᐊᖅᑲᐊᖏᒥᒃ, CH-13, ᐃᓄᖓᓄᑦ ᐱᔪᓇᐅᑎᖅᑲᐊᐅᔪᑐᒍᖅ ᖃᖓᓴᓂᑐᓇᐅᓂᖏᓐᓄᑦ ᑕᐃᓂ ᓄᓇᐅᑦ (ᑕᓂᕐᓴᖅ 2). ᑲᓇᑕᐅᑦ ᓴᖅᒪᑲᑎᖏᓐᓄᑦ ᓄᓇᕗᑎᑎᕐᕆᐅᓂᕆᖅᑕᐅᔪᑦ ᐃᓯᕗᖅᑐᓕ ᑕᐅᖅᓂ ᓂᑎᐊᑕ ᐱᖓᓐᓇᖅᑰᓯᔨᖓᓂᒃ ᐃᑭᕆᖓᖓ/ᑎᓪᓗᓂᔨᖓ ᑎᖏᐊᖅᑲᐊᐅᑦ, ᕿᑭᖅᑕᒋᔭᑦᓗ. ᐃᓄᖓᓄᑦ ᐱᕆᓂᑎᑎᕐᕆᐅᔪᑦ ᓄᓇᖏᑦ ᐊᐅᓪᓕᐅᑦᓗᑎᒃ ᑕᐃᒃᓂᓗᑎᖓ ᑭᖅᓪᓕᒥᒃ ᐃᑲᔪᖅᑎᑎᒃ ᐃᒃᓴᓴᓕᑎᑎᒃ ᐃᒃᓂᖅᑎᐅᑦᓗᑎᒃᓗ ᑕᒪᐃᓐᓄᒃ ᐃᓄᖓᓄᒃ. ᑲᓇᑕᒥ ᒥᓂᑕᐅᔪᖅ ᒪᖅᒪᓕᖏᓐᓄᑦ-ᐃᓄᐃᑦᓗ ᓄᓇᓕᑐᖅᑲᐃᑦ ᑲᓇᑕᒥ ᐱᓕᓐᓂᖅᑎᑎᕆᔨᓐᓇᖏᓐᓄᑦ ᐊᒻᒪᓗ ᐅᑭᐅᖅᑕᖅᑐᒥ ᐱᖅᓯᓕᐊᔪᓕᓐᓂᕐᓗᑦ ᑲᓇᑕᒥ, ᒪᓕᒃᑕᐅᑎᒃ *ᑲᓇᑕᐅᑦ ᐅᑭᐅᖅᑕᖅᑐᖅᓴᓗᐅᔪᓂᖅᓴᖅ ᓄᓇᐃᑦ ᒪᓕᓐᓄᖅ (ᓄᓇᕗᒻᒥ),* ᑎᐃᒥᐊᖅᑎᑎᕐᕆᑦ ᑕᒪᐃᓐᓂᕐᑎᐊᖅ ᓄᓇᐅᑦ ᐃᓗᐊᓂᑐᑐᓂᒃ ᐱᔪᓐᓇᐅᑎᓂᒃ.

ᑲᓇᑕᒥ ᐅᒪᔨᔪᓕᓐᓄᓕᒫᑦ ᑲᒪᔨᐅᕐᓂᑦ ᐊᐅᓪᓕᔪᕐᓂᑎᓂᕐᒥᑲᓗ ᑕᐅᖅᓂ ᑎᖏᒥᐊᖅᑲᐅᓂᕐᒥᓴᓗᒃ, ᐅᑦᓗᖏᓂᖅᓴᒃ, ᒪᓂᖅᓴᒃ, ᓇᓂᑐᐃᓐᓇᕆᔨᓗᒃᑭᑎᒃ, ᐊᒻᒪᓗ ᑎᖏᒥᐊᖅᑲᐃᐅᓂᖓᓴᓂᒃ ᑲᓇᑕᐅᑦ ᑲᖏᒪᐃᑦ ᓄᓇᖏᑦᓴᓂᑐᓂᒃ. ᐃᓄᖓᓄᑦ ᓄᓇᕗᑎᑎᕐᕆᐅᔪᓴᓂᑐᓂ ᑎᖏᒥᐊᖅᑲᐃᖅᑲᓇᑦ ᓄᓇᖏᑦᓴᓂ ᐊᐅᓪᓕᐅᑕᖅᑐᑐᓂ ᑭᖅᓪᓕᒥᒃ ᑲᓇᑕᒥ ᐃᓄᐃᑦ ᑲᑐᓴᖅᑲᐊᑎᖅᑲᓂᕐᓂᖅᓴᓂᑦ.

ᐃᓄᐃᑦ ᓄᓇᕗᑎᖏᓂᑦ ᐃᓯᓕᖕᒪᖅᑲᖅᓴᓂ ᐃᒪᐅᑦ ᑕᓚᐅᑦ ᐃᑎᓴᓂᖓᓴᓂᒃ ᕿᑯᖕᒪᓯᖓᓂᖅᓴᖓᓴᓂᒃ (*NA s.19.8.13*). ᑕᓚᐅᖏᑦ ᓇᑎᖅ ᐃᖅᑲᖅ ᐃᒪᐃᓕᓗ ᐅᖓᓕᑕᓂᑐᓂ ᐃᓄᐃᑦ ᓄᓇᕗᑎᖓᓕ ᐱᑎᕆᐅᓂᖅ ᑲᑕᐅᑦ ᑕᓚᐅᖅᓂ ᑭᖅᓪᓕᒥᒃ.

ᐃᖅᑲᓗᓕᓚᔨᔪᐊᑲᑯᑦ ᑕᓚᐅᖏᑎᐅᑕᓕᓚᔨᓕᓗ ᑲᓇᑕᒥ ᓂᐳᐊᓕᓚᐅᖅᑐᑦ ᓯᓴᖅᑰᓯᔨᐅᓂᖓᓴᓂᒃ ᐃᒪᖏᓂᒃ ᐊᖕᒍᓗᐊᓂ ᓴᓪᑦᑎᑦ ᕿᑭᖅᑕᒋᔭᐊᓂᒃ ᐊᖕᒍᓗᐊᓂᒃ ᐃᓕᐅᔨᒪᓂᖅᓴᓂᑦ, ᑕᐃᒪᐃᒪᑦ ᑕᒪᐃᓐ ᐱᑎᐊᖅᑲᐃᐅᓂᖅᓴᖅ ᑕᓚᐅᖅᓂᒃ ᐃᓄᐃᑦ ᑲᖏᒪᐃᑦ ᐅᖃᐅᓯᒪᓂᖅᓴᓂᑦ ᐱᖓᓐᓇᖅᑐᓂᒃ ᑎᖏᒥᐊᖅᑲᐃᐅᓂᖅᓴᖅ, ᕿᑭᖅᑕᒋᔭᐊᓂᒃ ᓄᓇᕗᑎᑎᕐᕆᐅᔪᓴᓂᑐᓂᒃ ᐃᓄᐃᑦ ᑲᖏᒪᐃᑦ ᐅᖃᐅᓯᒪᓂᖅᓴᓂᑦ ᑲᑎᒪᔨᖓᓂᒃ ᐃᓄᐃᑦ ᑲᖏᒪᐃᑦ ᐅᖃᐅᓯᒪᓂᖅᓴᓂᑦ



ᑕᑯᔭᕆᔨᖅ 2: ᐃᓄᐃᑦ ᓄᓇᕋᑎᑕᔭᖏᑦ ᐃᓗᐊᓂ ᑎᖅᒥᐊᖅᑎᐃᔭᐅᑦ.

1.4 ᐃᓚᓪᓯᐊᑎᑭᕆᐅᔪᓂ ᐱᓯᓂᑦᓗ

ᐱᑕᖃᖅᓴᖅ 9-ᓂᒃ ᒪᓕᒪᒃᓂᑦ ᐱᓯᓂᑎᕐᓂᓂᒃ ᑎᖏᑕᐃᖃᖅᓯᖏᖓ (ᐊᐃᒃᑐᖅᓭᒪᔪᓂᒃ ᑕᓂᑎᑎᕆᔪᓂ 3). ᐊᕿᑎᓕᓚᓪᓕᔪᐊᒃᓂᑦ ᒪᓪᑐᐃᓐᓇᖅᑐᖅ ᐊᓂᓂᑕᓕᓕᓚᖅᑐᒃ ᖃᐅᔨᓪᓕᐊᓂᖅ ᖃᐅᔨᓴᖅᑎᑦᑎᓐᓂᒃᑯᑦ <ᑦ ᓴᒥᑦ-Paul Smith (ᐅᒪᔪᓕᓚᓪᒃᓂᑦ ᓄᓇᒥᓗ ᖃᐅᔨᓴᖅᓯᖏᓐᓂᒃᑐᒃ, ᖃᐅᔨᓴᖅᓂᒃᑐᖅ ᓴᓇᖏᑐᑎᓕᓚᓂᖅ) ᐊᖏᑭᐅᒪᑎᑕᑎᐅᕐᑐᑐᒃᑯᑦ ᐊᐅᓕᕆᑎᓗᐊᖅᑯᓗᓗᓗ ᓄᓇᒥ ᑕᖏᒪᓯᖏᒥᒃ. ᓴᓗᖅᓴᓴᖏᓕᓴᐃᔪᓂᒃ ᖃᐅᔨᓴᖅᑎᐅᔨᓗᐊᖅᓴᓂ ᑕᐅᐅᓂ. ᓴᑕᒪᐅᓂᑦ 4 ᐊᑐᖅᑕᐅᖃᕆᑕᐃᓇᖅᑐᑐᑦ ᐱᓯᓂᑦᓗ; ᐃᓚᓪᓕᓪᓕᖅᑐᒃ ᑕᖏᒪᓯᖏᒃ, ᓴᓪᓗᐊᖅᑐᒃ, ᖃᐅᔨᓴᖅᖏᒃ, ᐊᓚᒪᓗ ᐊᓇᖏᒃ. ᑕᖏᒪᓯᖏᒃ ᖁᑭᖅᑕᒪᓗᓐᓂ ᐊᑐᖅᑕᐅᔨᓗᐊᖅᓴᐅᑐᖅ ᖃᐅᔨᓴᖅᓂᕐᓂᑦ ᒥᑎᓂᓂᒃ. ᐱᓯᓂᑦ ᖁᑭᖅᑕᒥ ᐃᓕᓴᖅᑐᖅᑯᑦ ᑕᐅᓕᕆᑎᓗᐊᖅᑐᒃ, ᖃᐅᔨᓴᖅᖏᒃᑐᒃ ᐃᓚᓪᓕᓪᓕᖅᑐᒃ, 7-ᖃᓪᓚᓂᒃ ᑕᓕᓴᐱᖏᓕᓂᒃ, ᐊᓚᒪᓗ ᐊᓇᖏᒃᑐᒃ. ᖃᐅᔨᓴᖅᑎᐅᔨᒍᑎᓕᒃ ᔪᑲᓂᑦ ᑭᐅᓂᒃᑯᕐᓂᑦ-Dr. Grant Gilchrist (ᐅᒪᔪᓕᓚᓪᒃᓂᑦ ᓄᓇᒥᓗ ᖃᐅᔨᓴᖅᖏᓴᓂᒃ, ᖃᐅᔨᓴᖅᓂᒃᑐᖅ ᓴᓇᖏᑐᑎᓕᓚᓂᖅ) ᑲᒪᔨᐅᔪᖅ ᖁᑭᖅᑕᒃᑯᓪᓕᒃᑐᒃ ᐱᓯᓂᑎᓂᒃ.

ᐊᖏᑭᐅᖅᓴᖅᑎᑦᑎᓂᒃ ᑕᓂᑎᑎᕆᔪᓂ 3: ᐃᓚᓪᓯᐊᑎᑭᕆᐅᔪᓂ ᐱᓯᓂᑦᓗ ᑎᖏᑕᐃᖃᖅᓯᖏᖓ


ᓴᓇᐅᓂᖏᑦ (ᐱᓯᓂᑦ)	ᖃᑲᓇᐃᓂᖏᑦ	ᐊᖏᑦᓂᖏᑦ ᐅᐸᑐᕐᑎᑲᓂᑦ Fᐃᑦᓂᑦ	ᖃᖏᒐ ᓴᓇᔨᐅᓂᓂᑦ	ᑲᒪᔨᐅᔪᓂ ᐱᓯᓂᑎᓂᒃ	ᓇᓂ
ᑕᖏᒪᓯᖏᒃ ᐃᓚᓪᓕᓪᓕᖅᑐᒃ - ᓄᓇᒋᕐᖏᒐᖏᓂᓂ	ᐱᐅᔪᖅ	16 x 24	2008	ᐊᖏᑎᓕᓚᓪᓕᔪᐊᒃᓂᑦ	63° 59' 14", -81° 41' 44"
ᓴᓪᓗᐊᖅᑐᒃ ᐃᓚᓪᓕᓪᓕᖅᑐᒃ - ᓄᓇᒋᕐᖏᒐᖏᓂᓂ	ᐱᐅᔪᖅ	12 x 16	2004	ᐊᖏᑎᓕᓚᓪᓕᔪᐊᒃᓂᑦ	63° 59' 14", -81° 41' 44"
ᐃᑲᖏᒃ ᐃᓚᓪᓕᓪᓕᖅᑐᒃ - ᓄᓇᒋᕐᖏᒐᖏᓂᓂ	ᐱᐅᔪᖅ	16 x 24	2013	ᐊᖏᑎᓕᓚᓪᓕᔪᐊᒃᓂᑦ	63° 59' 14", -81° 41' 44"
ᐊᓇᖏᒃ – ᓄᓇᒋᕐᖏᒐᖏᓂᓂ	ᐱᐅᔪᖅ	4 x 4	2008	ᐊᖏᑎᓕᓚᓪᓕᔪᐊᒃᓂᑦ	63° 59' 14", -81° 41' 44"
ᑕᖏᒪᓯᖏᒃ ᐃᓚᓪᓕᓪᓕᖅᑐᒃ – ᖁᑭᖅᑕᓂᓗᓐᓂᒥ	ᐱᐅᔪᖅ	16 x 24	2001	ᐊᖏᑎᓕᓚᓪᓕᔪᐊᒃᓂᑦ	64° 01' 47", -81° 47' 17"
ᖃᐅᔨᓴᖅᓯᖏᒃ ᐃᓚᓪᓕᓪᓕᖅᑐᒃ – ᖁᑭᖅᑕᓂᓗᓐᓂᒥ	ᐱᐅᔪᖅ	16 x 24	2010	ᐊᖏᑎᓕᓚᓪᓕᔪᐊᒃᓂᑦ	64° 01' 47", -81° 47' 17"
ᐊᓇᖏᒃ – ᖁᑭᖅᑕᒪᓗᓐᓂᒥ	ᐱᐅᔪᖅ	4 x 4	1998	ᐊᖏᑎᓕᓚᓪᓕᔪᐊᒃᓂᑦ	64° 01' 47", -81° 47' 17"

ᓱᓇᐅᓂᖅᓯᒪᓕᒃ (ᐱᖁᑎᒋᑦ)	ᖃᓄᐃᓂᖅᓯᒪᓕᒃ	ᐊᖏᕐᓂᖅᓯᒪᓕᒃ ᐅᓴᖅᑐᐅᑎᒃᑯᓕᒃ Fᐃᑦᑯᓕᒃ	ᖃᓄᖅ ᓴᓇᐅᔨᓂᑯᓕᒃ	ᑲᒪᔨᐅᔪᑦ ᐱᖁᑎᓂᒃ	ᓇᓂ
7-ᖏᒧᑦ ᑕᓕᖏᓚᐃᒃ ᖃᑎᐅᔨᕐᖁᐃᑦ - ᖃᑭᖅᖃᑕᐅᔪᖏᒥ	ᐱᐅᔪᖅ	4 x 4	1998-2012	ᐊᐊᖏᑎᓕᓗᔨᕐᔪᐊᒃᑯᓕᒃ	64° 01' 47", -81° 47' 17"
ᑐᖅᑯᐃᐃᓱᒪᓚᒃ ᖃᑭᔪᖁᑎᑎᑐ – ᑲᖏᕐᕆᖁᕐᖏᒥ ᓄᓇᒥ	ᐱᐅᔪᖅ	4 x 8	Unknown	OMNR	63° 57' 35", -81° 51' 17"

2 ᐊᕐᑎᓗᒃ ᐊᕐᑎᑎᐅᑕᖅᓂᒃᑐᓕᒃ

2.1 ᓄᓇᒥᐅᑕᐃᓕᒃ ᑕᓚᐅᖏᒥᓗ ᐃᒪᖏᐅᑕᐃᓕᑦ ᐅᒪᓴᖅᓴᐃᓕᑦ

ᖃᖅᖢᓴᐅᖅᑐᑦ ᑎᒃᓯᒥᐊᖅᖃᓴᒃᑯ ᐃᑕᓴᖅᖅᓂᖅ ᑕᓚᐅᒃᑯᓂᒃ ᐃᒪᖏᒥᒃ ᖃᖅᖢᓴᐅᖅᑐᑦ ᑕᓚᐅᒃᑯᓂᒃ ᐃᒪᖏᒥᒃ, 50 km ᑭᓪᓕᒋᓂᒃ ᑕᑭᓂᓕᒥᒃ, ᐊᓗᒪᓗ ᑕᒪᐃᐢᓂᒃᖢᓴᐃ ᐃᑕᑎᕐᓴᐅᑕᑎᓗᑎᒃ ᓄᓇᖅᓯ ᐊᖏᑎᑐᑦᑕᐅᓂᒃᑎᑦ ᖃᖅᖢᒃᑲᖃᖅᑯᓂᒃᑎᑦ ᑎᑭᑦᑐᖅᓯᓂᒃᑎᑦᑐᑦ, ᖃᖅᖢᒃᑲᖃᖅᑯᓂᒃᓴᖅ, ᑕᒪᐢᓐᓇ, ᐱᑕᖅᓇᐅᓕᒪᔪᖅ ᑭᓪᖅᓇᓂᒃ ᐊᖏᑎᑐᑦᑕᐅᓂᖅᑎᐢᓂᒃ ᓄᓇᒥ, ᐊᒥᓕᐊᓴᐅᓪᓚᑎᒃ ᐃᒃᑯᑐᑎᑦ ᑕᓕᓴᐃᑦ ᑕᒧᐊᖅ, ᐊᓗᒪᓗ ᖃᓄᖅᐢᓂᒃᖢᓴᐅᖏᑦᑐᓕᓕᒃ ᓕᓴᐅᓂᖅᓯ ᖃᖅᖢᒃᑲᖃᖅᑯᓂᒃᑐᑦ. ᑕᒪᖏᑕᒃᖢᓴ ᑎᒃᓯᒥᐊᖅᖃᓴᒃᑯ ᐱᓕᖅᓯᓂᒃᑐᑦ ᑕᒪᐃᓕᑦ ᑕᒧᐊᖅ ᐅᐊᖅᓇᐢᓂᒃᑐᑦ ᐃᓕᓴᐅᓂᒃᑎᑦ ᐊᓗᒪᓗ ᖃᓄᖅᐢᓂᒃᖢᓴᐃᓕᑦ ᑕᒪᐃᓕ



ᑕᑯᔭᕆᔭᖅ 3: ᓇᓗᑎᓴᐅᓪᓗᐊᑕᖅᑕᕐᓂᖅᓂᓐᓂᒃᓂᓄᑐᓂᒃ ᐊᐃᔭᑐᖅᑕᐅᓯᒪᓂᖅᓂᑦ ᑎᖅᒥᐊᖅᑎᖅᕿᐃᐅᓴ (ᒪᓕᒃᑕᐅᓯᒪᓪᓗᑎᒃ ᐱᓕᑦᓭᐅᖅᑐᑦ Fᐊᓐᑎᓐᓄ ᐊᓚᒪᓗ ᐃᓗᓐᓇ-Fontaine and Mallory, 2011).

2.2 ᑎᖅᑕᐊᓂ

ᐃᑲᑦᑐᐊᖅᑎ ᑎᖅᑕᐊᖅᑲᕐᓇᑲ ᐃᓂᓕᓂ ᑕᒪᓂ ᑎᖅᑕᐊᓂ ᓴᐳᖅᓂᐊᖅᑕᑕᐅᓯᒪᖅᓇᒪᓂ ᑲᓇᑕᒥ ᐊᓇᑲᑐᖅᓯᒪᓂᖅᒪᓂ 3-ᒥ ᐅᑭᐅᖅᑕᖅᑐᑦ ᐊᓇᑲᑐᖅᓯᒪᓂᖅᒪᓂ ᓇᕐᑐᑐᖅᑎᓂᖅᑕᖅᑎᑲ ᐊᒻᒪᓗ ᑭᖅᒪᓕᓂ (ᐊᐅᑎᑕᓚᓴᕐᔪᐊᑲᑯᑦ ᑲᓇᑕᒥ, 2013). ᑕᒪᕐᒥᑲ ᑕᓚᐅᕐᒥᓯᓪᓗᓂ ᓇᓇᒥᐅᑕᐃᓕᓂ ᐅᒪᔪᕐᑲᖅᓇᐅᖅᐳᖅᓂᖅᑎᑦ ᐊᒥᓯᐊᓗᓂᓂᑦ ᐱᑕᕐᑲᖅᓇᐅᔪᑦ ᑎᖅᑕᐊᑲᓴᕐᑲᑐᑐᓂᑦ. ᐅᖅᑕᑕᐅᓂᒪᒪᑕ 90 ᑎᒥᐊᖅᑲᖅᓇᖅᑭᓯᖅᓯᔪᑦ, ᐊᖅᐳᓪᓕᐊᖅᑎᓪᓗ ᓇᓕᐊᖅᑕᒃᓯᓯᑎᓂ ᑕᐅᐳᓂ ᑎᖅᑕᐊᖅᑲᖅᓇᖅᑕ.

2.2.1 ᑲᖓᐃᐃᓂ


ᐊᒥᓯᓂᖅᑲᐃᓂ ᐱᑕᖅᑲᓂᖅᑎᑦ ᒥᑭᔨᑎᐅᓂᖅᑲᐃᓂ ᑲᖓᐃᐃᓂ ᓇᓕᐊᖅᑕᒃᑐᑦ ᑕᐅᐳᓂ ᑎᖅᑕᐊᖅᑲᖅᓇᐃᐅᓂᖅᓯᒪᓂ ᖃᑲᖅᑲᐅᖅᑐᑦ, ᐃᓕᑎᕐᔪᐅᖅᓭ ᐊᒥᑎᓴᖅᓇᓇᖅᓭ ᓴᖅᑭᒋᓗᐊᖅᓂ. 2014-ᒥ ᑲᖓᐃᐃᓂ (ᓯᑲᓗ ᑕᒪᕐᒥᑲ ᒥᑭᔨᑎᐅᓂᖅᑲᐃᓂᓗ ᑕᐃᑲᑯᐊᓗᓂ ᑲᖓᐃᐃᓂ (Ross’ geese) ᖃᐅᑎᕐᑎᑎᓂ ᐊᒥᓯᑎᑎᑕᐅᖅᑐᑦ ᑕᒪᓂ 234,000 ᐅᓪᓗᖃᑲᖅᑐᑦ (ᐅ. ᓕᖅᓗᐊᖅ-J. Leafloor, ᓇᖅᒥᓂᖅ ᑐᓴᐅᒪᑎᑕᑎᓂᐅᓚᐅᓂᖅᓯᓂᖅ). ᑕᐃᑲᑯᐊ ᑲᖓᐃᐃᓂ (Ross’ geese) ᐊᒥᓯᐳᑕᐊᖅᓯᒪᔪᓯᒪᓕᐊᓗᐃᓂ ᑕᐃᒪᖅᑎᓂ 1970-ᖅᑎᓂᖅ (ᑯᐱᐸ, ᐱᓕᕆᔨᖅᑎᓪᓗ-Kerbes, et al., 2014), ᐊᒻᒪᓗ ᐱᔪᔾᑕᐅᓪᓗᑎᑐᓗ ᐊᒥᓯᓂᖅᑲᓴᒪᓕᐊᐅᓕᕐᓂᓂᖅᑎᓂ ᐱᑕᖅᑲᓯᔪᕐᓂᓂᖅᑎᓂᓇᓂ ᑲᖓᐃᐃᓂ ᖃᑲᐅᔨᔪᐅᓯᒪᓕᕐᓂ ᒪᓕᒃᑕᐅᓗᓂᑎᒃ ᓂᐅᖅᒥᐅᓂᖅᓴᖅᑕᐅᖅᑕᕐᓯᒪᓂᖅᑎᓂᓂᖅᑐᑦ. ᑲᖓᐃᓇᐃᓂ (Ross’ geese) ᐊᒥᓯᑎᑎᓕᐅᖅᑐᑦ ᐅᖅᑲᒋᓂ <1% ᓴᓴᓂᑎᓂᖅ ᖃᑲᐅᔨᒪᔪᐅᓗᑎᑎᓵ ᓂᐅᖅᒥᐅᑕᓕᖅᑕᐅᓯᒪᔪᐃᓕᐅᓂᐅᖅᓂᖅᑎᓂ, ᐊᖅᓴᒍᑕᒧᖅ ᓴᓴᓂᑎᓂᖅᓭ ᓂᐅᖅᒥᐅᑕᓕᖅᑕᐅᓯᒪᔪᐃᓕᐅᓂᐅᖅᓂᖅᑎᓂ, ᐊᖅᓴᒍᔪᑕᒪᖅ ᐱᑕᖅᑲᓯᔪᕐᓂᓂᖅᑎᓂ ᐃᓕᓂ ᖃᑲᐅᔨᔪᐅᓯᒪᓕᕐᓂ ᑲᖓᐃᐃᓂ ᖃᑲᐅᔨᔪᐅᓯᒪᓕᕐᓂ ᐱᑕᖅᑲᓯᔪᕐᓂᓂᖅᑎᓂ ᓂᐅᖅᒥᐅᑕᓕᖅᑕᐅᓯᒪᔪᐃᓕᐅᓂᐅᖅᓂᖅᑎᓂ, ᐊᖅᓴᒍᔪᑕᒪᖅ ᐱᑕᖅᑲᓯᔪᕐᓂᓂᖅᑎᓂ ᐃᓕᓂ ᖃᑲᐅᔨᔪᐅᓯᒪᓕᕐᓂ ᑲᖓᐃᐃᓂ ᖃᑲᐅᔨᔪᐅᓯᒪᓕᕐᓂ ᐱᑕᖅᑲᓯᔪᕐᓂᓂᖅᑎᓂ ᐱᑕᖅᑲᓯᔪᕐᓂᓂᖅᑎᓂᓂᖅᑐᑦ ᑲᖓᐃᐃ

ᑎᖏᒥᐅᓪ ᓄᑦᖢᒃᑐᓪ ᑐᓂᕆᐅᓄᓪ ᑕᐅᖅᓯᓗᓪ ᓂᕐᐊᓄᓪ ᐅᖏᓱᓕᑎᓯᒪᓪ ᓄᓪᓕᐊᕐᓗᓪ (M’Clure Point; Abraham & Ankney, 1986).


ᑲᖓᐃᓪ ᓄᓪᐊᖅᐸᖏᓂᖏᓐᓂ ᐊᐅᑦᓪᖢᑦᓪᓪᐊᓪᓂᕆᑦᑕᕆᑐᓪ ᓴᑦᓪᑎᑦᓪᓕᖅᑐᓪᓂᖅᑐᓪ ᕿᓪᓕᐅᕆᓂᕆᑲᕆᑎᓕᖅᑐᓪᒥ ᑕᒪᓂ ᐃᓗᓱᐅᑦ ᕿᑎᕆᑦᐸᓱᐊᓂ, ᓄᓪᐊᓪᒃᐅᕆᑐᓪᑕᓪ ᑲᖓᐃᓪ ᐊᐅᑦᓪᒃᓴᕆᑦᑕᕆᓱᑎᑲ ᑕᒧᖓ ᐃᓕᓱᐅᑦ ᐃᓗᓱᐅᑦ ᐅᕐᖓᓪᓄᓂ ᐱᑎᐊᓪᒃᓴᓱᐊᓂ. (Sutton, 1932).

ᕿᑲᕆᐅᕆᑐᑎᒥ ᑎᒥᐊᕆᑲᕆᕐᑕᐅᕆᑲ ᐱᓚᒪᓪᐊᓪᑐᖅᑲ ᑎᑭᑕᐅᓱᒪᖖᓂᕆᑲ ᓂᕆᓪᕆᓪᓄᓂ Atlantic brant. ᑯᑦᓴ-Cooch (1957) ᓪᓕᐅᑕᓪᓕᐅᔪᖅᑲ ᑕᒪᑯᓐᓪᓕᓱᑎᐅᓂᕆᑲ ᐊᒥᓱᓂᕐᑎᑦ 5,000 ᓂᕆᓪᕆᓪᓪᐃᓂ ᑎᖏᒥᐊᕆᑲᕆᕐᒥ, ᓪᓕᐅᑕᕆᑲᑕᐅᓱᒪᓪᑐᑎᑲᑎᖅ ᐱᑕᕆᑲᕆᓂᖏᓐᓂᑲ 3% ᐳᓴᓐᖏᓐᓂᑯᓂ ᓂᕆᓪᕆᓪᓪᐃᓂ ᑕᐃᓴᓱᒪᓂ ᐊᑦᑭᓐᑎᒥ ᑎᖏᒥᐊᓕᓪᓚᓂᓕᑦ ᑲᑎᒪᔨᖏᓕᓂ-Atlantic Flyway Council, 2011. ᑕᐅᙵᑯᐊ ᐊᒥᑐᑐᒥᐅᑦ ᑕᓪᐅᑲᓂᓂ Foxe Channel-ᒥᕆᑭᑕᑕᑌᓴᐃᑦ ᓄᓕᐊᕆᕐᕕᐅᖅᑲᑐᓴᖏᒪᑕ ᓂᕆᓪᕆᓪᓭᓄᑦ Brant-ᓄᑦ , ᓱᓴᖅᐸᓱᑯᑦ ᖃᐅᔨᓴᖅᑕᐅᑎᒃ ᐊᒥᑐᑐᒥᐅᑦ ᑕᓪᐅᑲᓂ ᐊᓘᒪᓗ Hᐊᑦᓴᐤᐸᑦ ᑕᓪᐅᑲᓂ ᑕᒧᖓ 1979-ᒥ, ᖃᐅᔨᓴᖅᑎᑦ ᑎᑎᕆᓪᕆᐅᖅᑐᑦ ᐊᒥᓱᓂᖅᐸᑦᕆᑎᐊᖓᔪᓂᒃ ᑕᑯᕈᑕᐅᓪᓂᕆᑦᑐᓂᒃ ᓂᕆᓪᕆᓪᓭᓄᑦ ᖃᖅᖏᓴᐅᖅᑐᑐᒥ (15.7 ᓂᕆᓪᕆᓪᓴᖃᓂᕆᑦᑐᓂᑦbrant/km-ᑭᓪᒥᑐᑕᒪᓴᖅ ᓄᓪᒥ; ᓴᓵᓐᓂ, ᐱᓕᓚᖃᑎᖅᑎᑦᓪᑐ Gaston, et al., 1986). ᑕᒪᒃᑯᐊ ᐊᒥᓱᓂᖅᑦ ᑐᓂᖏᐅᑎᓱᒪᔪᖅᓴᐅᑦᑭᓂ ᓂᕆᓪᕆᓪᓭᓄᑦ ᐊᒥᓱᓂᕆᓪᕆᓪᓪᐃᓂᒃᑐᓂᒃ ᑕᐅᙵᓂ, ᐱᔪᐊᑎᑦᓪᑐᐊ ᐊᑕᐅᓱᓂᓪᓕᔪᑦ ᐃᓄᖏᑦ ᑕᑯᕈᑕᐅᓪᓂᕆᑦᑐᓂᑦ (Reed, et al., 1980). ᓄᓪᒋᑦᑐᓪᓄᓂ ᖃᐅᔨᓴᖅᑕᐅᑎᒥᒃ ᐅᑦᓪᑐᖅᑲᐃᐅᓪᐊᖅᓂᖅᑎᑦ ᑕᐅᙵᓂ ᑎᖏᒥᐊᕆᑲᕆᕐᒥ, ᑎᑎᕆᓪᕆᐅᖅᓱᒪᔪᑦ ᑕᐃᓴᓱᒪᓂ ᐱᓚᒪᓪᐊᓪᓚᖅᒥᒃ ᐊᒥᓱᓂᔪᓂᒃᓱᓪᑦᑐᓪᓚᖅᑎᖓᓂᒃᓂ ᖃᒃᖏᑎᖅᓱᒪᑦᑐᓪᓚᖅᑐᓪᓂ ᐊᓂᔪᖅᑐᓪ 30-ᓂᒃ ᐊᖅᕿᔪᓂᒃ. , ᐅᑦᓪᑐᖅᑲᐃᐅᙵᖓᓂᖅᑎᑦ ᓂᕆᓪᕆᓪᓭᓄᑦ brant-ᓄᑦ ᐊᓂᔪᖅᓱᒪᔪᓂ 30-ᑎᓂᒃ ᐊᖅᕿᔪᓂᒃ (Sharp & Abraham, 2010; Nissley, 2016). ᑐᑭᓱᓭᕆᑎᐊᖓᑎᓪᓕᖅ ᓱᓕ ᖃᓄᓂᑎᑦ ᓂᕆᓪᕆᓪᓪᐃᓂ (Atlantic brant) ᐃᓴᓭᐅᔪᖅ ᐱᑕᖃᓱᐃᖏᓐᓇᖏᓐᓄᑦ ᑕᐅᙵᓂ ᖃᖅᖏᓴᐅᖅᑐᑐᒥ ᑎᖏᒥᐊᕆᑲᕆᕐᒥ ᐊᓘᒪᓗ ᓂᕆᓪᕆᓪᓭᓄᑦ ᐊᓱᐃᖏᓐᓂᓴᓭᐊᑦᓪᓕᖅᑐᓪᓂᕆᓪᕆᓪᓚᖅᓂᓴᐃᓪᓕᓂᕆᓪᕆᓪᓚᖅᐸᖓᓂᒃᑐᓪᓂ ᑕᓴᐸᓂ ᐊᒥᑐᑐᒥᐅᑦ ᑕᓪᐅᑲᖃᕆᑲᕆᓪᕆᓪᓚᖅᑐᓪᓂᖅᑎᓂ Foxe Channel-ᒥ. ᑕᒪᐃᓴᓂᒃ ᐃᓱᒪᑦᓗᑎᑦ, ᐊᒥᓱᓂᖅᑦ ᓂᕆᓪᕆᓪᓴᓕᑦ Atlantic brant ᑕᐃᒪᐃᓕᖏᓐᓄᐃᓴᓚᖅᑐᑦ ᐊᒥᓱᓂᖅᑎᓂᒃᓂᓂ, ᐊᖅᕿᔪᓄᓕᔪᓱᒪᔪᓂ ᕿᑭᓕᓂᓗᓪ ᐊᖏᑎᓄᓗᓪ, ᒪᓕᒃᑕᐅᑦᓗᑎᒃ ᐅᑭᐅᒃᖏᓂ ᖃᒃᖏᐅᑦᑕᒃᑕᕆᑲᕆᓪᕆᓪᓚᖅᓂᖅᑎᓂ (Dufour, et al., 2019).

ᓂᕆᓪᕆᓪᓭᓄᑦ (Atlantic brant) ᑭᖓᓂᐊᔪᑦ ᑕᐃᓂᓂ ᒪᕐᐳᓂᒃ ᐱᓇᓱᐊᑭᓱᓂᒃ ᑎᑭᐸᕆᓪᕆᓪᓚᖅᓕᓱᓪᑦ ᐊᓱᖏᑲᓂᒃ ᑲᖓᐃᓪᒃ (Nissley, 2016). ᑕᒪᒃᑯᐊ ᓂᕆᓪᕆᓪᓭᓄᑦ Atlantic brant ᓇᔪᐃᓐᓄᑕᒪᓱᖖᓂᕆᑲᕆᕐᒥ ᓱᓴᖅᐸᓱᐅᓂᕆᑎᒃ ᐊᒥᓚᓗ ᐅᑦᓪᑐᖅᑲᐃᐅᓪᓚᓂᒃ ᐊᑕᓵᕆᑐᒥᑎᒃ >ᖅᑐᓂᓕᓚᒃ 5 m ᑭᑕᓕᓚᓂᒃ ᖃᑯᙵᓱᓕᓕᓚᓂᒃ ᐅᑦᓕᓚᑲᐅᓂᕆᑦᑐᓂᒃ ᐃᒧᑦ ᐊᓗᑦ ᖃᑯᙵᓱᓕᓕᓚᓂᒃ ᐃᓕᒃᑯᓚᓐᓂᑦ. (Abraham & Ankney, 1986). ᐅᑦᓪᑐᖅᑕ ᑕᑯᐅᑎᔪᓂᓂᖅᑲᐃᐅᙵᖓᑐᑦ ᕿᓚᓂᒃ ᐃᓕᓚᓂᑕ ᖃᖅᖏᓴᐅᖅᑐᑦ. ᒪᓂᓚᓂᑦ ᐃᓂᓄᖅᑕᓂᖅᑐᑦ ᑕᓪᐅᑲᓂ ᕿᑭᐅᒃᖏᓂ



ᑕᐃᔾᔪᓴᖅ 4: ᐃᓄᐃᑦ ᖃᐅᔨᓚᑐᑎᖃᖅᓂᒋᓐᓂᓐᓄᓄᑦ ᑲᖏᐃᑦ ᑎᓕᒥᐊᖃᖅᓴᐅᓂᕐᒥᒃ (ᒪᓕᔪᑐᑦ ᑲᑕᑦ, ᐱᓕᕆᖃᑎᖅᓂᕐᓗ-Carter et al. 2018-ᒥ ᖃᐅᔨᔪᖅᓂᐊᓂᖅᓂᓐᓄᔪᓂ). ᐱᑕᖃᖅᓴᐅᓂᖅᓂᑦ ᓴᖅᑭᐅᒪᔨᓪᓂ ᑕᒪᐃᓐᓄᑦ ᐃᓄᐃᑦ ᖃᐅᔨᓚᑐᑎᖃᖅᓂᒋᓐᓂᓐᓄᔪᓂᔪᓂᒃ ᒪᓕᔪᓯᑎᑐᒃ 21-ᓂᔪ ᐃᓄᖅᓂᔪ ᐃᓕᖃᔪᓕᐅᖅᓚᑕ.



ᑕᔪᒃᑳᖅ 5: ᑲᖓᐃᑦ ᐅᓪᓗᖓᓐᓄᑦᑕ ᓇᓂᓐᓂᕆᓕᕆᓂᑦ ᑐᑭᓯᐅᖅᑕᐅᓗᓂᑐᑎᒃ 2014-ᒥ ᓴᓪᓕᐅᑦ ᖃᑭᖅᑕᓗᐊᓂᒃ ᖃᔪᓯᐅᑎᓪᓗᑎᑐᑎᒃ ᖃᒃᐅᔨᓴᖅᑕᐅᓂᖓᓐᓂᐊᓂᒃ ᐊᔨᓕᓯᐅᖅᑕᐅᓯᐅᖅᑐᓂᒃ.

ᐊᒥᓱᐊᓗᖅᑐᑦᑕᖅᑐᑦ ᓂᕆᑕᖅᓭᐃᓕᑦ cackling geese ᓄᓕᐊᖅᑕᕐᒥᔪᑦ ᑕᐅᖅᓂ ᑎᖏᐊᖅᑲᖅᓇᖏᑦ. ᐃᓐᓇᓴᖅ ᖃᐅᔨᓴᖅᑕᐅᓕᐅᖅᑐᑦ ᖃᐅᔨᓴᐅᓕᐅᖅᑐᑦ ᓂᕆᑕᖅᓭᐃᓕᑦ cackling geese ᐊᒥᓱᑭᓚᐊᖅᓯᒪᔪᑦ ᑕᐃᒪᖏᓂᒃ 1980-ᖏᑐᓂᒃ (Nissley, 2016; Sharp & Abraham, 2010). ᓂᕆᑕᖅᓭᐃᓕᑦ cackling geese ᑎᑭᕐᒃᑐᑦ ᐅᓪᓗᖅᓯᒧᓪᓗᑎᓪᓗ ᐱᐅᓯᖅᑲᖅᑐᑎᑦ ᑲᒧᖅᑎᑐᑦ. ᑭᓯᐊᓂᕆᑕᐅᖅ, ᐅᓪᓗᖅᑕᕐᓂᓂᖅᓴᐅᒋᔪᑦ ᑎᖏᐊᖅᑲᖅᓇᖏᑦ ᑕᒪᐃᓐᓂᒃ ᐱᐅᓯᖅᑲᖅᓇᐅᒋᔭᖅᓂᖏᓂᒃ ᑐᖓᔪᓐᓇᖅᑐᑦ. (Nissley, 2016).

2.2.2 ᒥᑲᓂ

ᒥᑲᓂ ᐃᓕᑎᕆᐅᔨᓪᓚᕆᐅᑉᓄᑦ ᐅᑭᐅᖅᑕᖅᑐᒥ ᐊᒋᑎᒍᑦ ᐊᒋᑎᑎᒥᐅᑕᓄᓪᓗ, ᐊᒥᓱᓂᖅᓴᓯᒪᓪᓗᑎᓪᓗ ᐱᑕᖅᑎᓂᕐᑎᑦ ᐅᑭᐅᖅᑕᖅᑐᒥ, ᐱᓗᐊᖅᑐᒥᒃ ᑲᓇᑕᐅᑦ ᐊᒋᓚᓗ ᐱᓕᕆᓇᖅᓴᓯᓚᐅᑉᓄᑦ ᐊᐅᓚᑦᑎᐅᑦ. ᒥᑲᓂ ᐱᓪᓚᕆᐊᓗᖅᒥᒃ ᓂᐅᖅᓵᕆᐅᕐᓄᑦᑐᓂ ᓴᓪᓕᒥᐅᑕᓄᓪᓗ (Henri, 2007; Henri, et al., 2018). ᖁᑭᖅᑕᓯᓗᖅᒥ ᐱᑕᓕᓚᐊᓗᖅᒥᒃ, ᐊᒥᓱᑎᑎᕐᒍᓂᒃ ᑕᐃᓗᓐ 8,000-ᖏᑦᓂᓱᓯᒪᔨᓂ ᐅᓪᓗᓂᒃ ᐊᖏᕿᔪᒥ 2005 ᐊᓪᓚᓗ 2006 (ᑕᓯᓴᐅᓴᖅ 1; Descamps, et al., 2012). ᒥᑎᖅᑲᓕᒃ ᐱᓗᐊᖅᑐᒥᒃ ᐱᑕᖅᑲᓕᐅᕐᖅ *(ᐅᑭᐅᖅᑕᖅᑐᒥᓂᒃ)* ᒥᑎᓂᒃ, ᐱᑕᖅᓱᓯᓚᓗᓚᓪᓗᑎᒃ ᑕᐃᒃᓯᐊ ᐊᔪᐅᓴᓴᖏᑦᑐᓂ ᐊᓯᖏᑦᑐᓂ ᕿᐊᑦᑕᓴᖅᑭᑦᑐᑦᑕᖅᑐᑦ ᐱᑕᖅᓱᓯᓚᐅᑉᓄᑦ ᐊᐅᒋᐊᖅᓴᓯᓚᐃᓕᑦ ᐊᒥᓱᑎᑎᒥᐅᑕᓄᓪᓗ ᐊᒥᓱᑎᑎᒥᐅᑕᓄᓪᓗᒥ ᐊᒥᓱᑎᑎᒥᐅᑕᓄᓪᓗᒥᓂᑦ. ᐊᒥᓱᓂᖅᓴᓯᓚᐅᑉᓄᑦ ᐊᓯᖏᑦᑐᓂ – ᒥᑲᓂ ᐱᑕᖅᑲᓕᖏᑦᑐᓂ). ᒥᑲᓂ ᑎᑭᖅᑲᑕᖅᐳᑦ ᐃᓱᓕᖅᑕᓱᓯᐊᑦ ᒋᑦ ᑕᕿᑭᖓᓂ ᑕᒪᐅᖓᓄ ᖁᑎᖅᑕᓕᖅᑕᓱᐊᓄᑦ ᔨᓪᓚᐃᑦ. ᐊᖏᔪᓐᓇᐃᓕᑦ ᓇᔨᕈᕈᐅᖅᓯᓂᑦ ᒥᑎᖅᑲᓕᖅᑐᖅ ᑕᒪᓕᖅ ᐱᑎᐊᒋᓐᓇᖅᑐᖅᓂᒃ ᑕᒪᓕᖅ ᑕᒪᓕᖅᑐᑦ (Abraham & Ankney, 1986). ᐊᖏᓇᓐᓇᐃᓕᑦ ᐱᐊᖏᖅᑎᓪᓗ ᐃᓄᓐᓄᓯᒪᓕᕆᖅᑎᓐᓇᑎᑦᑎᓂᕈᓐᓇᖅᑐᑦ ᑕᒪᓕᖅ ᖁᑎᖅᑲᓕᖅᑕᓱᐊᓂ ᐊᑲᓯᐅᑦ, ᐃᓕᖅᑎᑦ ᖁᑭᓚᐅᖅᑲᖅᓯᒪᓕᖅᑎᓐᓇᑎᑦᑎᓂᕈᓐᓇᖅᑐᑦ ᑕᒪᓕᖅ ᐊᐅᔪᓐᓇᖅᑐᖅᓂᒃ ᐊᑐᐊᓗᖅᑐᑦ. (Mosbech, et al., 2006).

ᒥᑲᑦ ᓇᒧᓴᐅᖅᑲᑦᑎᓂᖅᑎᑦ ᑎᖏᐊᖅᑲᖅᓇᖏᑦ ᖁᑦᓴᐊᖅᑎᑕᐅᓯᒪᓕᒧᑦ ᖁᓕᓴᐊᐅᑎᑕᐅᓯᒪᔪᓚᑎᒧᑦ ᓴᓕᒪᓐᖅᓂᑦ ᑕᐃᒪᖏᓂᒃ 2001-ᒥᒃ (Mosbech, et al., 2006). ᐄᐊᑦᓴᓄ ᖁᑭᖅᑕᒍᑦᓗ ᓄᓇᕗᐅᓕᓗ ᓂᐸᐃᒃ ᑕᕿᓂᑕ ᐃᖁᑕᕿᓴᓴ ᐱᓕᓚᐅᓐᔨᖓ ᐃᓚᓴᕿᕗᐅᖅᑲᑦᑎᓂᕐᓂᒧᑦ ᑎᑭᔨᓄᓇᐃᓚᓂᕐᑐᓂᓄᓪᓗ ᐊᒪᓴᕿᕗᐅᖅᑲᑦᑎᓂᕐᓂᒧᑦ. ᒥᑎᑦ ᐅᓰᓂᖅᑕᑦ ᑕᐅᓂᓄᓪᓗ ᑕᓚᐅᖅᓴᓂᓄᓪᓗ ᓂᐃᐸᑦ ᑲᓇᖅᓴᓂᓄᓪᓗ ᐱᐃᐸᑦ ᑲᐃᓐᓂᒃᓯᒪᒥ ᐊᒪᓚᓗ ᓂᐅᐃᓕᐅᓂᒃᑯᑦ, ᐅᓕᓗᓂᓱᑦ ᑕᐅᓂᓄᓪᓗ ᐱᓕᕆᓇᖅᓴᓯᓚᐃᓕᑦ. (Mosbech, et al., 2006).

ᑭᖓᓕᑦ ᒥᑲᓂ ᓄᓕᐊᖅᐸᒃᑭᓂᑦ ᑎᖏᐊᖅᑲᖅᓇᖏᑦ. ᓴᓂᐊᓂᑕ, ᑕᓚᒃᓂᐊ ᖁᑭᖓᓕᑦ ᒥᑲᓂ ᑕᐃᓕᓂᕐᖏᓪᓗᓯᒋᖓᓕᑦ, ᐅᓕᓗᓂᓱᑦ ᑕᐅᓂᓄᓪᓗ ᐱᓕᕆᓇᖅᓴᓯᓚᐃᓕᑦ ᐊᒪᓴᕿᕗᐅᖅᑲᑦᑎᓂᕐᓂᒧᑦ ᐃᓚᓴᕿᕗᐅᖅᑲᑦᑎᓂᕐᓂᒧᑦ ᐊᒪᓴᕿᕗᐅᖅᑲᑦᑎᓂᕐᓂᒧᑦ (2003-2013), ᖃᐅᖓᓄᑦᖓᓕᑦ ᐅᓪᓗᖅᑲᖅᐸᑦᑐᓚᐊᖅᑕᕐᓂᖅᓂᒃ ᓚᓕᖅᒥᔪᑦ ᒥᑎᓂᒃ. ᑭᓯᐊᓂᑕ, ᖁᑭᖓᓕᑦ ᒥᑲᓂ ᒥᔪᐃᓚᓕᑦᑎᓂᒃ ᐃᓚᓴᕿᕗᐅᖅᑲᑦᑎᓂᕐᓂᒧᑦᓂᓗ ᐅᑎᖅᑲᑦᑐᓚᐊᓂᕐᓂᖅᑎᓐᓇᑎᑦ ᐊᒪᓚᓗ

ᐅᑭᕕᑎᓴᒃᑲᓂᑦ ᐊᐸᐅᖏᑎᓂᐅᖏᑦ ᑕᐃᒃᑯᓇᖏᓂ ᐅᑭᐅᖅᑐᓂ ᓄᓴᖅᖢᓴᖏᓂ ᐊᑯᑭᑐᕋᕆᐅᓐ ᑲᓇᑕᓂ
(Janssen & Gilchrist, 2013).

2.2.3 ᓄᓴᖅᖢᓴᐅᒥᐅᑕᐃᑦ ᓄᓴᕆᓚᐊᓚᓂᐃᑦ

ᓄᓴᖅᖢᓴᐅᑐᓂᕆᓴᓂ ᓇᓂᑐᐃᓇᕎᑎᐊᖅ ᑲᓇᒥᑦ ᐅᐊᖏᓇᖅᖢᓴᖏᓂ ᐊᒥᓴᔨᓂᖅᓴᒪᓕᖅᑐᑦ ᑕᒪᐃᐧᓂ ᑲᓇᒥᑦ
ᐊᒥᐊᓕᑲᒥᐅᑦᓗ ᓄᓇᓗᓐᖏᓂᒃ (Gratto-Trevor, et al., 2011). ᖃᖅᖢᓴᐅᑐᑦᑎ ᑎᖏᒥᐊᖃᖅᖁᖏᒥ
ᐊᐸᐸᖏᓕᑐᑎᒪᓚᐅᕆᓂᒃ ᓄᓴᖅᖢᓴᐅᑐᓂᕆᔨᖓᕆᓂᒃ ᐱᑕᖃᖅᖁᐅᕆᖅ ᐱᐅᕆᑎᐊᓗᓂᓐᓗ ᑲᓇᖏᖅᖢᓴᐊᐧᓂ ᐅᑭᐅᖅᑕᖃᑐᑦ
ᓄᓇᖏᓂ ᓇᒪᒋᑎᐊᓂᖏᖅᓄᓂ ᑕᒪᐃᓄᖅ ᓴᐅᑦᓂᓄᓂ ᐊᒥᓴᔨᖅᖢᓴᖏᖅᓄᓂ ᑕᒪᐃᓄᖅᑎᑦ, ᑕᓐᓇ. ᐊᒥᓴᑎᑎᕐᔪᑦ
ᑕᓐᓇ. 36-ᖏᑦᑐᓴᓂ ᓴᐅᑦᐃᑦ/ᐊᑕᐅᕆᓭᒥ ᑭᒃᖢᓵᔪᑐᑦ km^2 ᑭᓪᓕᒋᑕᒻᒥᑎᒥ, ᖃᖅᐅᐸᓴᖅᑕᐅᑎᒪᒃ (Dickie, et al., 2014),
ᑭᓴᐊᓂᒋᑕᐅᖅ ᐊᐸᐸᖏᓂᓄᖅᓚᐅᓐᓂ ᐊᖏᑎᒥᒃ ᒪᓂᓇᓴᒧᒋᓕᖅᑐᑦ ᐊᕆᕌᔪᓂᒃ. ᑕᐃᒪᓕᓴᐃᓂᓇᖅᑕᐅᑦᖅ,
ᑐᐃᑎᕆᑎᕆᑭᐅᑦ/ᓄᓴᖅᓚᐊᓕ ᐊᒥᓴᔨᖅᖢᓴᖏᖅᓄᓂᒥᒋᑦ ᓄᓕᐊᖅᖢᓂᖏᓂᓄᖅ ᓇᓂᑐᐃᓇᖅ ᖃᐅᔨᓴᓕᓴᐃᓂᓇᖅᑕᐅᑦᖅ
ᑕᐃᒪᖅᑲᒃ ᐊᕆᕌᔪᐃᓕᓂ ᐱᑎᓕᓴᖏᓂᒃᑲᓂᖏᓂᒃ ᐃᓕᓴᕐᔪᐊᓂ 2000-ᖏᓂᓂ (14-18 ᑐᖃᐃᑎᕆᑎᑭᐅᑦ/ᓄᓴᖅᓚᐊᓂ/ᐊᑕᐅᓴᑕᒪᖅ km^2
ᑭᓪᓕᒋᑕᒪᖅ; Perkins, et al., 2007), ᑭᓴᐊᓂᓕ ᐊᒥᓴᔨᖅᖢᓴᖏᖅᓄᖅᑕᐅᓕᓚᐅᓐᓂ. ᑕᒪᒃᑯᐊ ᓄᓴᖅᓚᐊᕐᔪᐊᓂ ᓄᓕᐊᖅᖢᓴᖏᖅᓄᓂ
ᐊᐸᐸᖏᓴᖅᑕᕆᓂᕆᓂᑐᑦ ᐊᒥᓴᓂᒃ ᐊᕐᓇᒍᓂᒃ (6-ᐱᖓᓴᔪᖅᑐᓂᒃ-34-ᓄᑦ ᓄᓴᖅᓚᐊᕐᔪᐊᓂ/ᐊᑕᐅᓴᖅ ᑭᓪᓕᒋᑕᖅ km^2; Dickie, et
al., 2014). ᓴᓐᓪᓕᒥᐅᑦ ᐅᖅᑲᐅᓴᖅᑲᖅᓴᒪᔪᓂ ᐱᑎᓪᓕᐊᑐᓂᖏᓂᒃ ᓄᓴᖅᖢᓴᐅᑐᓂᕆᐅᑕᐃᐃᑦ, ᐱᑐᐊᖅᑐᒥᒃ ᑕᒪᒃᑯᐊ
ᐊᐅᖢᑐᒃᑐᓂ ᓴᐅᑦᓂᐃᑦ, ᓇᑭᓐᓂ ᐊᓴᖏᓂᓐᓗ ᓄᓴᕆᓚᐊᓴᓴᐅᓴᓂᒃ (Carter, et al., 2018). ᑕᐃᓵᓴᒪᑐᖅᑲᖅ
ᐊᒥᓴᐊᑐᖅᓭᓪᓕᐅᖅᑐᑦ ᐅᐱᖏᐃᒃᑯᓴ, ᑕᒪᒃᑯᐊ ᓄᓴᖅᓚᐊᕐᔪᐊᓪᓕᐅᑎᑦ ᑕᑯᐅᑎᓚᐊᐃᖅᑐᓂ ᖃᐅᔨᓴᖏᖏᖅᑐᒥᒃ ᓴᓐᓪᓕᒥᐅᑦ.

ᐊᓴᖏᑦᑕᐅᖅ ᓄᓴᖅᓚᐊᕐᔪᐃᓪᓕᐃᓂᑦ ᑎᖏᒥᐊᖃᖅᑎᒥᓴᓂ ᐊᒪᓪᓗ ᒪᑯᐊᑎᓴᐃᓂᑦ ᑎᖏᒥᐊᖃᖅᑎᒥᓴᓂ
ᐱᑕᖃᓂᖏᑦᓗᓂᓗᑎᒃ: ᓴᓐᖏᕐᔪᐊᖃᖅ/ᑐᑦᓪᓕᑎᕐᔪᐃᓂᑦ, ᖃᓂᓪᓂᓂᓪᓕᐊᕐᓂᒃ, ᑐᑦᓪᓕᑎᕐᔪᐃᓂᑦ, ᐊᒥᓴᓂᓂᖏᓂᖃᖅᑕᐅᖏᓂᒃᑲᓂᖏᓂᒃ ᑕᒪᒃᑯᐊ
ᓇᑭᓐᓂᑦ,ᐊᒪᓪᓗ ᖃᓂᓪᓂᓇᑦᓂ. ᓄᓴᖅᓚᐊᕐᔪᒃ-Purple sandpiper ᑕᑯᐅᑎᔪᐃᓇᓂᒃᖃᖅᑎᑭᐅᓂᓕ ᓄᓕᐊᖅᑕᓂᕆᓂᓕᓂ ᑎᖏᒥᐊᖃᖅᖁᖏᒥ.
ᑕᒪᓇᓇ ᓄᓇᖏᓂ ᐊᑐᖅᑕᐅᑎᔪᐅᓴᖅ ᓄᓕᐊᕐᓄᐃᑲᖏᓂᕆᓪᓗᑦ, ᓂᕐᑭᐅᓴᖅᓂᑭᐅᓴᓄᐃᓂᕆᓂᓪᓗᑦ, ᓚᓪᓄᐅᑭᓕᓪᓕᐊᓂᖏᓂᓄᓂ
ᓄᖅᖢᓄᒡᓄᐃᓂᐅᑲᖅᑕᓂᕆᓂᓪᓗᑦ ᑕᒪᒃᑯᐊ ᐊᒥᓴᑭᑎᓪᓕᐊᓕᓴᖅᑐᑎᓄᑦ ᓂᕐᖁᕐᓄᑦ ᓂᓴᕆᓚᐊᕐᔪᖓᓄᑦ red knot-ᓄᑦ (Lathrop, et
al., 2018). ᑭᐊᓂᓴᑎᐊᖅ, ᑐᖃᐃᑎᕆᑎᑭᐅᖅ ᐊᒪᓪᓗ ᑐᑦᓪᓕᑎᕐᔪ ᑕᒪᑭᓴᓂᖏᓂᒃᑲᓂᖏᓂᒃ ᑎᖏᒥᐊᖃᖅᖁᓴᒃᑯᓂ,
ᓚᓪᓄᐅᑭᓄᓂᖅᓴᓂᓕᓂᒃ.

2.2.4 ᐊᓂᖏᓂ ᐃᒪᐅᓕᕆᓂᐃᑦ ᑎᖏᒥᐊᓂ

ᖃᖅᖢᓴᐅᖅᑐᑦᑎ ᑎᖏᒥᐊᖃᖅᖁᓪ ᐅᖅᑲᐅᓴᖅᑲᖅᓴᒪᒃ ᑕᑭᓂ ᖃᖅᖢᓴᐅᖅᑕᐃᑦ, ᐊᒥᓴᓂᖅᑕᒡᒪᑕ ᐃᓯᖁᓂᓂᖏᓂᒃ
ᓄᓕᐊᖅᖢᓂᖏᓂᒃ ᑕᒪᒃᑯᐊᓂᓪᓗ ᐱᑕᖃᓂᖏᓂᒃ ᐃᒪᓕᐅᕆᓂᐃᑦ ᓴᓇᖅᑐᓄᑦ ᓴᓇᖅᑐᓂᓕᕆᓂᐃᓄᑦ ᐊᒪᓪᓗ ᐃᓂᖏᓂᒃ,
ᓚᓪᓄᐅᑭᓄᓂᖅᓴᓂᓕᓂᒃ. ᐊᓂᖏᓂᓂᒃ ᐱᑕᖃᓂᖏᓂᒃᑲᓂᖏᓂᒃ ᐃᒪᓕᐅᕆᓂᐃᑦ ᒪᓇᐅᓂᖏᓂᓗᓂ ᓄᓇᓕᒃ ᐊᒪᓪᓗ
ᑕᓂᐅᖅᓴᓂ, ᓂᓴᓗ ᑕᒪᒃᑯᓂᖏᓂ ᓇᐅᖅᓴᕐᔪᐊᓴᓂ, ᐃᒥᓴᑯᑕᐃᓚᓂᓴᓂᒃ, ᐃᒥᓴᑯᑕᐃᓚᓂᓴᓂᒃ ᐊᒪᓪᓗ ᐃᓂᖏᓂᒃᑲᓂᒃ, ᑕᒪᐃᓇᒃ ᒪᓴᐅᓂᒃ,

ᐊᓚᒐ ᐃᖃᐊᓴᕆᐃᐊᖅᑐᖅ/ᑎᕃᑎᕃᖅ, ᕿᓂᕕᖏᓂᖅ ᑐᑦᓯᔪᓂᒃ, ᐊᓴᑎᐊᖅᔪᓂᒃ,, ᖃᐅᖅᔪᓂᒃ ᐊᓚᒐ ᖃᐅᒧᒋᐊᕆᖅᓂᒃ. ᑕᒪᖅᑲ ᑕᒪᒃᑯᐊ ᓄᓕᐊᖅᑎᑦᓴᒃᑯᑦ ᑕᐅᔪᓂ ᑎᖏᒥᐊᖅᑲᕕᖏᒥ. ᖃᑭᖅᑕᓂᑐᖅᖏᒥ ᐊᒥᓂᔪᒃᑐᑦᑕᐅᖅ ᑕᒧᓂ ᐊᒥᓂᑎᑎᔪᒃᑐᑦ 200-300-ᖏᑎi ᐅᑦᑐᖅᑲᓂᖓᓪᓚᑕ ᐱᖁᓂᐅᓪᓗᑦ.

ᓴᖅᑦᕆᐅᑦ ᐅᖅᑲᐅᓂᖅᑲᓂᓪᓚᒃᑯᑦ ᐃᓂᓪᓚᑐᑎᑎᕐᕆᓂᒃ ᐊᒥᓂᔪᖏᓐᓄᑦᑕᖅᑕᕐᓂᔪᖏᓐᓂᒃ ᑕᒪᒃᑯᐊ ᑎᖏᒥᐊᑦ, ᐱᑐᐊᖅᑐᒥᒃ ᑕᒪᒃᑯᐊ ᐃᒥᖅᑯᑕᐃᓕᐅᑦᑐᓪ, ᖃᖅᖄᓴᑐᓪ ᑐᑦᓯᓪᓗᖅ (Carter, et al., 2018). ᖃᐅᔨᓴᖅᑕᐅᓂᒪᖏᑕ ᐊᒥᓂᔪᓂᑦ ᐊᓂᔪᖏᓐᓄᑦ ᐃᓕᖅᑐᓂᒃ ᑎᖏᒥᐊᑦ, ᐱᑐᐊᖅᑐᒥᒃ ᑕᒪᒃᑯᐊ ᐃᒥᖅᑯᑕᐃᓕᐅᑦᑐᓪ, ᖃᖅᖄᓴᑐᓪ ᑐᑦᓯᓪᓗᖅ (Carter, et al., 2018). ᖃᐅᔨᓴᖅᑕᐅᓂᒪᖏᑕ ᐊᒥᓂᔪᓂᑦ ᐊᓂᔪᖏᓐᓄᑦ ᑐᓴᕐᓯᓪᓗᖅ ᑎᖏᒥᐊᑦ ᐃᓗᐊᓂᑐᓪ ᑎᖏᒥᐊᖅᑲᓇᐃᖅᑐᓂ, ᖃᐅᔨᓴᖅᑕᐅᓂᒪᖏᑕ ᐊᒥᓂᔪᓂᑦ ᐊᓂᔪᖏᓐᓄᑦ ᑐᓴᕐᓯᓪᓗᖅ ᑎᖏᒥᐊᑦ ᐃᓗᐊᓂᑐᓪ ᑎᖏᒥᐊᖅᑲᓇᐃᖅᑐᓂ ᐊᒥᓂᔪᓂᑦ ᐊᓂᔪᖏᓐᓄᑦ ᑐᓴᕐᓯᓪᓗᖅ ᑎᖏᒥᐊᑦ, ᐱᑐᐊᖅᑐᒥᒃ ᑕᒪᒃᑯᐊ ᐃᒥᖅᑯᑕᐃᓕᐅᑦᑐᓪ, ᖃᖅᖄᓴᑐᓪ ᑐᑦ

ᑕᑯᐅᐅᓴᖅᓂᓂᑯᓐᓄᓂ ᐃᑲᑐᑐᐊᓱ ᑯᖓᓂᒃ ᑎᕆᐊᖅᑎᖏᕆ ᑕᒧᓂ ᐊᖅᑕᒧᐃ 2002 (Fontaine & Mallory, 2011). ᑕᒪᒃᑯᐊ ᐱᖓᓱᓕ ᐅᒪᔨᐃᑦ ᐊᐅᑭᖓᕐᑐᑦ ᐃᓕᓴᓂᕆᔨᖓᓕᒫᖅ ᐃᐱᒃᓴᕐᓂᔨᓕ ᓄᓇᖅᐱᐅᓂᕆᕐᖓᓯᖅᑐᓂᒃ ᑕᐅᖅᓯᓴᖅᓄᑎᒃ ᓴᓪᓕᐅᑦ ᑎᓪᓗᒍᑦ (Roe’s Welcome Sound-ᖓᓗᑦ), ᓴᓪᓕᐅᑦ ᖃᑭᖅᑕᒻᒧᐃᓄᑦ.

ᐅᑲᓪᓕ ᐱᑕᖅᑲᓰᕐᔪᓂ ᓄᓇᒥ >ᖅᑐᓂᖅᓴᐅᓂᖓᓕᓂ ᑎᕆᐊᖅᑎᖏᐅᑦ, ᖅᑲᓂᑯᖓᓗᓂ ᓄᓗᒻᒥᐅᑦ. ᑎᕆᐊᖅᑎᖏᐅᓂᖏᑦ, ᐱᑕᖅᖓᓗᑲᓰ ᖃᐅᑎᖓᓂᖅᓴᐅᓂᖓᓕᓂ ᓄᓇᒥ. ᐊᑕᐅᑲᖏᑦ ᑕᑯᐅᐅᓴᖅᑲᓕᐅᖏᒥᔪᖅ ᑎᓘᐊᖏᑦ ᑕᐅᖅᓯᖃᐅᓄᖓᒥ ᑕᒪᓕ 2015-ᒥ.

2.3.2 ᑕᓘᐅᖏᒥᐅᑕᐃᑦ ᐅᒪᔨᐃᑦ

9-ᖓᕐᓂ ᑕᓘᐅᖏᒥᐅᑕᐃᑦ ᐅᒪᔨᐃᑦ >ᐃᓕᓂ ᐱᑕᖅᑲᓰᓂ ᖃᖅᖢᓴᐅᖅᑐᑐᖏᒥ ᑕᓘᐅᖏᒥ ᐊᐅᓴᖅᑯᓂ. ᑕᐃᒪᐃᓂᖅᑐᓂᑐᖅ ᓄᓇᑐᓂᑐᑎᓂ, ᐊᐃᐹᓂ, ᓄᓇᐃᓂ, ᖃᐅᓇᓗᒍᐃᓂ, ᑐᐃᓪᓕᓇ ᐊᖏᓐᓂ ᐱᑕᖅᑲᓰᕐᓂᑐᑐᓂ ᓴᓪᓕᐅᑦ ᓴᓇᖅᑭᓯᑲᓂᖅᓴᓂᓂ ᑕᒪᒪᓇ ᑕᓘᐅᖏᒥ Hᐊᑦᓴᓄᐊᑦᒥ (Sutton, 1932; Bird, 1953; ᑭᓪᓕᓕᒥ ᐃᓄᐃᑦ ᑲᑐᑎᑎᑭᓂ ᓄᓇᕗᒻᒥᐅᑕᓂᒃ ᐅᓇᓗᒍᑦ; Stephenson & Hartwig, 2010).

ᑕᑯᐅᐅᖅᑕᖅᑲᓕᓴᐅᓂ ᐊᒥᓯᑦ ᑲᑎᖓᓴᐅᓂ ᖃᐅᓇᓗᒍᐃᓂ ᐊᒪᒪᓇ ᐊᐃᐹᓂ ᑕᐃᒪᖓᓂᒃ ᑎᕆᐊᖅᑎᖏᐅᒥ (Latour, et al., 2008; Allison, 1977; Irniurviit ACMC, 2017). ᖃᐅᓇᓗᒍᐃᓂ ᐃᑲᓐᓂᖓᓯᓯᖓᓂ ᖃᖅᖢᓴᐅᖅᑐᑐᖏᒥ ᐊᐅᑭᖓᑯᓂ ᐅᐃᓂᕐᒥᓂᒃ ᐱᓴᐃᓂᕆᒥᓂᒃ ᐊᒋᔭᖅᑎᓪᓗᒍᑦ. ᖃᖅᖢᓴᐅᖅᑐᑐᖏᒥᑕᐅᖅ ᓂᕆᐹᖅᓴᓯᐅᐃᐅᓪᓚᕐᓂ ᐊᒪᓚᓇ ᐃᓕᓴᖅᑕᖅᑲᓕᐅᖏᓪᓗᒍᑦ ᐅᓱᒋᐊᓕᒫᖅ whales (ᑭᓪᓕᓕᒥ ᐃᓄᐃᑦ ᑲᑐᑎᑎᑭᓂ ᓄᓇᕗᒻᒥᐅᑕᓂᒃ).

ᓄᓇᐃᑦ ᑕᐅᖅᓯᓴᖅᑕᑕᐃᓄᓇᓯᕗᖓᓂ ᑎᕆᐊᖅᑎᖏᐅᖅᑐᑦ. ᐱᓪᓚᕆᐊᓇᖅᑎᒃ ᐊᐅᑭᖓᑦ ᓄᓇᑌᓱᐅᐅᖅᑐᖅ ᐊᒪᓚᓇ ᐅᓪᓗᖅᑎᖏᐅᖅᑐᓄᓇᕗᑦ ᑕᐅᖅᓯᓄᓇᒧᑦ Foxe Channel-ᒥ ᓄᓇᐃᑦ ᖃᐅᔨᓴᖅᑎᑦᓴᓂᒃ ᑕᐅᖅᓯᓴᖅᑭᑦ ᐅᑭᐅᓯᓄᓇᓕᓴᖅᑐᓂᒃ ᓄᓇᐃᑦ ᑕᑯᐅᐅᖅᑕᖅᓴᖅᓂᖅ ᑕᐅᖅᓯᑎᑦᓴᖅᓂᖅ ᑎᕆᐊᖅᑎᖏᐅᒥ.

2.3.3 ᑕᓘᐅᖓᓂᖅᓴᖏᒥ ᑕᓯᖏᐅᖅᑕᐃᑦ ᑕᓘᐅᖏᒥᐅᑕᐃᓪᓗᒍ ᐃᖃᓗᓐᓂᕐᓂᐃᑦ

ᐊᒥᑎᓵᓗᐃᑦ ᑕᓯᓐᓂᕐᓂᐃᑦ, ᑕᓯᐊᖅᔪᐃᓪᓗᒍ, ᑯᓪᓗᒍ, ᑕᓯᐅᓴᖅᔪᐊᓪᓗᒍ ᑎᕆᐊᖅᑎᖏᐅᒥ ᐃᒪᖅᑲᐅᖅᓱᓂ ᐊᒪᓚᓇ ᑕᓘᐅᖏᒥ ᐃᒪᖅᓂ ᐊᒥᓯᓄᑦ ᐊᐅᑭᖓᑦᓯᓗᒍᑦ ᐃᖃᓗᓐᓂᕐᓂᐃᓕᓴᐅᓂᖓᓕᓂᒃ. ᑕᒪᐃᓐᓄᑦ ᖃᓄᐃᓂᕐᒥᑦ ᐊᐅᑭᖓᑦᓯᑎᓂᑦ ᐃᖃᓗᓐᓂᕐᓂᐃᓕᓴᐅᓂᖓᓕᓂᒃ ᐃᖃᓗᓐᓂᕐᓂᐃᓕᓴᐅᓂᖓᓕᓂᒃ ᐱᓕᕆᔨᒃᑯᐃᓂᖅᓴᓂᒃ ᐃᓕᓴᖅᑕᖅᑲᓕᐅᖏᓪᓗᒍᑦ ᐊᒪᓚᓇ ᓯᓚᑦᓴᖅᑕᐅᓂᖓᓕᓂᒃ ᐃᓕᓴᖅᑕᖅᑲᓕᐅᖏᓪᓗᒍᑦ ᐊᒪᓚᓇ ᓯᓚᑦᓴᖅᑕᐅᓂᖓᓕᓂᒃ ᐃᓕᓴᖅᑕᖅᑲᓕᐅᖏᓪᓗᒍᑦ ᐊᒪᓚᓇ ᑕᓘᐅᖏᒥ ᐃᒪᖅᓂ ᐊᒥᓯᓄᑦ ᐊᐅᑭᖓᑦᓯᑎᓂᑦ ᐃᖃᓗᓐᓂᕐᓂᐃᓕᓴᐅᓂᖓᓕᓂᒃ ᐃᖃᓗᓐᓂᕐᓂᐃᓕᓴᐅᓂᖓᓕᓂᒃ ᐱᓕᕆᔨᒃᑯᐃᓂᖅᓴ

ᒪᔪᖅᐸᒃᑐᑦ/ᓴᑐᐊᒃᑐᑦ ᐊᒻᒪᓗ ᑕᕆᐅᖅᑎᐅᑕᐅᖓᓂᓐᓄᖅᐸᒃᑐᑦ ᐃᖅᑲᓗᐃᑦ ᐊᐃᒍᖓᓂᑐᐃᓐᓇ, ᐃᓕᓴᖅᑲᓂᒃ ᑕᑦᓚᒥᓂᒃ 5 ᐊᒥᓱᖓᓗᐊᓕᕆᓂᖅᑕᐅᓂᒪᔪᒥᓂᒃ ᐅᒪᔪᓂᒃ, ᐱᑕᖅᑭᓐᓇᖅᑎᔪᓂᒃ ᑎᖓᒥᐊᖅᑲᐃᖓᒥ (ᐊᐃᒃᑐᖅᓂᒪᔪᓂᒃ ᑕᑯᑎᑕᑎᐅᔪᓂ 4; ᑯᑦ ᐊᒻᒪᓗ ᕿᐃᓴᑦ-Coad and Reist, 2004).

2.4 ᐱᑭᖅᑐᖅᑲᓂᖓᓕᒐ ᓄᓇ

ᓴᓂᓕᓚᐅᑐᐃᓚᐅᐅᔪᖅᑲᖅᓱᖅ ᖃᑯᓕᓪᓕᖓᓂᒃ ᖃᖅᖃᑉᕿᖓᓂᐅᔪᓂᒃ ᐊᐃᒃᓂᒪᔪᓂᒃ ᖃᑯᒧᖓᓕᓂᒃᓴᓂᑦ ᓂᐅᖅᓯᖓᑕ ᑕᐅᑭᖅᑐᑦ ᓂᐅᖅᑲᐅᑐᖓᓕᒐᓂᒃ ᐃᓐᓇᑐᖓᓕᒐᓂᒃ ᑕᒪᐃᓐᓇᖅᑐᑦ

ᓄᖅᓱᓯᒪᓕᕐᓂᐊᓂᒃ ᓇᓗᕐᓴᒃᑕᑐᕐᓂᖅᓱᖅᑐᓂᒃ, ᓄᖅᓱᓕᓴᕐᓂᖅᑐᑦ, ᐅᓗᓚᐊᓇᖅᑐᒋᓕᕐᓂᖅᑐᑦ ᓄᖅᓱᓕᓴᕐᕈᓇᖅᑐᓂᒃ, ᐊᐸᐅᖅᑎᑎᑐᒃ ᐃᓯᒪᓵᑕᐅᓕᕐᓂᖅᑐᑦ, ᐊᓪᓚᓗ ᐊᒥᓯᓇᖅᓴᖅᓴᕐᓕᐊᖅᑎᑦ.

ᑲᓇᑕᒥ *ᐅᒪᓴᐃᑦ ᐊᒥᓴᓯᖅᑕᐊᓕᕐᓂᖅᖁᖅᓄᓇᑦ ᒪᓕᑦᓴᖅᑎᑦ (SARA)* ᓴᖅᑭᑎᕆᑎᓕᐅᖅᓂᑦ ᑕᐃᔪᐅᓇᓂᒃ ᒪᓕᑦᓴᐊᖅᑐᖅᓂᖅᖁᖅᓄᓇᑦ ᓇᑎᐅᑎᓂᒃ 1-ᒥᒃ, ᐃᓕᓕᓚᔪᐅᓇᓂᒃ ᑎᑎᕆᖅᑕᐅᖅᑲᕕᕆᓂᖅᖁᖅᓄᓇᑦ ᐅᒪᓴᐃᑦ ᐊᒥᓯᖅᑕᐊᓕᕐᖅᑐᓄᓇᑦ. ᑐᑭᓯᐊᖏᑕᐅᖅᑲᒃᓕᐅᖅᓯᐅᖅᑎᑦ, ᒥᓂᕆᐃᑦ ᑲᑎᒪᔨᖅᑎᑦ ᑲᑎᒪᔨᖅᑐᓂᒃ ᑲᑐᓕᕐᑯᔪᓕᐅᖅᑐᖅᓂᒃ ᑲᑎᒪᔨᖅᑐᓂᒃ ᒥᓂᕆᓴᓚᓄᓕᕐᓗᑦ ᐅᒪᓴᐃᑦ ᐃᓕᓴᐅᔪᓚᐊᖅᑲᒪᖅᓱᑦ ᒪᓕᑦᓴᐅᑕᐅᓯᒪᔪᖅᓴᓗᑦ 1-ᓗᑦ ᑎᑎᕆᑕᐅᓯᒪᓕᐅᑎᒃ.

22-ᖅᑐᓂᒃᓴᑦ ᐊᐸᔨᖅᑎᑦᑐᑎᑦ ᐅᒪᓴᐃᑦ ᐊᒥᓯᖅᑕᐊᓕᕐᖅᑐᑦ (ᑕᐃᔪᐅᑦᓕᐅᑎᒃ ᑲᑎᒪᔨᕆᓭᓕᐅᑎᒃ ᑲᑎᒪᔨᖅᑎᑦ ᐊᒥᓯᖅᑕᐊᓕᕐᓂᖅᑐᓄᓇᑦ COSEWIC-ᑯᓘᓂᑦ-ᖃᐅᔨᔪᖅᑕᐅᓕᐅᖅᓯᐅᑎᒃ ᐅᖃᐅᓕᓘᓂᒃ ᐊᑐᖅᑕᐅᓕᓕᐅᑎᒃ ᐅᒪᓴᐃᑦ ᐊᒥᓯᖅᑕᐊᓕᕐᓂᖅᑐᓄᓇᑦ ᒪᓕᑦᓴᖅᑎᑦ) ᐃᓕᓚᔪᐅᓯᒪᔪᓐᓇᑦ ᐅᖃᐅᓕᓘᓂᒃ ᐃᓕᓴᐅᑐᐃᓘᐅᓚᐊᖅᖁᖅᓴᓐᓇᑦ ᒪᓕᑦᓴᖅᑎᒧᑦ ᑕᐃᑯᐊ ᑎᖅᒥᐊᖅᑲᖅᓱᖅᓴᓂ ᐱᑕᖅᑯᓯᑦ. (ᐊᐃᖅᑐᖅᓯᒪᔪᓂᒃ ᑕᑯᑎᑕᑎᔨᓪᓗᑦ 4).

ᐊᐃᖅᑐᖅᓯᒪᔪᓂᒃ ᑕᑯᑎᑕᑎᔨᓪᓗᑦ 4: ᐅᒪᓴᐃᑦ ᐊᒥᓯᖅᑕᐊᓕᕐᓂᖅᖁᖅᓄᓇᑦ ᑎᖅᒥᐊᖅᑲᖅᓱᖅᓴᓐᓇᑦ

ᑕᐃᔪᐅᓂᖅᓄᑦ ᐃᓄᖅᑎᑐᑦ ᖃᖅᐅᔨᕆᖅᑎᓄᓗᐃᓗ ᐅᕆᖃᖅᐅᔨᓕᐅᑎᓄᑦ ᑎᑎᕆᖃᖅᓯᒪᓂᖅᓄᑦ	ᖃᖅᓄᐃᑦᓱᖅᓂᖅᓄᑦ ᑲᑎᒪᔨᖅ ᐃᓕᓚᔪᐅᓯᒪᔪᖅᓴᓐᓇᑦ		ᐱᑕᖃᖅᓂᖅᓄᑦ ᐃᑲᑦᑐᐊᓕ ᑎᖅᒥᐊᖅᑲᖅᓱᖅᓴᓂ
	ᑲᑎᒪᔨᖅ ᑲᑎᒪᔨᕆᖅᓯᒪᓂᖅᓄᑦ ᖃᖅᓄᐃᑦᓱᖅᓂᖅᓄᑦ ᐊᒥᓯᖅᑕᐊᓕᕐᓂᖅᖁᖅᓄᓇᑦ ᐃᓕᓴᐅᑐᐃᓘᐅᓚᐊᖅᖁᖅᓴᓐᓇᑦ - COSEWIC-ᑯᑦ ᖃᖅᐅᔨᔪᖅᓯᒪᓕᐅᖅᓯᐅᑎᒃ	ᐃᓕᓴᐅᑐᐃᓘᐅᓚᐊᖅᖁᖅᓴᓐᓇᑦ ᐊᒥᓯᖅᑕᐊᓕᕐᓂᖅᖁᖅᓄᓇᑦ ᒪᓕᑦᓴᐅᑕᐅᓯᒪᔪᖅᓴᓗᑦ	
ᑎᖅᒥᐊᑦ			
Ivory gull ᖃᖅᑯᓯᓪᓗᐃᑦ *Pagophila eburnea*	ᓄᖅᓱᓕᓴᕐᓂᑦ	ᓄᖅᓱᓕᓴᕐᓂᑦ	ᓇᓗᓇᐃᖅᑕᐅᓯᒪᓕᕐᓂᑦ
Red knot ᑭᖅᑎᖅ ᓯᓚᕆᐊᓚᐅᖅ - ᓗᐊᓗᓗᐊᓪᓕᖅ *Calidris canutus rufa*	ᓄᖅᓱᓕᓴᕐᓂᑦ	ᓄᖅᓱᓕᓴᕐᓂᑦ	ᓇᓗᓇᐃᖅᑕᐅᓯᒪᓕᕐᓂᑦ
Ross's gull ᓇᐅᔨᖅ *Rhodostethia rosea*	ᐅᓗᓚᐊᓇᖅᑐᒋᓕᕐᓂᖅᓂᖅ	ᐅᓗᓚᐊᓇᖅᑐᒋᓕᕐᓂᖅᓂᖅ	ᓇᓗᓇᐃᖅᑕᐅᓯᒪᓕᕐᓂᑦ
Bank swallow ᖃᑦᓴᓂᐊᖅᓯᒃ *Riparia riparia*	ᐅᓗᓚᐊᓇᖅᑐᒋᓕᕐᓂᖅᓂᖅ	ᐅᓗᓚᐊᓇᖅᑐᒋᓕᕐᓂᖅᓂᖅ	ᓇᓗᓇᐃᖅᑕᐅᓯᒪᓕᕐᓂᑦ
Barn swallow ᑐᖓᔪᖅᑐᖃᕐᓇᖅᑐᑦ ᖃᑦᓴᓂᐊᖅᓯᒃ *Hirundo rustica*	ᐅᓗᓚᐊᓇᖅᑐᒋᓕᕐᓂᖅᓂᖅ	ᐅᓗᓚᐊᓇᖅᑐᒋᓕᕐᓂᖅᓂᖅ	ᓇᓗᓇᐃᖅᑕᐅᓯᒪᓕᕐᓂᑦ

ᑕᐃᔨᐅᓂᖅᓯᒪᔪᑦ ᐃᓄᐹᑎᑐᑦ ᖃᐅᔨᓴᖅᑎᓄᑕ ᐅᖃᐅᓯᐊᔪᑦ ᑎᑎᕋᖅᓯᒪᓂᖅᓯᑦ	ᖃᐅᓄᐃᑦᓴᓂᖅᓯᑦ ᐅᓇᑕᒥ ᐃᓕᑕᕆᔨᐅᒪᓂᖅ	
	ᐅᓇᑕᒥ ᐅᑎᒪᔨᕋᓱᒃᑖᓄᑦ ᖃᐅᓄᐃᑦᓴᓂᖅᓯᑦ ᐊᒥᓱᓕᕆᓂᖃᖅᑕᐅᓯᒪᔪᑦ ᐅᒪᔨᐃᑦ - COSEWIC-ᑯᑦ ᖃᐅᐅᔨᖃᖅᓯᒪᔭᖅᓯᑦ	ᐅᒪᔨᐃᑦ ᐊᒥᓯᖅᓗᓂᕆᑕᖅᓯᒪᓂᓇᓄᑦ ᒪᑕᒃᖅᑐᒃ
Harris sparrow ᖃᑯᔨᓇᐊᖅ ᕼᐅᓗᑉ *Zonotrichia querula*	ᐊᐱᐅᔾᓯᑎᑐᒥᒃ ᐃᓯᒪᓕᑕᐅᓂᖅ	ᒪᑕᒃᖅᑎᔪᖏᒃᖅᑎᑐᑦ ᓯᑕ	ᓇᓗᓇᐃᖅᑕᐅᓯᒪᔨᑦ
Red-necked phalarope ᐊᐅᓚᓗᑐᖅ ᓴᐅᖃᖅ *Phalaropus lobatus*	ᐊᐱᐅᔾᓯᑎᑐᒥᒃ ᐃᓯᒪᓕᑕᐅᓂᖅ	ᐊᐱᐅᔾᓯᑎᑐᒥᒃ ᐃᓯᒪᓕᑕᐅᓂᖅ	ᓇᓗᓇᐃᖅᑕᐅᓯᒪᔨᑦ
Short-eared owl ᒥᑭᓂᓂᖅᓴᖅ ᐅᓗᐱᓗ *Asio flammeus*	ᐊᐱᐅᔾᓯᑎᑐᒥᒃ ᐃᓯᒪᓕᑕᐅᓂᖅ	ᐊᐱᐅᔾᓯᑎᑐᒥᒃ ᐃᓯᒪᓕᑕᐅᓂᖅ	ᓇᓗᓇᐃᖅᑕᐅᓯᒪᔨᑦ
Peregrine falcon ᖃᑭᓇᐅᔨᐊᖅ/ᑲᔪᓗ *Falco peregrinus*	ᐅᓗᓕᐊᓇᖅᑐᑦᓯᑎᑐᑦ	ᐊᐱᐅᔾᓯᑎᑐᒥᒃ ᐃᓯᒪᓕᑕᐅᓂᖅ	ᓇᓗᓇᐃᖅᑕᐅᓯᒪᔨᑦ
ᓂᕆᔨᑎᑦ ᐊᒪᓚᓗᑎᑎᓕᖅᑐᑦ ᐱᐊᖏᕆᒥᓂᖅ			
Caribou, barren-ground population ᓇᐸᑐᖃᖅᓯᑎᑐᕆᒥ ᑐᖅᑐᐃᑦ *Rangifer tarandus*	ᐅᓗᓕᐊᓇᖅᑐᑭᓕᕆᓂᖅ	ᒪᑕᒃᖅᑎᔪᖏᒃᖅᑎᑐᑦ ᓯᑕ	ᓇᓗᓇᐃᖅᑕᐅᓯᒪᔨᑦ
Atlantic walrus (Central/Low Arctic population) ᐊᐃᕃᖅ (ᖁᑎᖅᓴᓪᓕᐊᓂ/ᐊᑦᑎᐅᑐᓂᖅᓯᓐᓂ ᐅᑭᐅᖅᑕᖅ ᐱᑕᖃᖅᓂᖅᓯᓐᓂ *Odobenus rosmarus rosmarus*	ᐊᐱᐅᔾᓯᑎᑐᒥᒃ ᐃᓯᒪᓕᑕᐅᓂᖅ	ᒪᑕᒃᖅᑎᔪᖏᒃᖅᑎᑐᑦ ᓯᑕ	ᓇᓗᓇᐃᖅᑕᐅᓯᒪᔨᑦ
Beluga whale (West Hudson Bay population) ᖁᓇᓗᒪᐃᑦ (ᐱᖏᒪᓇᖏᓂ ᕼᐊᓕᓐᓄᐃᑦ) *Delphinapterus leucas*	ᐊᐱᐅᔾᓯᑎᑐᒥᒃ ᐃᓯᒪᓕᑕᐅᓂᖅ	ᒪᑕᒃᖅᑎᔪᖏᒃᖅᑎᑐᑦ ᓯᑕ	ᓇᓗᓇᐃᖅᑕᐅᓯᒪᔨᑦ

ᑕᐃᔨᐅᓂᖅᑐᑦ ᐃᓄᐊᑎᑦ ᖃᐅᔨᓴᖅᑎᓄᑦ ᐅᖃᐅᓯᐊᒍᑦ ᑎᑎᕋᖅᓯᒪᓂᖅᑐᑦ	ᖃᐅᓄᐃᓕᓚᐅᓂᖅᑐᑦ ᑲᓇᑕᒥ ᐃᓕᓴᕆᐊᖃᒪᓂᖅ		ᐱᑕᖃᖅᓂᖅᑐᑦ ᐃᑲᑦᑐᐊᓕ ᑎᖅᒥᐊᖃᖅᐸᖅᓱᓕᒃ
	ᑲᓇᑕᒥ ᑲᑎᒪᔨᖏᓪᓗᓂᑦᑕᐅᓯᒪᔪᑦ ᐃᒪᓕᔪᐃᑦ - COSEWIC-ᓄᑦ ᖃᐅᔨᔪᖃᓯᒪᔭᖏᑦ	ᐅᒪᔪᐃᑦ ᐊᒥᓯᖏᓪᓗᐊᓕᕆᓂᖅᐹᓂᓄᑦ ᒪᓕᑐᖏᔪᑦ	
Bowhead whale (Eastern Canada-West Greenland population) ᐊᖏᕐᕿᑦ (ᑲᓇᖏᓇᖅᐸᓯᖏᓚᓂ ᑲᓇᑕᐅᔪ- ᑕᓪᐅᖏᑐᓂ - ᐊᑯᑭᑐᐃᑦᓗ ᐱᖏᓚᖏᓇᖅᐸᓯᖏᓚᓂ ᑕᓪᐅᖏᑐᓂ ᐊᖏᕐᕿᑦ) *Balaena mysticetus*	ᐊᔪᐅᖏᕐᕋᑐᒃ ᐃᓱᒪᓕᑕᐅᓂᖅ	ᒪᓕᑐᖅᑎᒍᖏᓚᖏᕐᕋᑦ ᓱᓕ	ᓇᓗᓇᐃᖅᑕᐅᓱᒪᓕᔪᑦ
Narwhal ᑐᐃᓕᑦ ᖃᓇᓗᖏᐃᑦ *Monodon monoceros*	ᐊᔪᐅᖏᕐᕋᑐᒃ ᐃᓱᒪᓕᑕᐅᓂᖅ	ᒪᓕᑐᖅᑎᒍᖏᓚᖏᕐᕋᑦ ᓱᓕ	ᐱᑕᖃᑐᐃᓇᓚᐊᓕᑦ
Polar bear ᓇᓄᖅ *Ursus maritimus*	ᐊᔪᐅᖏᕐᕋᑐᒃ ᐃᓱᒪᓕᑕᐅᓂᖅ	ᐊᔪᐅᖏᕐᕋᑐᒃ ᐃᓱᒪᓕᑕᐅᓂᖅ	ᓇᓗᓇᐃᖅᑕᐅᓱᒪᓕᔪᑦ
Ringed seal ᓇᑦᑎᑦ *Pusa hispida*	ᐊᔪᐅᖏᕐᕋᑐᒃ ᐃᓱᒪᓕᑕᐅᓂᖅ	ᒪᓕᑐᖅᑎᒍᖏᓚᖏᕐᕋᑦ ᓱᓕ	ᓇᓗᓇᐃᖅᑕᐅᓱᒪᓕᔪᑦ
Wolverine ᖃᑲᕐᓚᒃ *Gulo gulo*	ᐊᔪᐅᖏᕐᕋᑐᒃ ᐃᓱᒪᓕᑕᐅᓂᖅ	ᐊᔪᐅᖏᕐᕋᑐᒃ ᐃᓱᒪᓕᑕᐅᓂᖅ	ᐱᑕᖃᑐᐃᓇᓚᐊᓕᑦ
ᐃᖃᓗᐃᑦ			
Lumpfish ᖃᔪᖅᓱᖅᑐᖅ ᓂᐱᓴ *Cyclopterus lumpus*	ᐅᓗᓚᐊᓇᖅᑐᒋᑕᖃᓂᖅ	ᒪᓕᑐᖅᑎᒍᖏᓚᖏᕐᕋᑦ ᓱᓕ	ᐱᑕᖃᑐᐃᓇ-ᓚᐊᓕᑦ
Northern wolffish ᐊᑦᔪᐊᖅ/ᐊᑦᔪᐊᕆᓴᓗᒃ *Anarhichas denticulatus*	ᐅᓗᓚᐊᓇᖅᑐᒋᑕᖃᓂᖅ	ᐅᓗᓚᐊᓇᖅᑐᒋᑕᖃᓂᖅ	ᐱᑕᖃᑐᐃᓇᓚᐊᓕᑦ
Spotted wolffish ᑕᖅᓴᑐᖅᓂᖅᓴᖅ ᐊᑦᔪᐊᖅ/ᐊᑦᔪᐊᕆᓴᓗᒃ *Anarhichas minor*	ᐅᓗᓚᐊᓇᖅᑐᒋᑕᖃᓂᖅ	ᐅᓗᓚᐊᓇᖅᑐᒋᑕᖃᓂᖅ	ᐱᑕᖃᑐᐃᓇᓚᐊᓕᑦ
Atlantic wolffish ᑐᑭᒧᐊᑐᓂᒃ ᑎᒥᖅᒪᔨᑦ ᑕᖅᓴᓕᒃ ᐊᑦᔪᐊᖅ/ᐊᑦᔪᐊᕆᓴᓗᒃ *Anarhichas lupus*	ᐊᔪᐅᖏᕐᕋᑐᒃ ᐃᓱᒪᓕᑕᐅᓂᖅ	ᐊᔪᐅᖏᕐᕋᑐᒃ ᐃᓱᒪᓕᑕᐅᓂᖅ	ᐱᑕᖃᑐᐃᓇᓚᐊᓕᑦ

ᑕᐃᕆᐅᓂᖅᑎᑦ ᐃᓄᐊᑎᑐᑦ ᖃᐅᔨᓴᖃᑎᓄᑐ ᐅᖃᓕᕆᓂᖅᑎᑦ ᑎᑎᕋᖅᓯᒪᓂᖅᑎᑦ	ᖃᐅᓄᐊᑕᖏᓂᖅᑎᑦ ᑲᓇᑕᒥ ᐃᓕᑕᒪᔭᐅᒪᓂᖅ		ᐱᑕᖃᖅᓂᖅᑎᑦ ᐃᑲᑐᐃᐊᓕ ᑎᖅᑕᐊᖃᖅᐸᖏᓂ
	ᑲᑎᒪᔨᕐᓭᓂᖅᑎᑦ ᐃᓇᓗᒃᑐᑦ ᖃᐅᔨᓴᖃᑎᓄᑐ ᐊᑕᓇᓵᖅᓯᓂᖅᑐᑦ ᐅᒪᔨᐃᑦ - COSEWIC-ᓄᑦ ᖃᐅᔨᓴᖃᓇᓕᔪᖅᑎᑦ	ᐅᒪᔨᐃᑦ ᐊᑕᓇᓵᖅᑐᐃᓕᕆᓂᖅᑎᓄᑦ ᒪᑕᒐᖅᑐᓄᑦ	
Thorny skate ᑕᓗᐅᐸ ᓇᑎᐊᓂ ᖃᐅᑕᕋᓪᓕ ᑲᑭᑦᓭᐅᑎᖃᖅᑐᑐᒃᑯᑦ ᖃᐅᖅᑐᒍᑦ – ᐃᖃᐅᓚᐅᔪᓐᓇᖅᑕᐅᔭᖅᑎᑦᓱᑦ, ᖃᐅᓪᓗᓇᑎᑐᑦ ᑕᐃᔨᐅᔪᑦ *Amblyraja radiate*	ᐊᐱᐅᐅᖅᑭᑐᑐᒥᒃ ᐃᓕᓴᐲᑕᐅᓂᖅ	ᒪᑕᒐᖅᑎᒍᖏᓐᓄᑭᑐᑐᑦ ᓴᓪᓕ	ᐱᑕᖃᑐᐃᓪᓇᓗᐊᓪᓕᑦ

2.6 ᑕᒫᓂᖃᐅᑕᑐᖃᐅᔪᖅᑭᑦᑐᑦ ᐅᒪᔨᐃᑦ/ᖃᐅᔨᐱᖃᑭᐃᓗᓪᓚᑦ, ᐱᑭᖅᑐᓪᓗᓪᓚᑦ ᐱᑕᖃᐂᓂᖅᑭᑦ

ᓇᑭᑐᐃᓇᖅ ᐃᓄᐃᑦ ᐱᑕᓚᖃᑕᐂᓂᔪᓄᑦ ᓴᐳᑎᑦᑎᓗᑕᐅᔪᒃᓴᐳᑦ ᐱᑕᖃᑕᑭᓪᓚᖅᑭᔪᓂᓪᓚᑦ ᐱᑕᖃᖅᑲᓐᓂᕐᓗᑕᐅᔪᑦᑕᓂᑦ ᓄᓇᖅᑎᓪᓚ, ᐱᑭᖅᑐᓂᓪᓗᓪᓚᑦ, ᐅᒪᔨᓂᓪᓗᓪᓚᑦ, ᖃᐅᑦᐱᖃᑦᓂᓪᓗᓪᓚᑦ (Canadian Endangered Species Conservation Council, 2011). ᓄᓇᕗᑦ ᑲᖏᒪᐃᑦ ᑎᐅᐅᐊᖅᓯᓯᒪᓕᖅᑲᓴᐳᑦ 14-ᓂᐅ ᓄᓇᕆᑦᓂᐅ ᐱᑭᖅᑐᓂᐅ ᓄᓇᕗᒋᑦ ᐱᑕᖃᖅᑐᑐᖅᑎᑦᑕᐅᓇᖅᑐᖅᑐᑦᓂᐅ ᐱᑕᖃᑕᖅᑐᓂᐅ. ᑕᒪᐅᐅᐊ ᐱᑭᖅᑐᑦ ᓄᖏᑎᓚᖖᒋᓕᐊᓯᒪᓂᖅᑐᑦ ᐊᐱᐸᖅᑐᑦᑐᑎᐅᔪᓂᐅ ᐱᑭᖅᐸᐅᑐᑦᖃᐅᔪᓂᐅ ᓄᓇᕗᓴᒋ. ᖃᐅᔨᓴᒪᔪᐅᖅᑐᑦᑐᑦ ᑕᒫᓂᖃᐅᑕᑐᖃᐅᔪᖅᑭᑦᑐᑦ ᐱᑕᖃᑕᓴᖓᑎᒃ ᐃᑲᑦᑐᐃᓪᓇᓗᐊᓪᓕᑦ ᐱᑭᖅᐸᐅᑐᑦᖃᐅᔪᓂᐅ ᓄᓇᕗᓴᒋ. ᖃᐅᔨᓴᖃᑎᓄᑐ ᑕᒫᓂᖃᐅᑕᑐᖃᐅᔪᖅᑭᑦᑐᑦ ᐱᑕᖃᑕᓴᖓᑎᒃ ᐃᑲᑦᑐᐃᓪᓇᓗᐊᓪᓕᑦ ᐱᑕᖃᑕᖅᑐᓂᐅ.

3 ᐃᓕᓴᕆᔨᑐᖏᓪᓗᒍ ᐊᑐᓚᓱᐅᓕᖅᓯᓂᖅ ᓄᓇᒥᐅᑦᑕᐃ

ᑎᖅᒥᐊᖏᖅᓴᖅᒥ ᐱᑕᖏᖅᑎᑕᑎᓂᖅ ᐱᓱᐊᑎᖏᓵᐊᖅᓯᑦ ᐃᔨᓪᒪᑦᕆᐅᓂᖅᓯᓂᖅ ᓴᓯᓂᐊᖅᑕᐅᖏᓐᓇᕐᓂᖅ ᑎᖅᒥᐊᖏᓂᖅ ᓄᓇᕗᖅᓯᖅᑕᖅᓯᓂᖅᓴᓗ, ᑭᓱᐊᓂᒥ ᑕᐃᖅᑯᐊ ᐃᓄᐃᑦ ᐃᑲᔪᓱᐊᖅᑕᕐᑎᑲᖅᖁᓂᖅᓄᓂ ᐊᖅᑭᑕᕐᓂᓴᓂᑕᐅᑭᕐᓂ ᐃᓄᐃᑦ ᐃᓕᒪᕆᐅᑦᓭᓚᖅᓯᖅᓂᖅᓂ ᐊᐅᑎᖅᑭᖅᓄᓂ. ᐃᓄᐃᑦ ᖃᐅᔨᓴᓵᑐᖃᖅᑭᒃᑲᒪᑦ ᐊᑐᑎᓕᒪᓚᐊᓗᖓᒪᑕ ᖃᐅᔨᓴᓵᕆᐊᓂᖅᓯᓂᖅᓴᓗᒍ ᓄᓇᕗᒃ, ᐃᒪᖏᓂᖅ, ᓄᓇᕗᐅᑕᓚᕆᐅᔨᓂᖅᓴᓗ ᖃᐅᓂᕆᖅᑭᒃᑲᓂᖅᓂ, ᑐᖅᑲᐃᐅᓗᑎᑎᒃ ᑐᑭᓯᐊᖅᑭᖅᑑᑎᓂᖅᓂᖅᓯᓂᖅᓴᓗᒍ ᐃᓕᒪᓕᐅᕈᑕᐅᔪᓚᐊᖏᖅᓴᖅᑐᐊᖅᑐᓂᖅᑐᓂᖅᑐᖅ. ᑕᒃᑯᐊ ᑎᑎᕋᖏᓚᖅᓂᑕ ᐅᖏᖅᐅᔨᖏᖅᓴᐅᑦ ᑕᐃᓴᔨᓚᓂᐅᓕᐅᖅᑐᖅᓴᖅ ᐅᑦᓗᒥᐅᓕᖅᓂᖅᓴᓗ ᓄᓇᖅᓂᒃ ᐊᑐᖅᑕᕐᐅᖅᓯᖅᓂᖅᓂᖅᓴᖅ ᑎᖅᒥᐊᖏᖅᓴᖅᐃᓕᐅᖅ, ᐊᐅᓚᐊᓂᓗ ᓄᓇᖅᓂᖅ, ᐊᓪᓚᓗ ᖁᑎᕋᐅᐊᖅᑕᐅᖅᑕᕐᖅᓯᓚᖅᓂᖅᓂᖅᓴᖅ ᖃᐅᔨᓴᓵᓕᐅᕈᑕᖅᑐᖅᓴᖅ ᐱᑕᖃᖅᑐᖅᓴᖅ, ᑕᑯᔪᖃᐅᔨᐅᕐᑐᖅᑐᖅᓴᖅ ᑕᑯᔪᖃᐅᔨᐅᓚᖅᓯᓂᖅᓴᓗ ᐃᓕᒪᓕᐅᕈᑕᖅᑐᖅᓴᖅ ᐊᑐᖅᑕᕐᓂᖅᓴᖅ.

ᐊᐅᓪᓕᓐᓂᐊᕐᑲᖅᑐᑐᖅᑲᐃᓪᓕ, ᓄᓇᐃᓪᓕ, ᐃᓕᐃᓪᓗ ᑕᐃᔪᓱᐊᖅᑐᖅᓯᓂ, ᖃᐅᔨᓴᓵᓂᕐᓂᓗ ᓱᓕᒥᖅ, ᐅᓕᓴᐃᓪᓗ ᐱᐅᓱᓚᐊᖅᑕᕐᓂᖅᓂᖅ, ᐅᓵᑐᑎᑐᐃᓐᓇᐅᐅᖅᑕ ᐅᑦᑐᓴᓚᐊᐅᖅᑕᕐᓂᖅᓂᖅᓴᖅ, ᑕᓪᓕᐊᓱᓕᐅᕈᑕᖅᑐᖅᓴᖅ ᖃᐅᔨᓴᓵᑎᑐᐃᓐᓇᐅᐅᖅᑕ ᐃᓄᐃᑦᑐᖅᑕᐅᖅᑕᖅᑐᖅᓴᖅ, ᐱᕐᔪᓐᓇᖅᑐᖅᓴᖅ, (ᓱᓕᓴ ᓂᕆᑭᓂᖅ ᐱᑭᓴᐃᓐᓂᖅ, ᐃᖃᓴᓐᓂᖅ ᓴᓗᐅᓂᖅ, ᖃᓄᖅᑯᐃᑐᖃᖅᑐᖅᓴᖅᓂᖅᓂᖅᓴᖅ, ᐃᓄᐃᓱᓐᓂᖅᓴᓗ), ᑕᐃᔨᓱᓕᓂᒋᓪᓕᐃᓕᓴᓗ ᐃᓄᐃᖅᑕᕐᓂᖅᓯᓚᐅᕈᑕᖅᑐᖅᓴᖅ ᓄᓇᖅᓂᖅ ᑕᓪᓕᕆᖅ ᑕᐃᔨᓴᓵᐅᕈᑕᖅᑐᖅᓴᖅᑎᑦᓴᓗ ᐊᐅᑎᖅᑭᖅᑐᓂᖅᓴᓗ ᖁᑎᕋᐅᐊᖅᑕᐅᖅᑕᕐᓂᖅᓂᖅᓴᖅ ᐃᓄᐃᑦ ᓄᓇᕗᒃ ᐊᑐᖅᑕᕐᓂᖅᓴᖅᑎᑦᓴᓗ ᐃᓄᐃᖅᑕᕐᓂᖅᓂᖅᓴᖅ. ᐱᓪᓚᑕᐊᓚᐅᖅᓴᖅ ᐃᓄᐃᖅᑕᕐᓂᖅᓴᖅᐃᓕᐅᖅᑕᕐᓂᖅᓂᖅᓴᖅ ᐃᓄᐃᑦ ᓄᓇᕗᒃ ᐊᑐᖅᑕᕐᓂᖅᓴᖅᑎᑦᓴᓗ ᐃᓄᐃᖅᑕᕐᓂᖅᓂᖅᓴᖅ, ᐃᓄᐃᑦ ᑕᐃᔨᓴᓵᐅᕈᑕᖅᑐᖅᓴᖅ

ᑐᓴᓪᓗᕕᒃᑯᓪ ᒪᓕᑦᑲᖅᑎᑕᐅᕐᒪᑦ ᐊᒋᔪᓪ ᑎᑎᕋᖅᓯᒪᓂᖅᑎᓂᒃ 6, ᐃᓄᐃᑦ ᐃᑲᔪᓯᐊᕐᓴᖅᓄᑦ ᐊᓴᑎᕐᔨᑎᔪᓪ ᑎᑎᕋᖅᓯᒪᔪᕐᓇᖅᑐᖅᓯᒪᓕᖅᑐᑦ ᖃᑦᑕᕐᔪᐊᖅᑐᕐᓂᖅ ᓄᓇᒥᐅᑕᓚᐅᕐᔪᐊᓘᑦ ᐱᓚᓗᐅᔪᐊᓘᑦ ᐃᓄᖅᓄᑦ ᑎᖅᒥᐊᖅᑲᐃᐅᕐᓂᖅ ᐊᑐᓗ ᐅᓪᓗᓕᒫᖅᑐᑦ ᓄᓇᕗᒻᒥ. ᑕᓚᑦᑯᐊ ᐱᑕᖅᑲᑯᐅᐅᖅᓯᑎᑦ ᖃᐅᔨᓕᐅᔪᐊᓘᑦ ᐃᑲᔪᖅᓱᐃᓂᕐᒧᑦ ᐊᐅᓕᑕᐅᓂᖅᑎᓄᑦ ᐊᑐᓗ ᓴᐳᓂᐊᖅᑕᐅᓱᒪᕐᓇᓕᑦ ᓇᓂᑐᐃᓇᖅ ᓄᓇᕗᒻᒥ, ᐃᑲᔪᑕᐅᓗᑎᓗ ᑐᐱᓱᐅᓕᕐᑎᑕᐅᓇᖅᑐᓂᖅᓄᑦ, ᐊᓚᓗ ᑎᑎᖅᑯᑎᑭᕐᔨᐅᓂᐊᖅᑐᑐᒃ ᐱᑕᖅᑯᓂᐊᖅᑕᐅᓂᐊᖅᑐᑦ, ᐃᓕᖅᑯᓱᑐᖅᑎᒧᑦ ᐱᓚᓗᐃᐅᓂᐊᖅᑐᑐᒃ ᐃᓄᖅᓄᑦ, ᐃᑲᔪᑕᐅᓗᑎᓗ ᐱᓕᕐᓂᕐᔪᐊᑕᐅᓇᖅᑐᑦ, ᐊᑐᖅᑕᐅᔪᓇᖅᓯᒪᓂᖅᓄᑦ ᐃᓗᓕᑕᐅᓗᑎᓗ ᐱᓕᕐᓂᕐᔪᐊᑕᐅᓇᖅᑐᑦ, ᐊᑐᖅᑕᐅᓗᑎᓗ ᑕᒪᓐ ᓴᐳᓂᐊᖅᑕᐅᓱᒪᕐᓇᓕᑦ. ᐃᓂᖅᓄᑦ ᐃᑲᔪᑕᐅᓗᑎᓗ ᐱᓕᕐᓂᕐᔪᐊᑕᐅᓇᖅᑐᑦ ᐃᓕᓴᕐᔪᐊᓘᑦ ᐃᓄᖅᓄᑦ ᐃᓕᓴᕐᔪᐊᓘᑦ ᐱᓕᕐᓂᕐᔪᐊᑕᐅᓇᖅᑐᑦ, ᐃᓄᖅᓄᑦ ᐃᓕᓴᕐᔪᐊᓘᑦ ᐱᓕᕐᓂᕐᔪᐊᑕᐅᓇᖅᑐᑦ

ᐅᑭᐅᓚᓴᖏᓂᓂ ᑕᒪᓂ 1902-1903 ᐊᐢᓂᐊᕆᓂᐊᓗᖏᓪᓗᑦ ᖃᓂᒪᓂᐊᓗᖏᓪᓗᑦ ᑎᑭᑕᓂᓚᓗᖏᓪᓗᑦ, ᑐᕆᔪᖅᑐᐊᔪᓴᓕᑎᒃ ᑕᒪᖅᑎᕆᑐᓚᐅᖅ ᓴᖅᑭᓂᓚᐅᖅᑐᑦ. ᐊᖅᓯᐢᓂᐊᖅᑏᑦ ᐊᐢᓂᐊᕈᑎᓂᒃ ᑎᑭᐅᓴᐃᔨᕐᓂᐅᔪᖅᓴᐅᒥᒃ. ᐃᑲᔪᖅᑎᖅᑭᐢᓂᕆᒃ ᐃᓇᐢᓂᒃ ᑕᐅᓯᓂ, ᐊᖅᓯᐢᓂᐊᖅᑏᑦ ᐃᓇᐢᓂᒃ ᑎᑭᑎᑦᑎᓕᓕᐅᖅᑐᑦ ᓇᐅᔭᖅᑎᕆᐅᐢᓂᒃ, ᐅᒃᑯᐢᓴᖅᑎᕆᐅᓂᓗ ᑕᒪᓂ 1908-ᒥ ᓴᖅᓕᓂᑦ, ᓴᓇᔨᐅᑎᑕᐅᓕᖅᓴᑎᒃ.


1924-ᖁᑎᖅᓗᒍ, ᐄᐊᑦᓴᐢᐋ ᑲᒻᓯᓂᒃᑯᓂ ᓂᐅᕐᓕᖅᑎᒃᑯᓂ ᓄᓇᕗᑦᑎᑎᒃᑯᓂ ᓂᐅᕐᓕᓴᖏᓪᓂᒃ ᐊᒃᓯᑐᖅᓴᐊᖅᑎᓪᓕᐅᖅᑐᒃ ᓴᖅᓕᓂᑦ ᐃᓇᐃᓗᑐ (ᖃᑭᑲᒃᑰᓐᖏᑎᒃ [Baffin-ᒥᐅᑦ] ᓂᕆᐊᓂᖅᑎᐅᓂᒃ) ᓄᒃᑎᖅᑲᕈᑕᓇᖅᓴᓪᓚᒪᑕ ᓂᐅᕐᓕᖅᑎᒃᑯᓂᓇᓂ. ᑎᓚᒐᓂᐊᖅᑲᕈᓂᖅ ᐱᒻᓚᓚᐊᑐᓴᕈᑕᐅᔪᓂᓇᖅᑐᓂᖅᓗᑦ, ᓂᐅᕐᓕᐅᑎᒃᓴᓂᑦ ᐄᐊᑦᓴᐃᒃᑯᓂ ᐅᓇᕐᓕᖅᑎᒃᑯᓂ ᐅᓇᕐᓕᖅᑎᒃᑯᓂ (ᑕᑎᒥᒃ, ᖃᒃᓴᔪᓂᒃ, ᐊᓯᖏᓐᓂᒻᓗᓗ). 1940ᑦ ᐊᖅᓴᐃᐅᑦ ᐃᓯᖅᓴᓯᖏᓪᓂ, ᐊᒥᐊᑕᒪᒥᐅᑦ ᒥᑦᑕᖅᓴᐃᕈᑕᐅᕈᑕᓇᖅᑐᓂᑦ ᓯᕈᑎᑎᑦᐊᖏᓪᓂ ᓴᖅᑦᑦ. ᑭᖓᓂᐊᐃᑦ ᒪᓇᕈᑕᑦ ᓴᖅᓕᒪᓂᖅᓯᖏᓐᓄᑦ ᐊᐅᕈᑎᑕᓇᖅᑐᓂᑦ ᑕᒪᒃᑯᓂᖏᓪᓗ ᓯᕗᓪᓗ ᐃᕐᓂᐢᓂᐊᖅᑐᑦ, ᐋᑲᓂᐊᖅᑐᓂᑕᓚᓂᖅᑎᒃ, ᑭᓇᐅᔪᕐᓂᐊᖅᑐᑦ, ᑭᓇᔪᕐᓂᐅᕐᓂᖅᓴᓪᓗ ᐃᒃᖅᑎᑎᒍᑦ ᐊᑭᕐᓂᖅᑐᐃᔨᓂᒃ, ᐊᒥᓯᑦ ᓄᓇᒥᐅᑕᒪᔭᐃᑦ ᓯᕈᒪᒃᓴᑦ ᓴᖅᑦᑦ ᓄᓇᖅᓴᓂᑦ ᐱᓕᕐᓂᖅᑐᑦ.

ᓂᐅᕐᓕᖅᑎᖅᑲᕈᐅᓂᖅᓯᓪᓚᓂ ᓯᓂ ᓴᖅᑕᓂ, ᓴᖅᑕᓂᒥᐅᑦ ᐃᓄᓯᓴᖅᓴᖅᑕᕈᐅᓂᖅᑐᑦ ᓄᓇᒋᓴᖅᑕᓂᒥᓂᒥᑦ ᐊᐅᖅᓯᓄᖅᑕᓂᖅᑐᖅᓂᓪᓚᒥ, ᒪᑕᖅᑐᓂᓪ ᐱᑕᓴᖅᓂᖅᑐᓂᓪ ᐅᒪᔨᐃᑦ. ᐱᓗᐊᓯᐊᖅᑐᑦ ᓴᖅᑕᓂᒥᐅᑦ ᑕᓚᐅᖅᒥᐅᑕᓂᓪ ᓂᕆᐹᖅᓴᖅᓂᖅᓴᖅᑎᐅᖅᑕᖅᑐᓂᖅᑕᐃᓇᖅᓴᐅᖅ ᐃᓄᐃᑦᑎᐊᕐᓂᓄᑦ ᐃᓄᐃᑦᑎᐊᕐᓂᖅᑐᑦ ᐅᑦᑎᓂᓪ, ᖃᓄᐃᓗᑎᓂᓪ, ᐊᐃᐃᓚᓂᓪ (Bird, 1953; Collins, 1956). ᐃᓯᒪᒋᐅᐅᓕᓕᐅᖅᑎᑎᓪᓗᑎᓪᓗ ᓯᓂ ᖃᐅᐅᓴᖅᑎᓄᑦ ᑲᖓᐃᑦ, ᑕᓚᐅᖅᑎᐅᑕᓂᒃ ᐅᐃᐅᓂᒃ ᐊᖓᔪᓇᓯᓂᒃᐸᐅᒥᓂ ᑕᐃᒪᖓᑦ ᑎᑎᕐᓂᐅᓕᖅᑐᐅᓂᖅᓴᐅᓕᖅᑐᐅᓂᖅᑕᐃᓇᓕᖅᑐᒃ ᐅᑦᑎᓂᒃ ᐊᖓᔪᓇᓯᓂᒃᐸᐅᒥᓂ ᐃᓄᐃᑦᑎᐊᕐᓂᐅᓕᖅᑐᐅᓂᖅᑕᐃᓇᓕᖅᑐᒃ ᐅᑦᑎᓂᒃ ᓄᖅᑕᖓᖅᓴᓪᓗ

ᐃᓄᖅᑲᐃᑦᑐᓕᕆᔨᖅᑐᑦ ᐊᒻᒪᓗ ᐃᓐᓇᓴᒧᑎᖅᓴᖅ ᓄᓇᒋᐊᑦᑐᓕᕆᔨᖅᑐᑦ ᖃᖅᖢᓴᕆᔨᖅᑐᑦ ᑎᒥᐊᖅᑲᐃᖅᓱᓂ (ᑕᑯᔭᖅᓴᖅ 6). ᐃᓄᖅᑲᖅᐸᓕᕆᔨᖅᑐᑦ ᓄᓇᐃᑦ ᓄᓇᖏᐊᖅᓂᖅᑕᑎᓕᕆᔨᖅᑐᑦ ᖃᖅᖢᓴᕆᔨᖅᑐᑎᖓ ᐊᒥᓱᑎᓴᐃᑦ ᐃᓄᔭᓱᐃᑦ ᐊᒻᒪᓗ ᓂᕐᕿᑎᓂᔭ ᐳᓪᓕᓕᐊᖅᓕᓕᓂᑦ ᑎᒥᐊᖅᑲᐃᖅᓱᐸᑦ ᓄᓇᖅᑭᓂᔭ. (ᑐᑭᓱᑦᐊᖅᓕᕆᓱᒪᓕᕐᓂᖅ: ᐅᐃᓕᓴᒻ ᑭᐊᒻᐸ-William Kemp, ᑐᖓᓕᖅᔪᑦ ᐃᓕᓴᔪᓯᑐᖅᑎᒍ ᐱᕐᔪᑎᓕᓂᖅᑲᕐᖓᑭᓂᔭᓂᖅ ᑐᑭᓱᑦᐊᖅᓕᕆᔨᖅᑐᑦ, ᑐᑭᓱᑦᐊᖅᓕᕆᔨᖅᑐᑦ).

ᐅᐊᖅᓇᖅᑕᓱᖓᓕ ᖁᑭᓕᖅᑲᖅᓕᖅᖢ ᑎᖅᑎᐊᖅᑲᖅᓕᕆᔨᖅᐸᑦ ᐊᑕᓚᖅ ᓂᕈᖅᑕᓱᓕᓄᑦ ᐅᑭᓕᕆᔨᖅᓯᖅᓱᖓᓕᓄᑦ ᑐᐃᑐᓄᑦ ᐊᒻᒪᓗ ᑕᒪᓕᕆᔨᖅᓯᖅᓱᖓᓕᓄᑦ ᐅᐱᓕᕆᔨᖅᓯᖅᓱᖓᓕᓄᑦ ᑐᐃᑐᓄᑦ ᐊᑎᖅᑕᐅᕆᔨᖅᑐᑦ ᑕᒪᓐᓇ. ᑕᐃᒪᐃᓱᐊᖅᑐᑦ ᐃᓄᖅᑲᐃᖅᑐᑦ ᐃᓄᖅᑲᐃᖅᑐᑦ ᐊᑎᖅᑕᐅᕆᔨᖅᑐᑦ ᑕᒪᓐᓇ. ᑕᐃᒪᓐᓇᑦ ᐃᓄᐃᑦ ᑐᐃᑐᐃᓱᐊᖅᑐᑦ ᐃᓄᖅᑲᐃᖅᑐᑦ ᐊᑎᖅᑕᐅᕆᔨᖅᑐᑦ ᑕᒪᓐᓇ. ᓴᓪᓕᖅᑎᐅᕆᔨᖅᑐᑦ ᐃᓄᖅᑲᐃᖅᑐᑦ ᐊᑎᖅᑕᐅᕆᔨᖅᑐᑦ ᑕᒪᓐᓇ.


ᖃᒃᓗᓈᑎᑐᑦ ᑎᑎᖃᑎᒍᑦ	ᐃᓄᒃᑎᑐᑦ ᑎᑎᖃᑎᒍᑦᑕ	ᓯᓚᐅᓂᖅ/ ᖃᓄᐃᔪᓂᖅ-ᓂᖅᓗᓪᓚᕆᓂ	ᐃᓕᑕᕆᓚᐅᓯᒪᓕᕐᓂᖃᑦᑐᑦ ᑕᐃᔨᐅᓂᖅᑭᑦ	ᐅᓂᒃᑲᓚᐊᒃᖃᓂᕈᑎᑦ
Qikiqtakuluk	ᖁᑭᖅᑕᑯᓗᒃ	ᖁᑭᖅᑕᖅ		ᐅᓪᓗᖅᓴᖅᓄᑦ ᒥᑎᖅᓄᑦ
Itijujjuaq	ᐃᑎᓴᔪᐊᖅ	ᖃᖅᖃᓂᖏᓐᓂᖅ	ᐅᔨᕋᐃᑦ ᖃᖅᖃᓂᔨᑦ ᑎᓚᖅᑯᐊ	ᐊᖏᑭᓯᐊᓗᒃ ᑎᓚᖅᑯᖅ. ᓴᖅᑐᓂᖅᓱᖏᓪᓗᓗ ᑭᖑᓕ, ᖃᖅᖃᓂᔨᑦ ᐅᔨᕋᐅᓂᐊᓗᑦ ᐃᓯᐊᓂ
Nasipaariarvik	ᓇᓯᓵᓚᐊᖅᓴᒃ	ᖃᖅᖃᖅ		ᖃᐅᔨᓴᖅᓴᒃ. ᓴᖅᑐᓂᑐᒪᒥ ᐊᓯᖏᑭᓂᒃ ᑕᒧᓄ ᐊᖅᑎᒃᑐᑕᐅᔨᓂ ᓄᓇᒥ
Qakutajak	ᖃᑯᖅᑕᕐᒃ	ᖃᖅᖃᖅ	ᖃᑯᖅᑕᕐᕆᖅ ᖃᖅᖃᓂᔨᖅ	ᖃᒃᐅᓪᓗᑐᓂᒃ ᐅᔨᕋᓕᒃ ᐅᒃᑯᓯᓕᕐᔨᐅᓂᖏᓐᓂᓗ ᖃᑯᖅᓂᖅᓴᓂᒃ ᐅᔨᕋᖏᑦ
Qaqsauqtuuq	ᖃᖅᖅᓴᐅᖅᑐᖅ	ᓄᓕᐊᖅᓴᒃ		ᐱᑕᖃᖅᓂᕋᐃᓂᖅ ᒪᖁᑲᖏᓂᒃ ᖃᖅᖅᓴᐅᖅᓂᒃ ᑎᖏᒥᐊᖅᖃᖅᖅᓴᐅᓂᖅᑎ ᐃᓕᑕᕆᓚᐅᓯᒪᓕᕐᓂᖃᑦᑐᑦ, ᐊᓖᒪᓗ ᑕᓯᐅᔨᕋᐊᕐᓴᓂ, ᐊᒥᓯᐊᓗᐃᑦ ᐊᖏᓐᖑᕐᖑᑕᑦ ᑎᖏᒥᑦ ᐅᑎᖅᓯᒃᑕᑦ ᑕᒪᐅᖏᓐ ᐅᐱᖏᓗᐅᓴᒃᑯᑦ: ᑲᖓᐃᑦ, ᓂᕆᓪᓕᐃᑦ, ᒥᑲᑎᓗ. ᐃᓄᐃᑦ ᐱᑭᐅᑕᓚᐊᖅᓯᓪᓕᐅᖅᑕᖅ ᑕᒪᐅᖏᓐ.
Tajjaqturliq	ᑕᖏᖅᑕᕆᓪᓕᒃ	ᐅᔨᕆᓪᓂᔪᐊᕆᓗᐃᑦ		ᑐᒃᑕᓯᐅᕆᓇᑭᓪᓕᐅᖅᑕᖅ
Tasialuk	ᑕᓯᐊᓗᒃ	ᑕᓯᖅ	Tasialuk Lake	ᑕᓯᕆᒥᐅᓂᒃ ᐃᕆᓄᓗᓪᓚᑕᒪᓂ



ᑕᐱᓂᐊᖏᑎᒋᔪᓂ: ᑕᵃᵒᒐᐃᑲᓂ. 2014



ᑕᐱᓂᐊᖏᑎᒋᔪᓂ: ᓄᓇᐃᑦ ᐃᓄᒃᓄᑦ ᐊᑕᖅᑕᐅᓯᒪᓂᖅᓯᓂᖅᐳᖅ ᐱᑕᕆᐊᒃᒋᒪᑕᐅᖅᑕᑦ. ᐃᓄᑕᕆᔨᑕᖅᓂᓄᑦ 1976




ᑕᐱᓂᐊᖏᑎᒋᔪᓂ: ᓄᓇᕗᒻᒥ ᓄᓇᒃᓴᐊᑕᐅᖅᑎᑐ. ᓇᐅ-Riewe 1992

ᑕᯇᔭᕆᖅ 6: ᑕᐃᒪᓐᓇᓂᒃ ᐃᓄᐃᑦ ᐊᑐᖅᑕᕐᕕᓂᖅᓂᑦ ᓄᓇᒥᒃ ᓴᕐᓂᐅᑦ ᖁᑭᖅᑕᒧᑐᐊᓂᒃ.

3.3 ᐃᓐᓇᐅᓕᖅᑐᖅ ᐃᓄᐃᑦ ᐊᑐᖏᑦᑕᕐᓂᖅᐸᑦ ᓄᓇᒥᒃ

ᐅᓪᓗᒥᐅᓕᖅᑐᖅ, ᓴᓪᓕᖅᒥᐅᑦ ᓂᓪ ᓂᖅᑭᖏᓂᒃᓂᒪᓕᕐᑦ ᐊᒥᖅᓂᒃᓗ, ᖃᓄᖅᓂᒃᓗ ᐅᒪᔪᓂᒃ, ᐃᓄᒃᔪᐱᖏᓴᖅᑭᓐᓄᑦ ᐊᑐᖅᑐᓂᒃ. ᐃᓄᒃᔪᐱᖏᓂᒃ ᐊᖏᔪᓇᓂᕆᑦᑕᖅᓂᖅ ᐅᒪᔪᖅᖃᕆᐊᕐᓂᖅᓂᒃ ᒪᓕᒃᐸᒃᑐᑦ ᐱᓇᓂᐊᖅᐸᖓᓂᖅᐸᓂ ᓴᓪᓕᐅᑦ ᖃᑭᖅᑕᓗᐊᓂ. ᑕᒪᖅᓇ ᐱᖅᓂᐅᒃᓴᕐᓂᖅ ᐱᓕᕆᐊᓚᐅᖅᑕᖓᓂᒃ ᒪᖅᓂᑕᖅᑏᑕᖅᓂᕐᒥᒃ, ᑲᖏᖅᓂᐅᖅᑏᑕᖅᓂᕐᒥᒃ, ᐊᖏᔪᓇᓂᐊᖅᑏᑕᖅᓂᕐᒥᑦᓗ ᓄᓇᒥᐅᑕᓂᒃ ᐊᓚᒪᓗ ᑕᓚᐅᖅᒥᐅᑕᓂᒃ ᐅᒪᔪᓂᒃ, ᓂᑦᓗ ᓇᓄᖅᓂᒃ, ᑐᒃᑐᓂᒃ, ᓇᑎᖅᓂᒃ, ᖃᖅᓇᓗᑎᖅᓂᒃ ᐊᓚᒪᓗ ᐊᐃᕐᖅᓂᒃ (ᑕᔪᓴᓚᖅ 7). ᓴᓪᓕᖅᒥᐅᓕ ᐊᐅᓪᓚᕋᐃᓂᖅᓴᐅᑉᑐᑦ ᑕᐅᑉᖓ ᐅᖏᓴᓂᑐᒍᑦ ᖃᓄᖅᓴᐅᖅᑐᖓᒥᒃ ᑎᖅᒥᐊᖅᑏᖅᓴᓚᓄᑦ, ᑲᖏᖅᓂᐅᖅᓂᒍᑦ ᒪᓂᓕᕆᐊᖅᓂᒍᓕᓂᓪᓚᑉᓂ ᐱᓕᕐᑕᐅᓂᒍᑦ ᑕᒪᑲᔪᐊ ᖃᑲᓂᓂᓄᓴᓕᓕᓂᖅᑐᑦ ᓄᓇᖅᑐᓄᑦ (ᑕᔪᓴᓴᐃᑦ 4 ᐊᒻᒪᓗ 5).

ᑭᓴᐊᓂᓕᑕᐅᖅ, ᓴᓪᓕᖅᒥᐅᑦ ᓂᓪ ᐊᑐᖃᑦᑕᖅᐳᑦ ᑎᖅᒥᐊᖅᑏᖅᓴᖅᒥᒃ ᓄᓇᒋᑕᖅᑐᖃᐅᓂᒍᑦ, ᐃᓕᓂᑲᔪᑦᓗ ᐊᖏᔪᒪᐊᓂᒍᑦ. ᑕᖅᐸᒪᖃᓴᖅᓂᓄᓴᖅᑐᖅᓴᐅᑉᖓ ᐅᖅᑎᓂᖅᓄᖅᑐᖅ ᑲᓇᖅᓇᖅ ᑎᖅᒥᐊᖅᑏᖅᓴᖅᓯᐅᑦ, ᓂᓴᓱᑲᔪᑦ, ᓂᓪᓕᐊᓂᖅᓗᓪᓚᕆᐊᖃᓂᖅᓂᒃ ᐃᓄᒃᔪᐱᖏᓴᖅᑭᓐᓄᑦ ᐅᖃᓕᒫᓂᖅᓂᒃ ᐃᓄᒃᔪᐱᖏᓴᖅᑭᓐᓄᑦ (ᐃᓄᒃᔪᐱᑎᒃᓴᕐᓂᖅᓂᒃ ᐊᐅᓪᓚᕋᐃᓂᖅᓂᒃᓗ ᐊᐅᓪᓚᕋᐃᓂᖅᓂᒃ) ᐊᓂᖅᑎᓂᓗ ᑕᓚᐅᖅᑎᐅᑕᓂᒃ ᐅᒪᔪᖅᓂᐅᖅᑐᖅ.


ᐅᐱᖃᓐᓇᖅᑐᖅ
(ᐊᐅᓚᓐᓂᐊᓕᒫᑦᑐᖅᓯᓪᓗᔪ)
ᐊᐅᓴᖅᑎ
(ᑐᕐᐃᐃᓕᓂᖅᓭ)
ᐅᑭᐱᖅᑕᐅᖅᑎ
(ᓄᓇᕗᑦᓯᐊᓕᓂᖅᓭ)
ᐅᑭᐱᖅᑕᐅᖅᑐᑦ
(ᓄᓇᕗᑦᓯᐊᓕᓂᖅᓭ)
• ᑲᔪᓯᖅᓯᐅᓂᖅ ᐱᑭᐅᑕᓚᐊᓂᖅ (ᐅᐱᓇᖅᑐᒃᓯᒪᓂᖅ)
• ᓇᑦᑎᖅᓯᐅᓂᖅ ᐊᐃᕐᓂᕐᓂᐊᓂᑦᓗ
• ᓇᓐᓂᐊᖅᑕᑦᓂᖅ
• ᐃᖃᓗᒃᑖᐊᖅᓯᒪᓂᖅ ᑕᓯᑦᓂᖅ ᐊᓚᒪᓗ ᑯᓘᓂᖅ
• ᑎᓚᒋᓂᐊᑦᓂᖅ ᒥᑭᑎᑦᓂᐊᑦᓂᖅ
• ᓄᓇᕗᒻᒥᐅᑦᓂᖅ, ᑐᑐᓯᐅᑦᓂᑐᒃ, ᑲᔪᓯᖅᓯᐅᑦᓂᑐᒃᑲ ᐱᐅᓯᖅᑲᓱᒪᔨᖅᑐᑦ ᓴᖅᑭᑐᖅᑐᒥ ᐃᓚᖓᔪᒪᓂᖅ ᓴᓇᓕᖅᓯᓯᐅᑦᓂᖅ
• ᑕᓚᐅᑦᑎᐅᑕᓂᖅ ᐄᒪᔨᖅᓯᐅᓂᖅ ᐅᒥᐊᓂᖅ
• ᐸᐅᑦᓴᓕᖅᑕᑦᓂᖅ
• ᐅᑭᐱᖅᑕᐅᖅᑐᑦ ᒪᑐᐃᓐᓇᖅᑐᑦ, ᑐᑐᓯᐅᑦᓂᑐᒃᑲ ᓇᑦᑎᖅᑕᑦᓂᖅ
• ᑎᓚᒋᓂᐊᓂᖅ ᒥᑭᑎᑦᓂᐊᑦᓂᖅ
• ᑐᑐᓯᐅᑦᓂᑐᒃ, ᑐᑐᓯᐅᑦᓂᑐᒃ, ᓇᑦᑎᖅᑕ

4 ᑐᖏᓚᓴᐅᔪᓂ ᐊᓚᒪᓗ ᐱᓴᖏᑕᐅᔪᒪᔪᓂ

4.1 ᑕᐅᑐᓚᓴᐅᔪᓂ

ᑕᐅᑐᓴᖅᓱᖅᓯᓂ ᑎᖃᒥᐊᖅᑎᖓᒥ ᐃᒪᐃᓐᓇᒋᓚᔪᓂᖅ:

> ᖃᖅᖢᓴᐅᖅᑐᖏᒥ ᑎᖃᒥᐊᖅᑎᖓᖅ ᐊᐅᓪᓚᑕᐅᐉᖅ ᐊᑯᓂ ᓯᐉᓂᖅᓴᒪᒧᑦ ᐱᑕᖅᑲᐃᓐᓇᖏᓂᐊᕐᒪᑕ ᐊᓚᒪᓗ ᓴᓱᐃᓐᓇᖅᑕᐅᓯᒪᓂᐊᕐᒪᑕ ᓄᓇᖅᑎᑦ, ᐃᒪᐃᓕᓗ, ᐅᒪᔨᓕᒪᓄᑦ, ᐃᓄᖓᓄᓗ ᐱᔨᓇᐅᐅᑎᖅᓂᖓᓂᐊᓄᑦ ᐊᓚᒪᓗ ᖃᑭᒪᐃᐅᓯᒪᓂᖓᓄᑦ, ᐊᑐᖅᑕᑦᓗᑎᒃ ᐃᓄᐃᑦ ᖃᐅᐅᓚᓴᑦᖢᖅᓂᓂᒃ ᑕᒪᐃᓂ ᐃᓯᒪᒃᓴᑕᐅᑎᓂᖓᓂᐊᕐᒪᑕ ᖃᑭᒪᐃᐅᓯᒪᓂᖓᓂᐊᓄᑦ.

ᑕᒪᒃᑯᐊ ᐊᑯᓂ ᓯᐉᓂᖅᓴᒪᒧᑦ ᑕᐅᑐᓚᓴᐅᔪᓂ ᐊᐅᓪᓚᑕᑎᓂᔭᒧᑦ ᖃᖅᖢᓴᐅᖅᑐᖏᒥᐅᑦ ᑎᖃᒥᐊᖅᑎᖓᒪᑦ ᐱᖓᓱᓇᐅᓪᓚᖅᑐᖓᓂᐊᕐᒪᑦ ᐃᓕᓴᕐᓂᐊᓕᒫᖅ ᐃᓇᓯᓇᐅᑉᐸᓇᐊᖅᑐᖓᓂᐊᕐᒪᑦ ᐊᑐᖅᑕᑦᓗᑎᒃᓴᖏᓂᐊᕐᒪᑦ ᐊᐅᓚᑦᓯᐊᖅᑕᖓᓂᐊᕐᒪᑦ ᐱᓕᕆᔨᒃᑯᑦ:

- ᓴᓪᓂᐊᕐᑕᐅᓗᑎᒃ, ᐅᒪᔨᐃᑦ ᐅᒪᔨᓴᖅᑎᖓᐅᔪᓐᓇᖅᓂᒋᓗ ᓄᓇᐃᑦ, ᓴᓪᓂᐊᕐᑕᐅᓗᑎᒍᓗ ᑎᖃᒥᐊᑦ ᐊᒪᓗ ᐅᒪᔨᐃᑦ ᐊᒥᓱᖏᓪᓗᐊᓕᖅᑐᑦ;
- ᓴᓪᓂᐊᕐᑕᐅᓗᑎᒃ ᐃᓄᖅᑎᖓᐅᖅᓯᒪᓕᐅᖅᑐᑐᖅᑎᑦ ᓄᓇᖅᑐᑦ ᐊᒪᓗ ᐃᓕᓴᕆᔨᖅᑎᖓᔪᑦ ᐊᑐᖅᑕᐅᖅᓯᒪᓂᖅᑎᑦ (ᐊᑐᖅᓗᑎᒃ ᑎᑎᕋᖅᓱᒪᓂᖏᓐᓂ 2-ᒥ – ᐃᓄᐃᑦ ᐃᑲᔪᓱᐊᕈᓴᖏᓐᓂᓄ ᐊᖏᑭᑐᐃᑦ);
- ᐃᓄᐃᑦ ᐱᔨᓇᐅᑎᖏᓐᓂᒃ ᐊᑐᑭᓇᓗᑎᒃ ᐊᒪᓗ ᐊᖓᔪᓇᓱᖅᑕᑭᓇᓗᑎᒃ, ᐊᒪᓗ ᐅᓴᖅᑐᓗᑎᒃ ᐃᓄᐃᑦ ᐃᑲᔪᖅᑕᐅᖅᖾᓕᕐᖅᑕᕋᐊᖅᑎᖓᓂᐊᕐᒪᒃ (ᐊᑐᖅᓗᑎᒃ ᑎᑎᕋᖅᓱᒪᓂᖏᓐᓂ 2-ᒥ – ᐃᓄᐃᑦ ᐃᑲᔪᓱᐊᕈᓴᖏᓐᓂᓄ ᐊᖏᑭᑐᐃᑦ);
- ᐃᓕᓴᕐᕋᓴᐅᑎᒃ ᐃᓄᐃᑦ ᖃᐅᔨᓴᖅᑐᖏᓐᓂ ᐱᓕᕆᔨᐅᓂᖅᓴᖏᓐᓂᓄ ᐱᑕᖅᑲᐃᓐᓇᑯᓴᓂᖅᑐᑦ, ᐊᐅᓚᑦᓯᐊᖅᑕᖓᓂᐊᕐᒪᑦ, ᖃᐅᔨᓴᖅᑎᖓᔪᓐᓇᖅᓂᒋᓗ ᐃᓇᓯᓕᐅᖅᑕᐅᓂᖅᑐᓂᐊᕐᒪᑦ ᐃᓕᓴᕐᓂᐊᓕᒫᖅ;
- ᖃᐅᔨᓴᖅᑐᖏᓐᓂᖅᓴᓕᐅᑎᓗᑎᒃ ᐱᓕᕆᔨᐅᓂᖅᓴᖏᓐᓂ ᑎᖃᒥᐊᖅᑎᖓᔪᓄᑦ ᐱᑕᖅᑲᐃᓐᓇᑯᓴᓂᖅᑐᑦ, ᐊᐅᓚᑦᓯᐊᖅᑕᖓᓂᐊᕐᒪᑦ, ᖃᐅᔨᓴᖅᑎᖓᔪᓐᓇᖅᓂᒋᓗ ᐃᓇᓯᓕᐅᖅᑕᐅᓂᖅᑐᓂᐊᕐᒪᑦ ᐃᓕᓴᕐᓂᐊᓕᒫᖅ;
- ᖃᐅᔨᓴᖅᑐᖏᓐᓂᖅᓴᓕᐅᑎᓗᑎᒃ ᐱᓕᕆᔨᐅᓂᖅᓴᖏᓐᓂ ᑎᖃᒥᐊᖅᑎᖓᓂᒃ.

4.2 ᑐᖏᓚᓴᐅᔪᓂ ᐊᓚᒪᓗ ᐱᓴᖏᑕᐅᔪᒪᔪᓂ

ᐊᐅᓚᑦᓯᐊᖅᑐᑦ ᑐᖏᓚᓴᐅᔪᓂ ᖃᖅᖢᓴᐅᖅᑐᖏᒥᐅᖅᑐᖓᓂᒃ ᐱᑕᖅᑲᐃᓐᓇᖅᑎᖓᓚᖅᑐᖓᓂᒃ ᐊᖏᑭᑐᐃᑦ ᐊᐅᓚᑦᓯᐊᖅᑕᖓᓂᐊᕐᒪᒃ ᐊᓚᒪᓗ ᐊᑦᑭᐅᒪᑦᓯᐊᖅᑎᖓᓂᐊᕐᒪᒃ ᑕᐅᑐᓚᓴᐅᔪᓂᐅᖅᑐᓂᐊᕐᒪᒃ (ᐊᐃᒃᑐᑦᓯᐊᓂ ᑕᑯᑎᑦᓯᐊᓂ 6). ᐃᑦᓂᐅᑦᓴᒃᑯᓂ ᐊᐅᓚᑦᓯᐊᐅᑦᖢᖅᑕᐅᓂᖅᑐᑦ ᑲᑎᒪᔨᖅᑕᖅᑐᑦ ᑕᐅᑐᓚᓴᐅᔪᓂ ᑕᐅᑐᓚᓴᐅᔪᓂ ᐊᐅᓚᑦᓯᐊᖅᑐᑦ ᑐᑭᓯᓚᐅᖅᑐᑦ

ᐅᓗᓕᐊᓇᖅᑕᐅᔪᓐᓇᖅᑐᓂᒃ ᖃᐅᖅᓴᐅᖅᑐᒥᒃ ᑎᖅᑕᐊᖅᖃᕐᓱᓄᑦ (ᐃᓗᓕᖅᑎᑦ ᐊᕐᓴᐅᑐᖅᓯᒪᓂᖅᑎᓂᒃ 5.0 – ᐊᐅᓚᑦᑎᐅᐅᓂᖅᑎᓂᒃ ᐃᓱᒪᒃᓴᖅᓯᐅᑕᑎᒃᓴᖅᑎᑦ).

ᑐᖏᑎᓕᐊᓴᐅᔪᑦ ᐋᖏᑭᐅᒪᕐᒪᑦ ᒪᓕᒃᑐᓂᒃ ᐱᓕᓴᐅᓂᖅᑎᓂᒃ ᐊᐅᓚᑦᑎᐃᓕᑦ ᑲᑎᒪᔨᖏᑦᑕᐅᓂᖅ.

ᐊᕐᓴᐅᑐᖅᓯᒪᔪᓂᒃ ᑕᒪᑎᑦᑎᐅᔪᑦ 6: ᐊᐅᓚᑦᑎᓂᕐᒧᑦ ᑐᖏᑎᓴᓴᐅᔪᑦ ᐊᓕᒪᓗ ᐱᓴᓱᖅᑕᐅᔪᒪᔪᑦ

ᑐᖏᑎᓴᓴᐅᔪᑦ	ᐱᓴᓱᖅᑕᐅᔪᒪᔪᑦ
<u>ᑐᖏᑎᓴᓴᖅ 1:</u> ᓴᓇᓂᐊᖅᑕᐅᔭᕆᐊᑕᑦ ᓄᓇᐃᓪᓗ ᐃᒪᐃᓪᓗ, ᑕᒪᕐᒥᒃ ᐅᒪᔨᐃᓕᑦ ᐱᓪᔨᑕᐅᓂᖅᑐᓄᓐᓂ, ᐊᓕᒪᓗ ᐃᓄᓐᓄᐅᑎᖅᓂᖅᑕᑦ ᐊᓕᒪᓗ ᓄᓇᖅᐃᓇᓂᖅᑕᑦ ᐃᓄᐃᓕᑦ	ᐱᓴᓱᖅᑕᐅᔪᒪᔪᑦ ᐱᓴᓱᖅᑕᐅᔪᓐᓇᖅ 1.1: ᐱᓕᕆᓗᑎᒃ ᐊᔪᕐᓴᔨᒪᓂᖅ ᑎᖅᑕᐊᖅᖃᐃᐅᓂᖅᓯᓂᒃ ᑲᓇᑕᒥᓪᒪᑕ ᐅᓴᐃᑦᑐᐃᓂᖅ ᐱᓴᓇᓂᐅᑎᓂᒃ ᐊᐅᓚᑦᑎᐊᕐᓇᖅᓂᖅᑕᑦ, ᐊᓕᒪᓗ ᐊᖏᕿᔪᓕᒪᖅ ᓴᓇᑎᐅᓴᖅᓯᓇᖅᑕᑦ ᑕᒪᕐᒥᒃ ᐅᒪᔨᓕᒪᓂ, ᐱᓗᐊᕐᓗᑎᒃ ᐅᒪᔨᖏᑦᑕᐊᓕᖅᑕᑦ. ᐱᓴᓱᖅᑕᐅᔪᒪᔪᖅ 1.2: ᐱᔨᑎᓴᐃᓂᖅᓱᑦ ᐃᑲᔪᖅᔪᐃᖅᑕᖏᓂᖅᓱᓪᓗᑎᑦᓗ ᖃᐅᔨᓴᖅᑕᐅᓇᖅᑐᓂᒃ ᐅᖃᖅᖢᓂᒃ ᓇᐅᑎᖅᑐᖅᑕᐅᖅᑲᓪᓚᐊᓕᓴᐃᓂᒃ, ᐃᓕᖅᖢᓂᑐᐃᓪᓗ ᑎᑎᖅᑕᐅᖅᖢᓂᒃ ᐃᓄᐃᓕᑦ ᖃᐅᔨᓴᓕᕐᓂᖅᓱᒃ, ᖃᐅᔨᓴᓕᕐᔨᒪᑎᒃ ᐊᐅᓚᑦᑎᐊᓇᖅᓂᖅ ᑎᖅᑕᐊᖅᖃᐃᐅᓂᖅᓯᓂᒃ ᖃᐅᔨᓴᓕᖅᑐᖅᓯᓂᖅ ᖃᐅᔨᓴᓕᖅᓯᓂᖅ. ᐱᓴᓱᖅᑕᐅᔪᒪᔪᖅ 1.3: ᐱᓕᕆᖅᑎᑎᒋᓗᐊᖅᑐᓂᒃ ᐃᑲᔪᖅᑎᐅᖅᑎᑦᑎᓗᐊᖅᑐᒃ ᐃᓕᓴᐅᒡᒎᖅᑐᓂᒃ ᑎᑎᖅᔪᓂᑎᑦᑎᐅᔪᑦ (ᐃᓄᐃᓕᑦ ᐱᐅᔪᑐᖅᑎᖅᓯᓂᒃ ᐸᐸᑎᐃᓕᑦ/ᓄᓇᖅᑐᑦ ᓴᖅᓚᐃᓕᑦ/ᑐᐊᖏᓴᐃᓚᒃᑯᑦ) ᑎᑎᖅᑕᐅᖅᑐᖅᓯᓂᖅᓱᓂᒃ ᐃᓕᓴᒃᑯᓪᔨᑐᖅᓯᓂᒃ ᖃᐅᔨᓴᖅᑕᐅᖅᑲᓪᓚᐃᓕᑦ ᑎᖅᑕᐊᖅᖃᐃᐅᓂᖅᓯᓂᒃ ᐱᓴᓱᖅᑕᐅᔪᒪᔪᖅ 1.4: ᖃᐅᔨᓴᓕᐅᑎᐊᖅᓯᖏᓐᓂᐊᓕᖅᑕᐅᓂᖅᓱᑦ ᑎᖅᑕᐊᖅᖃᐃᐅᓂᖅᓯᓂᒃ ᐱᓕᕆᓗᑎᒃᓴᖅᓯᓂᒃ ᐊᐅᓚᑦᑎᐅᐅᓂᖅᓯᓂᒃ ᐱᐅᔪᑐᖅᑎᖅᓯᓂᒃ ᐊᓕᒪᓗ ᐃᓕᓴᒃᑯᓪᔨᑐᖅᓯᓂᒃ ᐊᐅᓚᑦᑎᐊᓇᖅᓂᖅᓯᓂᒃ ᐊᐅᓚᑦᑎᐅᔪᒪᔪᑦ ᐃᓄᐃᓕᑦ ᐱᓴᓱᖅᑕᐅᔪᒪᔪᑦ ᐊᐅᓚᑦᑎᐅ

ᑐᖏᓕᓴᐅᔪᓂ	ᐱᓴᓱᖅᑕᐅᔪᓚᔪᓂ
<u>ᑐᖏᓕᓴᐅᖅ 2:</u> ᐅᓴᕆᖅᑐᑎᐊᖏᓂᖓᓂ ᓄᓇᓕᓐᓂ ᖃᐅᐅᓚᒋᑕᐅᓴᓇᖏᖅᓂᖅᑎᑲᓂᖅ,ᐃᓕᐅᑎᑕᐅᓴᓇᖏᖅᓂᖅᑎᑲᓂᖅ ᐊᒻᒪᓗ ᐃᑲᔪᓯᐊᖏᖅᑕᖏᑕᓇᐊᖏᖅᓂᖅᑎᑲᓂᖅ	ᐱᓴᓱᖅᑕᐅᔪᒃᖏᖅ 2.1: ᓴᖅᑭᑎᑕᓗᑎᒃ ᑐᓴᐅᒪᖏᖅᑕᐅᑎᓂᖓᓂ <ᖏᓇᐅᑎᓂᒃ ᓴᓪᓕᒥᐅᑕᓂᒃ, ᑐᓴᐅᒪᓴᐅᔪᒃᖏᓂᒃ, ᑐᑭᓯᓴᐅᖏᓴᐅᔪᓂᑦᓗ ᐊᐃᑐᐃᔪᓐᓇᖏᓂᖃᖅᐳᑎᓂᐊᖏᖅᒪᑕ ᑕᒪᒃᑯᓂᖓᓗ ᐃᓄᖏᑎᖓᓐᓄᑦᓄᓂ, ᐅᖏᐅᓯᐅᖏᐅᑕᑭᓇᖏᓗᑎᒃ ᐃᓯᒪᓗᑕᐅᔪᑦ, ᑐᓴᖏᐊᕐᖏᖏᑕᖏᓗᑎᓗ ᖃᐅᐅᓴᖅᑕᐅᔪᒪᔪᓂ ᓯᑲᓪᓕᐅᖏᓴᐅᔪᓂᖓᓂᒃ. ᐱᓴᓱᖅᑕᐅᔪᒃᖏᖅ 2.2: ᐱᔨᒋᔭᐃᐊᓐᖓᓂᖅ ᓴᖅᑭᑎᑕᓂᖓ ᐱᑕᓕᐊᒃᖏᖅᑲᒡᖏᖅᑕᐅᔪᑦᓇᖏᓂᖓᓂᒃ ᐊᒻᒪᓗ ᐃᓯᒪᓂᖅᑐᑦᖏᖅᑕᐅᑕᖏᓇᖏᓂᖓᓂᒃ ᐱᔭᔨᑎᖏᖅᑲᖅᑐᓂ ᖃᐅᐅᓴᖅᑕᐅᔪᓴᖏᓂᖓᓂᒃ, ᓇᐅᑎᖅᑲᑐᑎᓂᖓ ᐊᒻᒪᓗ ᖃᐅᐅᓴᖅᑕᐅᔪᒃᖏᖅᑕᖏᓗᑎᓗ ᐱᑕᓕᐊᒃᖏᖅᑕᐅᔪᑦᓇᖏᓂᖓᓂᒃ. ᐱᓴᓱᖅᑕᐅᔪᒃᖏᖅ 2.3: ᓄᓇᑕᖓ ᐃᓄᐃᑦ ᖃᐅᐅᓴᒪᓂᖅᑲᐅᖏᖅᑕᖏᓗᑎᒃᑕᐅᔪᓗᑎᒃ ᐱᑕᓕᐊᒃᖏᖅᑕᐅᔪᓂᖓᓂᒃ ᑎᖓᒥᐊᖏᖅᖃᐊᖓᒥ.
<u>ᑐᖏᓕᓴᐅᖅ 3:</u> ᐃᓄᐃᑦ ᖃᐅᓂᐃᑲᓇᖅ ᐱᑕᓕᖏᓂᖓᓂᒃᑯᓂᖓᓂ ᐊᒃᑐᖏᑕᐅᔪᓚᑲᓱᐅᑎᒃᑲᐃᑦ ᐱᐅᓯᑐᖏᐅᔪᓂ ᐊᒻᒪᓗ ᐃᑕᖏᓂᖅᑲᖅᓯᐅᓂᖓᓂᒃ ᓄᓇᓕᖏᓂᖓᓂ	ᐱᓴᓱᖅᑕᐅᔪᒃᖏᖅ 3.1: ᐅᓴᕆᖅᑐᓴᓗᑎᒃ ᑕᒪᓴᓂᒃ ᐃᓄᐃᑦ ᒪᑕᓴᑎᐊᖏᖅᑕᓚᐊᖏᖅᓴᓂᖓᑦ ᐱᔨᓇᐅᐅᑎᖏᖅᑲᐊᖏᖅᓴᓂᖓᖅᑎᒃ ᐊᖏᕐᕕᐃᒪᓕᐅᑕᐅᔪᓂᖓᓂᒃ ᒪᑕᑐᓴᓂᒃ. ᐱᓴᓱᖅᑕᐅᔪᒃᖏᖅ 3.2: ᐊᑐᑕᖅᑎᓴᑎᓐᓕᓪᓕᐊᓂᖅᑐᐱᐅᓂᖅᓴᓂᒃ ᐱᐅᓂᖅᓯᓂᒃ ᐱᐅᓯᓚᐅᔨᓇᖅᑐᓂᒃ ᐊᒃᑐᖏᑕᐅᔪᓚᑲᓱᐅᑎᒃᑲᐃᑦ, ᓯᑎᖅᑎᖅᑕᐅᔪᑲᓯᐅᑕᐅᔪᓂᑲᓇᓂᓂ ᐱᐅᓯᑐᖏᐅᔪᓂ, ᐃᓕᑕᖅᑯᓯᐅᑎᓂᒃᑲᓇᓂᓂ ᐱᒻᒪᓚᐅᓂᖓᓂᒃ.	ᐱᓴᓱᖅᑕᐅᔪᒃᖏᖅ 3.3: ᐃᓕᐅᑦᖃᑕᖏᓗᑎᒃ ᐊᐃᓴᐅᑐᖏᓯᓚᐅᖓᓂᖅ ᐱᑕᓕᐊᖓᔨᓕᑕᐅᔪᓂᒃ, ᐱᑎᐊᖅᑎᑕᐅᔪᓯᐅᑲᓇᓂᓂ ᐃᓄᐃᖅᓱᑕᐅᔪᓂᖓᓂᒃ ᑎᖓᒥᐊᖏᖅᖃᐊᖓᒥᐅᑦ ᓴᓴᓂᐊᔪᓂᖅ ᐊᒻᒪᓗ ᐊᓐᓗᑲᐊᓂᑲᓐ ᓄᓇᕆᐊᑎᓂᖅᓂᒃ.

4.3 ᖃᐅᔨᓴᐃᖅᑕᕐᓂᖅ

ᑲᓇᑕᒥ ᐅᒪᔪᓕᕆᔨᒃᑯᑦ ᐊᖏᕐᓯᒍᒪᔪᓐᓇᖅ ᓇᐅᑦᑎᖅᑐᕆᐊᖅᑐᐃᖅᑕᑦᑎᓐᓄᑦ ᒪᓕᒃᓴᑎᒃ ᐊᑐᐃᓐᓇᐅᔨᓐᓇᕐᓂᖅᓂᒃ ᑭᓇᐅᔭᐃᑦ ᐊᐃᒪᓗ ᐱᓕᕆᔨᖃᖅᑐᓇᕐᓂᖅᓂᑦ. ᐃᖏᓂᐅᖏᖅᓴᒃᑯᑦ ᐊᐅᓪᓚᕆᑎᔨᐅᖃᖅᑕᐅᓂᕐᒧᑦ ᑲᑎᒪᔨᕿᓂᑦ ᖃᑎᖏᕐᐊᑭᒍᑦ>ᑦ ᐊᐅᓪᓚᕆᑎᓂᕐᒧᑦ <ᖏᓇᐅᑎᓂᒃ ᐊᖓᕆᖅᑕᐅᓪᓚᐅᖏᓐᓄᑎᒃ, ᑕᓪᒪᐃᑦ 5 ᐊᖏᕌᐃᑦ ᑭᖑᓂᐊᒍᑦ ᐊᓂᔪᖅᑐᓂᒃ ᑭᖑᓂᑲᓂᐊᒍᓪᓗ ᖃᑎᖏᕐᑎᑐᓂᒃ, ᓚᒋᑭᓇᐊᖅᑕᐅᖏᑲᑕᕐᓯᒪᓂᖓᓂᒃ, ᓚᒋᑭᓇᐊᖅᑕᐅᖏᑲᑕᖏᓐᓄᑐᓗ ᖁᔨᓂ 10 ᐊᖏᕌᐃᑦ ᓴᓴᖅᓯᒪᑕ. ᖃᐅᔨᓴᐃᖅᑕᐅᖏᑲᑕᓂᐊᖅᑐᑦ ᑎᑎᕋᖅᓯᒪᓂᕐᓂᑦ ᖃᑎᖏᕐᐊᖅᑕᐅᓗᑎᒃ ᐊᐅᓪᓚᕆᑎᓂᕐᓂᓂᒃ ᐱᓪᓚᐊᖅᑕᐅᓗᑎᒃ ᖃᐅᔨᓴᓴᐅᓯᒪᔪᓂᕐᓂᖅᓂᑦ ᐊᐃᒪᓗ ᖃᐅᔨᓴᐅᓴᐃᖅᑕᐅᓗᑎᓂᕐᓂᖅᓂᑦ. ᑲᓇᑕᒥ ᐅᒪᔪᓕᕆᔨᒃᑯᑦ ᐊᐃᒪᓗ ᐊᐅᓪᓚᕆᑎᔨᐅᖃᖅᑕᐅᓂᕐᒧᑦ ᑲᑎᒪᔨᕿᓂᑦ ᑕᐃᑲᓐᓂ ᓴᖅᑭᑎᓚᓂᐊᓚᖅᑕᐅᓗᑎᒃ ᐊᐃᑐᓴᓚᓂᐊᓚᖅᑕᐅᓗᑎᓂᕐᓂᖅᓂᑦ ᑕᒪᒃᑯᐃᓇᓂᐊᓚᖅᑕᐅᓗᑎᓂᕐᓂᖅᓂᑦ ᐊᐃᒪᓗ ᓴᓂᖅᖃᐃᓚᓂᐊᒃ ᐊᐃᑐᐅᓴᓚᓂᐊᓚᖅᑕᐅᓗᑎᓂᕐᓂᖅᓂᑦ ᑕᒪᒃᑯᐃᓇᓂᐊᓚᖅᑕᐅᓗᑎᓂᕐᓂᖅᓂᑦ ᐱᓕᕆᔨᓂᕐᒧᑦ.

5 ᐊᐅᓪᓚᕐᓂᐅᕐᒥᐅ ᐃᓱᒪᓕᕆᔨᓱᐅᑭᓚᕆᐊᑦ

ᐃᓐᓇᐅᔾᔨᖅ, ᐱᓕᒪᓇᐅᔾᔨᓂᐅ ᑐᐊᕕᕆᔨᓇᓚᖅᑐᑐᓂᓗᓐᓄᑦ ᐅᑐᓪᓚᐊᓇᖅᑐᑎᕆᖅᑭᕐᓚᖅ ᕆᐅᖅᓴᐅᖅᑐᑦ ᑎᓐᓕᐊᕆᐅᕆᕕᓐᓄᑦ. ᓄᓇᕆᐞᑎᖅᑭᑦ ᑕᐅᔨᓂ ᐊᐅᑐᖅᑕᐅᓂᕆᓕᕐᓕᒪᑕ, ᑎᓐᓕᐊᕆᐅᕆᕕᐅᓂᖅᓱᓗᓐ ᐃᓄᕆᐅᕆᔨᐃᑦᑐᑦᓗᓂᓂ. ᑭᓱᐊᓂ ᓂᓪ ᒪᑯᐊ ᐊᐅᑕᓪᓚᐅᕐᒥᑦ ᐃᓱᒪᓕᕆᔨᓱᐅᑭᓚᕆᐊᐅᑉᐳᑦ, ᐊᐅᑐᖅᓴᔨᓐᑕᕐᓂᑦ ᑎᓐᓕᐊᕆᐅᕆᕕᖅᑐᑎᒃ.

ᑲᖓᐃᑦ ᐊᒥᓱᓂᕐᑭᑦ ᓄᓇᐃᑦᓗ ᓀᔨᖅᑮᑕᕐᑭᑦ

ᑲᓇᑕᒥ ᐅᒪᔨᑦᓚᓕᕆᑯᑦ ᑎᓐᓕᐊᕆᐅᕆᕕᑕᐅᑕᖅᓴᒪᔨᓛᑦ ᑕᒪᓂ 1959-ᒥ, ᐊᒥᓱᓗᐊᖅᑮᑕᐅᕐᓄᑦᑎᑦᓗᑦᓓᑦ ᑕᐃᔪᓱᒪᓂ ᑎᓐᓕᐊᕆᐅᕆᕕᐅᕙᑕᐅᕐᖅᑕᑦ ᓄᓇᕐᑭᑦ, ᑲᓇᑕᐅᑦ ᐅᑭᐅᖅᑕᕐᖅᑐᐃᓂᕐᓂᒃ, ᑕᐃᔪᓱᒪᓂᓗ ᐱᑕᖅᑲᒪᐅᖅᓴᑎᑎᑦ ᑲᓇᑕᖅᑐᑎᒃ ᑲᑎᕐᓂᖅᑐᑦ 24,300 (Cooch & Barry, 1957) ᑎᓐᓕᐊᕆᐅᕆᕕᖅ ᕆᐅᖅᑭᕐᖅᑐᑦᓕᒃ ᐅᓐᒪᑕᓄᓐᒪᓚᑦ ᑭᓪᓕᕆᐅᕆᕕᐅᓂᓐᒪᑕ ᑎᓐᓕᐊᕆᐅᕆᕕᐅᖅ, ᐊᑕᐅᓱᐊᓗᓕᓕᖅᓴᓂ ᑲᔪᓱᐊᓗᓐᓂᖅ ᐅᑦᓗᕆᐅᕆᕕᐅᓕᓕᕆᓂᐅᓐᒪᑕ ᕆᐅᓕᑎᓐᓂᒃ (ᑕᐃᓲᖅᓚᐃᑦ 4 ᐊᒪᓚᓗ 5). ᐃᓐᓇᓴᓐᓂᖅᓯᒋᑎ ᕆᐅᐅᓕᐅᓚᖅᑕᐅᓚᑎᑲᖅ ᐊᒥᓱᓂᕐᓂᑦ ᐊᖅᓚᐃᑦ 2014-ᒥ, ᓇᓕᐅᑦᓴᖅᑕᐅᓕᐅᖅᑐᑦ ᐅᓐᒪᑕᓂᕐᑐᓂᓗᓂᑲᓂᑦ 289,700 ᑲᓐᒪᐃᑦ ᑦᑲᐃᓇᓂ ᒪᕐᕉᓂ ᑎᓐᓕᐊᕆᐅᕆᕕᐅᓕᖅᑐᖅᓂ (ᐅ. ᓂᐢᓗᐊᖅ-J. J. Leafloor, ᑲᓇᑕᒥ ᐅᒪᔨᓕᓚᐅᓕᑦ, ᓇᓕᒥᓂᖅ ᑐᐊᕆᐅᑎᑎᐊᕐᓂᖅ). ᓄᓇᕐᔨᐊᓕᒻᒪᒥ ᐃᓱᒪᒋᓕᑐᒋᑦ, ᑕᒪᑲᐃᐊ ᑲᓐᒪᐃᓗᐃᑦ ᐊᒥᓱᕈᕐᔨᐊᕆᓐᓂᕐᒪᖅᑕᑦ 50-ᓂᑲ ᐊᓂᔪᖅᑐᓂ ᐊᕐᕌᔪᓂᑲ, ᐃᓐᓇᐅᓕᖅᑐᕐᓗ “ᐊᒥᓴᓱᐃᐊᐊᓕᖅᓴᓂᑲ” ᑲᓇᑕᒥᓗ ᐊᒥᐊᑲᓕᑲᒥᐅᓕᓗ ᓄᓇᖅᑎᓱᓴᔨᔪᓕᑦ. ᓄᓀᓗᐊᑕ ᐊᑯᓂᓕᖅᓴᖅᑎᖅᑐᑦ ᑲᓐᒪᐃᑦ, ᑕᐃᑲᑯᐊᓗ ᑲᓐᒪᐃᑦ ᐅᓗᖅᑲᓱᐃᑦ ᓴᑦᓕᖅᑕᓀᓗᐊᓂ, ᐊᒥᓱᓂᑲᖅᓂᑎᓕᖅᑐᑦ 12.6 ᒥᓕᔾᓂᑲ ᐱᕈᐊᓂᑲᓕᔨᓂᑲ ᑲᓐᒪᐃᑦ (ᐊᓕᓴᐅᓴᑲᓴ, ᐱᓕᓐᖅᓂᖅᑎᓕᓗ-Alisauskas, et al., 2018a), ᑕᐃᑲᑯᐊᓗ ᐊᓱᖅᑎᑦ ᓂᓪᓕᓐᓄᐃᑦ (Ross-ᑭᓱᑦ) ᐊᒥᓱᓂᑲᑎᓕᖅᑐᑦ ᑕᐃᓂ 1.6 ᒥᓕᔾᓂᑲ ᐱᕈᐊᓂᑲᓕᔨᓂᑲ ᑲᓐᒪᐃᑦ, ᐊᓕᖅᔪᒪᐅᖅᑐᐃᓂᑲ 2006-ᒥᑲ 2015-ᔪᑦ (ᐊᓕᓴᐅᓴᑲᓴ, ᐱᓕᓐᖅᓂᖅᑎᓕᓗ-Alisauskas, et al., 2018b).

ᐊᒥᓱᓗᐊᓕᓐᓂᕐᑎᑦ ᐃᓱᒪᓗᑕᐅᓕᖅᑐᑦ ᐊᑲᑐᖅᑕᐅᔨᓐᓇᓂᕐᒧᓐᓄᑦ ᐅᑭᐅᖅᑕᖅᑐᒥ ᓂᓴᖅᑐᓱᐅᓂᕐᑎᑦ, ᐊᓕᑎᑲᑐᓂᕐᑎᑦᓗᓐ ᓄᓇᐃᑦ ᐅᒪᔨᓕᖅᑎᐅᑭᕐᓂᖅᑐᑦ ᐊᒥᓱᓗᐊᖅᑐᑦᓗᒥ ᑲᓐᒪᐃᑦ (Batt, 1997).. ᓴᓪᓕᐅᔨ ᓴᑭᖅᑮᓗᐊᓂ ᖃᐅᔨᓴᖅᑕᐅᓕᐅᖅᓯᒪᓂᖅᓂᒃ ᐅᓂᒃᑲᓕᐊᑦ ᐅᖃᐅᔾᔨᖅᑐᒃ ᑲᓐᒪᐃᑦ ᓂᓴᖅᑲᒪᔨᖅᑕᐅᑲᓕᓕᖅᓯᒪᓂᖅᓂᒃ ᐊᓕᑎᑲᑐᓂᐅᓂᖅᑐᓂ ᓂᓴᖅᑲᒪᔨᖅᑕᐅᓂᖅᑐᓂ, ᐊᒥᓱᓗᐊᖅᑐᑦᓗᒥ ᓄᓇᕗᑦ ᑲᓐᒪᐃᑦ (ᑲᑐ, ᐱᓕᓐᖅᓂᖅᑎᓕᓗ-Carter, et al., 2018; Fᐊᓂᐢ ᐊᒪᓚᓗ ᐃᓗᓐᓗ-Fontaine and Mallory, 2011). ᓄᓇᐃᑦ ᐊᓂᔨᓴᖅᓯᒪᓕᖅᑐᑦ ᑕᐃᒃᑯᒪᑐᓐᓇ ᐊᔨᔨᓴᖅᓂᖅᑎᑦᓗᐅᑎᓕᖅᑐᒃ ᕿᐊᑦᓴᓂᐢᐳᑦ ᑕᓂᐅᔭᕐᔪᐊᖏᓐᓇᑦ ᓄᓇᐃᑦ ᓂᓴᖅᑲᓂᕐᒧᐊᓂᖅᑐᓂᖅᑐᐃᓂᒃ ᐊᑦᑎᓂᒃᑑᓂᖅᑐᓂᓂᓂᖅᑐᐃᓂᒃ ᓄᓇᕐᑭᑦ (ᑲᑐ, ᐱᓕᓐᖅᓂᖅᑎᓕᓗ-Jano, et al., 1998; ᑯᑕᓂᐊᓐ ᐊᒪᓚᓗ ᔨᐊᐅᓗᓴᔨ-Kotanen and Jefferies, 1997). ᐊᐅᑎᓕᓚᔨᔪᐊᑲᑯᑦ ᐃᓐᓇᐅᔾᔨᖅ ᖃᐅᔨᓴᖅᑲᓕᓕᐊᓕᓂᖅᑐᑦ ᑎᓐᓕᐊᕆᐅᕆᕕᐅᓕᓕᓂᖅᑐᑦ ᓂᕆᖅᑎᖅᑕᐅᓕᐅᓕᓕᓕᐊᓂᖅᑐᓂᖅᑐᐃᓂᒃ ᐊᒥᓴᓱᐃᐊᐊᓕᖅᑐᑦ ᑲᓐᒪᐃᑦ ᓂᓴᖅᑲᒪᔨᖅᑕᐅᓂᖅᑐᓂᖅᑐᑦ, ᓂᓚᕕᐅᓗᐊᓕᑦᓂᖅᑎᖅᑐᑦ ᑐᓂᕕᐅᓂᖅᑐᓂᓗ, ᑐᐱᓂᔨᐅᔨᓚᓪᓗᑎᖅ ᖃᐅᔨᓴᖅᑕᐅᓂᖅᑐᓂᖅᑐᐃᓂᒃ ᐊᒥᓱᓗᐊᖅᑐᑦᓗᒥ ᐃᓐᓇᐅᔾᔨᖅ ᖃᐅᔨᓴᖅᑕᐅᓂᖅᑐᓂᖅᑐᐃᓂᒃ ᐊᑦᑎᓂᒃᑑᓂᖅᑐᓂᓂᓂᖅᑐᐃᓂᒃ ᓄᓇᕐᑭᑦ, ᓂᓚᕕᐅᓗᐊᓕᑦᓂᖅᑎᖅᑐᑦ ᑐᓂᕕᐅᓂᖅᑐᓂᓗ, ᑐᐱᓂᔨᐅᔨᓚᓪᓗᑎᖅ ᖃᐅᔨᓴᖅᑕᐅᓂᖅᑐᓂᖅᑐᐃᓂᒃ ᐊᒥᓱᓗᐊᖅᑐᑦᓗᒥ ᓄᓇᕗᑦ ᑲᓐᒪᐃᑦ (ᑲᑐ, ᐱᓕᓐᖅᓂᖅᑎᓕᓗ-Jano, et al., 1998; ᑯᑕᓂᐊᓐ ᐊᒪᓚᓗ ᔨᐊᐅᓗᓴᔨ-Kotanen and Jefferies, 1997). ᐊᐅᑎᓕᓚᔨᔪᐊᑲᑯᑦ ᐃᓐᓇᐅᔾᔨᖅ ᖃᐅᔨᓴᖅᑲᓕᓕᐊᓕᓂᖅᑐᑦ ᑎᓐᓕᐊᕆᐅᕆᕕᐅᓕᓕᓂᖅᑐᑦ ᓂᓴᖅᑲᒪᔨᖅᑕᐅᓂᖅᑐᓂᖅᑐᐃᓂᒃ ᐊᒥᓱᓗᐊᖅᑐᑦᓗᒥ ᓄᓇᕗᑦ ᑲᓐᒪᐃᑦ ᓂᓴᖅᑲᒪᔨᖅᑕᐅᓂᖅᑐᓂᖅᑐ

ᐅᑦᓗᓕᐅᕈᓐᓇᐃᓕᓕᓴᓐᓂᑎᓄᑦᓗ ᓄᓇᒥ (Flemming, et al., 2016). ᐊᓯᖏᑦ ᐄᒪᓪᐃᑦᑕᐅᖅ ᐊᐳᑐᖅᑕᐅᓯᒪᓪᓴᓐᓂᑦ ᓯᓚ ᑐᑭᓯᓇᑦᑎᐊᖏᒪᒪᑕ.

ᒥᑎᖅᑕᕆᔨᓂᖏᓐᓄᑦ

ᒥᔅᑦ ᐃᓕᓴᕆᐅᔪᓚᓇᐅᑐᑦ ᐅᑭᐅᖅᑕᖅᑐᒥ ᐊᕐᑎᓄᑦ ᐊᕐᑎᑎᒥᐅᑕᓄᑦᓗ ᐱᓚᓚᐊᓗᑦᓗᑎᑉᐳ ᐃᓄᕆᔪᑎᑲᓚᕆᐅᕐᓂᖓᓪᓗᑦ ᓂᕿᐳᕆᐅᓂᖓᓪᓗᑦ ᑲᓇᑕᐅᑦ ᐅᐊᖏᓐᓇᖏᓂᒃ ᐊᒻᒪᓗ ᐊᑯᑭᑐᖓᑎᐅᑦ ᐱᖏᓐᓇᖏᓂᒃ (Harms, 2011; Henri, 2007). ᐃᓱᒪᓗᑕᐅᕐᒃᑐᑦ ᒥᑎᖅᓱᐅᖏᐃᐅᓗᐊᖅᑕᕐᓂᖓᓂᑦ ᐅᑭᐅᖅᑯᑦ ᓇᔪᖅᓯᒃᑕᕐᓂᓐᓂ ᐊᑯᑭᑐᐃᑦ ᓄᓇᖏᓕ ᐱᖏᓐᓇᖏᓂᒃ ᑕᐃᒪᖏᓂᒃ 1990-ᖏᓐᓂᒃ, ᐱᔪᓪᑕᐅᓱᒪᑦᓗᑎᓚᓗ ᒪᑕᒪᑕᐅᕈᑕᐅᓱᒪᑕᕐᓂᖅ ᐊᖓᔪᓇᓱᒃᑕᐅᕆᔨᓇᓐᓂᖏᓐᓄᑦ ᒪᑕᒪᑦᓗᑐᐃᓐᓇᑦ ᑕᐅᓐᓂ. ᐊᕐᕌᒍ 2002-ᒥ, ᐊᑯᑭᑐᐃᑦ ᓄᓇᖏᓂᒃ ᑯᑕᕆᖅᑎᑕᐅᓕᑕᐅᖅᑐᑦ ᒥᑎᖅᑕᐅᕆᔨᓇᕆᐅᕈᕐᓂᐊᖅᑐᓗᒃ. ᖃᐅᖅᑕᓗᒍᒥ ᒥᑎᖅᕆᕐᓇᖅ ᑕᒪᑐᒥ ᑭᖓᓂᐊᒍᑦ ᒥᑎᖅᑕᕆᕆᓇᖅᑐᖅᑐᐃᓐᓇᑦᑐᖅᑐᑦ ᐊᖓᔪᓇᓱᒃᑕᐅᕆᔨᓇᓐᓂᖏᓐᓄᑦ, ᐃᓱᒪᓗᑕᐅᕐᓂᑦ ᓯᑕ ᖃᓄᐃᓕᓐᓂᖏᓐᓇᓕᕆᔨᓂᖏᓐᓂ ᒥᑎᖅᑐᑦ ᐊᓯᖏᑐᓂᒃᑕᐅᖅᑐᖅ ᐅᓗᓚᐊᓇᖅᑐᖅᑐᐃᓐᓇᑦᑐᖅᑐᑦ ᐊᖓᔪᓇᓱᒃᑕᐅᕆᔨᓇᓐᓂᖏᓐᓂᑦ.

ᐱᓚᓚᐅᓂᖅᓯᐅᒍᑕᐅᖅᑐᒥᒃ ᐋᑐᓂᐊᓂᐊᓗᑎᑦᓗᑦ ᑎᑭᑕᐅᒍᑕᐅᓂᖓᓂᑦ ᓄᓕᖓᓀᓗᓴᖅᑐᒥᒃ ᒥᑎᖅ ᐊᑐᒍᑕᐅᖅᑐᖅ ᖃᐅᖅᑕᓗᒍᒥ ᑕᐃᒪ 2005-ᐯᒋᓪᓗᔪ, ᑕᐃᒪᖏᓂᓪᓗ ᐱᑕᖅᐃᐅᓇᓕᖅᓯᓂ, ᓴᖏᓂᖓᓂᑦ ᐊᔨᐅᖏᓂᒃᓗᐊᖅᓯᑎᒃ. ᑕᒪᐅᓇ ᑐᖅᑯᐅᑭᐅᓇᖅᑐᐊᓗ ᓴᖏᓪᓕᐊᓕᓴᐃᒃ ᒥᑎᖅᓄᑦ ᐊᒥᓱᓕᐅᓂᒃᑕᕆᔨᓇᓐᓂᖅᑐᖅ ᑕᐃᒪ 33-56% ᐳᓴᐅᑎᖏᓂᔨᓇᖅᑐᒥ ᒥᑎᑦ ᐱᑕᖅᑲᓂᖏᓐᓂᒃᓂᒃ, ᐊᑐᖅᓂᖏᓐᓂ ᐱᖏᓕᓂ 3 ᐊᖅᑳᔪᐃᑦ ᑕᐃᒪᓕ 2006-2008-ᒥ (Buttler, 2009; Iverson, 2015; Descamps, et al., 2009). ᐊᒥᓱᓂᖏᓂ ᐊᖅᓇᓪᑐᓂᖏᓂ ᑐᖅᑯᖅᑕᑕᓇᐅᖅᑐᒥ ᓄᓕᖓᓂᐊᑐᖅᑐᒥᒃ ᐊᓐᓂᐊᖅᓂᖅᒥᒃ ᑕᐃᒪᖏᓐᓄᒃ ᐊᒥᓱᓕᐅᓂᖅᐸᓪᓕᐊᓕᕐᓂᐅᑦ ᐊᖅᑕᐃᒥᒃ 2009 (Iverson, 2015). ᐃᓄᐃᑦ ᓄᓇᕗᓂᓂ ᖃᓄᐃᓂᖓᓂᒃᓴᐃᑦ ᒥᑎᖅᓄᑦ ᐊᑯᖅᑕᐅᓱᒪᔪᓄᑦ ᑕᑯᓱᒪᕐᕕᑦ ᐊᒥᓱᓕᐅᓂᖅᓯᐅᓂᖓᓂᒃ ᑐᖅᑯᖅᑎᑦᓗᑎᓂᖅᓯᐅᓂᖅᓯᐅᒃ ᐊᐅᑕᑐᖅᑕᐅᓱᒪᔪᓂᖅᓯᐅᓂᖅᓯᐅᒃ ᐊᓐᓂᐊᖅᓂᖅᒥᒃ ᐱᔪᓂᒃ ᓯᕐᓗᓂᖅᓯᐅᒃ (Henri, et al., 2018). ᓴᓐᓕᕆᐅᑦ ᐃᓱᒪᖏᖅᑐᑦ ᒧᓐᓇᐅᑉ ᐊᓐᓂᐊᖅᓂᖅᖏᓂᖅᓯᐅᒃ ᖃᓄᐃᓕᑯᓚᖅᒥᒃ ᐱᔪᓐᓇᕆᔨᓇᖅ ᑕᐃᒪᐃᓂᖅᓯᐅᓂᖅᓯᐅᒃ ᐅᓐᓄᑦᑕᐃᓐᓇᖅᑐᖅᑐᒥ ᒥᑎᖅᓄᑦ ᐃᓚᖅᑕᐃᓐᓇᖅᑐᖅᑐᒥ ᐊᓯᖏᑐᓂᒃ ᓄᓇᕗᓂᒃ, ᖃᐅᖅᑐᖅᑐᖅᓴᐅᓕᕐᖀᑦᑐᑦ ᓄᓇᕗᓕᖅᓴᒃ. Hᐊᓄᓕᕐ-Henri (2007) ᑎᑎᕋᓕᐅᖅᑐᑦ ᑕᐃᒧᑦᓴᐃᓇᐅᑦᓂᐊᕐᓇᖅ ᐃᓄᐃᑦ ᖃᐅᐅᔾᔨᕐᑐᖅᑐᒥᒃ ᐱᔪᓯᑎᖅᑐᑦᑐᖅᑐᒥᒃ ᐱᐅᓯᑎᖅᑐᑦᓗ ᖃᐅᖅᑐᖅᑐᐃᓐᓇᑦᑐᖅᑐᒥᒃ ᐃᓕᓴᕆᔨᓂᖏᓐᓄᑦ ᐊᐅᓚᑦᑎᓐᓂᖏᓐᓄᑦ ᐊᓯᖏᑐᓄᑦ ᓄᓇᕗᓂᑦ, ᖃᐅᖅᑐᖅᑐᖅᓴᐅᓕᕐᖀᑦᑐᓄᑦ ᓄᓇᕗᓕᖅᓴᒃ. Hᐊᓄᓕᕐ-Henri (2007) ᑎᑎᕋᓕᐅᖅᑐᑦ ᑕᐃᒧᑦᓴᐃᓇᐅᑦᓂᐊᕐᓇᖅ ᐃᓄᐃᑦ ᖃᐅᐅᔾᔨᕐᑐᖅᑐᒥᒃ ᐱᔪᓯᑎᖅᑐᑦᑐᖅᑐᒥᒃ ᐱᐅᓯᑎᖅᑐᑦᓗ ᐄᕐᓗᑦ.

ᐃᓐᓄᐊᖅᓯᒪᔪᓂᖅ ᐊᖅᑕᐃᓂ, ᐊᕐᑎᑎᓕᕆᔨᓴᐅᑉᐳᑦ ᖃᐅᔨᔪᖅᑎᖏᓂ ᑎᑎᕋᖅᓯᒪᔪᑦ ᓄᓇᕗᓂᑦ ᓚᐅᑦᓂᖅᓯᐅᓂᖅᓯᐅᒃ ᒥᑎᑦ ᓚᐅᖏᓐᓂᒃᓂᒃ ᐊᒻᒪᓗ. 1997-ᐯᒋᓪᓗᔪ, ᓚᖓᔪᓄᑦ ᑎᑭᑕᐅᓯᒪᓪᓗᑎᒃ ᓄᓇᖓᓄᑦ, ᑕᒪᐃᐅᓪᓕᐊᓕᒫᖅᑎᑦᓗᑎᓂᖅᓯᐅᓂᖅᓯᐅᒃ ᓯᕐᓗᖅᑎᖅᑕᐅᓕᕆᔨᓂᖏᓐᓂ ᒥᑎᖅᑐᖅᑐᖅᓴᐅᓕᕐᖀᑦᑐᑦ. ᑕᒃᑯᐊᒃ ᓄᓇᒃ ᓂᓗᓕᕆᔨᓂᖏᓐᓇᖅᓱᒃ ᐅᖏᓪᓕᐊᓂᖅᓴᐅᓂᒃ 14,000-ᓂᒃ ᒥᑎᑦ ᓚᐅᖏᓐᓂᒃ ᐅᓪᓗᒥᒃ 7-ᓂᒃ ᐱᓇᓯᐊᕐᓯᒥ. (Gilchrist & Heise, 1997). In 2012 and 2015, ᓚᓂᒃᓯᐅᓴᐃᐅᓂᖅᓯᐅᓂᒃ ᐱᔪᓕᐅᕆᔨᓂᖏᓐᓂ ᐅᓪᓗᓕᕐᓂᐊᖅᓯᐅᓂᖅᓯᐅᒃ, ᐱᐊᕐᓇᖅᓴᐅᓂᖅᓯᐅᓂᒃ ᒥᑎᖅᑐᖅᑐᐃᓐᓇᕐᓂᖓᓪᓗᒥ. (Gilchrist & Jassen, 2015;

Dey, et al., 2018; Dey, et al., 2019). ᐊᒥᓱᓪᓗᓂᒃᑐᑦ ᐅᑦᓗᖅᖃᐃᐅᓪᓚᕐᓂᖅᑎᑦ ᐊᓪᓚᑕᒃᑕᐅᓱᒪᓪᓗ ᒥᔅᓄᑦ ᐊᒥᓱᑐᖅᑲᓂᑕᓚᕐᓂᖅ ᑭᖑᔪᓂᐊᔪᑦ ᓄᓇᕗᓂᐊᓕᕆᔨᖅᑐᒥᒃ ᐊᓂᓂᐊᖅᓂᖓᑦ ᑎᑭᑕᐅᑕᐅᕙᓕᖅᑎᑦᓴᑎᑦ, ᐊᓚᒪᓗ ᒪᓕᑐᖅᖃᖅᑎᖅᓯᓂᓄᑦ ᒥᑎᖅᑕᐅᓚᓕᖅᓯᓚᓂᓂᒃ ᐊᒥᓱᓂᖅᓂᓂᒃ ᐊძᑭᑐᖅᒥᑕᓂ.

ᓱᓕᐅᑦ ᐊᓱᔨᖅᑕᕐᓕᐊᓂᖏᓐᓄ

ᓱᓕᐅᑦ ᐊᓱᔨᖅᑕᕐᓕᐊᓂᖏᓐᓄᓚᓄᑦ, ᐃᖏᒥᔪᐃᖏᐅᐅᑎᓗᓕᐊᖅᓗ, ᐊᓱᖏᑎᓄᓕᓗ ᐱᔨᕈᑕᐅᖏᒥᔪᓄᑦ, ᑕძᔪᐅᓕᓕᓕᐊᖅᑲᕈᑕᖅᑐᑦ ᐊᓱᔨᖅᓱᒪᓂᖏᑦ ᓄᓇᐃᐊ ᐅᒪᓕᔪᑕᖅᑐᖅᑐᑦ ᖃᖅᖅᓴᐅᑕᖅᑐᑦ ᐃᓕᔪᖅᑲᖅᓯᓕᓕᐅᖅᑐᑦ. ᐊᓱᔨᖅᑕᕐᓕᐊᔪᑦ ᑕᕆᐅᕗᑦ ᐊᒪᓕᓗ ᑕᓱᖅᓂᓂᒃᑐᓂ, ᓄᓇᐅᑦ ᐃᓗᐊᔪᑦ ᖃძᐊᖓᐃᐅᓪᓚᑐᖅᑐᓂ, ᐊᐱᖅᑲᑕᐅᓪᓚᑐᖅᑐᓂ ᐊᒪᓕᓗ ᓂᓪᑕᓱᓕᖅᑐᓂᓂᒃ ᖃᐅᔨᓴᖅᑕᐅᖅᑕᖅᓱᒪᓕᒫᒃ ᓇᓂᑐᐃᐅᓪᓗᖅ ᐅᑭᐅᖅᑕᖅᑐᑦᑎᓗ ᐅᓪᓚᕐᓇᖅᑐᖅᑐᓂ, ᖃᒦᓇᐃᑕᓕᑦᑕᐊᓂᖏᓐᓂᑐᓗ ᐅᓪᓚᕐᓇᐃᑦ ᑕᒪᒃძᓇᓂᖅᑲᑕᖅᑕᑐᑐᖅᑐᒃᑦ. (CAFF, 2010). ᓱᓪᓗ ᐅᒃᑐᑎᑕᓗᒍ, ᖃᑭᒥᐅᓂᖅᑲᓴᖅ ᑐᓕᐃᔨᖅᓵᒥᓂᖅᑲᓴᐅᓕᑦᓂᖏᓐᓄᓚᓄᑦ ᕼᐊᑦᓴᐊᓚᐸᑦ ᑕᕆᐅᕆᑐᐊᔪᑦ, ᓇᓇᐃᑦ ᑎᑭᑕᐅᑐᖅᑲᓴᐅᓕᕆᔨᖅᑐᑦ (Lunn, et al., 2016) ᖃᐅᔨᔨᐅᖅᑕᖅᑲᓱᒪᓗᑎᒃ ᑎᑎᕆᑕᐅᕙᓱᒪᔪᓂᒃ ᓇᓇᐃᐃᐅ ᐊᒥᓱᓂᖅᑲᐅᓕᕆᓂᖏᓐᓄ ᑎᖓᒥᐊᖅᑲᖏᓐᓇᓇᐊᖅᑕᓕᕆᖅᑐᑦ (Iverson, et al., 2014) ᑎᑭᓴᓕᓂᖅᑲᐅᖅᑲᑕᓕᕆᖅᓱᑎᒃ ᑎᖓᒥᐊᑦ ᐅᑦᓗᖅᖃᖏᓐᓄᓂᓂᑦ ᒪᓐᓂᓂᒃᑐᒃ ᓂᓚᔭᒃᑲᖅᓱᐅᖅᓂᓂᒃᑦ ᑕᐅᓐᓂ ᓴᑦᓕᐅᑦ ᖃᑭᖅᑑᓗᐊᓂᒃ ᐊᒃᑕᑐᖅᓴᐊᖅᑎᒥ (Smith, et al, 2010). ᐅᓇᖅᓂᖅᑲᐅᓂᓐᓄ ᓱᓕᐅᑦ ᐱᔨᕈᑕᐅᓕᕆᓕᓴᖅ ᒥᑎᑦ ᓄᓇᕗᑦᑕᓕᕆᓂᖏᓐᓄᓚᓄᑦ ᐊᖅᓴᒪᓗ ᐊᒥᓱᓂᖏᓐᓄ ᒪᓐᓂᑑᓚᓐᓂᖏᓐᓄᓚᓄᑦ ᐃᓐᒥᓄᑦ ᐊᒥᓱᓄᖏᓐᓂᕐᓂᖅᓯᓕᕆᔨᖅᑐᑦ (Dey, et al., 2018).

ᐊᓱᖏᑐᓂᓚᑕᐅᖅ ᖃᐅᔨᔨᐅᓱᒪᓕᓴᖅᑐᑦ ᓄᓇᒥ ᐱᐅᓱᓚᔨᐅᖅᑕᖅᑐᓂᓚᓄ ᑎᖓᒥᐊᖅᑲᖅᓯᓐᓄᖓᑦ, ᓱᓪᓗ ᐸᓂᓗᐊᖅᑕᕐᓕᓂᓂᖏᓐᓄᑐᓂᓚᓄᑦ ᓄᓇᐃᐃᑦ, ᐃᓚᖅᑲᓱᓕᕆᓂᖏᓐᓄᑐᓂᓚᓄ ᐊᓂᓂᐊᖅᑕᖅᑐᑦ ᐊᒪᓕᓗ ᓄᓇᐃᐃᑦ ᐃᓕᓴᐅᑦ ᐊᒥᓱᓂᖅᑲᓱᒪᓕᕆᓂᖏᓐᓄᓚᓄᑦ ᓄᓇᐃᐃᓂᒃᑐᓂᓚᓄ

ᐊᓯᖏᑐᓂᓗ ᐱᕕᖅᓴᓕᕆᔨᓄᑦᑕᕐᕕᐊᓂᖅ ᐊᐅᓪᓚᐅᓂᒃᓴᓂ ᑎᖏᑕᐊᖅᖃᕕᖅ. ᑲᔪᓯᑎᐊᕈᑐᓄᖅᑕᐊᕈᑎᑎᒃ ᐊᐅᓪᓚᕆᐊᔾᐅᓂᖏᑐᓂ ᑎᖏᑕᐊᖅᖃᕕᖅᓯᖏᓐᓄᑦ ᑐᖅᓚᕕᖅᑕᐊᖅᖃᕕᖅᓴᓕᓂ ᖃᓄᐊᓕᓴᐅᑎᖅᑕᐅᔪᓐᓇᖅᑐᑦ ᑕᒪᐃᔨᓄᖅ ᐊᑐᖅᑕᕐᓗᑎᒃ, ᐊᐃᓯᔾᔨᑕᐅᖅᑕᕐᕕᖅᑐᑐᔨᓂ ᖃᓄᐊᓕᓴᐅᑎᖅᑕᐅᔪᓐᓇᖅᓯᒪᔨᓂ.

ᐃᓄᐃᑦ ᑭᓇᐅᔪᑐᐃᓇᐃᑦ ᐱᓕᕆᐊᖅᓂᕐᓂ

ᐱᓕᕐᓕᐊᔪᒪᑐᐃᓇᓕᖅᓴᓐᓂ ᓄᓇᒥᐅᑕᓂᐅᒃ ᐊᒍᑦᖁᐃᐃᓴᖅᑭᒪᑦ ᓄᓇᒥᒃ ᐊᑐᑭᓇᐃᓕᓴᐅᑎᖅᑕᐅᔪᓐᓇᖅᓯᒪᓕᕐᖅᓯᐅᑎᓪᓗ ᐅᒥᐊᑦᔪᐊᖅᑯᑦᖃᕐᓂᐊᕆᐊᓇᓂᖅᓯᓄᐅᒃ ᓄᓇᒥᑦ Δᖅᑭᑦᕋᐃᖅᓴᓂᒃ ᑕᐅᖓᓇ ᐊᒥᕐᑐᑦ ᑕᓪᓕᐅᖅᓯᒪᔪᑦ Foxe Channel-ᑯᑦ ᐊᐅᓪᓚᑎᓂᖅᑭᓂᒃ ᐅᓴᕋᖅᓂᐊᑦᓂᒥᒃ. ᑕᒪᓐᓇ <ᑦᓇᐅᑕᐅᓯᒪᓂᖅᓯ ᑭᖓᔪᖅᓇᐊᖅᑕᐅᔪᓚᐅᐊᖅ ᑕᒪᐅᖅᓯ ᐊᑦᕋᒧᒪᑦ 2025-ᒧᑦ, ᓯᖖᓂᓴᒥ ᐅᖅᓯᐊᓗᖅᒥᒃ ᑯᕕᓯᐊᔪᒪᓕᕆᓂᖅᓴᓐᓂ ᐅᓗᓚᐊᓇᕐᑕᐅᖖᖅ ᑎᑭᐅᒪᔪᓕᕆᓂᖅ ᑎᖏᑕᐊᖅᖃᕕᖅᓴᖅᓯᒪᓂ.

ᓴᓪᓕᖏᐅᑕᒡᑕᐅᖅ ᐅᖅᖁᐅᓯᖁᖅᓯᒪᔪᑦ ᓴᓇᓴᐅᖅᑯᓴᓂᖅ ᒪᖅᑲᖅᓯ ᕿᐊᖅᑎᓂᒃ ᐃᐊᑭᑕᒃᑯᑦ ᐊᖅᑯᖅᓯᓯᒪᔪᓕᕆᓂᖅᑲᖅᓯᓂᒃ ᑎᖏᑕᐊᖅᖃᕕᖅᓴᖅᓯᒪᒃ. ᓯᖖᓪᓕᓂᖅ ᐊᖅᑯᑎᑎᖅᓴᐅᓇᓴᖅᑐᖅ ᐋᖅᑭᐅᒪᓇᓴᖅᑐᖅ ᐊᑭᐅᐊᑯᓗᓂ ᑭᓴᕐᖁᖅᓴᖅᓯᐊᕐᓇᓂᒃ, ᑕᐅᖏᓐᓄᑐᐊᓗᖅᓯᒃ ᑭᓄᐊᑎᑐᐊᓗᖅᓯᒃ ᐊᑭᐅᓯᑎᑐᐊᓗᖅᓯᒃ ᓄᓇᒥᑦ ᑭᓴᕐᖁᖅᓴᖅᓯᐊᕐᓇᓂᒃ ᑭᓴᕐᖁᖅᓴᖅᓯᐊᕐᓇᓂᒃ ᑎᖏᑕᐊᖅᖃᕕᖅᓴᖅᓯᒪᓂ. ᐃᓕᓴᕆᓂᖅᓴᖅᓯᒪᓂᒃ ᑎᖏᑕᐊᖅᖃᕕᖅᓴᖅᓯᒪᓂ ᑕᐅᖏᓐᓄᑐᐊᓗᖅᓯᒃ ᑎᖏᑕᐊᖅᖃᕕᖅᓴᖅᓯᒪᓂ ᐊᑭᐅᐊᑯᓗᓂ ᑭᓴᕐᖁᖅᓴᖅᓯᐊᕐᓇᓂᒃ ᑎᖏᑕᐊᖅᖃᕕᖅᓴᖅᓯᒪᓂ ᐊᑭᐅᓯᑎ

ᐊᐃᔅᑐᖏᓱᓪᓗᒍᓂᖅ ᑕᑯᑎᑎᑦᓴᓪᓕᒃ 7: ᓇᐃᐧᓴᖅᑕᐅᔾᔨᓱᓪᓕᒃ ᐊᐅᓪᓚᑎᓂᖅᑭᖅᓂᒃ ᐃᔨᓪᒪᔭᖏᖅᐅᔾᔨᕈᑎᔭᖏᖅᓴᐅᑦᓗ ᐊᓪᓚᓗ ᖃᓄᐃᓕᓐᐅᕈᑎᔭᖏᖅᓴᖅᑭᖅᓂᒃ

ᐊᐅᓪᓚᑎᓂᕆᑎᓂᒃ ᐃᔨᓪᒪᔭᖏᖅᓴᖅᑭᐅᕈᑎᔭᖏᖅᓴᐃᑦ	ᒪᓪᓕᖅᑐᑦ ᑐᖓᓚᐅᖅᑐᐅᔾᔨᓂᒃ/ᐱᔭᕆᖏᖅᑕᐅᔾᔨᒪᔨᓂᖅᓂᓗ ᐊᐃᔅᑐᖏᓱᓪᓕᒃ ᑕᑯᑎᑎᑦᓴᓪᓕᒃᑎᓪᓗᒍᓂᖅᓴᐅᓂᖅᓂᒃ 6-ᒥᑦᑐᓂᒃ	ᐊᐅᓪᓚᑎᓂᕆᓪᓗᑦ ᖃᓄᐃᓕᓐᐅᕈᑎᔭᖏᖅᓴᐃᑦ
ᐱᓪᓚᓐᓂᖅ ᐊᔨᔭᓪᓚᖅᓴᓂᖅ ᑎᖅᑎᐊᖏᖅᓴᐃᐅᓂᖅᓴᓂᒃ ᑲᓇᑕᒥᒪᒃ ᐅᒪᔭᐅᑕᐅᓗᑎᒃ ᐱᐅᓂᖏᓐᓄᑦᓴᓂᖅᑭᑦ ᐊᐅᓪᓚᑎᓂᖅᓴᐃᐅᓂᖅᑭᑦ ᐊᓕᖅ ᐊᖏᔪᒪᓂᒪᖅ ᓴᓐᑎᕐᓯᐅᓂᖅᑭᒃ ᑕᒪᖏᖅᑐᒃ ᐅᒪᔭᓚᓂᒪᑦ, ᐱᓗᐊᖏᖅᑎᑦᓗ ᐅᒪᔭᖅᑭᑦ ᐊᒥᓱᖅᑭᓗᐊᓚᖏᖅᑐᑦ.	1.1, 1.2, 1.3, 1.4	6.1 ᐱᐅᔨᓴᒪᒥᔪᑦ ᓄᓇᒥᐅᑕᓂᒃ ᐊᐅᓚᑎᓂᖏᖅ 6.2 ᐅᒪᔭᓚᓴᓂᖏᒃ ᐊᓕᓗᓐ ᐅᒪᔭᖏᖅᓴᐃᐅᓕᖅᓂᖅᑭᖅᓂᒃ ᐊᐅᓚᑎᓂᖏᖅ 6.3 ᓇᐅᑎᑎᖅᑐᐃᓴᖅᑐᖏᖅᓴᐃᐅᖏᖅᑕᓂᖅᑭᑦ ᖃᓄᐅᔨᖏᖅᓴᐃᐅᕈᖅᑭᑦᓗ 6.4 ᐃᓄᖏᑎᑎᑭᓴᐅᔨᓂᒃ ᖃᓄᐅᔨᓗᑎᑎᓂᖅᓴᖅᑕᐃᓴᖅᑕᖏᖅᖢᓂᒃ ᐊᓕᓗᓐ ᖃᓄᐅᒪᔨᓴᒃᑕᐃᓕᖅᓂᒃ ᐊᐅᓚᑎᓂᕐᓂᖅ.
ᑲᖓᐃᐃᑦ ᐱᔨᓭᑕᐅᓂᖅᑭᑦ ᓄᓇᐃᑦ ᓴᖅᓯᑎᖅᑕᐅᓂᖅᓴᓄᑦᓴᓂᖅᑭᑦᓗ ᐊᓕᓗᓐ ᐊᓂᖅᑭᓂᒃ ᐅᒪᔭᓐᓄᑦ ᐊᑯᖅᓂᓂᒪᔨᓇᓕᖅᓂᖅᑭᒃᓄᑦᓗ	1.2	6.2 ᐅᒪᔭᓚᓴᓂᖏᒃ ᐊᓕᓗᓐ ᐅᒪᔭᖏᖅᓴᐃᐅᓕᖅᓂᖅᑭᖅᓂᒃ ᐊᐅᓚᑎᓂᖏᖅ 6.3 ᓇᐅᑎᑎᖅᑐᐃᓴᖅᑐᖏᖅᓴᐃᐅᖏᖅᑕᓂᖅᑭᑦ ᖃᓄᐅᔨᖏᖅᓴᐃᐅᕈᖅᑭᑦᓗ
ᓄᓕᖅᓂᐊᓚᐅᖅᑐᒃ ᐊᓐᓂᐊᓂᐊᓚᐅᑐᖅ ᓴᓂᑭᑕᕐᖅᓴᑦ ᐊᐅᑐᖅᑐᐊᓚᐅᑐᖅ ᐱᓂᐅᓂᖅᓴᖅ ᒥᑋᑦ ᐱᑕᖅᓂᖅᓴᐅᓂᖅᑭᑦ	1.2	6.2 ᐅᒪᔭᓚᓴᓂᖏᒃ ᐊᓕᓗᓐ ᐅᒪᔭᖏᖅᓴᐃᐅᓕᖅᓂᖅᑭᖅᓂᒃ ᐊᐅᓚᑎᓂᖏᖅ 6.3 ᓇᐅᑎᑎᖅᑐᐃᓴᖅᑐᖏᖅᓴᐃᐅᖏᖅᑕᓂᖅᑭᑦ ᖃᓄᐅᔨᖏᖅᓴᐃᐅᕈᖅᑭᑦᓗ
ᓇᓄᐃᑦ ᓂᓴᐳᓴᖅᓴᐅᖅᐸᖓᓂᖅᑭᑦ ᒥᑎᑦ ᐱᑕᖅᓴᓂᖅᑭᖅᓂᒃ	1.2	6.2 ᐅᒪᔭᓚᓴᓂᖏᒃ ᐊᓕᓗᓐ ᐅᒪᔭᖏᖅᓴᐃᐅᓕᖅᓂᖅᑭᖅᓂᒃ ᐊᐅᓚᑎᓂᖏᖅ 6.3 ᓇᐅᑎᑎᖅᑐᐃᓴᖅᑐᖏᖅᓴᐃᐅᖏᖅᑕᓂᖅᑭᑦ ᖃᓄᐅᔨᖏᖅᓴᐃᐅᕈᖅᑭᑦᓗ

ᐊᐅᓪᓚᕐᓂᕆᓐᒃ ᐃᓱᒪᓕᕆᔨᐅᕙᑎᓕᕆᓂᑦ	ᒪᓪᒃᑐᑦ ᑐᕆᕆᐃᓐᓂᐅᔭᕐᓂᖅ/ᐱᓕᕆᖏᑕᐅᔪᓚᐅᖅᓗᓂ ᐊᓘᐃᑐᖏᔨᒪᔪᓂ ᑕᔪᑎᑦᑎᐊᕐᓇᓕᕆᓂᖅ 6-ᒥᑦᑐᓂᒃ	ᐊᐅᓪᓚᕐᓂᕆᔨᑦ ᖃᓄᐃᓪᓕᐅᕙᑎᓕᕆᓂᑦ
ᓴᓪᓕᕆᐅᑦ ᐃᓱᒫᒐᕐᓂᕆᓂᖏᑦ ᖃᐅᔨᓴᖏᑕᐅᖏᑦᑕᓂᓕᕆᓂᖅ	2.1	6.4 ᐃᓄᖅᔪᑎᑎᕆᐅᔪᓂᒃ ᖃᐅᔨᒪᑎᑦᑎᓂᕐᖏᓴᐅᓚᐅᖏᓂᖅ ᐊᐅᒪᓗ ᖃᐅᒪᕆᐅᓴᕆᓚᐅᓕᓂᒃ ᐊᐅᓪᓚᕐᓂᕆᓂᖅ.
ᓴᓪᓕᕆᐅᑦ ᐃᓱᒪᓕᕆᓂᖅᓂᕆᓂᖏᑦ ᐊᑐᖏᑕᐅᓂᕐᖏᓚᐅᖏᓂᖅᓂᕆᓂᖅᓂᖅ ᐃᓄᐃᑦ ᖃᐅᔨᒪᕗᑦᓯᐅᑕᖅᓯᒪᓂᖅ ᖃᐅᔨᓴᖏᑕᐅᓂᐊᖏᑐᓂᒃ ᐊᒻᒪᓗ ᐊᐅᓪᓚᕐᔪᐅᖏᑕᐅᔪᑦ ᐃᓱᒪᓕᐅᕈᑕᐅᔪᓚᐅᓕᓂᒃ	2.1	6.4 ᐃᓄᖅᔪᑎᑎᕆᐅᔪᓂᒃ ᖃᐅᔨᒪᑎᑦᑎᓂᕐᖏᓴᐅᓚᐅᖏᓂᖅ ᐊᐅᒪᓗ ᖃᐅᒪᕆᐅᓴᕆᓚᐅᓕᓂᒃ ᐊᐅᓪᓚᕐᓂᕆᓂᖅ. 6.3 ᓇᐅᑦᑎᖏᑐᐃᓴᖏᑐᔾᔨᖏᓚᐅᖏᑦᑕᓂᕐᓂᑦ ᖃᐅᔨᓴᕐᔨᖏᓐᓂᑦᓗ
ᓴᓪᓕᕆᐅᑦ ᐃᓱᒪᓕᕆᓂᖅᖏᑦ ᐃᑲᕆᐊᖏᓴᐅᑎᑦᑕᐅᓂᕐᖏᓚᐅᖏᓂᖅᓂᕆᓂᖅᓂᖅ ᐃᓕᐅᐅᓂᖏᓚᐅᖏᑦᑕᕐᓗᑎᓗ ᑭᓇᐅᔭᓕᐅᖏᓚᐅᖏᑦᑕᖏᑐᓂᒃ ᐊᓱᖏᑐᓂᓗ ᐱᓵᖅᓯᕐᕕᐅᔪᓚᐅᖏᑦᑕᖏᑐᓂᒃ ᐊᓱᖏᑐᓂᓗ ᐊᐅᓪᓚᕐᓂᕆᓂᖅᓂᖅ ᑎᖏᒥᐊᖏᓴᐅᒥᒃ	2.2	6.4 ᐃᓄᖅᔪᑎᑎᕆᐅᔪᓂᒃ ᖃᐅᔨᒪᑎᑦᑎᓂᕐᖏᓴᐅᓚᐅᖏᓂᖅ ᐊᐅᒪᓗ ᖃᐅᒪᕆᐅᓴᕆᓚᐅᓕᓂᒃ ᐊᐅᓪᓚᕐᓂᕆᓂᖅ.
ᐊᐅᑐᖏᑕᐅᓱᒪᓂᖅᓂᑦ ᓱᓪᒪᐅᑦ ᐊᓱᓴᖏᑦᓯᓪᓚᐊᓂᖅᓯᒪᓄᑦ ᑎᖏᒥᐊᖏᓴᐅᒥᒃ ᐊᐅᓪᓚᕐᓂᕆᓂᖅᓂᖅ	1.2	6.3 ᓇᐅᑦᑎᖏᑐᐃᓴᖏᑐᔾᔨᖏᓚᐅᖏᑦᑕᓂᕐᓂᑦ ᖃᐅᔨᓴᕐᔨᖏᓐᓂᑦᓗ
ᐊᒥᓱᓂᖏᓚᐅᓪᓚᕐᓇᖏᓂᖅᓂᑦ ᐅᒥᐊᑦᔾᐊᑦ ᐅᓱᓴᐅᔾᔨᑦ ᑕᐅᐅᓇ ᐊᒥᑦᑐᑦ Foxe Channel-ᓄᑦ ᐅᓗᓚᐊᖏᓂᕐᓂᓗ ᐅᖏᓱᐊᓗᓂᒥᒃ ᓄᐃᓱᔾᔨᖏᓂᕐᓂᓴᓄᑦ	1.2, 3.2, 3.3	6.3 ᓇᐅᑦᑎᖏᑐᐃᓴᖏᑐᔾᔨᖏᓚᐅᖏᑦᑕᓂᕐᓂᑦ ᖃᐅᔨᓴᕐᔨᖏᓐᓂᑦᓗ 6.4 ᐃᓄᖅᔪᑎᑎᕆᐅᔪᓂᒃ ᖃᐅᔨᒪᑎᑦᑎᓂᕐᖏᓴᐅᓚᐅᖏᓂᖅ ᐊᐅᒪᓗ ᖃᐅᒪᕆᐅᓴᕆᓚᐅᓕᓂᒃ ᐊᐅᓪᓚᕐᓂᕆᓂᖅ.

ᐊᐅᓪᓚᕐᓂᕆᓂᖅ ᐃᓱᒪᖏᖅᓯᐅᑕᑎᖏᓐᓄᑦ	ᒪᓕᖅᑐᑦ ᑐᕆᓚᐅᖅᑕᐅᔪᓂᖅᓯᐅᑕᐅᔪᒪᔪᓂᒃᓗ ᐊᓘᖅᑐᖏᓯᒪᔪᓂ ᑕᔾᐅᑎᖏᓐᓄᖏᖅᓯᒪᔪᓂᖅ 6-ᒥᑦᑐᓂᖅ	ᐊᐅᓪᓚᕐᓂᕐᒧᑦ ᖃᓄᐃᓕᓴᐅᑕᑎᖏᓐᓄᑦ
ᐄᐊᓐᑕᖏᔾᑦ ᐊᖏᖏᓯᐅᖏᑕᐅᔪᒪᓂᖅᑭᑦ ᐃᓗᐊᓂᓗ ᐊᑭᐊᓂᓗ ᑎᖅᑎᐊᖏᖅᖃᓘᓐᓃᖅᐸ	2.3, 3.1, 3.2, 3.3	6.4 ᐃᓄᖏᔾᔪᑎᑎᒃᐅᔪᓂᖅ ᖏᐅᕗᒪᑎᑦᑎᓂᖏᖅᓴᐅᓚᐊᖏᖅᖏᓂᖏᖅ ᐊᓯᒪᓗ ᖏᐅᒪᔾᐅᓴᓚᐊᑕᖅᓂᖅ ᐊᐅᓚᑦᑎᐊᖅᓂᖏᖅ.
ᐊᒥᓱᓂᖏᖅᓴᐅᓴᖅᓂᖅᑭᑦ ᐃᓚᐅᔪᓴᖅᑕᑦ ᐊᐅᓪᓚᐹᑦᓴᐅᑕᖏᓂᖅᒥᑦ ᐊᑭᓕᑭᐅᖅᑕᑦ/ᐊᖓᔪᓇᓱᑎᐊᑭᐅᓴᔪᓴᖅᑕᑦᓗᓴᓐᓂᑦ ᑎᑭᑕᐅᓴᓵᔾᓂᖏᖅᓴᐅᑕᓇ ᑎᖅᒥᐊᖏᖅᖃᓘᓐᓃᖅ	2.3, 3.1, 3.2	6.4 ᐃᓄᖏᔾᔪᑎᑎᒃᐅᔪᓂᖅ ᖏᐅᕗᒪᑎᑦᑎᓂᖏᖅᓴᐅᓚᐊᖏᖅᖏᓂᖏᖅ ᐊᓯᒪᓗ ᖏᐅᒪᔾᐅᓴᓚᐊᑕᖅᓂᖅ ᐊᐅᓚᑦᑎᐊᖅᓂᖏᖅ.
ᐅᒥᐊᔪᖏᓴᓕᓵᓚᖅᐸᑦ ᓂᐅᑕᐅᔪᓂᖅᓯᐅᑕᐅᔪᓂᖅᓴᐅᓚᐊᖏᖅᓂᖏᖅ ᐳᓚᕆᑎᓂᖅ ᑲᓇᖏᖅᓴᐅᓴᖅᑐᖃᓕᓴᐅᑕᐅᔪᓂᖅᓴᐅᓚᐊᖏᖅᓂᖏᖅ ᖏᖅᖏᓚᐅᖅᓴᖅᑐᓕᕆᓂᖅ	1.2, 3.1, 3.2, 3.3	6.4 ᐃᓄᖏᔾᔪᑎᑎᒃᐅᔪᓂᖅ ᖏᐅᕗᒪᑎᑦᑎᓂᖏᖅᓴᐅᓚᐊᖏᖅᖏᓂᖏᖅ ᐊᓯᒪᓗ ᖏᐅᒪᔾᐅᓴᓚᐊᑕᖅᓂᖅ ᐊᐅᓚᑦᑎᐊᖅᓂᖏᖅ.

6 ᐊᐅᓪᓗᕆᐅᔪᓂᖅ ᖃᓄᐃᓕᓚᐅᕈᑎᒃᓴᐃᓪ

ᑲᓇᑕᒥ ᐅᒪᔨᓪᓚᕆᐊᒃᑯᓪ ᐃᓕᓴᕐᓂᕐ ᐃᖏᓕᕐᕕᖏᓐᓂᖅ ᐊᐅᓪᓗᕆᐊᐅᖏᖃᑦᑕᐅᓂᖅᓂᓪ ᑲᑎᒪᔨᕋᓱᒃᑐᕐ ᐊᐅᓪᓗᕆᑎᓂᐊᖏᖅᑐᓂ <ᖏᓇᐅᑎᓂᒃ ᐊᖏᕐᕋᑕᐅᓪᓚᐅᖏᓪᓗᑎᒃ, ᓂᕈᒪᓂᒃ ᐱᓗᐊᖏᖅᑐᒥᓪᓗᒥ ᖃᓄᐃᓕᓴᕐᕕᖏᓐᓂᖅᓂᒃ: 6.1-Γ, ᐱᐅᓂᓴᒥᔪᓂ ᓄᓇᒥᐅᑕᓂᒃ ᐊᐅᓪᓗᕆᑎᓂᖅ 6.2-Γ ᐅᒪᔨᓪᓚᕆᓂᒥᑦᑐᒃ ᐊᓯᓗᓐ ᐅᒪᔨᖏᖃᕐᕕᐅᕙᓂᖓᓂᕐᓂᒃ ᐊᐅᓪᓗᕆᑎᓂᖅ, 6.3-Γ ᓇᐅᓂᕆᖏᑐᐃᓴᖏᑐᖃᕐᕕᐅᖏᖃᑕᑎᓂᕐᓂ ᖃᐅᔨᓴᖃᕐᕕᐅᕙᓂᖓᓂᕐᓂᓪᓗ ᐊᓯᓗᓐ 6.4-Γ ᐃᓄᓴᐃᑎᑎᕐᐅᔨᓂᒃ ᖃᐅᔨᓚᑎᓂᑎᓂᕆᕐᓴᐅᔪᐊᖏᖃᓄᖅᓂᖅ ᐊᐅᓪᓗᕆᓚᖅᓂᒃ ᐊᓯᓗᓐ ᖃᐅᔨᓴᐅᕐᓴᐅᔪᐊᓪᓚᖅᓂᒃ ᐊᐅᓪᓗᕆᑎᓂᖅ. ᑕᓚᐅᔪᐊ ᖃᓄᐃᓕᓴᕐᐅᑎᓂᐊᖏᖅᑐᓂ ᐃᑲᔪᑕᐅᓂᐊᖏᖃᑦᑕᓂᖅ ᐊᖓᒍᓚᔪᓐᓇᖅᓂᖓᓪᓗᒥ ᑕᐅᑐᓚᓂᕆᐊᔪᓂᒃ, ᐅᖁᓚᓂᕆᐅᔪᐊᓂᒃ ᐊᓯᓗᓐ ᐱᓴᓱᖏᑕᐅᔪᓚᔪᓂᒃ ᑕᐅᔪᓚᐅᓂᖅ ᐊᐅᓪᓗᕆᑎᓂᖅ <ᓇᐅᐅᑎᓂᒃ, ᐅᔨᓴᖏᑐᑎᐊᖅᑎᓪᓗᓐᓄ ᐊᐅᓪᓗᕆᑎᓂᖓᓂᒃ ᐃᓕᓚᒃᓴᖏᑐᐃᓴᖏᑐᕐᕕᐅᕙᓂᖓᓂᕐᓂᒃ. ᑲᓇᑕᒥ ᐅᒪᔨᓪᓚᕆᐊᒃᑯᓪ ᐊᑐᓕᖏᑐᐃᓴᖏᑐᖃᑦᑕᓂᖅ ᑕᒪᒃᑯᐊ ᖃᓄᐃᓕᓴᕐᕕᐅᕙᓂᖓᓂᕐᓂᒃ, ᐊᖏᖅᑎᒋᒪᒃ ᐊᑐᐃᓐᓇᕐᓂᖓᓂᕐᓂᒃ ᐱᓕᕆᔨᐅᓂᐊᖏᖃᑦᑕᓂᖅᓂᒃ, ᐊᑐᖏᓪᓗᓐ ᐊᑐᐃᓐᓇᕐᓂᖓᓂᕐᓂᒃ ᐱᓕᕆᔨᐅᓂᐊᖏᖃᑦᑕᓂᖅ

2.1.6). ᑲᓇᑕᒥ ᐅᒪᔨᓕᓗᐸᒋᑦ ᐃᖏᓂᐅᕙᖅᓯᒋᐅᓂᖅᓴᓗ ᐊᐅᓕᕐᑎᓴᐅᖏᑦᑐᖅᑕᐅᓂᖅᓴᓗ ᑲᑎᒪᔨᖁᓕᖅᑭᓂ ᑐᓴᕐᓯᐃᐸᓂᑦ ᐃᓐᓇᐸᒋᓂᑐᓂᐸ ᐱᔨᑎᖏᖅᑐᑐᓂᐸ ᐃᓐᖏᑐᒃᓯᐃᐅᓂᖅᑐᑐᖅᑐᖏᖅᑐᓂᐸ, ᐃᓐᖏᓐᓄᑦᓗ ᐱᓕᒪᓕᐅᔨᓂᐸ ᐃᓕᓴᐸᒋᓂᖅᑐᒥᔪᑦ ᓄᓇᕗᒻᒥᖅ, ᖃᐅᔨᓴᒪᔭᐅᓕᖅᑐᑐᓂᐸ ᖃᐅᔨᓴᑎᐊᖅᑕᐅᓕᐅᔪᓂᕐᓂᓕᕐᓂᐸ ᒪᓕᕐᓂᑎᑐᓂᐸ 6.4-ᒥᐸ 6.7-ᒧᑦ ᐃᓐᐃᑦ ᐃᑲᔪᐊᖅᓴᓕᕐᓂᐸᓂᑦ ᐊᖏᕐᕕᑎᓂᐸ, ᐱᔨᓴᒪᔭᐅᓕᖅᑐᑐᖅᑐᓂᐸ ᑎᓕᐊᖅᑕᐅᓕᐅᔪᓂᕐᓂᓕᕐᓂᐸ.

ᑲᓇᑕᒥ ᐅᒪᔨᓕᓗᐸᕐᓂᑦ ᐃᖏᓂᐅᕙᖅᓯᒋᐅᓂᖅᓴᓗ ᐊᐅᓕᕐᑎᓴᐅᖏᑦᑐᖅᑕᐅᓂᖅᓴᓗ ᑲᑎᒪᔨᖁᓕᑦ ᓄᓇᕗᒻᒥᖅ ᖃᐅᔨᓴᖁᓕᖅᑭᓂᐅᓂᖅᑐᑐᖅᑐᖏᖅᑐᓂᐸ, ᖃᐅᔨᓴᒪᔨᐅᖅ ᐊᑯᐅ ᐱᓕᓕᐊᖅᑕᐅᓕᐅᔪᓂᕐᓂᓕᕐᓂᐸᓂᑦ, ᑎᑎᕋᖅᑕᐅᓂᖅᑐᑐᖅᑐᖏᖅᑐᓂᐸᓗ, ᑐᓴᕐᓯᐃᐸᓂᑦ,ᓴᓗᐅᓂᖅᑐᑐᖅᑐᖏᖅᑐᓂᐸᓂᑦᓗ ᓄᓇᕗᒻᒥᐅᑕ

(4) ᓇᓚᐅᓂᔪᖅᓯᒪᓂ, ᑐᑭᓯᐊᖏᓐᓄᐃᑦᑐᕆᔨᓗᑎᒃ ᐃᓄᐃᑦ ᐃᓂᒪᓕᑕᐅᔨᖏᓪ ᐊᑐᑎᓂᑦᑎᐊᖅᑐᑦᑐᒃ ᐊᖏᕐᐅᒪᑦᑎᐊᖅᑐᑦᓯᓗᓪᓕ ᐊᐅᑕᑦᑎᓴᓇᐊᖅᑎᓂᖏᓐᓂᒃ ᐅᒪᔨᔨᓂᖅᑎᓄᐅᕙᖅᑐᑦᑐᒃ ᐊᓚᒪᓗ ᐅᒪᔨᔨᓂᖅᓄᐃᐅᕙᖅᑐᑦᑐᒃ ᓄᓇᒃ ᑎᓐᓕᐊᖅᑎᓂᖅᓴᕐᑐᑦᑐᒃ.

6.1.3 ᐃᓄᐃᑦ ᑕᐅᕙᓂ ᑕᐃᔪᓯᑐᖅᑎᑦᑕ ᓄᓇᒧᑦ ᐃᒪᖅᓄᑦᓗ

ᐱᓱᓴᖅᑎᖅᑎᑕᐅᕐᓂᑦ ᐊᑕᐃᑦ *ᐃᑲᔪᐊᓴᖅᓯᒪᓄᑦ ᐊᖏᕐᕈᑎᑎᒍᑦ (ᑎᑎᖅᑲᖏᖓᓂ 2.1.6-ᒥ)* ᑎᑎᕋᖅᑕᒪᔨᐊᖅᑎᓂᖏᖓᓂᒃ ᐃᓄᖓᓄᑦ ᑕᐅᔪᓯᑐᖅᑲᓱᐅᔨᓂᒃ ᓄᓇᒧᑦ ᐃᒪᖅᓄᑦᓗᓄᐃᐅᕙᖅᑐᑦᑐᒃ ᐃᓄᐃᑦ ᑕᐅᔪᓯᑐᖅᑲᓱᐅᔨᓂᒃ ᑕᒪᑦᑯᐊ ᐊᐅᑕᑦᑎᓴᐅᑎᓂᖏᖓᓄᑦ ᓴᓗᐅᓂᖅᑕᐅᓯᒪᔪᖓᑎᐅᑎᑕᐅᕐᓂᒃ. ᓄᓇᒧᑦ ᑕᐃᔪᓯᑐᖅᑐᓄᑦ ᑎᑎᕋᖅᑕᐅᑕᐅᓱᒪᔨᓂ ᒪᕐᐃᕐᖅᑕᖅᓯᓗᒃ ᓴᓗᐅᓂᖅᑎᑎᖅᑕᐅᓯᒪᔨᓂ ᑐᓴᐅᒃᑐᓂᒃ (2006-ᒥᓗ 2012-ᒥᓗ). ᑲᓇᑕᒥ ᐅᒪᔨᓕᒪᓕᕐᑐᑦ ᐃᔅᓯᐅᖅᓱᑯᓗ ᐊᐅᑕᑦᑎᓴᐅᖅᑎᑕᐅᕙᑦ ᐊᑐᑕᖅᑎᑎᒍᑦᓕᓕᐊᓯᐃᓐᓇᔅᓯᐊᖅᑎᒥᔨᑦ ᐊᓯᕐᑐᓂᒃ ᖃᐅᔨᓴᒪᐅᓯᕐᑐᓂᒃ ᑲᑎᕋᖅᑕᐅᓯᒪᔨᖅᑕᐅᓯᕐᑐᓂᒃ ᐊᐅᑕᑎᑕᐅᓯᐊᖅᑕᒥᑦ ᑎᑎᕋᖅᑕᐅᑕᓂᑎᒍᑦᓕᓕᐊᓯᐃᓐᓇᔅᓯᐊᖅᑎᒥᔨᑦ ᖃᐅᔨᓴᒪᐅᓯᕐᑐᓂᒃ ᑲᑎᕋᖅᑕᐅᓯᒪᔨᖅᑕᐅᓯᕐᑐᓂᒃ ᖃᐅᔨᓴᒪᐅᓯᕐᑐᓂᒃ ᑕᐃᔅᓯᐊᖅᑕᒥ.

6.2 ᐅᒪᔨᓕᒪᓂᖅᑎᒃ ᐊᓚᒪᓗ ᐅᒪᔨᖅᑲᓇᐅᒍᖏᓂᖅᓯᒪᓂᒃ ᐊᐅᑕᑦᑎᓂᖅᓴᖅ

6.2.1 ᐊᓯᔾᔨᖅᑕᐅᓂᖅᓱᒃ ᐃᓕᑕᒪᔭᐅᓯᒪᔨᔪᖅ ᖃᐅᓄᐃᑦᑐᑐᓂᖅᓱᓂ ᓴᓗᐅᓂᓯᐊᖅᓯᒪᒍᓇᐅᓂᖅ

ᐃᓐᓇᐅᔨᔪᖅ ᑕᒪᓐᓇ ᖃᖅᖢᓴᐅᖅᑐᑎᒥ ᑎᖅᑕᐊᑦᑎᓇᖅ ᓴᓗᐅᓂᖅᑕᐅᓯᒪᓂᖅᓱᒃ ᐊᑐᑎᓂᖅᑎᑎᐊᖅᑐᑦᑐᒃᓴᓇᓗᐊᖅᓯᓂ ᑎᖅᑕᐊᑦᑎᓇᐅᓂᒍᑦ, ᐊᓯᖅᑎᑦᑕᐅᖅ ᐅᒪᔨᐃᑦ ᐊᒥᓯᖅᑕᓗᐊᓕᖅᑐᑦ ᐊᓚᒪᓗ ᐅᒪᔨᐃᑦ ᐃᑲᔪᖅᑕᐅᖏᓪᓕᕐᓂᐅᓇᖅᓴᐅᓇᖅᑐᑦ ᐊᑦᕿᔪᓕᒫᓇᖅ ᓴᓗᐅᓂᖅᑕᐅᓯᒪᔨᖅᑕᐅᓯᕐᑐᓂᒃᓇᓇᐅᓂᖅᓯᓄᑎᓂᑎᒃ. ᑲᓇᑕᓕᒫᒥ ᐅᒪᔨᖅᑎᓂᐅᑕᐅᓂ ᐱᐅᓂᖅᓴᓂᒃ ᓴᓗᐅᓂᖅᑕᐅᓯᒪᔨᖅᑕᐅᓯᕐᑐᓂᒃᓇᓇᐅᓂᖅᓯᓄᑎᓂᑎᒃ ᑕᒪᑎᓂᒃ ᐅᒪᔨᐃᑦ ᐊᓚᒪᓗ ᐃᓐᓇᐅᔨᓂᖅᓴᐅᓂᖅᓴᓂᒃ ᓴᓗᐅᓂᖅᑕᐅᓯᒪᔨᖅᑕᐅᓯᕐᑐᓂᒃᓇᓇᐅᓂᖅᓯᓄᑎᓂᑎᒃ ᐊᑐᑎᓂᖅᑎᑎᐊᖅᑐᑦᑐᒃ ᐃᓄᐃᑦ ᐃᑲᔪᖅᑕᐅᖏᓪᓕᕐᓂᐅᓇᖅᓴᐅᓇᖅᑐᑦ ᐊᑦᕿᔪᓕᒫᓇᖅ ᓴᓗᐅᓂᖅᑕᐅᓯᒪᔨᖅᑕᐅᓯᕐᑐᓂᒃᓇᓇᐅᓂᖅᓯᓄᑎᓂᑎᒃ

ᐃᓱᒪᒃᖃᓱᐅᑎᖃᖅᓂᐊᖅᑐᑦ ᐃᓕᑕᒫᖅᐅᓱᒪᓕᓂᒃᓴᓂᒃ ᐊᓱᔪᖅᑕᐅᖅᑯᓲᐅᓂᒃᓴᓂᒃ. ᑭᒃᑕᖃᖅᖁᐅᓂᖏᑕᐅᖅ ᐊᓱᔪᖅᑕᐅᓂᐊᖏᓐᓇᑦ ᐅᖅᐅᓱᐅᐊᓪᓗᓐᓄᑦ ᐅᒪᑭᖅᐅᓂᒃᓴᓂ.

6.2.2 ᐊᒥᓱᓕᐊᑕᖅᓂᖅᑎᑦ ᑲᖏᐃᑦ

ᐊᒥᓱᓕᐊᖅᑐᓴᒪᓗᐊᓕᑕᖅᓂᖅᑎᑦ, ᓇᒍᑎᑭᓕ, ᐊᕐᑎᑎᑭᓴ ᐊᐅᑐᖅᓱᕐᑕᐊᓂᖅᑎᑦ ᑲᖏᐃᑦ ᓂᕿᑭᖅᖁᓱᐅᓕᐊᖅᑕᑕᖅᓂᖅᑎᑦ ᐃᓕᓕᐊᓂ ᖃᖅᖁᖅᐅᖅᑐᑦ ᑎᖅᑭᐊᖅᖁᖃᕐᓂᖅᑕᒪᐊᖅᑭᓕᑦ, ᑭᓱᐊᓂ ᐅᒪᓕᔪᑕᓗᓴᐅᑦ ᑕᒪᐃᓐᓇᖅᑐᑦ ᐃᓱᒪᒃᖃᓱᐅᑎᖃᖅᓂᖅᑎᑦ ᐊᐅᑕᑕᐅᔪᓐᓇᖅᓂᖅᑐᓂᓂᐅᑦ ᖃᓄᐃᐃᑕᖅᓂᖅᑐᓂᓂᐅᑦ, ᐊᓖᓗᓐ ᐊᑐᖅᑕᐅᔪᓐᓇᖅᑐᑦ ᖃᓄᐃᐃᑕᐅᑭᐅᖅᓂᓕᑦ. ᑲᖏᐃᑦ ᐊᒥᓱᓕᐊᑕᖅᓕᒃ ᖃᐅᔨᓴᖅᑕᐅᓱᒪᔪᓂᓗ ᑎᑎᕋᖅᓱᒪᑕᖅᑐᑦ ᐊᐅᑐᖅᑕᐅᑕᓂᖅᑎᑦ ᓄᓇᕗᑦ (ᑲᑐ, ᐱᑕᓗᖅᑎᖅᑭᑦᓗ-Carter, et al., 2018; Fᐃᓴᖅ ᐊᓖᓗᓐ ᐃᓗᓐᓇ-Fontaine and Mallory, 2011). ᐃᓱᒪᖅᓱᑕᐅᓗᑎᒃ ᖃᖏᐃᖅᓱᐅᕐᓇᐅᔪᓐᓇᖅᓱᓱᒪᐅᑦ ᑲᓇᑕᒥᓗ ᐊᒥᐊᓕᑲᒥᐅᓗ ᓇᓴᓗᐊᑕ ᕿᑎᑎᖅᓯᓱᐊᖅ ᑲᖏᐃᑦ, ᑕᐃᒪᖓᑦ 1999. ᑲᖏᖅᑕᐅᖅᑲᑕᓂᖅᑦ ᐊᒥᓱᑦᖅᓱᒪᒪᐊᖅᓱᑎᒃ, ᑲᖏᐃᑦ ᐊᒥᓱᓴᖅᓯᓕᓕᐊᖅᓯᑎᑐᓂᖅᓱᑎᒃ ᑲᖏᖅᑕᐅᑲᓂᖅᑦ ᐊᒥᓱᓴᖅᓯᓕᓕᐊᖅᓯᑎᑐᓂᖅᓱᑎᒃ ᑕᒪᐃᐃᑲᓗᐊᖅᑎᓯᒪᔪᑦ, ᑲᖏᐃᖅᑐᓂᖅᓴᐅᓕᐊᖅᓯᑎᑐᓂᖅᓱᑎᒃ ᑲᖏᖅᑕᐅᖅᓯᓕᓕᐊᖅᓯᑎᑐᓂᖅᓱᑎᒃ ᑲᖏᐃᑦ ᑲᖏᐃᖅᑐᓴᖅᓯᓕᓕᐊᖅᓯᑎᑐᓂᖅᓱᑎᒃ ᑲᖏᐃᑦ ᐊᒥᓱ

6.2.3 ᒥᑎᖅᑲᕆᓪ

ᐱᓴᒋᐅᖅᑲᒋᑕᑭᓐᓇᖅᑐᑦ ᑕᒪᐃᓄᐊ ᓄᑦᓗ ᓄᖅᓂᐊᒍᑦᑐᓴᑦ ᐊᑭᓐᓂᐊᖅᓂᑦ ᑎᐹᒋᐅᖅᖁᓂᖅ, ᓇᓄᐃᑦ ᓂᕆᑭᓴᐅᓄᑦᓯᐊᕐᑐᖅᒋᖅᑐᑦ, ᐊᖓᔪᓇᓄᒋᐅᑎᑎᖅᒋᖅᑐᑦ, ᓄᓇ, ᓄᕆᑦᓇᖅᑐᑦᓴᓐᓂ, ᐊᖅᑎᑎᕐᒋᑦ ᓂᕆᐹᖅᓴᖅᑲᐃᐅᓇᓐᓂᕐᓂᓗ ᑲᔪᓄᓪᓕᖅ ᓄᓐᓂᐅᑦᑐᐅᓴᓚᐊᖅᑲᖅᑐᑦ ᖃᐅᔨᓴᖅᒋᐅᖅᖁᓂᖅᓯᓐᓂᖅ ᐊᒻᒪᓗ ᓇᐅᑦᑎᖅᑐᑦᑕᐅᖅᓴᖅᑐᑦᑲᖅᖢᓪᓕᐹᒃ ᒥᑎᓴ. ᑕᒪᐃᓄᐊ ᐱᓴᒋᐅᖅᖁᓂᕐᓂᑦ ᐱᓪᒪᓚᐅᓗᓐᓂᖅ ᐊᐅᓪᓚᑎᓂᑦᒪᑦ ᖃᐅᔨᓴᖅᑲᓂᕐᒋᐅᖅᓴᓂᒪᒋᑦ ᒥᑎᓴ ᐊᒥᓱᓄᖅᓂᓗᓐ ᑕᒪᓐᓇᖅᑐᑦᑕᐅᖅᓴᖅᑐᓂᒪᒋᑦ ᐊᒥᓱᓄᖅᓂᓗᓐ ᑲᑎᒪᓂᖅᑐᑦ. (Dey, et al., 2018; Iverson, 2015; Provencher, et al., 2014). ᐊᖓᕐᓂᖅᓯᑎᐊᓚᐅᐅᓪᓗᑎᒃ ᐊᑕᐅᓱᐅᐅᓪᓂ ᑎᖓᒥᐊᖅᑲᓴᐅᓄᖅᓂᑦ ᑲᓇᑕᐅᑦ ᐅᑭᐅᖅᑕᖅᑐᖅᓯᓱᐅᓄᖅᓯᓂᓄᑦ, ᐊᒋᓄᓕᐊᓪᓗ ᑐᖓᓯᒥᒃ ᐊᑐᔨᓴᖅᑲᖅᓯᓱᒃ ᖃᐅᔨᓴᖅᑲᖅᑐᑦᑕᐅᖅᓴᖅᑐᓂᒪᒋᑦ ᐊᒥᓱᓄᖅᓂᓗᓐ ᑐᖓᓯᒥᒃ ᐊᑐᔨᓴᖅᑲᖅᑐᑦᑕᐅᖅᓴᖅᑐᓂᒪᒋᑦ ᐱᓴᒋᐅᖅᖁᓂᕐᓂᑦ ᐊᒥᓱᓄᖅᓂᓗᓐ ᑐᖓᓯᒥᒃ ᐊᑐᔨᓴᖅᑲᖅᑐᑦᑕᐅᖅᓴᖅᑐᓂᒪᒋᑦ ᐊᒥᓱᓄᖅᓂᓗᓐ ᑲᑎᒪᓂᖅᑐᑦ

ᐱᓕᓀᐊᓐᓇᕐᓂᓐᓄᑦ, ᐅᒪᔨᐃᑦ ᑎᑭᑐᓐᓇᓐᓂᕐᓂ ᐱᑕᖅᓕᑐᓐᓇᓐᓂᕐᓂ ᐊᒥᓱᑐᖅᐸᓕᐊᓂᐊᖅᑐᒃᓴᐅᑦ. ᑲᓇᑕᒥ ᐅᒪᔨᓕᓀᐃᑦ, ᑐᑭᓯᓂᐊᕐᕕᖅᖅᓯᑎᒃ ᐃᓕᓂᐅᕐᐅᓇᕐᓂᓐᓄᑦ ᐊᐅᓪᓚᑎᑦᑐᖅᑕᐅᔨᓂᒃ, ᖃᐅᔨᐃᓕᓐᐅᐸᑎᓐᓄᑦ ᐸᖅᓇᐅᔨᓯᒪᓂᐊᖅᑐᑦ ᓇᐅᑦᑎᖅᑐᖅᓯᒪᓂᐊᖅᑐᑦ ᖃᐅᔨᓴᐅᔨᔪᖅᐸᑦ ᑕᒪᓂᕆᐅᑕᐅᖅᓴᖅᑐᑦ ᑎᑭᑎᐅᔨᓂᖅᓯᒪᓂᐊᖅᑐᑦ.

6.2.6 ᐊᖏᓇᓯᒃᐸᓐᓂᖅ

ᐃᓕᓐᐅᕐᕕᒃᓂ ᐊᐅᓪᓚᑎᑦᑕᐅᔨᖅᑕᐅᔨᔪᑦ ᐃᓕᓚᑦᐅᔨᓂᓐᓄᑦ ᐊᖏᓇᓯᒃᐸᓐᓂᖅ, ᐃᓇᓐᓄᑦ ᓂᕆᐊᖅᑐᓐᓄᑦ ᐊᖏᓇᓯᒃᐸᓐᓂᖅ, ᐃᓇᓐᓄᑦ ᓂᕆᐊᖅᑐᓐᓄᑦ ᐃᓇᒃᐸᔪᑎᑎᓐᓄᑦ, ᓇᓇᒧᓐᓄᑦ ᐃᓕᓐᓂᐊᕐᓂᖅᑐᓐᓄᑦ, ᐃᓇᓐᓄᑦ ᐃᓕᓐᓂᐊᕐᓂᖅᑐᓐᓄᑦ, ᐃᓇᓐᓄᑦ ᐃᓕᓐᓂᐊᕐᓂᖅᑐᓐᓄᑦ ᐊᐅᓪᓚᑎᑦᑐᖅᑕᐅᔨᓂᒃ ᐃᓕᓐᓂᐊᕐᓂᖅᑐᓐᓄᑦ ᐊᐅᓪᓚᑎᑦᑐᖅᑕᐅᔨᓂᒃ ᐊᐅᓪᓚᑎᑦᑐᖅᑕᐅᔨᓂᒃ ᐃᓕᓐᓂᐊᕐᓂᖅᑐᓐᓄᑦ

ᑐᖏᓚᐅᔨᓗᑎᒃ ᓱᓇᓪᓚᓂᖅᓴᓪᑎᑦ ᑕᒪᒃᑯᐊ ᐱᓪᓚᓂᖅᓴᓪᓂᒃ, ᑕᒪᒃᑯᐊ ᐊᖅᑎᖓᓂᖅᖢᓵᕐᒪᖅᑎᑕᐅᔨᑦ ᖃᐅᓪᓕᐊᓂᖅᓯᐅᑦ ᐊᔨᑭᓐᓇᓱᒋᐅᔭᑦ ᐃᓕᓂᒃᑯᑦ (ᓱᓕᓗ ᐅᒪᔨᓪᓚᓂᖅᓯᐅᑦ ᓴᓴᐃᓂᖅᓴᓪᓂᒃ) ᐊᖏᓗ ᐃᓱᒪᒃᖃᓱᐅᑭᑕᐅᓂᔪᓂᐊᖅᑐᑦ ᐱᔨᑦᓇᓱᐊᖅᑐᑦ ᐱᓪᓚᓂᖅᓴᓪᓂᖅᑐᑦ.

6.2.8 ᑕᓚᐅᖅᑎ ᐅᒥᐊᖅᑐᓂᖅ/ᐅᒥᐊᖅᓯᐊᖅᑲᓂᖅ

ᑲᓇᑕᒥ ᐅᓪᓗᓯᓚᓐᓂᒃᑯᑦ ᐃᖅᓂᐅᖅᓴᐅᑯᓕᓪᓕᑦᓗ ᐊᐅᓪᓕᖅᑎᓂᐅᐅᖅᑲᑕᐅᔨᑦ ᐊᐅᓪᓕᖅᑎᓂᐅᐅᒋᑦ ᑕᓚᐅᖅᑎᑦ ᐅᓪᓗᓯᖅᑲᖅᓴᐅᑯᓕᓪᓕᖅᑐᓂᒃᑯᑦ ᐃᑲᑐᑐᖅᑎᑦ ᑎᑎᕋᖅᑲᖅᓴᖓᓕ, ᑐᑭᓯᓂᐊᖅᓴᐅᑯᓕᖅᑲᑕᖅᓂᐊᖅᑐᑦ ᑕᒪᒃᑯᐊ ᑲᑲᒥ, ᑲᑲᑕᐅᑦ ᐅᑭᐅᖅᑕᖅᑐᑐᒍᓕᒃ ᐊᓴᒃᑐᖅᓯᒪᓂᖏᓐᓂᒃ, ᐊᖏᓗᓕ ᓄᓇᕐᓂᓂ ᒪᓕᒪᐅᔨᓂᒃ ᐊᖏᓗᓕ ᐱᓪᓚᓂᖅᓴᐅᓕᖅᑐᓂᒃ, ᐊᖏᓗᓕ ᓄᓇᕐᓂᓂ ᐃᖅᑲᓪᓕᐊᓂᖅᑎᐅᓕᖅᑐᓂᒃ ᐊᖏᓗᓕ, ᐊᖓᔪᓇᓱᑎᑦ >ᐃᐅᓂᒃ. ᑕᒪᓐᓇ ᒥᒃᓪᑎᑭᑕᐅᓂᐊᖅᓯᒪᓕᑦ ᐊᒃᑐᖅᑕᐅᔨᓯᒪᔪᓐᓇᖅᑐᓂᒃᑯᒃ ᑕᓚᐅᖅᑎᐅᑕᐃᑦ ᐅᒪᔨᔪᐃᑦ >ᐃᐅᓕᓪᓕ ᐅᒥᐊᖅᑐᓂᖅᓴᐅᑯᓕᓪᓕᖅᑐᓂᒃ ᐃᒪᖅᓂ ᐃᓗᐊᓂ ᑎᓕᒥᐊᖅᑲᖅᓴᖓᓕᖅᑐᓂᒃ ᐃᒪᖅᓂ ᐊᑭᐊᓂ ᑎᓕᒥᐊᖅᑲᖅᓴᐅᑯᓕᖅᑐᓂᒃ.

ᐊᖅᖁᒍᓕᒪᖅᓱᒃ, ᐅᒥᐊᖅᓯᐊᑦ ᐃᒪᓐᓇᐃᓕᓚᐅᕆᓚᐊᖅᑲᖅᓱᓕᒃᑐᑦ:

(1) ᒪᓕᕈᐊᖅᑲᖅᑐᑦ ᑕᐃᒃᑯᓂᓐᒃ ᐅᑭᐅᖅᑕᖅᑐᑐᒍ ᐅᒥᐊᖅᓯᐊᑦ ᓱᑕᖅᑎᓚᖅᑕᐃᓕᓚᕆᓚᐊᖅᑲᖅᓴᖓᓕᖅᑐᓐᓇᓄᑦ ᒪᓕᑎᓪᓗᑐᖏᑦ ᐊᑎᓯᒃ ᐅᑭᐅᖅᑕᖅᑐᑐᒍᓱᒃ ᐃᒪᖅᓂᑎᖅᑐᓐᓇᓄᑦ ᓱᑕᖅᑎᓚᖅᑕᐃᓕᓚᕆᓚᐊᖅᑲᖅᓴᖓᓕᖅᑐᓐᓇᓄᑦ ᒪᓕᑎᓪᓗᑐᖏᑦ *(1985)*;
(2) ᐅᒥᐊᖅᓯᐊᕐᓂᒃ ᐃᓱᒪᒃᖠᓪᓕᐊᖅᑲᑕᖅᓴᖓᓕᖅᑐᓐᓇᓄᑦ ᐃᒪᓕᑐᑯᖏᓐᓇᓄᑦ ᑯᐃᓱᑕᑕᐃᓕᓚᕆᓚᐊᖅᑲᖅᓴᖓᓕᖅᑐᓐᓇᓄᑦ, ᐊᓱᔪᐅᖅᑕᐅᑕᐃᓕᓚᖅᑕᓐᓄᑐᓂ ᐃᒪᐃᓕᑦ ᐅᒥᐊᖅᓯᐊᕐᓂᒃ ᐊᑕᑦᒎᑦᑕᐅᑕᖅᑐᑦ ᐅᕝᕙᓪᓕᐊᓂᖅᑐᓐᓇᓄᑦ ᐅᕝᕙᓪᓕᐊᓂᖅᑐᓐᓇᓄᑦ ᑯᐃᓱᑕᑕᐃᓕᓚᕆᓚᐊᖅᑲᖅᓴᖓᓕᖅᑐᓐᓇᓄᑦ, ᐊᓱᔪᐅᖅᑕᐅᑕᐃᓕᓚᖅᑕᓐᓄᑐᓂ ᐃᒪᐃᓕᑦ ᐅᒥᐊᖅᓯᐊᕐᓂᒃ ᐊᑕᑦᒎᑦᑕᐅᑕᖅᑐᑦ ᐅᕝᕙᓪᓕᐊᓂᖅᑐᓐᓇᓄᑦ

ᐊᖏᕆᑎᐅᓂᖅᓴᐃᑦ ᐅᒥᐊᕋᐊᑦ ᐅᓗᓚᐊᓇᖅᓄᖅᑐᑦ ᐃᖅᑲᓚᑕᑐᓄᖅᓂᖅ ᐅᕐᓀᓗᓂᒋᑦ ᑭᐃᕋᖅᓄᖅ ᐅᓴᖏᓗᒋᓂᐊᓕᖅ >ᖅᑐᒍᓂᖅ <ᖏᓂ ᖃᐅᔨᖅᓴᐅᖅᑐᑦ. ᐅᒥᐊᕋᐊᑦ ᓂᕈᐊᖏᓄᑦ >ᓕᕆᕆᑎᓄᑦ ᑭᓴᖅᓭᒪᖅᑕᕆᐊᖃᖅᑐᑦ ᓭᓕᑦᑐᖅᓭᒪᓕᖅᖃᑦ, ᑎᑭᓴᖅᐸᓚᓗᑎᒃ ᒥᑭᕋᑎᐅᓂᖅ ᐅᒥᐊᖏᓕᓂᖅ (ᓭᕐᓗ >ᑦᓕᓕᖅᓂᖅ ᓭᑎᓴᖅᓂᖅ).

6.3 ᓇᐅᕆᓂᖅᑐᐃᓴᖅᑐᕐᓂᐊᐅᖅᖃᕆᑕᓂᖅᓭᕆᑦ ᖃᐅᔨᓴᕐᓂᐊᐅᓕᖅᖢᖅᓂᖅᓭᕆᑦ

ᐃᕐᑭᒪᐅᑕᕆᐊᖅᑐᐃᓂᖅ ᐊᑐᑎᓂᖃᕆᑎᐊᖅᑐᐃᓂᓚᑐ ᓇᐅᕆᓂᖅᑐᐃᓴᖅᑐᖃᑕᕐᓂᖅᓭᕆᑦ ᐱᓕᓚᐅᖅᓴᑦ ᑲᑐᖅᐅᓴᖅᓚᐊᕐᓂᖅᖃᓄᑦ ᐊᖅᑐᓂᖅᓭᕆᑦ ᐱᓕᓚᐅᖅᓴᑦ ᐊᐅᓚᑦᑐᖅᓴᐅᖅᑐᑦ, ᐱᓕᓚᐅᖅᓯᐅᓴᑐᓂᒃ ᐃᓕᒪᑐᓂᑦ ᓂᖃᐅᔨᖅᑕᐃᑦ ᐊᓯᖏᑦᓗ ᐅᓴᖅᓭᒪᓕᖅᖢᖅᓂᖅᓭᕆᑦ ᐱᓕᓚᐅᖅᓴᑦ ᐃᓕᓴᕐᓂᐊᐅᓕᖅᖢᖅᓂᖅᓭᕆᑦ ᑕᕐᓴᑦᓯᐅᖅᓴᑦ ᐱᓕᓚᐅᖅᓴᑦ ᐊᒥᓱᓕᕆᓂᖅᓭᕆᑦ ᐃᓕᓴᕐᓂᐊᐅᓕᖅᖢᖅᓂᖅᓭᕆ

ᑲᓇᑕᒥ ᐅᒪᔨᓕᕆᔨᖏᑦ ᐃᓕᓴᕐᐳᑎᑦᑕᐅᔨᓪᓗ ᐊᐅᓚᑦᑎᔨᐅᑉᑲᒪᐅᔨᓕᕆᓂᖅ ᑲᑎᒪᔨᖏᑦ ᖃᐅᔨᔪᖃᑦᑕᓕᐊᖅᑐᑦ ᑐᓴᕐᕋᐅᓴᖅᑕᐅᓴᒪᔨᓂᖅ ᐅᔨᓴᖅᑐᑦᑎᐊᕆᐊᖃᖅᓯᒪᓂᖅ ᒪᓕᓐᓴᒪᑦᑎᐊᕆᐊᖃᖅᖏᓐᓂᒃᓴᖅ ᐊᐅᓚᑦᑎᓂᕐᒧᑦ ᑐᓵᔪᕆᐅᕐᓂᖅᓴᖅ ᐊᖏᒪᓗᐊᖅ ᐱᓕᕆᔨᓴᖅ. ᐃᓕᓴᕐᕋᐅᓴᒥᑦ ᐊᐅᓚᑦᑎᔨᐅᑉᑲᒪᐅᔨᓕᕆᓂᖅᖏᑦ ᐃᑲᔪᖅᓴᐃᓂᒋᖏᑦᑐᑦ ᐱᓕᕆᐊᖃᖅᓯᒪᓂᖅᓴᖅ ᐊᖏᕐᕌᐊᑐᖓᓕᒃ ᐱᔪᕆᐊᖃᖅᑎᓕᖃᖅᑐᑦᑕᒃ ᓇᓪᓕᐊᓂᒃᑯᑦᑐᑦ ᐱᓕᕆᔨᓴᖅᑐᑦ ᐃᓕᓴᕐᓂᖅ, ᐊᒪᓚᑐ ᐱᒪᓚᐅᔨᓴᖅ ᕿᑲᓚᕈᑕᐅᓕᖃᖅᑐᑦᑕᒃᑐᖅ ᐃᓕᓴᕐᓂᖅ ᐅᕙᓪᓕᐊᓂᒃᑯᑦᑐᑦ ᓇᔨᓴᓕᒃᑕᕐᕕᖅᑐᑦ. ᐃᓕᓴᕐᕋᐅᓴᒥᑦ ᐊᐅᓚᑦᑎᔨᐅᑉᑲᒪᐅᔨᓕᕆᓂᖅᖏᑦ ᐃᐅᔨᒪᔨᓕᕆᓂᖅᓴᖅ ᐊᐅᓚᑦᑎᔨᐅᑉᑲᒪᐅᔨᓕᕆᓂᖅᓴᖅ ᐃᓕᓴᕐᓂᖅ ᐅᕙᓪᓕᐊᓂᒃᑯᑦᑐᑦ ᐃᓕᓴᕐᓂᖅᓴᖅ ᐅᕙᓪᓕᐊᓂᒃᑯᑦᑐᑦ ᐅᔨᖅᑕᕐᓂᒃᑯᑦᑐᑦ. ᐃᓕᓴᕐᕋᐅᓴᒥᑦ ᐊᐅᓚᑦᑎᔨᐅᑉᑲᒪᐅᔨᓕᕆᓂᖅᖏᑦ ᐱᔪᕆᐊᖃᖅᓯᒪᓂᖅᓴᖅ ᖃᐅᔨᔪᖃᑦᑕᓕᐊᖅᑐᑦ ᐅᓂᒃᑳᖅᑐᑦᑕᐅᓴᖅᑐᑦ ᐃᓕᓴᕐᓂᖅᓴᖅ ᐊᐅᓚᑦᑎᔨᐅᑉᑲᒪᐅᔨᓕᕆᓂᖅᖏᑦ ᐱᔪᕆᐊᖃᖅᓯᒪᓂᖅᓴᖅ ᐃᓚᐅᕐᓴᐅᔪᖅᑐᑦᑕᐅᓴᖅᑐᑦ

ᐊᐪᓂᐊᖅᑐᒥᓗᓂᑎᒃ ᐱᓴᐁᑎᖅᖅᑐᓂᒃ, ᐅᑐᓗᐊᓇᖅᑐᖅᑭᓇᓂᓂᕆᑎᓪᓗᓯᓂ ᐃᓱᒪᓕᑕᐅᔪᓂᒃ. ᐃᓕᓴᕐᓄᓂᒃᑕᐅᖅ ᐱᓵᖅᐸᓚᕐᓯᓂᒃ <ᕐᓇᐅᑎᖅᑭᓯᓂᑐᓂᒃ ᐱᓕᓚᐊᖅᑭᓇᓂᕆᔨᓪᓗᕐᔪᓂ ᑕᐃᒪᓴᕐᓂ ᐱᕮᕐᐸᖅᑎᓂᑎᖅᑎᓂᖅᑐᓂᖅ ᐃᓇᓴᓂ ᖃᐅᔨᒪᔨᐅᐸᓚᐊᖅᖅᑐᓂᒃ ᖃᐅᔨᑎᓂᑎᖅᑕᓂᓴᕐᓂᓯᓂᒃ ᐊᓚᒪᓗ ᖃᐅᔨᒪᔪᑐᖅᖃᕐᓂᓯᓂᒃ ᐃᓇᓴᓂ ᓄᓇᒪᔪᓂᒃ, ᐅᖅᐅᔨᓯᔨᖅᑕᑭᓇᓗᓀᓗ ᐃᓱᒪᓗᑕᐅᔨᓂᒃ ᓯᔭᕐᓂᐅᑎᑕᐅᔨᓂᖅᑐᓂᓕᕆᔨᒃ ᓇᐃᔅᓯᓗᓐᓂᑐᒃ. ᑕᒪᓐᓇ ᐃᓕᖅᑭᓐᓇᓚᔭᖅ ᓄᓇᖃᓐᓂᒃ ᐃᓇᓐᓂᒃ ᑲᑎᒪᑎᑎᖅᑕᓂᕆᑎᓪᓗᒃ ᐃᓕᓐᓂᐊᑭᑕᐅᔨᖅᑎᑎᓂᐊᖅᑕᓂᓴᓗ ᑲᑎᒪᓂᖅᓂᓂ. ᐃᓕᓂᐅᔨᖅᓴᓂᓂ ᐊᐅᓕᑎᑦᓯᓂᖅᑐᓂᒃ ᑲᑎᒪᓴᕋᓂᖅᑐᓂᒃ ᐃᓕᓐᓂᐊᑭᑕᐅᔨᖅᑎᑎᓂᐊᖅᑕᓂᓴᓗ ᐊᓯᒋᓕᐊᖅᑐᓂᒃ ᐃᓕᓐᓂᐊᑭᑕᐅᔨᖅᑎᑎᓂᐊᖅᑕᓂᓴᓗ

ᐊᓗᒪᓗ ᐃᓄᐃᑦ ᐱᐅᓯᑐᖅᑎᒍᑦ ᐱᑎᓂᖏᓐᓂᒃ ᐊᓗᒪᓗ ᐃᓕᓴᐃᓯᑐᖅᑕᓴᖅᓯᒪᓂᒃ (ᐃᓄᐃᑦ ᐃᑲᔪᓯᐊᖅᓯᒪᓂᐅᓄᑦ ᐊᖏᑕᕆᑦ ᐃᓗᓕᓂᖏᓂ 6.8.4) ᖃᑎᖓᕕᐊᖅᑕᐅᖅᑕᖅᑐᓂᐊᖅᓯᒪᔪᓗ ᑲᓇᑕᒥ ᐅᒪᔪᓕᕆᔨᖏᓐᓄᑦ, ᐅᓪᓴᖅᑐᑎᑎᐊᖅᓂᐊᖅᓯᒪᖓᒪᑕ ᑕᓪᒪᖅᓯᒪᔪᕐᓗᒃᖏᓂᖅᓂᑕ ᐅᒪᔪᓕᕆᔨᓂᒃ ᐅᒪᔪᓕᕆᔨᓂᒃ ᓄᓇᕗᑦᐅᓂᒃᖏᓂᒃᓂᑕ ᖃᐅᔨᓴᒪᑎᑎᑕᐅᑦ, ᐱᑕᖅᖃᖅᑕᖅᓂᖏᓂ ᑐᑭᓯᐅᑦᑕᐅᔪᓂᒃ ᓴᓇᔨᐅᓯᒪᓂᖏᓐᓂᒃ (ᐃᓄᐃᑦ ᐃᑲᔪᓯᐊᖅᓯᒪᓂᐅᓄᑦ ᐊᖏᑕᕆᑦ ᐃᓗᓕᓂᖏᓂ 6.8.5).

ᑲᓇᑕᒥ ᐅᒪᔪᓕᕆᓴᖅᔪᑦ ᑕᔪᐊᖅᓴᓂᓕᒫᓂᒃ ᖃᐅᔨᓴᒪᔪᓂᕐᑎᑎᐊᕈᑎᓂᓂ ᖃᖅᖄᓴᐅᖅᑐᑦᑐᒃ, ᐊᑐᖅᑕᕐᓂᐊᖅᓂᑦᑐᐊᖅᓯᒪᓂᖅ ᓂᐅᒥᐊᖅᑎᓂᓂᒃ ᐊᑐᖅᑕᕐᓂᐊᖅᓯᒪᓂᖅ ᐃᓕᓴᐃᓯᑐᖅᑕᖅᑐᓂᒃ ᑕᓪᓕᖅᑐᓂᒃ ᐊᑐᖅᑕᕐᓂᐊᖅᓯᒪᓂᖅ ᐃᓕᓴᐃᓯᑐᖅ

6.4.4 ᓂᐅᕐᑕᐊᕐᓂᐅᖅ ᐊᒥᓱᐊᕐᖁᕐᓂᖅ

ᒪᓕᖅᑕᐅᔪᓗᑎᒃ ᑕᐃᒃᑯᐊ ᐃᓄᐃᐃᓐ ᐃᑲᔪᐊᕐᓇᖅᑕᑭᑎᑎᒃᓴᖅᓯᓄᓐ ᐊᖏᕐᕕᑦᓂ, ᑐᖓᓕᒃᑯᓐ ᐊᐅᓪᓚᕆᔨᐅᒋᑐᑦ ᑭᓇᐅᔨᓂᐊᑎᕐᑎᐅᔪᓂᖅᑐᒃᓴᓂᒃ ᐊᔪᕐᑭᓐᓂᖅᓴᐃᔨᐊᑎᒃᓴᓂᒃ ᐃᓄᖅᓂᒃ ᓴᓪᓕᓂ ᓂᐅᕐᑕᐅᔪᓗᑎᒃ ᐱᓪᓚᔨᐅᔪᓯᓐᓇᖅᓂᖅ ᐊᓚᒪᓗ ᐊᑐᑎᖅᑎᐊᕐᑭᓯᓇᖅᑐᓂᒃ ᓄᓇᓕᖅᓂᒃ ᑐᖓᓕᒃᑯᓐ ᐊᔪᕐᑭᓐᓂᖅᓴᐃᔨᐊᑎᒃᓴᓂᒃ. ᑕᒪᒃᑯᐊ ᐱᔪᓂᑎᕆᔨᑦᓴᓂ ᐃᓕᓴᖅᓂᐊᖅᑐᑦᓂ ᐊᔪᕐᑭᓐᓂᖅᓴᐃᔨᐊᑎᒃᓴᓂᒃ, ᐃᓗᓕᒥᖅᓴᑦᑕᐅᔪᓯᓐᓇᕐᓂᓕᕆᓂᒃ ᐃᓕᐅᔨᓪᓚᕆᔨᐅᒋᑐᑦ ᐱᔪᓂᑎᕆᐃᑦᓴᓂᒃ ᐅᓂᒃᑯᐅᑦᓂᒃ; ᐱᓕᕐᓂᐊᑎᓂᑎᓗᑎᒃ ᓄᓇᓕᖅᓂᒃ ᐱᔪᓂᑎᕆᔨᐅᒋᑐᑦᓂᒃᑐᓂᒃ ᑎᑎᕋᐅᓂᖅᓴᐃᔨᐊᑎᒃᓴᓂᒃ ᑎᑭᓕᓚᐅᖅᓴᓂᒃ; ᐊᓚᒪᓗ ᑎᑎᕆᔨᑦᓴᓂᒃ ᐊᔪᕐᑭᓐᓂᖅᓴᐃᔨᐊᑎᒃᓴᓂᒃ ᐊᖏᕐᕕᑦᓂ ᐃᓄᖅ

ᑎᑭᓯᒪᔨᖅ ᓄᓇᕗᕋᓪᓴᐊᖏᒪᔪᓂ ᑕᖅᓯᖏᐅᐅᖅᓂᔭᖅ, ᐊᐅᓚᕋᑎᔨᐅᖏᕆᑕᐅᓂᔭᖅ ᑲᑎᒪᔨᕋᓪᒐ ᐅᖃᐅᖅᔨᐅᖅᓂᐊᕐᓂᖅ ᑎᑭᓯᒪᔨᐸᑉ >ᓚᕋᕋᑎᑦᑲ ᓇᒪᒋᔭᐅᖃᕐᓂᖅᓱᖅᓂᖅᓄᑦ ᐅᔨᒪᔨᖅᓚᓚᑦᓴᓂ ᐊᖏᕐᖅᑕᐅᓂᐊᕐᓯᒪᖅᓯᒪᓂ ᖃᐅᔨᔾᔪᐊᕐᓴᑎᑦ ᐊᑐᖅᑕᐅᔨᓐᓄᑦ ᐊᔾᔨᑎᑎᖅᓂ (ᐃᓄᐃᑦ ᐃᑲᔪᓯᐊᓕᖅᖢᓂᓐᓄᑦ ᐊᖏᕐᕕᒃᓴᓂ 5.5.5).

ᓄᒌᓂᑲ ᓄᓇᒦᖏᐅᔨᖅᑐᖅᑕᖅᓯᐸᑲᓂᑲ ᑕᖓᒪᔨᓴᓂ ᑕᐅᖅᓂ ᓄᓇᖅᓯᓂ ᑎᖅᒥᐊᖅᑎᖏᐅᑦ ᐱᔨᓐᓇᔭᓂᖅᑎᖅᑎᑕᐅᓯᒪᕐᒉᒃᓵᓂ ᐊᑐᖅᑕᐅᔨᓐᓇᔭᓂᖅ ᐊᖏᕐᕕᒃᓴᓂᐊᑎᑦ ᐊᖏᖅᑕᐅᓯᒪᔪᑎᑎᓂᑦ ᐊᖏᕐᕕᒃᓴᓂᐊᑎᑦ ᐊᖏᕐᕕᒃᓴᓂᐊᑎᑦᑲ ᑲᑎᒪᔨᖅᑎᖏᐅᑦ ᐱᔨᓐᓇᔭᓂᖅᑎᖅᑎᑕᐅᓯᒪᕐᒉᒃᓵᓂ ᓄᓇᖅᓯᓂᑦᑎᖅᑎᑕᐅᓯᒪᕐᒉᒃᓵᓂ ᐊᖏᕐᕕᒃᓴᓂᐊᑎᑦ ᐊᖏᕐᕕᒃᓴᓂᐊᑎᑦᑲ ᑲᑎᒪᔨᖅᑎᖏᐅᑦ ᐱᔨᓐᓇᔭᓂᖅ

6.4.7 ᒪᓕᑕᕐᓂᐊᓕᒃ ᒪᓕᒃᑕᐅᑲᕐᓂᖅᑕᕐᓂᖅᓯᒪᓕᒃ

ᒪᓕᑕᕐᓂᒃ, ᒪᓕᑕᓪᓗᑐᓂᒃ, ᐊᖏᑭᕈᑎᓂᒃᓗ ᐱᓱᐊᑎᖃᖅᑐᑎᒃ ᖃᓄᖅᓂᐊᖅᑕᐅᓯᒪᖏᖅᓂᒃ ᓄᓇᕗᒻᒥ ᒪᓕᒃᑕᐅᕐᓂᔪᓐᓇᖅᓂᖅᖃᖅ ᐱᓘᒪᓕᐊᔪᕗᖅ ᐊᐅᓕᕐᑎᓂᕐᒧᑦ ᑕᒪᒃᑯᓂᖅᓯᒪᔪᖅᑕᐅᓯᒪᓕᐅᕐᓂᒃ. ᑲᓇᑕᒥ ᐅᐃᒪᓱᓕᓚᐅᖅᑯᑦ ᐊᓘᒪᓗ ᐊᐅᓕᕐᑎᐅᕐᖃᖅᑕᐅᓂᕐᒧᑦ ᑲᑎᒪᔨᖏᑦᑑᒃ ᒪᓕᒃᑕᐅᑎᕐᑎᓂᕐᓯᒪᔪᖅᑕᕐᒥ ᒪᓕᑕᓂᒃ ᖃᐅᔨᒪᑎᕐᑎᖃᖅᑕᕐᒥᓂᒃ ᐃᓇᖅᓂᒃ ᖃᐅᔨᒪᔭᐅᓚᖅᑐᓂᒃ, ᐃᓕᓴᓂᐊᖅᓯᒪᓕᑕᐅᓪᓗᓂᒃ ᓄᓇᕗᒻᒥ ᒪᓕᒃᑕᐅᑎᕐᑎᓂᖅᓯᒪᓕᖅᓂᖅᓂᒃ ᐃᓇᖅᓂ, ᒪᓕᑕᓪᓗᑐᓂᒃ ᐊᓘᒪᓗ ᐊᑐᐊᑎᓪᓗᑲᑦᑐᖅᓯᒪᓕᐅᖅᓯᒪᓕᒃ ᑕᐅᖅᓂᖅ, ᐊᓘᒪᓗ ᑐᖏᑎᓪᓗᑲᑦᑐᖅᓯᒪᓕᐅᖅᓯᒪᓕᒃ ᐱᓕᕐᓂᐊᖅᑕᐅᓯᒪᓕᐅᖅᓯᒪᓕᒃ ᐊᓯᐊᓂᓪᓗᑲᑦᑐᖅᓯᒪᓕᐅᖅᓯᒪᓕᒃ ᑕᐅᖅᓂᖅ ᓇᓗᓇᖅ.

ᐊᐅᓕᕐᑎᓂᕐᒧᑦ ᖃᓄᐃᓕᓕᐅᕈᑎᒃᓴᐃᓕᑦ ᒪᓕᒃᑕᐅᑎᕐᑎᓂᕐᓂᖅ ᒪᓕᑎᕐᓂᒃ ᐃᓄᒃᑐᐊᖅᑎᖓ ᑎᒃᓚᕐᑎᖅᓴᖏᓐᓄᑦ ᒪᓕᒐᓂᖅ ᐊᓕᖅᑐᐊᖅᑎᖓ:

- ᐊᑐᐃᓐᓇᐅᑎᑕᐅᖅᑲᒃᑕᕐᓗᑎᒃ ᑕᒃᔪᐊᐊ ᐊᐅᓕᕐᑎᑦᓴᕐᑕᐅᓂᕐᒧᑦ <ᕐᓇᐅᑎᖏᑦ;
- ᐊᑐᓕᖅᑎᑕᐅᓂᖏᓐᓂᒃ ᑐᓴᐅᒪᓕᖅᑲᑕᐅᑎᓂᕐᒧᑦ <ᕐᓇᐅᑎᓕᐊᓕᓚᖏᑦ ᓴᓇᔨᐅᓪᓚᔪᓂᖏᑦ, ᒪᓕᒃᓪᓗᑎᒃ ᑐᑭᓯᓂᐊᕐᓇᐅᓪᓚᔪᓂᖏᓐᓂᒃ ᓴᖅᓕᕐᒥᐅᑦ;
- ᐃᓕᓴᕐᖃᐃᓪᓚᓕᓗᑎᒃ ᑎᑎᕐᖃᖅᖃᖅᑐᓂᒃ ᖃᐅᔨᒪᑎᑦᑎᐊᕐᑕᐅᒃ ᐱᓘᒪᓕᐅᓂᖏᓐᓂᒃ ᑎᑭᓴᕐᓇᐅᑦᓯᒪᔪᖅᑕᐅᓂᖏᓐᓂᒃ ᑎᖏᕐᐊᓂᖅᓇᖅᑐᓂᖏᓐᓂᒃ;
- ᐅᔾᔨᖅᑐᑦᑎᐊᕐᓂᖅ ᖃᐅᔨᒪᑎᑦᑎᐊᕐᑕᐃᑦ ᐊᑐᐃᓐᓇᐅᓂᖏᓐᓂᒃ ᐃᓄᖅᑲᕐᓇᐅᑦᓯᒪᔪᖅᑕᐅᓂᖏᓐᓂᒃ ᑎᑭᓴᕐᓇᐅᑦᓯᒪᔪᖅᑕᐅᓂᖏᓐᓂᒃ ᐃᓚᒋᑭᓐᓇᓂᖏᓐᓂᓐᓂᒃ ᐃᓄᖅᑐᐊᐸ;
- ᐅᔾᔨᖅᑐᑦᑎᐊᕐᓂᖅ ᑎᑎᕐᓯᖅᑎᑦ ᖃᐅᔨᒪᓂᖏᓐᓂᒃ ᐱᓕᕐᓂᐊᖅᑕᐅᓯᒪᔪᖅᑕᐅᓂᖏᓐᓂᒃ ᐃᓗᐊᓂᓐᓂᑕᐊᕐᑕᐅᒃ ᐃᓄᖅᑐᐊᐸ ᑎᑎᕐᑎᖅᓴᖏᓐᓄᑦ;
- ᐅᔾᔨᖅᑐᑦᑎᐊᕐᓂᖅ ᑭᓇᐅᔭᖅᑎᑦ ᖃᐅᔨᒪᓂᖏᓐᓂᒃ ᐱᓕᕐᓂᐊᖅᑕᐅᓯᒪᔪᖅᑕᐅᓂᖏᓐᓂᒃ ᐃᓗᐊᓂᓐᓂᑕᐊᕐᑕᐅᒃ ᐃᓄᖅᑐᐊᐸ ᑎᑎᕐᑎᖅᓴᖏᓐᓄᑦ;
- ᐅᔾᔨᖅᑐᑦᑎᐊᕐᓂᖅ ᑭᓇᐅᔭᖅᑎᑦ ᐃᓗᐊᓂᑦ, ᓇᖏᑦᓂᓪᓗ ᐱᓕᕐᓗᐊᓂᑦ, ᖃᐅᔨᒪᔭᕆᐊᖃᖅᓯᒪᔪᖅᑕᐅᓂᖏᓐᓂᒃ ᐱᓕᕐᓂᐊᖅᑕᐅᓯᒪᔪᖅᑕᐅᓂᖏᓐᓂᒃ ᐃᓕᖏᓂᒃ ᐱᓕᕐᓗᓇᐅᓂᕐᒧᑦ ᑎᖏᕐᐊᓂᖅᓇᖅᑐᓂᖏᓐᓂᒃ ᐃᓄᖅᑐᐊᐸ ᑎᖏᕐᐊᓂᖅᓇᖅᑐᓂᖏᓐᓂᒃ;
- ᐱᓱᒋᐊᓂᐃᓂᖅ ᑭᓇᐅᔭᖅᑎᑕᐅᓂᒃ ᐃᓄᖅᑕᐅᓚᐅᖅᑎᑦᓯᒪᔪᖅᑕᐅᓂᖏᓐᓂᒃ ᒪᓕᒐᓂᒃ ᓯᕆᐃᐊᓂᒃ ᑎᖏᕐᐊᓂᖅᓇᖅᑐᓂᖏᓐᓂᒃ ᐃᓕᓴᕐᓇᐅᑦᓯᒪᔪᖅᑕᐅᓂᖏᓐᓂᒃ ᐅᖅᑯᓂᖅ ᐃᓕᓴᕐᓇᐅᑦᓯᒪᔪᖅᑕᐅᓂᖏᓐᓂᒃ ᐃᓄᖅᑐᐊᐸ ᑎᖏᕐᐊᓂᖅᓇᖅᑐᓂᖏᓐᓂᒃ ᑎᑎᕐᑎᖅᓴᖏᓐᓄᑦ

6.4.8 ᐊᕐᓇᖅᑐᓂᒃᑯᓐᓂᒃ ᖃᓄᐃᓕᓚᐅᕐᓂᐊᕐᓂᖅ ᐱᑎᐊᖅᑎᑕᐅᓯᒪᓕᕐᓂᖅᓗᓪ

ᓯᓚᑎᒥᐅᑕᕆᔨᐅᔪᓂᒃ ᐃᒃᐱᒋᔨᐅᖃᖅᑕᕐᕋᔭᓂᖅ ᐃᓯᒪᒋᔨᐅᓗᓂᒃᑯᓐᓂᕐᓂᖅᖢᐊᐅᓚᓗᐊᖅᓯᑎᒃ ᑕᒃᑯᓄᓂᖅᓗᒍ ᐊᐅᓚᕐᑎᓂᖅᓗᒍ ᐃᓕᓴᐅᒪᓂᖅ ᐊᐅᓚᕐᑎᓯᐅᖃᖅᑎᑦᓗᓂ ᐃᖅᑲᓇᐃᓐᓂᒃᑯᓐᓂᕐᓂᖅᓗᒍ ᐊᐅᓚᕐᑎᓯᐅᖃᖅᑎᑦᓗᓂ ᑎᒥᒋᐊᖃᖅᑎᖅᑐᖅ ᐊᐅᓚᕐᑎᓯᐅᖃᖅᑎᑦᓗᓂ ᑎᒥᒋᐊᖃᖅᑎᖅᑐᖅ ᐊᓯᖏᓐᓂᒃᓗᓪ ᐅᒪᓕᔨᓚᓂᕐᒥᖅ.

ᐃᓕᓴᐅᕐᓇᒃᑯᓂ ᐊᐅᓚᕐᑎᓯᐅᖃᖅᑕᐅᓂᕐᓗᓂ ᑲᑎᒪᔨᕆᓇᖅᑐᓂ ᐃᓚᐅᕐᖃᖅᑕᕐᐃᓐᓇᕐᓂᖅᓗᓂ, ᐃᓚᐅᕐᓴᐅᓕᔨᓯᑎᓪᓗᓐᓂᐅᓂ ᐃᓗᓚᒋᓂ ᐃᒪᓐᓇᖅᑐᖅ. ᑕᒪᐃᑯᐊ ᒪᑯᓂᖓᓕᕐᔪᓐᓇᖅ ᐃᓚᖃᖅᑕᕐᐃᓐᓇᕐᓂᖅᓗᒍ. ᑕᒪᐃᑯᐊ ᒪᑯᓂᖓᓕᕐᔪᓐᓇᖅ ᐃᓚᖃᖅᑕᕐᐃᓐᓇᕐᓂᖅᓗᒍ:

- ᖃᐅᒥᖅᑕᐊᖅᑕᐅᓂᕐᓂᐅᓂᒃ ᐱᓚᓗᐊᖅᓴᓕᐅᔪᓪᓚᕐᓂ, ᐅᖅᑲᐅᔨᐅᔨᓪᓚᕐᓂ ᑕᐃᒃᑯᓇᓐ ᓄᓇᕗᑦ ᐅᓇᓕᕐᓂᖅᓗᒍ ᐊᐅᓚᕐᑎᓯᐅᖃᖅᑎᑦᓗᓂ ᓴᓇᓕᐅᔪᓪᓚᕐᓂᓗ ᖃᐅᔨᓴᓴᖃᓕᕐᑎᖅᑎᑦᓗᓂ ᒪᓚᑦᓗᓂ, ᐊᐅᖅᑕᐊᖅᑕᐅᔨᓂᐊᕐᑎᐅᒃ ᐅᒪᓚᔪᐃᓐ ᐅᒪᓚᔪᐃᓐ ᐊᐅᓚᕐᑎᓯᐅᖃᖅᑎᑦᓗᓂ ᓄᓇᕗᑦ ᐊᐅᓚᕐᑎᓯᐅᖃᖅᑎᑦᓗᓂ ᓄᓇᕗᑦ ᐊᐅᓚᕐᑎᓯᐅᖃᖅᑎᑦᓗᓂ ᓄᓇᕗᑦ *ᐊᐅᓚᕐᑎᓯᐅᖃᖅᑎᑦᓗᓂ ᓄᓇᕗᑦ*; ᐊᓚᓗᓪ;
- ᓄᓇᕗᒻᒥ ᓄᓇᐃᑦ ᐊᑐᑎᐅᓴᖏᓐᓂᒃ <ᖁᓇᐅᑎᑕᓚᓂᕐᓗᓂ ᐊᓚᓗᓪ ᓯᕗᓂᐅᓴᒥ ᐋᑭᑭᑎᐊᖅᑕᐅᔨᓴᓇᕐᓂᖅᓂᐅᓂᒃ ᐱᔭᔪᑎᖅᑲᖅᑐᖅ.

ᐊᓯᖏᓐᓂᒃᑯᐅᑕᐅᖅ ᐱᑎᐊᖅᑎᑕᐅᓯᒪᔨᓴᓇᕐᓂᖅᓗᒍ ᐊᓚᓗᓪ ᐱᐃᓐᓴᐅᖅᑲᐅᔨᑕᓴᓇᕐᓂᖅᓗᒍ ᐃᓭᓂᒃᑯᓂ ᓴᖅᑭᑦᑐᖅᓗᒍ ᐃᓯᓂᒃᑯᓂ ᐊᓯᖏᓐᓂᒃᑯᓐᓂᕐᓂᖅᓗᒍ

7 ᐊᖏᑭᑕᐅᓯᒪᔨᕆᐊᓂᖅ ᐱᓕᕆᐊᒃᓴᐅᑦ, ᐊᔪᖅᓯᖅᑎᑕᐅᔪᑦ ᐱᓕᕆᐊᕐᐳᐅᔪᖅᓯᓂᖏᑦ ᐊᒻᒪᓗ ᐊᔪᖅᑎᑕᐅᔪᑦ ᑎᑭᓯᒪᕐᐳᐅᔪᖅᓯᖏᓐᓄᑦ

ᑲᑎᒪᐅᑕᐅᖅᑕᑐᓂ ᑎᖏᑎᐊᓕᕆᓂᖏᑦ ᒪᓕᑐᓕᐊᕆᔨᐅᖅᑕᖅᑐᑦ ᒪᓕᑐᐅᑭᑕᐅᖅᓯᒪᓕ ᑐᖏᓪᓗᐊᐅᓂᖏᑦ ᓴᓇᓴᖅᑐᑦ ᐱᓇᓱᐊᒋᒪᖅ, ᐅᑦᓗᓂᖅᓴᖅᖢᓗ, ᒪᐃᓂᖏᓐᓂᓗ, ᓯᔪᖅᑎᖅᑕᐅᔪᖅᑐᕆᐊᖅᑕᖅᓯᖅᑎᓂᖅ, ᐸᓴᓴᖅᑕᐅᔪᖅᑐᕆᐊᖅᑕᖅᓯᖅᑎᓂᖅ, ᐊᒻᒪᓗ ᐃᓂᖅᑕᐅᔪᖅᑐᕆᐊᖅᑕᖅᓯᖅᑎᓂᖅ. ᑕᐃᒃᑯᐊᓗ ᑲᑎᒪᒪᐅᑕᐅᖅᑕᖅᓂᑦ ᑎᖏᑎᐊᓕᕆᓂᖏᑦ ᒪᓕᑐᓚᐊᕆᔨᐅᖅᓯᒪᔪᖅᑐᖅᑐᑦ ᒪᓕᑐᓴᓪᓗᕆᔨᐅᖅᑕᖅᑐᑦ ᑐᑭᖅᑕᐅᓂᖅᓯᒪᓂᖅᑐᓂ ᐊᑐᖅᑕᐅᔪᖅᑐᕆᐊᖅᑕᖅᑐᑦ ᐊᕐᓇᖅᑐᖅᑐᑦ ᒪᓕᑐᓴᓪᓗᕆᔨᐅᖅᑕᖅᑐᑦ ᑲᑎᒪᒪᐅᑕᐅᖅᓯᒪᓂᖅᑐᑦ ᑲᑎᒪᒪᐅᑕᐅᖅᑕᖅᓂᑦ ᒪᓕᑐᓴᓪᓗᕆᔨᐅᖅᑕᖅᑐᑦ ᑲᑎᒪᒪᐅᑕᐅᖅᑕᖅᓂᑦ ᐊᔪᖅᑎᖅᑕᐅᔪᖅᑐᕆᐊᖅᑕᖅᑐᑦ ᐊᒻᒪᓗ ᑐᖏᓪᓗᐊᐅᓂᖏᑦ ᒪᓕᑐᓴᓪᓗᕆᔨᐅ

ᑎᖏᐊᖅᓴᖅᓄᓂ ᒪᓕᒐᓪᓗᑕᓗᐅᔪᓂ ᒪᓂᒋᓂ ᐊᒡᓂ ᑎᑎᕋᖅᓯᒪᔪᓂᒃ ᐱᔪᓐᓇᖅᑎᕋᖅᑕᒥᓪᓚᕐᓂᑦ, ᑭᓯᐊᓂ ᐊᖏᕐᕋᐅᔪᓯᒪᓪᓗᓐᓄᑦᑐᓂᒃ ᐱᓕᕆᐊᖓᔪᓇᖅᑐᖅᓂᒃ ᐱᔪᓐᓇᐅᑎᖅᕋᓗᑎᒃ (ᖃᐅᔨᒪᓕᐅᓂᐊᖅᓴᖅ ᐃᓄᐃᑦ ᐊᖓᔪᖅᓯᓕᕆᓂᖅᓯᓂᖅ ᐊᐅᓚᓯᓂ ᐃᓐᖏᕋᐅᔪᓯᒪᓕᖅᓯᓂ ᒡᒃᑯᓕᐅᔪᓯᒪᓕᕐᓂᑐᑦ ᑲᒪᔨᓄᖅ, ᐅᖃᐅᔨᔪᐅᖏᓂᒃᔪᓯᒪᓕᒐ ᐱᔪᓐᓇᐅᑎᑎᖃᕋᓂᖅᓯᓂᖅ ᖃᐅᓪᓂᓂ);

- ᑭᓇᐃᔪᓇᖅ ᑎᖏᐊᖅᔪᐅᑭᓇᖅᓯᐃᖅ.
- ᑭᓇᐃᔪᓇᖅ ᔪᕆᖅᑎᓚᓂᐊᖅᓯᒪᓕᖅ ᑎᐃᔪᖅᑲᕈᑕᓚᐊᖃᖅᕆᓂᓗᓄᑦᓂ ᑎᖏᐊᑦ ᐅᑦᓗᖅᓯᓂᒃ.
- ᑭᓇᐃᔪᓇᖅ <<ᕆᑎᓂᖅᓂᐊᖅᓯᒪᓕᖅ ᐅᒪᔪᒥᒃ ᔪᓕ ᑎᖏᐊᖅᓯᒃ, ᑎᒥᕕᓂᖅᓄᑦᓗᓄᑦᓂ, ᐊᒥᖅᓄᑦᓗᓄᑦᓂ, ᒪᓂᖅᓄᑦᓗᓄᑦᓂ ᑎᖏᐊᑦ.
- ᑭᓇᐃᔪᓇᖅ ᖃᑭᐅᔪᔨᖅᐸᖅᓂᐊᖅᓯᒪᓕᖅ ᐊᖓᔪᓇᔪᓂᓕᒫᓂᑦ ᐊᑐᑎᓕᖅᑎᒃ ᐱᖃᑎᓂᖅ ᐱᔪᒪᓕᖅᓯᒪᓕᖅ
- ᑭᓇᐃᔪᓇᖅ ᑎᔪᐊᓇᔪᒪᔨᒥᓂᖅ ᐅᒃᑎᒥ ᖃᑭᒥᖃᒥᓂᓗᓄᑦᓂ ᐃᐊᖅᑐᕆᑎᖅᓄᑦᑎᓂᓗᓄᑦᓂᖅ ᓇᒧᑐᐃᓇᐅᖅᑐᒥᒃ
- ᑭᓇᐃᔪᓇᖅ ᐱᓕᕆᓂᖅᓂᐊᖅᓯᒪᓕᖅ ᔪᓂᑯᑎᓇᖅᑐᒥᒃ ᑎᖏᐊᖃᖅᓂᒃ, ᒪᓂᖅᓄᑦᓂᓗᓄᑦᓂ, ᐅᔨᒪᓇᖅᑕᖅᓄᑦᓂᓗᓄᑦᓂ ᑎᖏᐊᑦ ᓄᓇᓇᖅ.

7.1 ᐊᐅᓚᖃᑎᒌᓄᑦ ᐊᖏᕐᕋᐅᔪᓯᒪᔭᕆᐊᓂᑦ

7.1.1 ᐊᖏᕐᕋᐅᖃᕋᐅᖏᓂᑦ ᐱᔪᓐᓇᐅᑎᑐᑦ

ᐊᑐᕋᐅᐅᓂᖏᓐᓂ *ᑎᖏᐊᖅᓴᖅᓄᓂ ᒪᓕᒐᓪᓗᑕᓗᐊᓂᑦ* ᐊᒻᒪᓗ ᐅᐸᑐᕋᖅᑐᖅᓴᓐᓂᑦ, ᒥᓂᔅᑕᐅᔪᕋᖅ ᐊᐅᑎᓕᓚᓂᖅᓴᓗᓪᓚᕐᓂ ᐊᖏᕐᕈᓐᓇᖅᓴᖅ ᐊᔪᖃᓐᓇᖅᑎᑕᐅᔪᓂᖅ ᐱᓕᕆᐊᖓᔪᓇᖅᑐᖅᓂᒃ ᐊᑐᖅᓗᓂᒥᓂ *ᑎᖏᐊᖅᓴᖅᓄᓂ ᒪᓕᒐᓪᓗᑕᓗᐊᓂᑦ* ᑐᓂᓯᔪᓐᓇᖅᓯᓂᒃ. ᑕᒪᓐᓇ ᐱᓕᕆᓂᖅᓗᓂᑦ ᑐᑭᖓᒪᔪᖅᓴᖅ ᐊᖓᔪᒪᔨᓚᐊᖃᖅᑎᖅᓴᖅ ᐊᑕᐅᔪᓯᖅᑎᒃ ᑕᒪᐦᑭᔪᓂᓕᒫᓂᑦ ᐊᑐᑎᖅᓄᑦᓂᐊᖅᓯᒪᓕᖅᑐᒃ ᐊᒻᒪᓗ ᖃᐅᔨᓴᖅᑎᑕᐅᔪᓂᖅᓯᒪᓕᖅᑐᒃ ᒪᓕᒐᓪᓗᑕᓗᐊᖅᓯᒪᓕᖅᑐᒃ ᐊᐅᑎᓕᓚᓂᔨᓴᓕᐊᑦᓂᖅᓯᒪᓕᖅᑐᒃ ᐊᐅᑎᓕᓚᓂᐊᖅᓯᒪᓕᖅᑐᒃ ᐊᑐᖅᓗᓂᒥᓂ ᐃᓕᓴᕆᔨᖃᖅᑎᑕᐅᔪᓂᖅᓯᒪᓕᖅᑐᒃ ᐱᓕᕆᐊᖓᔪᓇᖅᑐᖅᓂᒃ ᐊᐅᑎᓕᓚᓂᐊᖅᓯᒪᓕᖅᑐᒃ ᐊᐅᑎᓕᓚᓂᐊᖅᓯᒪᓕᖅᑐᒃ ᐊᑐᖅᓗᓂᒥᓂ ᐃᓕᓴᕆᔨᖃᖅᑎᑕᐅᔪᓂ

- ᐊᓯᖏᐃᓕᓐᓂᒃ ᐱᓕᕆᐊᖓᔪᐊᓘᓐᓄᑦᓂᒃ ᐃᓱᒪᓕᖅᓯᐅᑎᓕᐅᕐᓂᒃ ᐃᓱᒪᓕᖅᓯᐅᑎᖑᓕᕐᓂᖅ; ᐊᒻᒪᓗ
- ᐊᐅᑐᖅᑕᐅᓗᐊᓵᔪᒥᑎᓕᖅᓴᐃᑦ ᐃᓱᒪᓕᖅᓯᐅᑎᕕᓱᒪᔨᑎᒃ ᐊᒻᒪᓗ ᐊᖏᕈᑕᐅᓱᒪᔨᑎᒃ.

ᒪᓕᑐᖃᖅᑎᑕᐅᓂᖏᓐᓂ ᖃᓄᐃᓕᓐᓄᑦᓴᓚᐊᕐᖃᖅᖃᖅᑎᑕᐅᓂᖏᓐᓂᓪᓗ ᐊᐅᓪᓚᕆᑎᓂᖏᓐᓂᒃ ᐱᓕᕆᐊᓚᐅᓂᖏᓐᓂᒃ, ᒥᓂᔾᑕᐅᐸ ᐃᓱᒪᓕᖅᓯᐅᑎᖑᓕᕐᓂᖅᑕᐅᓚᐊᕐᖃᖅᖃᖅᑐᖅ ᓴᓐᓄᓂᐅᓂᖏᓐᓂᒃ ᐊᐅᑐᖅᑕᐅᓗᐊᓵᔪᒥᑎᖑᓕᕐᓂᖅᓄᑦᓗ ᑎᓐᒥᐊᑦ, ᓇᔨᖅᐸᒃᑕᖏᓪᓗ ᓄᓇᖏᓪᓕ, ᑕᒪᒃᑯᐊᓗ ᐃᓕᓴᐅᓱᒪᔨᓇᓐᓇᕐᓂᒃ ᐱᔨᓇᐅᑎᑑᓚᐅᓂᖏᓐᓂᒃᓄᑦᓗ ᒪᓕᑐᖃᖅᑎᑕᐅᓗᐊᓵᔪᒥᑎᓕᖅᓴᐃᑦ. ᑐᐱᓯᐊᖅᑐᑎᓕᐅᓂ ᐅᖃᐅᓯᐅᓯᒪᔨᑦ ᑎᑎᕋᖅᓯᒪᔨᑦ 7.1.7-ᖓᓂᓗ ᑐᐱᓯᐊᖅᑎᐊᕆᐅᓂᕐᒪᔪᐃᓐᓇᑦ ᐱᔨᓇᐅᑎᑦ ᐊᑐᖅᑕᐅᔾᔨᖏᓐᓂᒃᓂᖅ.

7.1.2 ᓄᓇᕗᓂᒃ ᐃᓄᐃᑦ ᐱᓕᕆᔨᓇᖏᓐᓂᑦ ᐱᔨᓇᐅᑎᕕᖃᖅᓯᓐᓂᑐᖅ

ᓄᓇᕗᓂᒃ ᐃᓄᐃᑦ ᐱᔨᓇᐅᑎᕕᖅᑲᓂᑐᑦ ᑎᓐᒥᐊᑦᑕᖃᖅᓴᖓᓄᐊᑦᑕᑕᐅᓇᖏᓐᓇᖅ, ᐊᐳᓪᓴᕆᓚᐊᖅᑕᕐᑎᒃ ᐱᔨᓇᐅᑎᕕᖃᖅᓯᓐᓂᑐᓴᓐᓄᓪᓗ, ᒪᓂᒪᓐᖓᓂ ᐱᓕᕆᐊᖅᑕᐅᓇᖏᓐᓇᕐᓂᒃ.

- ᐊᖏᔪᓚᐅᔨᓂᖏᓐᓂᒃ ᑕᒪᐅᐊᖅᑐᑦ 60-ᓂᒃ ᐳᔨᒪᓕᖅᑐᑦᓂᒃ ᐅᓴᑕᕐᑐᑎᓂᒃ ᐅᓂᒃᑯᐊᑎᓂᒃ ᑎᓐᒥᐊᑦᑕᖃᖅᓴᖓᓂᒃ ᐊᒻᒪᓗ ᖃᓄᐃᓕᐅᓇᖅ ᐱᔨᓇᐅᑎᖓᓂᒃ ᐃᓄᐃᑦᑦ ᓄᓇᖏᑎᑎᕐᓴᖓᓂᒃ ᑎᓐᒥᐊᑦᑕᖃᖅᓴᖓᓄᐊᑦ ᐃᓗᐊᓂᑕᑐᓂᒃ (*ᓄᓇᕗᓂᒃ ᓄᓇᕆᓂᖅᓯᒪᔨᑦ ᐊᖏᕈᑕᐅᓇᖏᓐᓇᕐᓂᒃ 19.9.4* ᐊᒻᒪᓗ *ᐃᓄᐃᑦ ᐃᑲᔪᐊᕐᕕᖏᓐᓂᓄᑦ ᐊᖏᕈᑕᐅᓂᑦ ᐃᓗᓕᕐᓂᑦ 5.4*), ᐊᒻᒪᓗ
- ᓴᖅᑭᑎᑎᕐᓂᖅᓄᑦ ᓄᑖᑎᓂᒃ ᓄᓇᓕᕆᔨᒪᔨᐅᓂᐊᕐᖃᖅᑐᑎᓂᒃ ᑕᖅᓱᖅᓴᕐᕕᖏᓐᓂᒃ (ᒪᓕᑐᖃᓗᐊᕈᑎᒃ ᑕᐃᑲᔪᓂᖅᓱᒪᓕᖅᓯᐅᑎᖑᓕᕐᓂᖅᓄᑦᓗ ᐃᓗᐊᓂᕐᕕᐅᓂᖅᓄᑦ 6.4.5, *ᐃᓄᐃᑦ ᐃᑲᔪᐊᕐᕕᖏᓐᓇᕐᓂᒃ ᐊᖏᕈᑕᐅᓂᑦ ᐃᓗᐊᓂᕐᕕᐅᓂᖅᓄᑦ 5.5-ᒥ* ᐊᒻᒪᓗ ᑕᐃᑲᔪᓂᖅᓱᓚ, *ᓄᓇᕗᓂᒃ ᓄᓇᕆᓂᖅᓯᒪᔨᑦ ᐊᖏᕈᑎᓂᖅ*).

ᐃᓄᐃᑦ ᐃᑲᔪᐊᕐᕕᖏᓐᓇᕐᓂᒃ ᐊᖏᕈᑕᐅᓂᑦ (ᐃᓗᐊᓂᕐᕕᐅᓂᖅᓄᑦ *5.3*) ᐃᓄᖅᓄᑦ ᐱᔨᓇᐅᑕᐅᕕᖅᑕᑦ ᑎᑎᕋᖅᑕᕐᕕᒃᓴᓇᓐᓇᕐᓂᖅ ᑕᐅᖅᓱᖓ ᐊᐅᓚᐅᕐᓴᓗᑎᒃ ᐊᖏᔪᓚᐅᕐᓂᖅᓯᒪᔨᓕᕐᓂᖅᓄᑦ ᑎᓐᒥᐊᑦᑕᖃᖅᓴᖓᓂᒡᒎᖅᓇᓐᓇᕐᓂᒃ ᐃᑲᖏᕈᑕᐅᓇᖏᓐᓇᕐᓂᒃ ᐃᖅᑲᓇᐃᑦ ᑎᓐᒥᐊᑦᑕᖃᖅᓴᖓᓂᒡᒐᖅ ᐊᒻᒪᓗ ᖃᓄᐃᓕᐅᓇᖅ ᐱᔨᓇᐅᑎᖓᓂᒃ ᐅᕙᓪᓕᐊᓂᒃᓴᓇᓐᓇᕐᓂᖅ ᐊᐅᑐᖅᑕᐅᓗᐊᓵᔪᒥᑎᓕᖅᓴᐃᑦ. ᐱᔨᓇᐅᑎᖓᓂᒃᓴᖅ ᑎᑎᕋᖅᑕᕐᕕᒃᓴᓇᓐᓇᕐᓂᖅ ᑕᐅᖅᓱᖓ ᐱᔨᓇᐅᑎᖓᓂᒃ ᐊᒻᒪᓗᓯᐅᑎᓕᕐᓂᑦ, ᐱᓕᕆᔨᓇᐅᑎᓕᕐᓂᑦᓪᓗ ᑐᐱᓯᐊᖅᑐᑎᓕᐅᓇᖓᓂᒃ ᐊᒻᒪᓗᓯᐅᑎᓕᕐᓂᑦ ᐱᓴᔨᑎᖓᓂᒃᖃᖅᑐᖅ ᐊᐅᓚᐅᕐᓴᓗᑎᒃ ᐱᔨᓇᐅᑎᖓᓂᒃ.

ᐃᓄᓕᒫᐅᖅᑕᐅᑐᓂ ᑐᓂᔾᔨᐅᓯᒪᔨᓇᓐᓂᑕᓂ ᑕᒪᐅᐊᖅᑐᑦ ᓄᓇᕗᑦ ᑎᑎᕋᖅᑕᕐᕕᒃᓴᓇᓐᓇᕐᓂᖅᓄᑦ ᐊᒻᒪᓗ ᒪᓕᑐᖃᖅᑎᑕᐅᓕᕐᓂᖅᓄᑦ, ᑐᓂᔾᔨᐅᓯᒪᓪᓗᐊᑕᐅᑎᓪᓗᓂᒃ ᐃᓄᓕᒫᐅᖅᑕᐅᓗᐊᓵᔪᒥᑎᓕᖅᓴᐃᑦ ᐊᓇᓯᐊᓂᒃᓴᓇᓐᓇᕐᓂᖅ ᐱᔨᓇᐅᑎᖓᓂᒃᓴᓇᓐᓇᕐᓂᖅ ᐊᓇᓯᐊᓂᒃᓴᓇᓐᓇᕐᓂᖅᓄᑦᓗ ᐃᓄᖅᑎᖓᓂᒃᖃᖅᑐᒥᒃ ᐃᓄᖅᑎᓪᓚᕆᖃᖅᑐᒥᒃ ᐱᓯᒪᓪᓗᐊᑕᐅᓂ, ᒪᓕᑐᖃᖅᑎᑕᐅᓕᕐᓂᖅᓄᑦ ᓄᓇᕗᓂᒃ ᓄᓇᕆᑕᐅᓂᑦ ᐊᖏᕈᑎᓂᖅ 5.7.35-ᒥᑦᑐᑦ.

7.1.3 ᓄᓇᕗᒻᒥᐅᑦ ᐃᓄᐃᑦ ᐱᓕᕆᔨᒋᐊᓂᖅᓯᓕᕐᓂᑦ ᑭᓱᑐᓂ ᐱᔪᓐᓇᐅᑎᖃᕐᑎᑐᖅ

ᑭᓇᐅᔨᓕᕆᐅᕐᑎᕆᔨᓂᒃ ᐱᓕᕆᔨᕆᐊᖃᐅᔪᑦ ᐱᓕᕆᔨᔨᒋᐊᕐᕈᓐᓇ ᐱᔪᓐᓇᐅᑎᖃᕆᐊᖃᖅᑐᑦ ᑎᒍᒥᕐᕆᖃᖅᓯᒪᔪᖅ ᓄᓇᕗᒻᒥ ᐃᓄᒃᓄᑦ ᓇᒍᑎᓂᖅᓴᐅᔨᓂᒃᓴᓐᓂᑦ. ᑕᐃᒪᓕᕆᓂᐊᕆᖃᖅᑎᑎᕐᓂᕐᒧᑦ ᑭᓱᑐᓂ ᐊᐅᓪᓚᑦᑭᓕᕆᓂᖅᑐᑦ ᐊᖏᓪᓗᓂᐊᕐᓂᐅᓚᑎᑐᖅ ᐊᖏᓪᓗᓂᓗᑎᓂᒃ ᐃᓕᒃᑲᓕᐊᕐᓂᒃᓴᓐᓂᑦ. ᓄᓇᕗᒻᒥ ᐃᓄᐃᑦ ᐊᐅᓪᓚᑦᑭᓕᕆᓂᖅᓴᒃᑐᑦ ᐱᔪᓐᓇᐅᑎᖃᕆᐊᖃᖅᓯᒪᓕᕆᓂᖅᓴᐅᑎᖃᖅᓯᒪᔪᖅ ᑭᓱᑐᓂ ᐃᓄᒃᑲᓕᐊᕐᓂᖅᓴᖅᑎᑦᑎᓂᑦ ᐊᖏᓪᓗᓂᐊᕐᓂᐅᓚᑎᑐᖅᓴᖅᑎᑦᑎᓂᑦ, ᐃᓕᒃᑲᓕᐊᕐᓂᖅᓴᐅᑎᖃᖅᑐᑦ ᐊᐅᓪᓚᑦᑭᓕᕆᓂᖅᓴᐅᑎᖃᖅᑐᑦ ᑭᓱᑐᓂ ᐃᓄᐃᑦ ᐊᐅᓪᓚᑦᑭᓕᕆᓂᖅᓴᐅᑎᖃᖅᑐᑦ. ᑕᑯᓗᑎᒃ ᐅᖃᐅᓯᖅᑐᑦ ᐊᖏᓪᓗᓂᐊᕐᓂᐅᓚᑎᑐᖅᓴᐅᑎᖃᖅᑐᑦ (ᓄᓇᓗ ᐊᐢᑎᑎᒃ ᑕᑯᓂᐊᕆᖃᖅᑎᑎᕐᓂᕐᒧᑦ, ᖃᐅᔨᓚᐊᕈᑲᔪᓂᓂᕐᒧᑦ, ᐊᓱᖏᑐᓂᑦᓗᓗ) ᓄᓇᕗᒻᒥ ᐃᓄᖕᓂᒃ ᐱᔪᓐᓇᐅᑎᖃᖅᑎᑎᕐᓂᕆᐅᓇᖅᓴᖅᑐᑦ. ᑐᑭᓱᓂᐊᕐᓇᒪᓕᕐᓂᑦ ᐃᓕᓴᖏᓂᑦ ᑎᑎᕋᖅᑭᓱᓂᖅᓴᖏᓂᑦ ᐱᔪᓐᓇᐅᑎᓕᓚᔪᓂᑦ 7.1.7-ᒥ ᑐᑭᓱᐊᕋᑎᐊᕈᓐᓇᐅᓇᖅᓴᓐᓂᕐᒧᑦ ᐱᔪᓐᓇᐅᑎᓕᓚᔪᓂᒃ ᐱᓕᕆᓂᖅᓴᖅᑐᑦ.

7.1.4 ᐃᓄᒃᓕᓚᐅᖏᓂᑐᓄᓂ ᐱᓕᓚᐊᖅᑭᓐᓇᕆᓂᖏᓂᑦ ᐱᔪᓐᓇᐅᑎᖃᖅᖏᓂᒃᓗᑎᑐᖅ


ᑕᒪᐃᓐᓂᕎᑎᐊᖅᑐ ᐱᓕᕆᐊᖓᔪᓐᓇᖅᑐᑦ, ᐃᓕᓴᖅᑭᓱᑎᑦᓗ ᑕᐃᒃᑯᓂᖅᓴ ᓄᓇᒪᑦ ᑕᐅᕐᕉᖅ ᑎᑎᕈᓱᓕᐊᓐᓇᐅᑎᖃᖅᑐᑦ ᐊᒻᒪᓗ ᓄᓇᖏᑉᐳᑐᓂᐅᓐᓇᖅᓴᓂᖏᓂᒃ ᑲᔪᓯᑎᑕᐅᓇᓴᐅᑎᒃ ᐱᔪᓐᓇᖅᑎᑕᐅᔨᓗᑎᒃ, ᐱᔪᓐᓇᐅᑎᖃᕆᐊᖃᖅᓯᓂᐊᕐᓂᖅᑐᑦᑐᖅᑕᐅᕐᕉᓂ ᓱᓕ ᐃᓄᒃᓕᓚᐅᖏᓂᑐᓂ, ᑎᓪᓗᒍᑦ ᓂᕆᐹᒃᑲᓚᔨᐅᕈᑐᑦ ᐃᓄᐃᑦ ᓄᓇᖏᑉᐳᓕᕐᓂᓴᑦᑎᓪᓗ (ᑕᒪᓐᓂ ᐃᓕᓐᓂᒃ ᐊᒃᑐᐱᓚᒪᑦ ᑎᑎᕋᖅᑭᓱᑐᓂ, ᐱᔪᓐᓇᐅᑎᖃᖅᑎᑎᕐᓂᕐᒧᑦ ᐊᒻᒪᓗ ᖃᐅᔨᓴᓇᓕᖅᑐᖅ ᐊᐅᓚᑦᑭᓕᕆᓂᖅᓴᐅᑎᖃᖅᑐᑦ ᐊᒻᒪᓗ

7.1.6 ᐱᓕᕆᔨᓐᓇᖅᑐᖅ ᐊᖏᖅᑕᐅᓂᒪᔪᓂ ᐱᔨᔪᓐᓇᐅᑎᖃᖅᑎᓗᑎᑐᖅ ᐃᓄᖅᓯᐅᓂ ᐱᖅᒋᐱᑎᕐᓯᐅᔪᓂ ᓄᓇᓕᒫ

ᑕᐃᒃᑯᐊ *ᑲᑎᓴᒪᕆᑕᐅᖅᓯᖏᓐᓄᓄᓂ ᑎᖏᐊᓕᓚᓴᖅᓱᓕᒍᓂ ᒪᓕᑦᑐᐃᑦ, ᐊᓴᒪᓗ ᒪᓕᑦᑐᓗᑐᓂᓕᒍᓂ ᐊᑐᖅᑕᐅᕆᐅᓂ ᐃᓄᖅᓯᐅᑎᖃᖅᑎᓗᑎᑐᖅ ᐱᖅᒋᐱᑎᕐᓯᐅᔪᓂ ᐱᖅᒋᐱᑎᕐᓯᐅᔪᓂ ᓄᓇᓕᒫ ᑎᖏᐊᓕᖅᑐᖅᓯᖏᓐᓄᓄᓂ.* ᒥᓂᔨᑕᐅᔪᖅ ᐊᐅᑎᓕᓚᓴᕐᔪᐊᑲᓴᖅᓂᓄᓂ ᑐᓂᓯᑕᐅᔪᓂ ᐱᖅᒋᐱᑎᕐᓯᐅᔪᓂ ᓄᓇᓕᒫ, ᑐᐱᓯᓂᐊᑕᑕᐅᓕᐅᔨᓐᓇᑎᓄ ᑭᐅᓂᓕᒥ ᐃᓄᓐᓂᑦ ᐱᓕᕆᔨᓐᓇᐅᑎᓂᐊᖏᓐᓄᓄᓂ ᐊᖏᖅᑕᐅᓂᒪᔪᓂ ᐃᓄᖅᓯᐅᑎᖃᖅᑎᓗᑎᑐᖅ ᓄᓇᓕᒫ, ᑐᐱᓯᓂᐊᑕᑕᐅᓕᐅᔨᓐᓇᑎᓄ ᑭᐅᓂᓕᒥ ᐃᓄᓐᓂᑦ ᐱᖅᒋᐱᑎᕐᓯᐅᔪᓂ ᓄᓇᓕᒫ, ᑭᐅᓂᓕᒥ ᐃᓄᓐᓂᑦ ᐱᓕᕆᔨᓐᓇᐅᑎᓂᐊᖏᓐᓄᓄᓂ ᐃᓄᖅᓯᐅᑎᖃᖅᑎᓗᑎᑐᖅ ᓄᓇᓕᒫ

ᑲᓇᑕᒥ ᐅᒪᔨᓕᕆᔨᖅᑯᓂ ᑐᓂᓯᔪᓐᓇᐧᐃᖓᑕᓂ ᐱᔪᓐᓇᐅᑎᓂᒃ ᐱᔪᓐᓇᐅᑎᑕᕋᓯᐊᖅᑐᒍᓂ ᐅᒃᑐᖅᓯᒪᔨᒍᓂ, ᑭᓯᐊᓂ ᐱᐅᔨᓂᒃ ᒪᓕᕐᑎᐊᖅᓂᕋᖅᑕᐅᓯᒪᔪᑎᑎᒃ ᑐᑭᓕᐅᖅᑕᐅᓯᒪᓕᕐᑐᒍᓂᒃ ᑐᓂᓴᐅᓯᒪᔨᔪᓕᕐᓂᐅᓂᒃ ᑭᓯᐊᓂ, ᐸᖅᓴᓇᖅᑕᐅᓴᓇᐊᖅᓴᖅᑎᓕᖅᑕᐅᔪᑎᑎᒃ ᑭᓯᐊᓂ, ᐅᖃᓕᕆᔨᒃᑯᑦ ᐅᖅᓯᓯᒪᔨᔪᓕᕐᓂᐅᓂᒃ. ᐊᓚᒪᑕᐅᖅᑲ ᓯᓗ, ᐅᖅᑲᐅᔾᔨᑎᐊᖅᑎᑎᐅᖅᑲᖅᑕᖏᒪᑕ ᒥᓂᔪᒧᑦ ᑕᒪᐃᓐᖅᑲᖅᑐᓂ ᐸᖅᓴᐅᑎᓕᐊᓇᒃᐅᕐᓚᐅᖅᑐᓂ ᐊᓚᒪᓗ ᐊᐅᓕᐅᕐᓴᓐᖓᓕᓐᓂᓂᓂ. ᐃᖅᓂᐅᖅᓇᐃᓕᑎᑕᐅᖅᑲ ᐊᐅᓕᖅᑎᑎᓇᐅᖅᑲᖅᑕᐅᓂᖅᔪᑦ ᑲᑎᒪᔨᖅᑕᕐᕕᖅᑎᓕᖅᑕᐅᓂᖅᑎᒍᓂ

ᖃᒃᖏᕐᓂᖅᓯᒪᔪᖅ ᑎᒥᐅᔨᖏᓐᓂ ᐊᓯᖏᑐᓂᒃ ᓯᓚ, ᓯᕗᓪ ᐅᒃᑐᑎᑦᓚᐃ, ᖃᐅᔨᓴᖅᑭᓚᕐᔪᐊᕐᖃᖅᓯᓂᐅᔨᖅᑕᕐᓂ ᐱᓯᒪᔪᓚᐅᖃᖅᑎᑕᐅᔪᓐᓇᖅᑐᑦ ᖃᐅᔨᓴᖅᑭᐊᓚᖕᖓᑦ ᓚᐃᔪᓐᓇᓯᑕᕐᕕᐅᒃᑐᓂᒃ ᓄᓇᕗᒻᒥ ᖃᐅᔨᓴᖅᑭᖏᓐᓇᖅᑐᓂᒃ ᐅᕙᓪᓕᐊᓂᒃ ᐅᑎᓪᓗᒍ ᐅᒃᑐᓂᒃ ᓴᖅᑭᒪᔭᖓᓂᒃ. ᐅᖃᖅᑭᐅᐅᔪᓚᐅᖃᖅᑎᑕᒃᑐᓂ ᐊᐃᔪᑐᖅᓯᒪᓂᑯᓘᓐᓃᓕᓂᒃ ᓄᓇᕗᒻᒥᓄᓇᓕᒫᓂ ᐱᓕᕆᔨᒃᑲᓪᓚᕆᔨᒃᑲᓪᓚᕆᐊᓘᓐᓃᑦ ᑐᑭᓯᐊᓂᒃ ᐱᓇᓱᐊᕐᓂᖅ.



ᑕᒪᔭᒃᓴᖅ 8: ᑎᓪᓗᒍᖃᖅᑎᑕᒃ ᐱᓇᓱᐊᕐᑕᕆᔨᓴᓕᐊᑭᑎᓂᒃᑎᒪᒃᑕ ᐅᒃᑐᖅᑭᑐᓂ ᖃᓄᐃᓕᑕᐅᖅᑕᐅᔪᓐᓇᖅᑐᓂᒃ.

8 ᓄᓇᖅᑐᑦ ᑎᖅᑲᐊᖏᖅᖢᐅᔪᑦ ᖃᓄᐃᓕᓕᐅᔪᖅᓯᖅᑐᓂᓄᑦ ᑕᐃᔨᐅᓂᖅᑐᑦ ᐃᓕᓴᕆᐊᓕᓂᖅᑦᓗ

ᖃᖏᕿᐅᖏᖅᑐᑎᖅ ᐊᓐᓂᐊᕐᓇᖅ ᐃᓕᓴᕆᐊᓕᖅᑐᑦ ᑎᖅᑲᐊᖏᖅᖢᐅᔪᖅᓯᖅᑐᓂᑲᑦ; ᑭᓱᐊᓂᓕ, ᐊᖏᖅᔪᓕᐊᓐᓇᖅ ᓴᐅᓂᐊᖏᖅᑕᐅᕐᓚᓐᖓᑭᑕᓂ ᐅᓪᓗᖏᓐᓂᑦᑐᓂᓕᕆᐊᓂᓐᓄ ᓄᑕᖅᑕᐅᓂᐊᕐᓇᓂᓐᓄ ᐱᐅᓂᖏᓕᕆᐊᓂᓐᓄ ᑲᑎᖅᑐᐃᓐᓇᖅᑐᑦ ᐅᓪᓗᑕᓕᐊᒥ ᐃᓕᓴᕆᖏᖅᖢᓚᐅᑕᐅᓂᓄ ᐱᐅᓂᖏᓕᕆᐊᓂᓐᓄ ᐃᓕᓴᕆᐊᓂᖅᑐᑦ.

ᓯᕐᓂᓕᖏᑦ ᑕᐃᔨᓱᓕᓂ ᑎᒃᑯᐊᖏᖅᑕᐅᓚᓕᑦ ᑎᖅᑲᐊᖏᖅᖢᐅᔪᖅᓯᖅᑐᑦ ᓴᓕᒪᓕᐅᔪᖅᓕᑦ ᐃᓕᓴᐃᓕ ᓴᓂᐅᓂᐊᖏᖅᑕᐅᑎᐊᖏᓂᖅᑎᓂᓄᑦ, ᑲᖏᖅᑐᑦ ᐃᓕᓴᕆᖏᖅᖢᐅᔪᖅᓯᖅᑐᓂ 60-ᖓᔪᓂᓂ ᐊᖏᖅᔪᓂᓂ, ᓴᓂᐅᓂᐊᖏᖅᑕᐅᔨᓚᐊᖏᖅᖢᓂᖅᑎᓂᓄᑦ ᐊᓯᕐᓂᖅᓯᓕᖅᓂᓂ ᑎᖅᑲᐊᖏᓂᓂ, ᐊᓯᖅᑐᑦ ᐃᓕᓴᕆᖏᖅᖢᐅᔪᓕᖅᑐᑦ ᐊᓚᓚᓗ ᐃᓕᓴᐃᓕ ᐃᓕᖅᑐᑦ ᐊᑎᓕᕐᓂᖅᑐᓄᑦ, ᐊᓚᓗ ᐃᓄᐃᓕ ᐃᓕᖏᖅᑯᓯᑎᖅᑲᖅᑐᓂᖅᑐᑦ ᐱᖏᓂᖅᑐᐃᓐᓇᖅᑐᑦ ᐊᓯᔪᖏᖅᑐᓕᕐᓂᖅᓯᓯᒪᓂᖅ.

ᑕᐃᔨᓱᓕᓂ 1959-ᖓᔪᑎᓪᓗᒍ, ᑲᓇᑕᒥ ᐃᓕᓴᕐᓚᓴᖅᑭᑦ ᓴᖅᑭᑎᑎᓚᐅᖏᖅᓯᒪᖓᓚᑦ ᑎᖅᑲᐊᖏᖅᖢᐅᔪᖅᓯᐊᓕᕐᓂᖅᑐᒥᑦ, ᑲᓇᑕᒥ ᒪᓕᕐᓚᐅᕈᑲᓇᖅᑐᓂᑦ ᐱᔨᓄᓇᐅᑐᐃᓇᖅᑐᓴᓕᖅᓚᓕᑦ ᑲᓇᑕᒥ ᓄᓇᓂ ᐊᑐᑎᖏᖅᓂᖅᑐᓂᓄ ᓇᔨᖏᑕᐅᓕᖕᖏᓂᕐᑐᓄ ᐃᓕᓴᕆᖏᖅᖢᐅᔪᖅᓯᐅᔨᓂᖅ. ᑕᐃᔭᓕᐊ *ᑲᓇᑕᒥ ᐃᓕᔨᓕᓚᓂᕐᒧᑦ ᒪᓕᕐᓚᐃᑦ* ᒥᓵᓪᑕᐃᑦ ᑲᑎᖅᑐᐃᓐᓇᖅᑐᑦ ᑲᑎᕕᒃᑐᓕᕐᓂᖅᑐᑦ ᑕᒧᓂᑐᖅᖢᓂᖅᓯᒪᓕᕐᓂᖅ. ᐅᖃᓕᒫᒃ 1973-ᖓᔪᓕᖅᑎᓪᓗᒍ. *ᒐᖅᓕᐊ ᒪᓕᕐᑎᖓᓂ* ᐱᔨᓄᓇᐅᑕᐅᕐᓴᓂ ᐊᖏᖅᑭᔨᓇᓂᕐᒥᑦ, ᐊᐅᓚᓂᑎᔨᓇᓂᕐᒥᑦ ᐊᓚᓚᓗ ᓴᓂᐅᓂᐊᖏᖅᑕᐅᔨᓇᓂᖅᑎᓂᖅᑐᓄᑦ ᑲᓇᑕᒥ ᐃᓕᓴᕆᖏᖅᓱᓂ, ᖃᐅᔨᓴᓂᖅᑕᐅᔨᓇᓂᖅᑎᓂᖅᓯᓂᓄᑦ ᐅᓴᓇᓂᖅ ᑐᓴᐅᓂᖅᑐᓂ ᑐᓐᖓᓂᖅᑕᐅᔪᓕᖅᑐᑦ ᐃᓕᔨᖓᓂ, ᐱᑕᖅᑲᐃᓐᓇᐊᖏᖅᓯᖅᑎᓂᖅᓯᓂᓄᑦ ᐅᓴᓇᓂᖅᑎᓂᖅᓯᓂᓄᑦ ᐱᑕᖅᑲᑕᐅᓂᖅᑐᑦ.

ᓄᓇᐅᓂᖅ ᐊᑕᐅᔨᕐᒪᓇᓐᓂᖅᓴᓂ ᓴᖅᑭᑎᑎᐊᓐᓇᐊᖏᑕᐅᓂᕐᒧᑦ ᒪᓕᕐᓂᓂᑦ ᐱᑕᖅᑲᓚᐊᖏᖅᖢᓂᖅᑐᑦ ᐊᑕᐌᓂᑐᓂᑦ ᑎᒃᑯᐊᖏᖅᑕᐅᖏᖅᑕᐅᓂᕐᒧᑦ ᐃᓕᓴᕆᖏᖅᓱᓕᒃ. ᖃᖏᕿᐅᖏᖅᑐᑎᖅ ᐊᓐᓂᐊᕐᓇᖅ ᓴᐅᓂᐊᖏᖅᑕᐅᓕᖕᖏᓂᕐᑐᓄ ᐃᓕᓴᕆᖏᖅᖢᓂᖅᑐᑦ ᐃᓕᓴᕆᐊᓂᖅᑐᑦ ᐱᑕᖅᑲᑕᐅᓂᖅᑐᑦ ᐱᔨᓄᓇᐅᑐᐃᓇᖅᑐᓴᓕᖅᓚᓕ

ᐊᐃᔭᑐᖏᓪᓗᕆᓂᔭᖅ ᑕᔨᑎᕐᑎᓴᕆᓂ 8: ᒪᓕᓚᓴᐅᔪᓂ ᖃᓄᐃᓕᖅᑕᐅᓚᓴᓚᐊᖏᖅᖢᓂᖅᓄᑦᓂ ᑎᑦᔨᐊᖏᖅᑕᐅᔪᖅᓀᓂᐊᕐᓂᑐ ᖃᖏᖏᓴᐅᖏᖅᑐᑦ ᑲᓇᑕᓕᒫᒥ ᐅᒪᔪᖏᖅᓃᐃᓕᓂᒧᑦ

ᒪᓕᓚᐃᑦ ᖃᓄᐃᓕᓕᖅᓴᓚᐊᖏᖅᖢᓂᒧᑦ ᑲᓇᑕᓕᒫᒥ ᐅᒪᔪᖏᖅᓃᐃᓕᓂᒧᑦ (ᐊᕐᕈᑎᓕᓚᔪᔪᐊᖅᑯᑦ 2017b)	ᖃᖏᖏᓴᐅᖏᖅᑐᑦ ᖃᓄᐅᔪᓂᖏᖅᑕᐅᓕᕆᓂᖅ ᖃᓄᐃᓕᓕᖅᓴᓂᖅᓂᖅ ᒪᓕᓚᓕᓚᓴᐅᔪᓂᑐ ᒪᓕᖅᓂᖅᓂᖅ
1. ᑕᒪᐅᓗ ᓄᓇᖅᓴ ᐱᑕᓕᒃ ᒥᕈᓂᖏᓐᓂᒃᖢᑕᐅᔪᒥᒃ 1% ᔪᓂᖅᑎᖅᑯᓕᑐᓂᒃ ᑲᓇᑕᒥ ᐅᒪᔪᑯᑎᖅᖢᓂᒃ ᐊᔨᐅᑲᑎᕐᖏᖅᖢᓂᓚᐅᑯᓂ ᐅᒪᔪᓂᒃ ᑎᐹᐸᑐᑐᓂ ᑎᖅᑎᐊᖅᑐᓂ ᐅᒪᔪᓂᒃ ᐱᖓᓱᓚᐅᑯᓂ ᑕᐃᒃᑯᓂᖅᓴ ᐅᒪᔪᓂᒃ ᐊᒥᓱᖅᑎᓚᐅᑯᓂ ᐅᒪᔪᓂᒃ ᐊᒥᓱᖅᑎᓚᐅᑯᓂ ᐱᖓᓱᓚᐅᑯᓂ ᐅᒪᔪᓂᒃ ᐊᒥᓱᖅᑎ

	✓ ᐱᓚᒪᓕᒧᑦᓴᖅ ᖃᐅᔨᒪᓂᖅᑐᓄᑦᓗ ᐅᑦᑐᖅᑎᓂᖅᓂᓗ ᐊᐃᕕᖏᓄᑦ (ᖃᐅᑎᖅᑕᓕᕆᓂ/ᐊᑦᑎᐅᑐᑐᖅᑕᐅᓂᖅᓂ ᐅᐸᐅᖅᑕᖅᑐᑎ ᐊᐃᕕᖏᓄᑦ, ᐅᐊᖅᓇᖅᓚᓂᓗ ᐱᖅᑐᖅᓇᖅᓚᓂ ᕼᐊᐸᖅᐱᑦ ᐊᐃᕕᖏᖅᑐᓄᑦ)
3. ᑕᒪᐃᓇ ᓄᓇᖅᑐ ᐃᓕᑕᒪᔭᐅᓕᕆᔨᖅ ᐱᑕᖃᖅᓚᐊᖃᖅᓯᒪᓚᖅᓂᖅ ᑎᑎᕋᖅᑕᐅᓕᕆᔨᓄᑦ ᑎᖅᑎᐊᖏᓄᑦᓗᐃᓐᓂᑦ ᐊᓕᕆᔨᓄᑦ ᐊᒥᓕᖅᑎᓗᐊᓕᖅᖢᑎᓄᑦ, ᐊᐸᔭᑦᓴᖅᑎᓄᑦᓄᓗᐃᓐᓂᑦ ᑕᒪᒃᑯᐊ ᐅᒪᔨᐟᖅᑎᒋᔨᓄᑦ.	ᐊᑐᖅᑎᑐᖅ
4. ᑕᒪᐃᓇ ᓄᓇᖅᑐ ᐊᐸᐅᖅᑎᓕᖅ ᐅᐃᓇ, ᐊᒥᓕᖅᑎᑐᑦᔨᐊᖅᓯᓕᖅᖢᑎᓄᑦ ᑕᑯᐸᐅᓚᐃᑦᑐᑎᓄᑦᓗ ᐅᒪᔨᖃᓕᓕᖅᑐᒥ ᑕᒥᓂᒃ ᐅᒪᔨᐃᑦ ᐊᕕᑐᖅᑎᑦᓴᖅᑐᑦᓂᖅ, ᐅᖏᐟᓓᓇᓂᓂᒃ ᐱᑕᖃᖅᓚᐊᖃᖅᓯᒪᓚᖅᓂᖅᓄᑦ ᐃᓕᓴᕐᕕᒃᑐᖅ ᐱᑕᖃᖅᓚᐅᐸᑦᑐᖅᓯᒪᓚᖅᓂᖅᓄᑦ ᐃᓚᒋᑦᑐᖅᓯᒪᓚᖅᓂᖅᓄᑦ ᐊᐅᓚᑦᑐᖅᓯᒪᓚᖅᓂᖅᓄᑦ ᑕᑯᐸᐅᓚᐃᑦᑐᑎᓄᑦᓗ ᐃᓚᒋᑦᑐᖅᓯᒪᓚᖅᓂᖅᓄᑦ ᐊᑐᐊᓂᖅᓄᑦ

ᐃᓕᖅᑕᕐᓂᑦ ᑕᒪᐅᑯᐊ ᖃᑯᓪᓗ ᒪᓕᑕᒫᕐᓂᑦ ᐅᒪᔨᖅᖃᓕᓕᐅᕈᓐᓇᕆᑕᖅᑕᕐᓂᓐᓇᑦ, ᑕᒪᓐᓇ ᖃᖅᖢᓴᐅᖅᑐᖅᑎᒋᑦ ᑎᓯᒥᐊᖅᖃᖅᓱᓕᒃ ᐊᑯᓂᐊᓗᓄᑦ ᑲᒪᒋᐅᖅᑕᖅᑕᕐᕕᒃᓴᓪᓚᕆᔨᒃᑯᑦ ᖃᐅᔨᓴᖅᑕᕐᕕᖅᑕᕐᓂᓐᓇᑦ, ᐱᓗᐊᖅᑐᒥᒃ ᐱᔨᔪᑎᖅᖃᖅᑐᓂᒃ ᑲᖏᖅᓯᓂᒃ 9-ᒥᑦᑐᑦ ᓇᐃᓰᖅᓴᒪᔨᓂᒃ ᐅᓂᒃᑲᖅᓯᐅᑦ ᖃᐅᔨᓴᖅᑕᕐᕕᓴᒪᓂᕐᓂᓂᒃ ᐊᖏᓗ ᓇᐅᑦᑎᖅᑐᖅᑕᕐᕕᐅᖅᑕᖅᑕᕐᕕᓴᒪᓂᕐᓂᓂᒃ ᐊᑐᖅᑕᕐᕕᓴᒪᓗ, ᐊᑐᐃᓇᐅᓗᓄᒃ ᐱᔨᓇᐅᑎᓕᓪᓕᖅᑲᒃ ᑎᑎᖅᑐᖅᑕᕐᕕᐅᔨᔪᑎᖅᑐᖅᑕᕐᕕᓴᒪᓂᕐᓂᓂᒃ ᐊᖏᓗ ᑎᑎᕆᑐᖅᑕᕐᕕᓴᒪᔨᓂᒃ ᖃᑭᖅᑕᐊᖅᑕᕐᕕᖅᑕᕐᕕᓴᒪᔨᓂᒃ ᖃᐅᔨᓴᒪᔨᖅᑕᕐᕕᐅᓂᒃ.

ᐊᓚᐅᑐᖅᓴᒪᔨᓂᒃ ᑕᒧᑎᑎᑎᔨᓂᑦ 9: ᓇᐃᓰᖅᑕᕐᕕᓴᒪᔨᓂᑦ ᖃᐅᔨᓴᖅᑎᑎᕐᕕᐅᓕᕐᕕᖅᑐᑦ ᓇᐅᑦᑎᖅᑐᐃᔪᖅᑐᑦᑕᖅᑕᕐᕕᓴᒪᔨᓂᓪᓗ ᑎᓯᒥᐊᖅᖃᓚᐅᖅᓯᒪᔪᒃ

ᐊᖏᖅᓯᒪᔪ	ᖃᐅᔨᓴᖅᑎᑦ ᐊᑎᑭᓴᖅᑎᑦ (ᑎᒥᖏᒧᑎᖅᑎᓪᓗᓂᒃᓂᑦ)	ᐱᔨᔪᑎᖅᑐᖅᑕᕐᕕᖅ
1952	ᓗᑌ-Loughrey (ᑲᓇᑕᒥ ᐅᒪᔨᓕᑌᓵᓂᑦ ᐊᐅᑎᓕᑌᓴᖏᐊᖅᑯᑦ)	ᑎᓯᒥᓴᑦ ᖃᐅᔨᓴᖏᓂᖅ
1955	ᑯᑦᓴ-Cooch (ᑲᓇᑕᒥ ᐅᒪᔨᓕᑌᓵᓂᑦ ᐊᐅᑎᓕᑌᓴᖏᐊᖅᑯᑦ)	ᑎᓯᒥᓴᖅᑯᑦ ᖃᑯᓇᐅᑕᐅᓪᓗᑎᒃ ᐃᒪᐅᓂᖏᑦᓴᑦ ᑎᓯᒥᐊᑦ ᖃᐅᔨᓴᖅᑕᕐᕕᐅᓂᓯᕐᓂᑦ
1956-1957	ᐱᐅᓇ-Barry (ᑲᓇᑕᒥ ᐅᒪᔨᓕᑌᓵᓂᑦ ᐊᐅᑎᓕᑌᓴᖏᐊᖅᑯᑦ)	ᑲᖓᖏᓂᒃ ᖃᐅᔨᓴᖏᓂᖅ
1966	ᑯᐱᓴ-Kerbes (ᑲᓇᑕᒥ ᐅᒪᔨᓕᑌᓵᓂᑦ, ᐊᐅᑎᓕᑌᓴᖏᐊᖅᑯᑦ)	ᑲᖓᖏᓂᒃ ᖃᐅᔨᓴᖏᓂᖅ
1972-1973	ᑯᐱᓴ-Kerbes (ᑲᓇᑕᒥ ᐅᒪᔨᓕᑌᓵᓂᑦ, ᐊᐅᑎᓕᑌᓴᖏᐊᖅᑯᑦ)	ᐊᓕᓕᐅᖅᑕᕐᕕᐅᓗᑎᒃ ᖃᐅᔨᓴᖅᑕᕐᕕᐅᓂᓯᒪ ᑲᖓᓴᖅᖃᓚᒃ
1979	ᓀᓪᓂ-Reed, Dupuis (ᑲᓇᑕᒥ ᐅᒪᔨᓕᑌᓵᓂᑦ, ᐊᐅᑎᓕᑌᓴᖏᐊᖅᑯᑦ)	ᐊᓕᓕᐅᖅᑕᕐᕕᐅᓗᑎᒃ ᖃᐅᔨᓴᖅᑕᕐᕕᐅᓂᓯᒪ ᑲᖓᓴᖅᖃᓚᒃ
1979-1980	ᐃᐳᑕᒪᕼᐃᓕ, ᐊᖕᑭ-Abraham, Ankney (ᐊᑲᓐᑎᐅᓕᑌᓵᒥ ᒥᓂᔪᕆᖅᖃᓚᒃ ᓄᓇᒥᐅᑐᑐᓕᑌᓴᖏᐊᖅᑯᑦ, ᔨᑦᓂᑭᓕᑎᖕᓂᓂᒃ ᐱᖓᓱᓇᖓᓚᑕ ᐊᑲᓐᑎᐅᓕᑌᔨᑦ)	ᑲᖓᐃᐅᑦ, ᓇᐅᑲᐃᐅᑦ, ᐃᒥᖅᑯᑕᐃᓕᐃᐅᑦ ᖃᐅᔨᓴᖅᑕᕐᕕᐅᓂᓯᕐᓂᑦ
1987	ᖃᐅᔨᓴᓕᐅᖕᓂᑕᖅ	ᓄᓇᖕᓂᓂᒃ ᖃᐅᔨᓴᖅᑎᑎᖅᑐᖅ
1992	ᑲᐅᐃᓪ-Caswell (ᑲᓇᑕᒥ ᐅᒪᔨᓕᑌᓵᓂᑦ, ᐊᐅᑎᓕᑌᓴᖏᐊᖅᑯᑦ)	ᑲᖓᖏᓂᒃ ᖃᐅᔨᓴᖅᑎᑎᖅᑐᖅ
1997	ᑯᐱᓴ-Kerbes (ᑲᓇᑕᒥ ᐅᒪᔨᓕᑌᓵᓂᑦ, ᐊᐅᑎᓕᑌᓴᖏᐊᖅᑯᑦ)	ᐊᓕᓕᐅᖅᑕᕐᕕᐅᓪᓗᑎᒃ ᖃᐅᔨᓴᖅᑕᕐᕕᐅᓂᓯᕐᓂᑦ ᑲᖓᓴᖅᖃᓚᒃ
1997-2002	ᑎᐅᓪᑯᓗᓂᑦ, ᕕᓪᓕᓐᓂᑦ, ᒧᓗᓇ-Gilchrist, Wayland, Mallory (ᑲᓇᑕᒥ ᐅᒪᔨᓕᑌᓵᓂᑦ, ᐊᐅᑎᓕᑌᓴᖏᐊᖅᑯᑦ)	ᒥᑎᓂᑦ, ᓄᑕᖏᓇᖅᑐᖅᑎᖅᓂᓯᕐᓂᑦ ᐃᒪᐃᓪᓗ ᐃᓕᓴᖅᓯᐅᑕᐅᓇᖅᑐᓂᓯᕐᓂᑦ
1999-2001	ᕿᓄᓵ-Bazin (ᑲᓇᑕᒥ ᐅᒪᔨᓕᑌᓵᓂᑦ, ᐊᐅᑎᓕᑌᓴᖏᐊᖅᑯᑦ)	ᑲᖓᖏᓂᒃ ᖃᐅᔨᓴᖏᓂᖅ
1999-2019	ᑎᐅᓪᑯᓗᓂᑦ-Gilchrist ᖃᐅᔨᓴᖏᓂᖅ ᐊᑎᓪᓗ ᓴᓇᖏᑎᐅᓴᓕᓂᓯᕐᓂᑦ, ᐊᐅᑎᓕᑌᓴᖏᐊᖅᑯᑦ)	ᑎᓯᒥᐊᑦ ᖃᐅᔨᓴᖅᑕᕐᕕᐅᓂᓯᕐᓂᑦ (ᒥᑎᓂᑦ, ᑕᓚᐅᖏᒥᓄᑦ ᑎᓯᒥᐊᑦ,

ᐊᖏᕕᒋ	ᖃᐅᔨᓴᖃᑎᒋᔭᖅ ᐊᑎᑕᓯᖏᓐᓂᒃ (ᑎᒥᖏᑎᖏᓐᓂᒃᓴᓇᓂᒃ)	ᐱᔪᓐᖃᑦᑕᕐᓂᖅ
		ᓯᓚᖃᖅᐸᓯᐅᓂᖏᓕᓴᐃᓂ ᑎᖏᐊᑦ, ᓄᓇᒋᓴᐃᓂ ᑎᖏᐊᑦ), ᐊᑲᓂᐊᓕᕆᔨᐅᕙᖅᑐᑦ, ᓯᕆᖏᓇᖅᑐᖃᖏᓂᖏᓐᓂᒃ, ᐅᓄᐃᑦ.
2002-2003	ᔭᑭᓂᓪ-Perkins (ᔪᓂᒪᓯᑎᖓᓪᓚᑦ ᒋᓂ)	ᓯᓚᖃᖅᐸᓯᐅᓂᖏᓕᓴᐃᓂ ᑎᖏᐊᑦ ᖃᐅᔨᓴᖃᑎᒋᐅᓂᖏᓐᓂᒃ
2002-2020	ᓕᖅᓚᐊᖏᓪ-Leafloor (ᑲᓇᑕᒥ ᐅᒪᔪᓕᓗᓐᓄᑦ, ᐊᐅᑎᓕᓗᓐᓄᕐᔪᐊᖅᑯᑦ)	ᑲᖓᐃᐃᑦ ᖃᐅᔨᓴᖃᑎᒋᐅᓂᖏᓐᓂᒃ
2003-2019	ᓯᒥᑦ-Smith ᖃᐅᔨᓴᖏᓂᖅ ᐊᓚᒪᓗ ᓴᓇᖏᑎᐅᖅᓕᓕᓗᓐᓂᒃ, ᐊᐅᑎᓕᓗᓐᓄᕐᔪᐊᖅᑯᑦ)	ᓯᓚᖃᖅᐸᓯᐅᓂᖏᓕᓴᐃᓂ ᑎᖏᐊᑦ ᖃᐅᔨᓴᖃᑎᒋᐅᓂᖏᓐᓂᒃ
2005-2008	ᑲᐃᐅᑦ-Caswell (ᑲᓇᑕᒥ ᐅᒪᔪᓕᓗᓐᓄᑦ, ᐊᐅᑎᓕᓗᓐᓄᕐᔪᐊᖅᑯᑦ)	ᑲᖓᐃᐃᑦ ᖃᐅᔨᓴᖃᑎᒋᐅᓂᖏᓐᓂᒃ
2007-2009	ᐱᓂᓴᓪ-Pienitz (ᔪᓂᒪᓯᑎᑎ ᓕᖅᑦ)	ᓄᓇᖅᐱᐅᓂᖏᓐᓂᒃ ᐃᒪᓕᓂ ᖃᑯᐱᑦᖃᖏᓂᖏᓐᓂᒃ ᐊᓚᒪᓗ ᐃᒪᐃᑦ
2008-2013	ᒐᓴᑎᓪ-Gaston ᖃᐅᔨᓴᖏᓂᖅ ᐊᓚᒪᓗ ᓴᓇᖏᑎᐅᖅᓕᓕᓗᓐᓂᒃ, ᐊᐅᑎᓕᓗᓐᓄᕐᔪᐊᖅᑯᑦ)	ᑕᓗᐅᖏᓪᓯᖓᓂᖏᓐᓂᒃ ᑎᖏᐊᑦ ᖃᐅᔨᓴᖃᑎᒋᐅᓂᖏᓐᓂᒃ
2011	ᓕᖅᓚᐊᖏᓪ-Leafloor (ᑲᓇᑕᒥ ᐅᒪᔪᓕᓗᓐᓄᑦ, ᐊᐅᑎᓕᓗᓐᓄᕐᔪᐊᖅᑯᑦ)	ᐃᒪᖏᐅᓯᓂ ᑎᖏᐊᑦ
2014-2015	ᓂᓴᓪ, ᐅᐃᓕᔭᓴᓴ-Nissley, Williams (ᔪᓂᒪᓯᑎᑎ ᑎᐊᓪᓚᐅᑦ)	ᑲᖓᐃᐃᑦ ᖃᐅᔨᓴᖃᑎᒋᐅᓂᖏᓐᓂᒃ
2014-2020	ᖃᐅᔨᓴᓪ-Rausch (ᑲᓇᑕᒥ ᐅᒪᔪᓕᓗᓐᓄᑦ, ᐊᐅᑎᓕᓗᓐᓄᕐᔪᐊᖅᑯᑦ)	ᓯᓚᖃᖅᐸᓯᐅᓂᖏᓕᓴᐃᓂ ᑎᖏᐊᑦ ᖃᐅᔨᓴᖃᑎᒋᐅᓂᖏᓐᓂᒃ
2017-2019	ᔪᖏᓂ-Gurney ᖃᐅᔨᓴᖏᓂᖅ ᐊᓚᒪᓗ ᓴᓇᖏᑎᐅᖅᓕᓕᓗᓐᓂᒃ, ᐊᐅᑎᓕᓗᓐᓄᕐᔪᐊᖅᑯᑦ)	ᑎᖏᐊᑦ ᓯᕆᖏᓇᖅᑐᖃᖏᓂᖏᓐᓂᒃᓂᒃ ᖃᐅᔨᓴᖃᑎᒋᐅᓂᖏᓐᓂᒃ

ᑲᓇᑕᓕᒪᒥ ᐅᒪᔪᖏᖃᕆᐊᖓᔪᖅᓯᑦ ᐃᑲᔪᓂᖏᓐᓂᖏᓂᐊᖅᑐᖅ ᓴᐳᓂᐊᓂᖏᓐᓄᒪᑦ ᐃᑕᖅᑯᓯᑐᓗᓴᐅᔪᓂᒃ ᑕᐃᒪᖓᑦᓗ ᐱᓕᓗᐅᑎᑕᐅᕙᖅᑐᖅᑐᒃ ᐃᓄᖓᓂᒃ, ᑕᒪᑯᔪᐊ ᓯᑦᓗ ᐃᓗᐃᖃᖏᓐᓄᑦ, ᓄᓇᑭᔅᐅᑕᓕᓇᓂᖏᓐᓂᒃ ᐊᓚᒪᓗ ᐱᐅᓯᑐᖅᑎᖅᑎᒍᑦ ᐃᓄᐅᑦᖃᐃᖅᑎᖓᓂᒃ, ᐃᖏᓂᐅᖃᐃᑯᔪᐊ ᐊᐅᑕᓭᑎᓕᐅᖅᑕᑐᖃᖏᓐᓂᒃ ᑲᑎᒪᓕᓭᓂᒃ ᐱᐅᓂᖅᓴᓂᒃ ᐱᓕᓗᐅᑎᑕᐅᕙᖅᑐᖅᑐᒃ. ᐱᑕᖅᑭᑕᐅᓂᖏᓐᓂᒃ ᐱᔪᓪᑕᐅᓯᒪᓂᖏᓐᓂᓪᓗ ᑕᐃᑯᔪᐊ ᐃᓄᐃᑦ ᐃᑲᔪᓯᐊᖅᓴᖏᓐᓄᑦ ᐊᖏᑭᑭᐸᑦ ᐱᔪᓐᖃᑦᑕᕐᓂᖅᓴᖏᓐᓂᒃ ᐊᐅᑕᓭᑎᓕᐅᖅᑕᑐᖃᖏᓐᓂᒃ ᐃᓄᖓᓂᒃ ᑕᒪᐃᓕᓴᐅᓂ

ᐃᕐᓂᐅᕐᖁᓴᖅ ᐊᐅᓪᓚᕐᑎᐅᕐᑕᐅᓂᕐᒧᑦ ᑲᑎᒪᔨᕆᓪᕈᕐᑐᑦ ᑕᒪᐃᓐᓂᕐᑎᐊᖅ ᑕᒪᐃᓂᖅᓴᐃᖅ ᐃᑲᔪᖅᓴᐅᕐᓂᕐᒪᑦ ᐊᓯᔨᕆᑕᐅᕐᓂᐃᖅᑐᓂᖅ. ᐊᐅᓪᓚᕐᑎᐅᕐᑲᑕᐅᕐᔾᑦ ᑲᑎᒪᔨᕆᕈᑦ ᑎᑎᕆᕈᕐᑲᑐᑦ ᑲᓇᒥᑦ ᐅᒪᓪᓚᔨᖅᑯᓐᓂᕐᒪᑦ, ᑐᓴᖏᓚᖅᑯᓐᓂᕐᒪᑦ, ᐊᓘᓚᓗ ᑭᓯᓕᕐᓂᒥ ᐃᓄᐃᓂ ᑲᑐᔨᕐᓴᑎᑎᕐᑐᓴᓂ (ᑕᒪᓂ ᔪᓕᐃ 06, 2015-ᒥ) ᐃᓕᓕᓚᖁᐅᓪᓚᔨᒃᑐ ᑐᓴᕆᕈᑎᓂᖅ ᐱᓇᓱᐊᔨᓕᑎᖅ ᑲᑲᓇᖅᑕᐅᓕᖅᑐᓂᖅ ᐃᓕᓕᓚᖁᐅᓪᓚᔨᓂᖅ ᑕᐃᖁᐅᓪᓚᔨᓂᖅ ᐃᓕᓕᓚᖁᐅᓪᓚᔨᓂᖅᓗᓐᓗ. ᐃᓗᓕᓂᕐᓕᑦ 13-ᒥ ᐃᓄᐃᓂ ᐃᑲᔪᕆᐊᖅᓴᕐᓄᓂ ᐊᖏᕐᕈᑎᑦ ᐅᓂᖅᑲᓂᕐᒪᑦ ᖃᓄᐃᓕᓕᐅᕈᑕᐅᓱᒋᓂᖏᕐᓂᖅ. ᐃᓕᓴᖅᓴᓐᓂ, ᑕᒪᖅᑯᓂᖅᑕᕐᓗᒍᓗᑎᒃᓴᐅᔪᓕᖅᑐᓂᖅᓴᐅᑎᖅ ᖃᐅᔨᓴᕐᑎᐅᕐᑕᐅᓴᖅ ᑐᓴᕆᕈᑎᓂᖅᓴᐅᑎᖅᓴᓂᖅᓗᓐᓗ ᑐᓴᕆᕈᑎᓂᖅᓴᐅᑎᖅᓴᓂᖅᓗᓐᓗ ᑐᓂᔭᕐᓂᒃᑯᑦ ᐃᓄᐃᓂ ᑲᑎᒪᔨᕆᕈᑎᓂᖅᓴᐅᑎᖅᓴᓂᖅᓗᓐᓗ ᑐᓴᕆᕈᑎᓂᖅᓴᐅᑎᖅᓴᓂᖅᓗᓐᓗ ᐃᓕᓴᕐᓂᐊᕐᓗᒍ ᑲᑎᒪᔨ

9 ᐊᐢᓂᖅᑕᐅᑦᑕᐃᓕᒪᑎᑦᑎᓂᖅ, ᐊᐢᓂᐊᖅᑐᓕᕆᓂᖅ ᐊᓪᓚᓗ ᐅᓗᕆᐊᓇᖅᑐᓕᕆᓂᖅ

ᐊᐳᑎᓕᕆᓂᕐᒥᒃ ᑐᐊᐃᔭᓇᖅᑐᖃᖅᖢᑦ, ᐊᑎᑐᖅ ᐅᖃᖅᖁᐅᓂᐊᖅᓴᑦ ᐅᑯᐊ
ᑲᓇᑕᒥ ᐊᐳᑎᓕᕆᓂᕐᔪᒪᑦ ᑐᐊᐃᔭᓇᖅᑐᓕᕆᔨᑯᑦ ᖃᐅᐅᔨᑎᑎᐃᔪᖏᑦ
ᓄᓇᕗᑦᑎᒍ ᓄᓇᕐᒥᒍ:
1-867-920-8130.

ᑲᓇᑕᒥ ᐅᒪᔨᓕᕆᔨᖏᑦ ᓴᓂᐢᓂᐊᖅᑎᐅᓂᐊᖅᑐᑦ ᐃᐢᓂᐊᓕᕆᑯᑯᐅᔭᕆᐊᖅᖃᓄᖅᓂᐢᖏᐢᓂᐢᓂᒃ ᐊᓪᓚᓗ ᐅᑐᐊᓇᖅᑐᖃᖅᑲᖏᓂᐢᖏᐢᓂᐢᓂᒃ ᐃᓄᓕᒫᓂᒃ, ᖃᐅᐅᔨᒪᑎᑦᑎᖃᑦᓴᑎᒃ ᓄᓇᒧᖓᐅᓕᖅᑐᑦᑕᓂᖅᖢᑦᓴᓐᓂᒃ ᐅᑐᐊᓇᖅᑐᓂᖃᖅᑐᒥᒃ, ᐊᐢᓂᐊᓕᕐᑕᐅᔨᓇᖅᑐᑐᓄᓴᓐᓂᒃ, ᐊᑐᑦᓴᑎᒃ ᑐᑭᓕᐅᑦᑎᐊᑐᐃᐢᓇᖅᑐᒥᒃ ᐊᑐᖅᑕᐅᔨᓇᖅᑐᒥᒃ. ᐊᓪᓚᒪᑕᐅᖅ, ᑲᓇᑕᒥ ᐅᒪᔨᓕᕆᔨᓕᒃᑯᑦ ᐱᓕᕆᔨᖏᑦ ᓴᓇᔨᖏᑦᓴ ᑕᒪᐃᐢᓂᕐᑐᐊᖅ ᐅᓴᖅᑐᑦᑕᑐᐊᖅᑐᑦ ᐃᓕᓴᕆᔨᓂᖅᑐᐊᖅᑐᓂᒃ ᐅᑐᐊᖅᑐᖃᖅᑐᑦᑕᐃᓕᒪᓕᕆᓂᖅᑐᓂᒃ, ᐅᑐᐊᓇᖅᑐᖃᖅᑐᓂᒃᑐᓂᒃᑐᓂᒃ ᐃᓐᓇᑦ, ᐱᓕᕆᖃᑎᒌᖏᑦᑐᓂᒃ ᐊᓪᓚᓗ ᐃᓐᓇᑦᓴᓐᓂ ᐊᔪᖏᑦᓴᓐᓂᒃ ᖃᐅᔨᓴᕆᔨᑎᑎᐃᔪᖏᑦᓴᓐᓂᒃ, ᐊᑐᖅᑕᐅᔨᓇᖅᑐᖃᖅᑐᑦᑕᐃᓕᒪᓕᕆᓂᖅᑐᓂᒃ ᐃᓐᓇᑦᓴᓐᓂᒃ ᐊᑐᖅᑕᐅᔨᓇᖅᑐᑐᓄᓴᓐᓂᒃ ᐃᓐᓇᑦ, ᐱᓕᕆᖃᑎᒌᖏᑦᑐᓂᒃ ᐊᓪᓚᓗ ᐃᓐᓇᑦᓴᓐᓂᒃ ᖃᐅᔨᓴᕆᔨᑎᑎᐃᔪᖏᑦᓴᓐᓂᒃ, ᐊᑐᖅᑕᐅᔨᓇᖅᑐᖃᖅᑐᑦᑕᐃᓕᒪᓕᕆᓂᖅᑐᓂᒃ ᐃᓐᓇᑦᓴᓐᓂᒃ ᐊᑐᖅᑕᐅᔨᓇᖅᑐᑐᓄᓴᓐᓂᒃ, ᐅᑐᐊᓇᖅᑐᓂᖃᖅᑐᒥᒃ ᐃᓐᓇᑦᓴᓐᓂᒃ ᐊᓪᓚᓗ ᐃᓐᓇᑦᓴᓐᓂᒃ ᐊᔪᖏᑦᓴᓐᓂᒃ ᖃᐅᔨᓴᕆᔨᑎᑎᐃᔪᖏᑦᓴᓐᓂᒃ ᐅᑐᐊᓇᖅᑐᖃᖅᑐᑦᑕᐃᓕᒪᓕᕆᓂᖅᑐᓂᒃ, ᐸᑦᓇᓱᒪᑦᑎᐊᑦᑕᑦᓴᑎᒃ ᐊᓪᓚᓗ ᐃᓐᓇᑦᓴᓐᓂᒃ ᐊᔪᖏᑦᓴᓐᓂᒃ ᖃᐅᔨᓴᕆᔨᑎᑎᐃᔪᖏᑦᓴᓐᓂᒃ. ᑭᓱᓕᕆᔨᖏᑦ ᐃᓐᓇᑦᓴᓐᓂᒃ ᐊᑐᖅᑕᐅᔨᓇᖅᑐᑐᓄᓴᓐᓂᒃ

9.1 ᓄᓇᕗᒻᒥᔪᖅᑐᑦ ᐊᕐᓂᐊᓕᕆᔨᒃᓴᓂᖅ ᒥᑎᓄᑦ

ᐊᖏᑕᖅᑭᑎᐊᖅᑎᒃ ᐊᕐᓂᐊᖅᑕᕆᓚᐅᓕᕆᔨᓂᖅ ᓄᓇᕗᒻᒥᔪᖅᑐᑦ ᑎᑎᖅᑲᓇᓕᑦ ᐊᑐᓕᕆᔨᖅᑐᖅ ᑕᐅᓐᓂ ᖃᐅᔨᖅᑕᓗᖏᒃ ᑕᒪᓐ 2005-ᒥ, ᐱᑕᖅᐃᓐᓇᓕᖅᓯᓂᓗ ᑕᐃᒪᖏᓂᖅ ᐊᖏᕿᔪᑕᒪᖅ. ᐃᓄᖏᓄᑦ, ᐊᐃᓂᐊᓕᕆᔨᑕᐅᔪᓐᓇᖅᓂᖏᑦ ᓄᓇᕗᒻᒥᐊᓕᔪᖅᑐᑦ ᐊᑎᑎᕐᑐᐃᓐᓇᖅ. ᑭᓯᐊᓂᓕᕆᔨᖅ, ᑕᒪᖏᒃ ᐃᓄᐃᑦ ᐅᔪᓴᖅᑐᑎᐊᕆᐊᖅᑲᖅᓱᒃ ᐊᐅᓚᑦᑎᓂᖏᓐᓂᖅ ᖃᐅᓄᐃᑐᑐᐃᓐᓇᖅᓂᖅ, ᑎᑭᓐᓄᓚᔪᓂᓕᓐᓂᖅ, ᑕᐃᒃᑯᓂᓂᐊᓕᖅᑐᖅ ᖃᐅᔨᓴᖅᑭᑐᖅ ᑎᑎᖅᑲᓇᓕᑦ ᐅᓂᐅᔪᓂᖅ ᐊᐃᓂᐊᖅᑐᓂ, ᑐᑭᓯᒪᔪᓂᒪᓂᓕᒃ ᖃᐅᔨᓴᖅᑭᑐᖅ ᑕᒪᒃᑯᐊ ᑎᑎᕐᑕᑦ. ᓄᑖᖅᓯᓪᓚᔪᓂᖅ ᖃᐅᔨᑎᑎᐊᖅᑲᒃᖃᓂᓕᒪᔭᐃᓕᑦ, ᐅᖃᖅᖃᐃᐅᓂᐊᕐᓇᖅᑐᑦ ᐅᑯᐊ bᓇᑕᒥ bᑐᔅᓴᖅᑲᑎᑦᑦ ᐅᒪᔪᓕᓚᓂᕐᒧᑦ ᐊᓐᓂᐊᖅᑐᕐᖃᖅᖢᓪᓕᐊᓂᒧᑦ Canadian Cooperative Wildlife Health Centre ᐅᓪᓗᒥ 1-800-567-2033.

9.2 ᐊᓄᐃᑦ ᑐᖅᑯᑕᐅᓪᓚᕐᓂᑦ

ᑭᓇᑐᐃᓐᓇᖅ ᑐᓴᑯᔪᖅ ᐊᓄᕐᒃ ᑎᒥᐊᖅᑎᖃᖏᓐᓂᒥ, ᐱᔨᔪᑎᖅᑲᖅᓱᓂ ᑐᐊᕆᖃᖅᑐᑐᒃ ᐱᐅᖅᑯᕆᐅᓪᓚᕐᓂᑐᑭᓯᓂᑦ, ᐅᖅᑲᐅᓯᖅᑎᓚᐊᖅᑲᖅᓴᓐᓂ ᐊᔪᖅᓇᑭᓐᓂᐊᖅᑐᑦ. ᐱᓗᐊᖅᑐᒃ ᐅᑯᐊ ᐅᖅᑲᐅᑎᕐᐅᔨᕆᐊᑯᓂ ᓄᓇᕗᒻᒥᔪᖅᑐᑦ ᑲᑎᒪᔨᖅᑎᑦᓯᐊᓂᖅ (867-925-8823) ᐊᓚᒪᓗ ᑕᐃᒃᑯᐊ ᐊᐅᑎᓕᓚᓕᔪᐊᒃᑯᓂ ᓯᓚᐅᑦᓗ ᐊᓯᔨᖅᑮᑦᓕᐊᓕᓚᓕᖅᑐᓂᒃ ᐅᒪᔪᓕᓚᓕᖅᑯᓂ ᐱᔪᓐᓇᐅᑎᓕᖅᑐᓂᒃᑯᑦ (ec.nupermisscf-cwspermitnu.ec@Canada.ca). ᓴᒃᑯᓕᐅᓂᖅᑐᓂ ᑐᓂᔨᐅᓗᑎᒃ ᐅᒪᔪᓕᐃᑦ ᑎᑎᖅᑲᓂᖅᑎᑦᓗᓂᑦ, ᒪᓕᑐᖅᓂᒃ ᓯᕆᐃᔪᖅᑲᖅᑎᑦᓗᒍ, ᐱᔨᑯᓕᐅᓂᖅᑐᑦᑐᒥᓗᓂᑦ ᑐᑭᓯᒪᔪᓕᖅᑲᖅᑎᑦᓗᓂᑦ ᒪᓕᑐᖅᓂᒃ ᐃᓄᐃᑦ 5.6.55-ᖅᑐᓂᓕᑐᓂᒃ, ᓄᓇᕗᒻᒥ ᓄᓇᕐᓂᓪᓕᒍ ᐊᖏᑕᑎᓂ. ᓇᓪᓕᐅᑲᑯᒪᓴᓂᕐᕆᐊᖅᑐᑦ ᐊᐃᐃᓕᑦ ᐊᖏᔪᓯᑲᑎᑦ ᑲᑐᔨᖅᑎᑦᓯᐊᓂᖅ ᓇᓂᓄᑐᖅᑭᑦᓂ ᑎᒥᐊᖅᑎᖃᖏᓐᓂᒥ, ᒪᓕᑎᓄᑎᒃ ᐃᓄᓕᖅᑐᖅ *12.3-ᒥᑐᑐᓂᒃ, ᐃᓄᐃᑦ ᐃᑲᔪᑲᖅᑭᑐᓄᓂ ᐊᖏᑕᑎᓂᒃ*. ᐃᓄᐃᑦ ᑎᒥᐊᖅᑎᖃᖏᓐᓇᒪᑦᓯᐅᔨᓪᓚᕐᓂ ᐊᔪᑕᓂᕐᓴᖅᑲᑕᐅᓯᒪᔪᓕᖅᑎᖃᖅᑐᑦ ᓇᓄᖅᑕᑕᓇᓂᕐᑐᓂᒃ ᐅᑭᐅᖅᑕᑕᓇᓂᕐᒥᑦ ᐊᓚᒪᓗ ᖃᐄᐅᔨᓴᖅᑎᑦᓯᓂᕐᒥᑦ ᐅᓇᓕᖅᑐᖅ ᐅᓇᓂᓕᖅᑐᖅ ᐅᐅᓂᖅᑐᐃᐃᓕᖅᑲᓂᒃ ᐃᓕᓐᓂᒥᓂᖅ.

10 ᐊᑐᓪᓂᔪᒃᑯᓪᓗᒍᓂᖅ

ᐊᐅᓪᓚᑕᐅᓂᖏᓐᓂᖅ ᐊᑐᖅᑕᐅᔭᕆᐊᓯᒃᓴᓪ ᒪᓪᓕᖏᓪᓚᑕᐅᔪᓂᖏᓐᓂᖅ ᑎᖅᑎᐊᓪᓚᑕᓂᖅᓱᒍᓪ ᒪᓪᓗᑕᐊᓪᓚᑕᓂᖅ ᐊᒃᑎᐊᖅᑐᓂᖅ ᓴᖅᑭᑐᖅᑐᓂᖅ ᐊᔪᑎᓪᓚᕆᐊᖃᖅᑯᓂᖅ ᐅᓪᓗᕆᓚᐅᔭᖅᓯᒪᓂᖅ ᐃᓕᓴᓪᓚᕆᓂᖅ ᓴᖅᑭᑐᖅᑐᓂᖅ ᐊᐅᓚᓂᑎᐅᑎᑕᐅᔪᓂᖅ ᓴᖅᑭᑐᖅᑐᓂᖅ ᐃᓕᓴᓪᓚᕆᓂᖅ ᐊᐅᓚᓂᑎᐅᑎᑕᐅᔪᓂᖅ ᓴᖅᑭᑐᖅᑐᓂᖅ ᐃᓕᓴᓪᓚᕆᓂᖅ ᐊᑐᖅᑕᐅᔪ

ᐅᕐᓂᓗᓄᓕ ᓇᖏᑦᓂᖅ ᐅᖅᑲᐅᑎᓴᐅᓗᑎᑲ ᓇᑦᑕᐊᖏᓐᓄᑐᓇᖅ ᐃᖅᓂᐅᖅᓱᐊᕐᓂ ᐊᐅᓚᑦᑎᓴᐅᖅᑕᐅᓂᖅᓗᑦ ᑲᑎᒪᓴᕆᓇᖏᓐᓄᓕ ᓴᑦᑕᓂᓂ.

11 <ᕿᓇᐅᑦᑎᑦ ᐊᑐᓕᖅᑎᑕᐅᓂᖅᓯᒋᑦ

ᑲᓇᑕᒥ ᐅᒪᔨᓕᓗᓐᓂᑦ ᐊᒻᒪᓗ ᐃᕿᓂᐅᕿᖏᓐᓄᑦ ᐊᐅᓪᓚᑎᓕᐅᕿᑦᑐᑕᐅᓂᕿᒧᑦ ᑲᑎᒪᓐᕿᓭᓪᓕᐊᖅᑐᑦ ᐊᐅᓪᓚᑎᓂᕿᒧᑦ ᐊᐅᓪᓚᑎᑕᐅᓂᖅᑐᑦ ᐊᐅᓪᓚᑎᓂᕿᒧᑦ <ᕿᓇᐅᑎᓂᖅ ᑕᒥᓂ ᐊᑯᓂᐅᓂᖅᕿᓂᐊᖅᑐᒧᑦ ᕿᑯᓪᓗᓄᑦ ᐊᕿᕋᔪᓄᑦ. ᐊᑐᓕᖅᑎᑕᐅᓂᖅᑎᑦ ᒪᓪᓂᓂᐊᖅᑐᑦ ᐊᕿᕋᔪᑕᒥᖅ ᐱᓪᓚᑦᑎᓕᖅᑐᓂᓂᖅᑐᓂᖅ <ᕿᓇᐅᑎᑎᓯᒪᓕᖅᑐᓂᖅᑐᓂᖅ ᐊᒻᒪᓗ ᐱᓪᓚᑦᑎᓂᕿᒪᕐᓇᓂᖅᑎᑦᑐᓂᖅ ᐊᒻᒪᓗ ᐲᓇᐅᔨᖅᓴᖅᑕᐅᓯᒪᓂᖅᑐᓂᖅ. ᑲᓇᑕᒥ ᐅᒪᔨᓕᓗᓐᓂᑦ ᐊᒻᒪᓗ . ᐊᐅᓪᓚᑎᓂᐅᕿᖏᓐᓄᑦᑕᐅᓂᓪᓕᑦ ᑲᑎᒪᓐᕿᓭᓪᓕᐊᖅᑐᑦ ᐊᑐᖏᑦᑕᕐᓂᖅᓴᐅᓂᐊᖅᑐᑦ ᐃᓕᑭᒋᕿᖏᓐᓄᑦᑕᕐᓇᖅᑕᖅᑎᓂᖅᑐᓂᖅᑐᓂᖅ ᐊᐅᓪᓚᑎᓂᖅᑐᖏᓐᓄᑦᑐᖅ.

ᐊᐅᓪᓚᑎᓂᐅᕿᖏᓐᓄᑦᑕᐅᓂᓪᓕᑦ ᑲᑎᒪᓐᕿᓭᓪᓕᐊᖅᑐᑦ ᐊᑐᓂᖅᑎᑎᓂᖅᑐᑦ ᒪᓪᓂᓂᐊᖅᑐᑦ ᑕᒪᐃᓐᓄᑦ ᐱᓪᓚᑦᑎᓂᕿᒪᕐᓇᓂᖅᑎᑦᑐᓂᖅᑐᑦ ᐃᓕᓴᕐᓂᐊᓕᓂᓪᓕᑦ (ᐃᓄᐃᑦ *ᐃᑲᓴᐊᕿᖏᓐᓄᑦᓂᓪᓕᑦ ᐊᓐᑎᑕᑎᑦ ᐃᓗᓪᓚᓐᓂᑦ 2.1.9*). ᑲᓇᑕᒥ ᐅᒪᔨᓕᓗᓐᓂᑕᐅᕿᖅ ᕿᑲᐅᐱᐊᖅᑐᑕᓂᐅᐊᖅᑐᓂᖅ ᐊᑐᓂᖅᑎᑎᑕᐅᓂᖅᑐᖏᓐᓄᑦᑐᖅ <ᕿᓇᐅᑎᑦ ᑕᓪᓕᒪᐃᑦ 5 ᐊᕿᕋᔪᐃᑦ ᐊᓐᑎᑕᑎᑕᐅᓪᓚᐅᕿᓂᖅᑎᓂᖅᑐᑦ ᐊᒻᒪᓗ ᑭᖓᓂᐊᓕᑦ ᕿᑯᓪᓗᔪᑦ 10 ᐊᕿᕋᔪᐃᑦ ᐅᓴᕿᖓᑕᒃ, ᒪᓕᖅᓗᓂᖅ ᐊᐅᓪᓚᑎᓂᓪᓕᑦ ᐱᓪᓚᑦᑎᓂᕿᒪᕐᓇᓂᖅᑐᖏᓐᓄᑦ ᐃᓕᑕᒪᓴᐅᓯᒪᔪᓂᖅᑐᑦ ᑕᐃᑲᓂᐊᐸᖅᓯᒪᓂᖅᑐᖏᓐᓄᑦ ᑕᑯᑎᑕᑎᐱᔪᑎᕿᖏᓐᓄᑦᑐᖅ ᑕᐃᑎᑦ 10-ᒥ ᐃᓗᓪᓚᖅᑎᓂᖅᑐᑦ. ᑕᒃᑯᐊ ᑎᑎᕿᖅᓯᒪᓂᖅᑎᑦ ᐊᐅᓪᓚᑎᓂᓪᓕᑦ <ᕿᓇᐅᑎᓂᖅ ᐊᓯᔾᔨᓗᑕᐅᕝᕕᖓᖅᑎᑦᑐᓂᖅ ᐊᕿᕋᔪᑕᒥᖅ ᐱᓪᓚᑦᑎᓂᕿᓇᓂᖅᑕᕐᓂᖅᑐᖏᓐᓄᑦ <ᕿᓇᑕᐅᐊᖅᑐᓂᖅᑐᑦ, ᐃᑲᔪᕿᓂᖅᑎᖏᓪᓕᑕ ᓴᕿᑭᑎᑕᐅᓯᒪᓂᖅᑐᓂᖅᑐᑦ ᓄᑎᖅᑎᔪᑎᑎᑭᐅᓂᖅᑎᑦ ᓯᐅᑦᑐᑕᐅᓯᒪᓂᖅᑐᓂᖅᑐᑦ.

ᑲᓇᑕᒥ ᐅᒪᔨᓕᓗᓐᓂᑕᐅᓂᓪᓕᑦ ᑲᑎᒪᓐᕿᓭᓪᓕᐊᖅᑐᑦ ᐊᐅᓪᓚᑎᓂᐅᕿᖏᓐᓄᑦ

ᐊᓘᓵᖅᓯᓯᒪᔪᓂᖅ ᑕᑯᑎᑎᓴᓪᓚᓂ 10: ᑕᒡᓕᒪᓄᑦ ᐊᖏᕐᓯᒍᓄᑦ ᐊᐅᓚᓯᑎᓂᖏᓐᓇᑦ ᕿᖏᐅᑎᖅᓴᖅᓂᓂᖅ ᐊᑐᓵᖅᑎᑎᓂᐅᓕᒡᓕᐊᓂᖅ

ᐱᓕᕆᐊᓂᖅ	Year				
	1	2	3	4	5
ᐅᖃᐅᓴᖃᑕᖏᓂᖅ ᒥᓂᑕᕆᓂᖅ ᐊᓂᑎᓕᕆᓂᖏᓐᓄᑦ, ᓇᓱᒪᓴᓯᖏᓐᓄᑦ, ᑕᒪᐃᓂᖅ ᐱᓪᓚᕐᓂᖏᓐᓄᑦ ᐊᐅᓚᑕᖏᓂᖏᓐᓄᑦ ᕿᖏᐅᑎᖅᓴᖅᓂᓂᖅᓯᒪᓕᒃ; ᐅᓴᓴᖅᑐᑎᐊᖏᔪᓂᑎᓂ ᐃᓕᓴᕆᓂᖅᓴᖅᑐᑦ ᖃᓄᐃᓕᓴᕆᓂᖅᓴᖅᑐᑦ ᐅᖃᐅᓯᐅᓯᒪᓪᓚᑦ ᑲᑎᒪᔨᓴᕐᓂᖅᓴᖅᑐᑦ ᐃᓚᖓᑕᐅᓪᓗᐃᓚᐃᓕᓂ ᐃᓚᐃᓕᑦ ᖃᐅᔨᓴᕐᓂᖅᓴᖅᑐᑦ (ᐃᓚᐃᓕᑦ ᐃᑲᔪᓯᐊᓴᖅᓂᖅᓚᐃᓕᓂ ᐊᖏᕐᕈᑎᑦ ᐃᓗᓕᖅᓂᖅ 3.3).	X	X	X	X	X
ᐱᔨᑦᓴᖅᑕᐅᓴᖅᑕᕆᓗᓂᑐᖅ ᓴᖅᓚᐃᓕᓕᐅᕈᑕᐅᓂᖏᓐᓄᑦ ᐊᐅᓚᓯᑎᐅᕐᓯᖅᑕᐅᓂᖏᓐᓄᑦ ᑲᑎᒪᔨᓴᕐᓂᖅᓴᖅᑐᑦ ᐃᓚᐃᓕᑦ ᐃᑲᔪᓯᐊᓴᖅᓂᖅᓴᖅᑐᑦ ᐃᓚᐃᓕᑦ ᐃᑲᔪᓯᐊᓴᖅᑕᕐᕕᒃᓴᖅᓂᖅᓚᐃᓕᓂ ᐃᓚᐃᓕᑦ ᐃᑲᔪᓯᐊᓴᖅᓂᖅᓴᖅᑐᑦ ᐃᓗᓕᖅᓂᖅ

ᐱᓕᕆᐊᓂᖅ	Year				
	1	2	3	4	5
ᐃᓕᐅᖅᑕᕐᓗᑎᒃ ᐊᐃᒃᑐᖅᓯᒪᓂᖅᑭᓂ ᐱᓕᕆᐊᖏᓪᓕᓕᐊᖅᑐᓂᒃ ᐱᑎᐊᖅᑎᑕᐅᓕᖅᑐᓂᒃᓗᓐᓄᑦ ᐃᑲᔪᖅᓯᐃᓂᕐᒧᑦ ᓴᓇᓂᐊᓕᖅᑕᐅᓂᖏᓐᓄᒃ ᑎᖅᑎᐊᖅᖃᐃᐅᕐᓴ, ᓄᓇᖅᑭᓂᖓᓗ ᐊᐅᓗᐊᓂᑐᑐᓂᒃ.	X	X	X	X	X
ᑎᑎᖅᑯᓂᖅᑲᐃᓇᓕᓐᓂᒃ ᐃᓚᓇᐅᔪᖅ ᐱᑕᖅᖅᓂᖅᓂᒃ ᐃᒃᓗᑕᓪᓚᑦ ᑎᖅᑎᐊᖅᖃᐃᖅᑎ, ᐱᓕᕆᖅᑎᑎᒋᓗᑦ ᐊᐃᐃᓴᑦ ᐊᐃᒍᓇᔭᑎᑦ ᑲᑐᔾᔨᖅᑎᑎᒋᓕᑦ	X	X	X	X	X

11.1 ᐊᐅᓪᓚᕐᒪᓂᖏᑦ ᐸᖏᓐᓄᐅᑎᑦ ᐋᖅᑭᒋᐊᖅᑕᐅᓐᓄᐊᖅᓯᓂᖅᑐᑦ

ᑲᓇᑕᒥ ᐅᒪᓚᑦᑕᓱᓕᒃᑯᑦ ᐋᖅᑭᒋᐊᖅᓯᓚᖅᑐᖅᓯᖅ ᐊᐅᓪᓚᕐᒪᓂᖏᑦ ᐸᖏᓐᓄᐅᑎᓂᒃ ᖃᖅᖏᒐᔪᐃᖅᓯᖅ. ᑭᓇᐯᔪᑐᐃᓇᐃᓐᓇᑦ ᓴᖅᒪᓕᒃᓴ ᐃᓇᓴᓪᓗᓂ ᐊᐅᑐᖅᑕᐅᓯᒪᓚᓂᒃ ᑕᐯᔪᓐᓇᐅᑎᓂᒃ ᐊᐅᓪᓚᕐᒪᓂᖏᑦ ᐸᖏᓐᓄᐅᑎᓄᑦ ᑐᐯᓯᖅᑭᓇᖅᑐᑦ ᐱᕈᐊᖅᑎᑕᐅᕐᓯᔪᓚᐅᑎᒃ ᐋᖅᑭᒋᐊᖅᑕᐅᕐᓯᔪᓚᐅᖅᓱᒃ ᐅᖅᖃᖅᓴᐅᔪᓂᖅᑎᒃ ᑲᓇᑕᒥ ᐅᒪᓚᑦᑕᓱᓕᒃᑯᑦ (ᐃᓄᐃᑦ ᐃᑲᓴᓯᐊᕐᓴᖏᓐᓄᑦ ᐊᖏᑕᐅᓴᑦ *ᐃᓗᓕᑦ 3.7.1*). ᐊᐅᓪᓚᕐᒪᓂᖏᑦ ᐸᖏᓐᓄᐅᑎᑦ ᑭᖑᓂᐊᒍᓐᓇ ᐋᖅᑭᒋᐊᖅᑕᐅᕐᓯᒪᓂᖏᓐᓂ ᑎᑎᕋᖅᑕᐅᕐᓯᒪᓚᓴ ᐊᖏᕋᖅᑕᐅᕐᓴᓗᕋᖅᑐᓂᖅ> ᓄᓇᕗᒻᒥᐅᑦ ᐅᒪᓚᑦᑕᓱᓕᒃᑯᑦ ᐃᓚᖓᔪᖅᑐᑦ.

ᑕᐃᒃᑯᐊ ᐃᑎᓂᐅᕐᓴᐃᒃᑯᑦ ᐊᐅᓪᓚᕐᒪᓕᐅᕐᖏᓐᓂᖅᑐᑦ ᑲᑎᒪᓕᕋᓱᒃᑐᓂ ᐱᓕᕆᖅᓯᓂᐊᖅᑐᑦ ᖃᓄᐃᓕᐅᕐᓂᖅᑐᑦ ᐋᖅᑭᒋᐊᖅᑕᐅᕐᓯᒪᓂᖏᓐᓂᖅᑐᖅᓯ ᐋᓐᓂᖅᑕᐅᕐᓯᒪᓚᓴᖅᑐᑦ ᐊᓯᔪᓐᓇᖅᑕᐅᕐᓂᖏᓐᓂᑦᓴᓂᒃ ᐊᐅᓪᓚᕐᒪᓂᖏᑦ ᐸᖏᓐᓄᐅᑎᑦ. ᑲᓇᑕᒥ ᐅᒪᓚᑦᑕᓱᓕᒃᑯᑦ ᑲᑎᒪᓕᕋᓱᒃᑐᓂ ᐊᓯᔪᓐᓇᖅᑕᐅᕐᓂᖏᓐᓂᑦᓴᓂᒃ ᖃᑎᒌᖅᑕᐅᕐᓂᖏᓐᓂᑦᓴᓂᒃ ᐸᖏᓐᓄᐅᑎᑦ. ᐊᓯᔪᓐᓇᖅᑕᐅᓚᑦ ᖃᑎᒌᖅᑕᐅᕐᓂᖏᓐᓂᑦᓴᓂ ᐊᖏᕋᖅᑕᐅᕐᓂᖏᓐᓂᒡᓗ ᒪᓕᖏᓐᓂᖅᑕᐅᖅᑐᖅ ᐃᓗᓕᑦ *5.3.34(c)* ᐊᒻᒪᓗ *5.3.16* ᒪᓕᑐᖏᓐᓂᖅᑐᖅ ᐃᓗᐊᓂ ᓄᓇᕗᒻᒥ ᓄᓇᕆᔨᓴᑦ ᐊᖏᕋᐅᑎᑎᒍᑦ.

11.2 ᐊᐅᓪᓚᕐᒪᓕᐅᕐᓯᒪᓂᖏᑦ ᑎᓕᖅᐅᓯᒪᓚᓂ

ᐊᐅᓪᓚᕐᒪᓕᐅᕐᓂᖅ ᑲᓇᑕᐅᐸ ᒥᓂᑦᑕᖑᑦ ᐊᐋᑎᓕᓚᕐᓯᒪᓂᖏᑦ ᒪᓕᖅᑕᐅᓗᑎᒃ *ᑲᑎᒪᓕᐅᑕᐅᐋᖅᓯᖏᓐᓂᑦ ᑎᑭᓪᓚᕐᓯᓚᓂᖏᑦ ᒪᓕᓚᐃᓕ ᐊᒻᒪᓗ ᑎᑭᓪᓚᕐᒥᔪᖅᓴᓂᑦ ᒪᓕᒪᓗᑐᐃᓕᑦ*, ᐱᓕᕆᓇᐅᑕᐅᓂᖏᓐᓂᖅᑐᖅᓯ ᐊᐋᑎᓕᓚᕐᓯᒪᓚᐅᖅᑐᑦ ᐱᓕᕆᓇᐅᑕᐅᓂᖏᓐᓂᖅᑐᖅᓯ ᐊᐅᓪᓚᕐᒪᓂᖏᑦ ᐸᖏᓐᓄᐅᑎᑦ

12 ᐱᓕᕆᖅᑎᑦᑎᐊᑭᓐᓇᖅᓂᖅ

ᑲᔪᓯᓐᓇᑦᑎᐊᖅᓂᐊᑭᑎᑎᒃ ᑕᒃᑯᐊ ᐊᐅᓪᓚᑦᑎᓂᖅᓗᒍ <ᖅᓇᐅᑦᑐᑦ, ᑐᖏᑕᖅᑭᒪᓕᖅᓂᐊᖅᓂᒃᑐᑦ ᐱᓕᕆᖅᑎᑦᑎᐊᑭᓐᓇᖅᓂᕐᓂᖅᓂᒃ ᑐᑭᓯᓂᐊᑭᑕᐅᕐᓂᖅᓂᕐᓂᖅᓂᒃᓗ ᑲᓇᑕᒥ ᐅᒪᔪᓚᐅᔾᔨᑯᑦ, ᖃᓪᓗᓂᑦᑐᐅᓪᓗ ᐃᓄᖅᑎᑦᑎᓂᖅ, ᐊᔪᕐᓗᒍ ᑲᑎᒪᔨᖅᑕᕐᓂᖅ ᐱᓕᕆᖅᑎᑦᑎᐊᑭᓐᓇᖅᓂᕐᓂᖅᓂᒃ ᑐᑭᓯᐅᑎᖅᑕᕐᓂᖅ ᐱᓕᕆᖅᑎᑦᑎᐊᑭᓐᓇᖅᓂᕐᓂᖅ, ᐃᓄᖅᑎᑦᑎᐊᑭᓐᓇᖅᓂᕐᓂᖅᓂᒃᓗ ᑲᑎᒪᔨᖅᑕᕐᓂᖅᓂᒃᓗ ᐱᓕᕆᖅᑎᑦᑎᐊᑭᓐᓇᖅᓂᕐᓂᖅᓂᒃᓗ ᑲᑎᒪᔨᖅᑕᕐᓂᖅᓂᒃᓗ ᐱᓕᕆᖅᑎᑦᑎᐊᑭᓐᓇᖅᓂ

ᐊᓚᒐ ᑲᑎᒪᖅᓯᐊᕈᑕᐅᓕᖅᑐᓂᒃ ᐊᓚᒐ ᐃᓕᖅᑲᓪᓗᑎᒃ ᑕᒪᐅᔪᓐ ᑲᓱᓕᔪᓐᓇᖅᑐᑦ ᑲᒪᒋᔭᐅᓕᖅᑐᑦ ᓄᓇᕗᒻᒥ ᐅᒪᓴᓂᒃ, ᑕᐃᔪᓂᖏᓐᓇ ᒪᓕᕐᐊᖅᑲᓪᓗᑎᒃ ᓄᓇᕗᒻᒥ ᓄᓇᕆᓂᖅᓯᒪᔪ ᐊᖏᕐᕉᑎᓂᒃ. ᐃᓕᓴᐅᓂᒃ ᐱᓗᐊᖅᑐᑦ ᑐᓴᕐᓴᖅᓯᒪᓕᕐᓂᖅ ᓇᓕᒧᑐᓄᖅᑐᓂᒃ ᐅᒪᔨᓕᖏᓂᖅᑎᑦ ᐊᐅᓚᕋᐅᓂᖅᖢᓂᒃ, ᒪᓕᕐᐊᖅᑐᑦᑐᓂᒃ ᒪᓕᑎᒪᐃᔨᓂᒃ, ᐊᒻᒪᓗ ᐊᑐᖅᑕᕆᐊᖅᑐᕐᓂᒃ ᖃᓄᓪᓚᓄᓂ ᖃᐅᔨᓴᕐᓂᕆᔨᕈᑎᑐᓂᒃ ᐊᒻᒪᓗ ᐃᓄᐃᑦ ᖃᐅᔨᒪᔭᑐᑦᑐᓂᒃ, ᐃᓅᒪᑎᔭᑐᓂᒃ ᐊᒻᒪᓗ ᐱᔨᑕᑎᖅᑕᐅᓂᒃ ᐃᓄᐃᑦ ᓄᓇᕗᒻᒥᑦ ᐃᓄᓐᓂᒃ ᐊᐅᓚᕋᐅᓂᖅᑎᑦ ᐃᓄᐃᑦ ᓄᓇᕗᒻᒥᑦ.

ᓄᓇᕗᓂ ᑲᕈᒪᐃᑦ ᐱᓕᕆᓚᖓᑕ ᐃᓕᖅᑯᓯᓕᕆᓂᖅᓗᑦ ᐱᒻᒪᓚᐅᑎᑕᐅᔨᓂᓗ ᐃᓕᑕᕆᔨᐅᓯᒪᔨᓕᕆᓚᐅᔭᓂ ᐱᓕᕆᐊᖅᑯᓂ ᑲᓇᔨᐅᖅᑲᑕᓂᖓᓂᕐᓂᓂᖅ ᐊᑐᓕᖅᑎᑕᐅᕈᓐᓇᖅᑐᓂᓗ ᐊᑐᐊᖏᐃᓂ, ᐊᑐᑎᓴᒻᒪᓂ ᐱᓕᕆᓴᒻᒪᓂ ᐊᒻᒪᓗ ᐱᔨᑎᖅᑎᑐᑎᓂ ᐃᓄᖅᓂᒃ. ᑕᒪᓴᐃᑦ ᐱᔨᑎᖅᑎᓯᒪᔪᓂᒃ ᐊᓕᓴᕐᓂᖅᓴᓕᖅᑐᓂᖅᓂᖅ ᐃᓕᓴᕐᓯᓕᖅᑐᓂᒃ, ᐅᖅᓴᐅᔨᓕᖅᑎᑕᐅᔪᓂᒃ, ᐱᒻᒪᓕᕆᐅᓂᖅᓴᓕᖅᑐᓂᒃ ᐊᒻᒪᓗ ᖃᐅᔨᓂ ᐱᓕᕆᓚᖓᑕᓂᕐᓂᓂᖅ ᐃᓕᓴᕐᓯᓕᕆᓂᖅᓴᓕᖅᑐᓂᒃ, ᐅᖅᓴᐅᔨᓕᖅᑎᑕᐅᔪᓂᒃ, ᐱᒻᒪᓕᕆᐅᓂᖅᓴᓕᖅᑐᓂᒃ ᐊᒻᒪᓗ ᖃᐅᔨᓂ ᐱᓕᕆᓚᖓᑕᓂᕐᓂᓂᖅ ᐃᓕᓴᕐᓯᓕᕆ

13 ᐅᖃᓕᒫᓂᐊᕆᔨᒪᔭᖏᓂᒃ ᑐᑭᓯᐊᖃᖅᐳᑎᔨᓂᐅ

Δ>ᓪᕼᐊᒡ-Abraham, D. M. & Ankney, C. D., 1986. *Summer birds of East Bay, Southampton Island, Northwest Territories,* s.l.: Canadian field-naturalist 100(3):180-18.

ᐊᓂᔭᐅᓴᓴ-Alisauskas, R. T., Dufour, K. W. & Leafloor, J. O., 2018a. Midcontinent lesser snow goose Chen caerulescens caerulescens. In: A. D. Fox & J. O. Leafloor, eds. *A global audit of the status and trends of Arctic and Northern Hemisphere goose populations..* Akureyri: CAFF International Secretariat.

ᐊᓂᔭᐅᓴᓴ-Alisauskas, R. T., Dufour, K. W. & Leafloor, J. O., 2018b. Ross's goose Chen rossii. In: A. D. Fox & J. O. Leafloor, eds. *A global audit of the status and trends of Arctic and Northern Hemisphere goose populations..* Akureyri: CAFF Internation Secretariat.

ᐁᓂᔭᵒ-Allison, L., 1977. *Migratory bird sanctuaries in the Northwest Territories - a background paper,* s.l.: Unpubl. rep., Can. Wildt. Serv., Edmonton. 3 vols. 370 pp. .

ᐊᖅᓪᓗᖏᒥ ᑎᖃᐊᐅᓂᓱᓂᕐᒧᑦ ᑲᑎᒪᔨᑦ-Atlantic Flyway Council, 2011. *Atlantic Brant Management Plan,* s.l.: s.n.

Bᐁᑦ-Batt, B. D., 1997. *Arctic ecosystems in peril: report of the Arctic Goose Habitat Working Group.,* s.l.: Arctic Goose Joint Venture Special Publication..

Bᐅᑦ-Bird, J., 1953. *Southampton Island,* Ottawa: Edmond Cloutier, Queen's Printer and Controller of Stationery.

ᕿᑦᓗ-Buttler, I., 2009. *Avian cholera among arctic breeding common eiders somateria mollssima: temporal dynamics and the role of handling stress in reproduction and survival,* Ottawa, Ontario. : M. Sc Thesis. Department of Biology, Carleton University. 83 pp.

CACNMP, 1990. *A review of the boundaries of Bird Sanctuaries in the Northwest Territories,* Ottawa, ON: s.n.

ᓴ>ᵒᓂᐊᖅᑎᑦ ᐅᑭᐅᖅᑕᖅᑐᒥ ᐅᒪᔪᓂᒃ ᐱᑭᖅᑐᓂᓗ-CAFF, 2010. *Arctic Biodiversity Trends - Selected indicators of change.,* Akureyri, Iceland: CAFF International Secretariat.

ᑲᒃ>ᑦ-Campbell, M. & Boulanger, J., 2015. *Preliminary Report on The Long-Term Abundance Fluctuations of the Southampton Island Caribou Herd - 1978 -2015,* Arviat, NU: Nunavut Department of Environment, Wildlife Research Division.

ᑲᓇᑕᐅᓂ ᓄᖕᒍᓕᖅᑐᑦ ᐅᒪᔪᐃᑦ ᓴ>ᵒᓂᐊᖅᑎᑦ ᑲᑎᒪᔨᖏᑦ-Canadian Endangered Species Conservation Council, 2011. *Wild Species 2010: The General Status of Species in Canada,* s.l.: National Status Working Group.

ᑲᖏᑐ-Carter, N. A. et al., 2018. *Inuit knowledge about light geese in the Kivalliq region, Nunavut.,* s.l.: s.n.

ᑯᑦ-Coad, B. W. & Reist, J. D., 2004. *Annotated list of the Arctic Marine Fishes of Canada.,* s.l.: Canadian Manuscript Report of Fisheries and Aquatic Sciences 2674.

ᑲᓕᓐ-Collins, H., 1956. *Archaeological investigations on Southampton and Coats islands, N.W.T.* , s.l.: National Museum of Canada Bulletin No. 42:82-89.

ᑯᑦ-Cooch, F. & Barry, T. W., 1957. *Proposed Migratory Bird Sanctuaries Southampton Island, Keewatin.,* s.l.: CWSC 3498.

ᑎᑲᒪᐸ-Descamps, S. et al., 2009. *Costs of reproduction in a long-lived bird: large clutch size is associated with low survival in the presence of a highly virulent disease,* s.l.: Biology Letters 5.

ᑎᑲᒪᐸ-Descamps, S., Jenouvrier, S., Gilchrist, H. G. & Forbes, M. R., 2012. Avian Cholera, a threat to the viability of an arctic seabird colony?. *PLoS ONE,* 7(2), p. e29659.

Ṅ-Dey, C. J., Semeniuk, C. A. & Gilchrist, H. G., 2019. Climate change impacts on arctic seaduck colonies are mediated through changes in predator behaviour (in revision). *Proceedings of the Royal Society.*

Ṅ-Dey, C. J. et al., 2018. *Forecasting the outcome of multiple effects of climate change on northern,* s.l.: Biological Conservation 220.

DΔP-Dickie, M., Smith, P. A. & Gilchrist, H. G., 2014. The importance of survey timing on shorebird density estimates at East Bay, Nunavut, Canada. In: s.l.:s.n., pp. 394-401.

Dᐅᐳᐊᖅ-Dufour, K. W., Dickson, K. M., Castelli, P. M. & Leafloor, J. O., 2019. Atlantic Brant, Branta bernicla hrota.. In: A. D. Fox & J. O. Leafloor, eds. *Global Audit of the Status and Trends of Arctic and Northern Hemisphere Goose Populations (Component 2: Population accounts).* Akureyri: s.n., pp. X-Y.

Dᐅ>Δ-Dupuis, P., 1979. *Comments on the 1979 photographic survey of Lesser Snow Geese of the Eastern Canadian Arctic,* Ste-Foy, QC: Canadian Wildlife Service.

ᐊᖏᑎᓪᓗᓴᖅᓯᐊᐱᑯᑦ, 2016. *Environment and Climate Change Canada's input to the Nunavut Planning Commission regarding Key Habitat Sites for migratory birds in the Nunavut Settlement Area, May 2016 revision.,* Yellowknife, NT: Canadian Wildlife Service.

ᐊᖏᑎᓪᓗᓴᖅᓯᐊᐱᑯᑦ bᓇᑕᒥ-Environment Canada, 2013. *Bird Conservation Strategy for Bird Conservation Region 3 Prairie and Northern Region: Arctic Plains and Mountains.,* Yellowknife, Northwest Territories.: Canadian Wildlife Service, Environment Canada.

ᖃᓪᒥᖅ-Flemming, S. A., Calvert, A., Nol, E. & Smith , P. A., 2016. *Do hyperabundant Arctic-nesting geese pose a problem for sympatric species?,* s.l.: Environ. Rev. 24.

ᐸᵒṄᵒ-Fontaine, A. & Mallory, M., 2011. *Detection and Classification of Land Cover Classes of Southampton Island, Nunavut, Using Landsat ETM+Data,* Prairie and Northern Region: Occasional Paper No. 119, Canadian Wildlife Service.

ᑌᑕᵒGaston, A. J., Decker, R., Cooch, F. G. & Reed, A., 1986. *The distribution of larger species of birds breeding on the coasts of Foxe Basin and northern Hudson Bay, Canada.,* s.l.: Arctic Vol. 39 No. 4.

ᑎᐅᖅᑯᓗᒃ-Gilchrist, G. & Heise, K., 1997. *Studies on the Demography and Reproductive Ecology of Common and King Eiders Breeding in East Bay, Southampton Island—Second Annual Report 1997,* s.l.: Canadian Wildlife Service.

ᑎᐅᖅᑯᓗᒃ-Gilchrist, G. & Jassen, M., 2015. *East Bay Island Migratory Bird Research, 2015 Field Season Report,* s.l.: Environment Canada.

ᔪᖏᑐᖅ-Gratto-Trevor, C. et al., 2011. *Trends in Canadian shorebirds.,* Ottawa, ON.: Canadian Biodiversity: Ecosystem Status and Trends 2010, Technical Thematic Report No. 13. Canadian Councils of Resource Ministers.

Hᐊᐅᓴᓂ-Harms, N., 2011. *Avian Cholera in the Eastern Canadian Arctic: Investigating Disease Origins and Reservoirs,* s.l.: InfoNorth.

Hᐊᓄᓕ-Henri, D., 2007. *The Integration of Inuit Traditional Ecological Knowledge and Western Science in Wildlife Management in Nunavut, Canada: The Case of Avian Cholera Outbreaks among Common Eider Ducks in the West Hudson Strait and North James Bay Area. M.Sc Thesis, Oxford University*.

Hᐊᓄᓕ-Henri, D. A., Jean-Gagnon, F. & Gilchrist, G. H., 2018. *Using Inuit traditional ecological knowledge for detecting and monitoring avian cholera among Common Eiders in the eastern Canadian Arctic.,* s.l.: Ecology and Society 23(1): 22.

ᐃᓄᓕᓪᓚᕐᔪᑐᖅᑯᓕ-INAC, 1976. *Inuit Land Use and Occupancy Project, 1974 (Milton Freeman Research Ltd.),* s.l.: Department of Indian and Northern Affairs Canada.

ᐃᖃᓂᐅᕙᓱᑯᓕ-Irniurviit ACMC, 2017. *Site Visit Report,* s.l.: s.n.

ᐊᐃᕐᓴᓐ-Iverson, S. A., 2015. *Quantifying the demographic and population impact of avian cholera on northern common eiders in the face of ancillary threats and changing environmental circumstances.,* Ottawa, Ontario.: Ph. D. Thesis. Carleton University. 228 pp.

ᐊᐃᕐᓴᓐ-Iverson, S. A. et al., 2014. *Longer ice-free seasons increase the risk of nest depredation by polar bears for colonial breeding birds in the Canadian Arctic.,* s.l.: Proc. R. Soc. B.

ᐊᓄ-Jano, A. P., Jeffries, R. L. & Rockwell, R. F., 1998. *The detection of vegetational change by multitemporal analysis of LANDSAT data: the effects of goose foraging.,* s.l.: J. Ecol.

ᐊᓐᓴᓐ-Janssen, M. & Gilchrist, H. G., 2013. *East Bay Island Migratory Bird Research - 2013 Field Season Trip Report and Update to Collaborators,* Ottawa, ON: Unpublished Annual Summary Report.

ᑯᐱᓴ-Kerbes, R. H., Meeres, K. M. & Alisauskas, R. T., 2014. *Surveys of Nesting Lesser Snow Geese and Ross's Geese in Arctic Canada, 2002 – 2009.,* U.S. Fish and Wildlife Service, Washington, D.C. and Canadian Wildlife Service, Ottawa, Ontario: Arctic Goose Joint Venture Special Publication.

ᑯᐱᓴ-Kerbes, R. H. et al., 2006. *Surveys of Nesting Mid-continent Lesser Snow Geese and Ross's Geese in Eastern and Central Arctic Canada, 1997-98,* s.l.: Environment Canada, Canadian Wildlife Service, Prairie and Northern Region.

ᑭᖅᑕᒥ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑦ ᓄᓇᕗᒻᒥᐅᑎᖓᑦ-Kivalliq Inuit Association Mapping Tool. [Online] Available at: http://www.kivalliqinuit.ca/environ2.html [Accessed 2014].

ᑯᑭᔾ-Kokelj, S. V. et al., 2005. *The Influence of Thermokarst Disturbance on the Water Quality of Small Upland Lakes, Mackenzie Delta Region, Northwest Territories, Canada,* s.l.: Permafrost and Periglac. Process. 16.

ᑯᑕᓂᐊᖅ-Kotanen, P. & Jefferies, R. L., 1997. *Long-term destruction of sub-arctic wetland vegetation by lesser snow geese.,* s.l.: Écoscience 4: 179-182.

ᓕᑐᕋᐠ-Lathrop, R. G. et al., 2018. *Mapping and modeling the breeding habitat of the Western Atlantic Red Knot, Calidris canutus rufa, at local and regional scales.,* s.l.: PANGAEA.

ᓪᑐᐊᕋ-Latour, P. B. et al., 2008. *Key Migratory Bird Terrestrial Habitat Sites in the Northwest Territories and Nunavut (3rd Ed.),* s.l.: Occasional Paper No. 114. Canadian Wildlife Service.

ᓪᖅ-Lunn, N. J. et al., 2016. *Demography of an apex predator at the edge of its range: impacts of changing sea ice on polar bears in Hudson Bay,* s.l.: Ecological Applications 26(5).

ᒪᓂᖏ-Manning, T., 1942. *Remarks on physiography, Eskimos, and mammals of Southampton Island,* s.l.: Can. Geog. Jour. 24 (1): 17-33.

ᒪᓂᖏ-Manning, T., 1967. *A report on the transfer of barren-ground caribou from Coats Island to Southampton Island, NWT,* Ottawa: CWSC Rept. 1143, Canadian Wildlife Service .

ᒧᑦᐱᐊᑲ-Mosbech, A. et al., 2006. Year-round movements of northern Common Eiders Somateria mollissima borealis breeding in Arctic Canada and West Greenland followed by satellite telemetry.. *Ardea,* 94(3), pp. 651-665.

ᓂᓪᓕ-Nissley, C., 2016. *Assessing the impact of Lesser Snow Goose and Cackling Goose competition on breeding Atlantic Brant,* s.l.: University of Delaware.

ᓄᓇᕗᑦ ᓄᓇᐃᑦ ᐊᑐᑎᒃᖏᑐᓂᑦ ᐸᖅᓇᐃᓕᑦ ᑲᒥᔪᓇᒃᑯᑦ-Nunavut Planning Commission, 2000. *Keewatin Regional Land Use Plan,* s.l.: s.n.

ᓄᓇᕗᑦ ᓄᓇᐃᑦ ᐊᑐᑎᒃᖏᑐᓂᑦ ᐸᖅᓇᐃᓕᑦ ᑲᒥᔪᓇᒃᑯᑦ-Nunavut Planning Commission, 2012. *Draft Nunavut Land Use Plan - Options and Recommendations,* s.l.: s.n.

<ᐅᑯ-Parker, G., 1974. *A population peak and crash of lemmings and snowy owls on Southampton Island, NWT,* s.l.: Can. Field-Nat. 88:151-156.

<ᐅᑯ-Parker, G. R., 1975. *An investigation of caribou range on Southampton Island, Northwest Territories,* Ottawa, ON.: Canadian Wildlife Service Report Series No. 33.

>ᑭᓴ-Perkins, D., Smith, P. A. & Gilchrist, H. G., 2007. *The breeding ecology of Ruddy Turnstones in the eastern Canadian Arctic,* s.l.: Polar Record 43, 135-142.

ᐱᑭᕆᐊᓐᓂᖅ-Provencher, J. F. et al., 2014. *Mercury and marine birds in Arctic Canada: effects, current trends, and why we should be paying closer attention,* s.l.: Environ. Rev. 22.

ᓱᒃ-Reed, A., Dupuis, P., Fischer, K. & Moser, J., 1980. *An aerial survey of breeding geese and other wildlife in Foxe Basin and northern Baffin Island, Northwest Territories, July 1979.,* s.l.: Canadian Wildlife Service Progress Notes No. 114.

ᓇᐅ-Riewe, R., 1992. *Nunavut Atlas,* s.l.: Canadian Circumpolar Institute and Tungavik Federation of Nunavut.

ᕿᓴ-Ross, M. V., Alisauskas, R. T., Douglas, D. C. & Kellett, D. K., 2017. *Decadal declines in avian herbivore reproduction: density-dependent nutrition and phenological mismatch in the Arctic,* s.l.: Ecology 98(7).

ᓴᐅᕕ-Sharp, C. M. & Abraham, K. F., 2010. *Status of Atlantic Brant breeding at East Bay, Southampton Island: 30 years later.,* Peterborough, ON.: Poster to the North American Arctic Goose Conference.

ᓂᒥᑦ-Smith, P. A., Elliott, K. H., Gaston, A. J. & Gilchrist, H. G., 2010. *Has early ice clearance increased predation on breeding birds by polar bears?,* s.l.: Polar Biol..

ᓴᒍᑦ-Smol, J. P. & Douglas, M. S., 2007. *Crossing the final ecological threshold in high Arctic ponds,* s.l.: PNAS.

ᐅᓴᐃᔨᑦ ᑲᓇᑕᒥ-Statistics Canada, 2016 Census of Population. *Coral Harbour, HAM [Census subdivision], Nunavut and Nunavut [Territory] (ᐊᐅᔭᑐᓂᒪᓴᓂ ᑕdᑎᕋᑎᕐᓴᑦ). Census Profile. 2016 Census.,* Ottawa.: Statistics Canada Catalogue no. 98-316-X2016001.

ᓴᑏᕐᓴᕐ-Stephenson, S. A. & Hartwig, L., 2010. *The Arctic Marine Workshop: Freshwater Institute,* Winnipeg, Manitoba: Canadian Manuscript Report Fisheries and Aquatic Science 2934.

ᓴᑏᕐᓴᕐ-Stephenson, S. M. & McCormick, K. J., 1986. *Draft East Bay Bird Sanctuary Management Plan,* Yellowknife, NT: Canadian Wildlife Service - Western and Northern Region.

ᓴᑕᐟ-Sutton, G., 1932. *The Exploration of Southampton Island, Hudson Bay,* s.l.: Memories of the Carnegie Museum, Vol. 12.

ᐅᖅᕼᐊᐅᑦ-Urquhart, D. & Schwcinsburg, R., 1984. *Polar bear: life history and known distribution of polar bear in the Northwest Territories up to 1981,* Yellowknife: Unpubl. Rept., Dept. Ren. Res., GNWT, Zone C, p.30-35.

ᕿᐟ ᓴᑐᐟ-Van Stone, J. W., 1960. *Notes on the Economy and Population Shifts of the Eskimos of Southampton Island.,* Toronto, ON.: Department of Anthropology, University of Toronto.

ᑎᑎᕐᒃᐃᓕᓴ ᐃᓕᓴᒪᓂᖅᓯᓐᓂ

ᑎᑎᖅᑲᐃᑦ ᐃᓕᓴᖅᓂᖓᓂ A: ᒪᓕᑐᖅᑎᒍᑦ ᐅᓂᓴᑲᖅᑕᐅᕐᑎᐊᖅᓂᖓᓂ ᓄᓇᐃᑦ ᓇᓵᓂᖓᓂ

ᒪᓕᑐᖅᑎᒍᑦ ᐅᖅᑲᐅᓯᐅᓯᒪᓕᒪᑦ ᐃᓴᐸᐊᖅᑕᐅᓯᒪᓐᓂᑎᖅ ᓇᖕᒥᓂᖅ ᑎᑎᕆᖅᓯᒪᓂᖓᓂ ᑕᐃᖅᑯᓇᖕᑐᑦ ᑎᖅᒥᐊᖅᑎᖃᖅᓄᓇᓄᑦ ᒪᓕᑐᓐᓂᑐᕆᔨᐸᓂᖅ (*C.R.C., c. 1036*; 1958) ᐃᒪᖓᓇᐃᓕᖕᖏᒪᔪᓂᖅ:

“ᖃᖅᑲᓇᐅᑲᑐᒃᑯᑦ ᑎᖅᒥᐊᖅᑎᖃᖅᓄᖅ


ᑕᒪᖅᑎᖅ ᐃᓕᑭᐅᕐᔪᓂ ᓄᓇᖅᓯᓄᓂ ᓴᓪᓕᐅᑦ ᖃᑭᑲᒡᓗᐊᓄᓂ ᐃᒪᖖᑭᓄᓪᓗ ᐄᐊᑦᓴᐃᓐᓇ ᑕᓯᐅᕐᔪᐊᖏᓪᓗᐊᓂᒃ, ᐊᐃᖅᑐᑲᖅᓯᒪᓕᖅᖏᑦ, ᓄᓇᕗᐊᖅᑭᑦ, ᐃᓕᑭᐅᕐᐅᓂᖓᓂ ᐱᓗᐊᖅᑐᑦᑕᖅ ᐅᓂᓴᑲᖅᑕᐅᕐᓄᑦ ᐃᒪᖓᓇᐃᓕᖕᖏᓚᓂᕆᐊᖃᓪᓗᓂᒃ:

*ᐱᑭᐊᖅᐃᓚᖅᑲᓂ ᑕᒧᐊᓂ ᖃᑎᖅᑐᐊᓂᒃ ᖃᐅᔨᓴᖅᐃᐅᓂ ᐱᖅᔨᑎᖅᑲᖅᐃᐅᑎᓂᒃ ᔪᐊᑦ ᓄᑭᐊᓂᒃ Gore Point-ᒥᒃ, ᑕᒪᓐᓇ ᐃᓕᑕᓴᐅᓯᒪᔪᖅ ᖃᐅᔨᓴᖅᐃᓚᐅ ᑕᑭᓂᖅᑲᖅᓯᓂ 15-Fᐃᑦᓂᒃ ᐱᖓᓱᐃᓕᓂᖅᑲᖅᓯᒪᔪᒃ ᖃᐅᔪᖅᒥᒃ ᒪᑭᑐᐊᖅᑲᖅᑕᒥᒃ ᑲᓇᓯᒪᔪᖅ, ᑕᒪᓐᓇ ᑐᑭᓯᓚᐅᖅᑕᒥᒃ ᑐᑦᓯᓖᓂᓂ ᐅᐊᖓᓇᖅᑕᒪᑕ ᐱᖓᓱᓇᖅᑐᐱᖓᓚᐅᖅᑕᒥᒃ ᑕᐅᑭᖓᓖ ᑎᑭᓗᐊᓂᖅᑐ

ᐱᑎᐊᖏᖅᑲᓕᓂ ᑕᒧᖅ ᖃᑎᖅ ᖃᓄᐊᓂᖏᑦ ᐱᖏᓂᖏᑦᑐᑦ ᐱᖏᓂᖏᓕᒃ ᓄᓇᕗᒻᒥᐅᑦ - *M'Clure Point*-ᒥᒃ ᑕᒪᓐᓇᑐᒥᒃ ᑐᑭᓯᐊᖅᑐᒥᒃ 64°4'-ᒥᒃ ᐊᓚᒐᓂ ᑲᓇᓗᐊ ᓴᓂᒍᑦ 81°17'-ᒍᑦ, ᐃᓕᑕᒪᓕᐅᕙᓚᐅᖅᓯᒪᓕᒃ ᖃᓄᐊᓂᖏᓕᒃ ᑕᑯᓂᖃᖅᑐᒥᒃ 15-Fᐃᑦᓂᒃ ᐱᖓᓱᐊᓕᕆᖃᖓᓚᐅᒥᒃ ᖃᐅᔨᖅᑐᒃ ᒪᑯᑕᐊᒥᒃ ᓴᓇᓕᓚᐅᑦ; ᑕᒪᖓᑦ, ᐅᐊᖓᓇᖅᑲᓱᖓᑕ ᑲᔪᓱᓂᑎᒃ ᓂᑎᐅᓂᖅᐸᓪᓴᖓᓄᑦ ᐃᓱᐊᓄᑦ ᒥᑯᔪᓄᑦ ᑕᐃᔪᑎᖃᖓᓕᒫᑦ ᖃᓄᖅᑕᒧᑦ ᑕᐃᔪᑎᖃᖓᓕᒫᑦ ᑐᑭᓯᒪᔭᖅ 64°8'-ᒍᑦ ᐊᓚᒐᓂ ᑲᔪᓱᑎᓂᔭ ᓴᓂᒍᑦ 81°34.5'-ᒍᑦ; ᑕᒪᖓᑦ ᑲᔪᓱᑎᓪᓂᔭ ᐱᖓᓇᖅᑲᓱᖓᓄᑦ 82.30'-ᒍᑦ; ᑕᒪᖓᑦ ᑲᔪᓱᑎᓕᕐᓂᔭ ᓂᑎᐅᓄᑦ 64°4'-ᒍᑦ; ᑕᒪᖓᑦ, ᓴᓇᓕᒍᑦ ᓂᑎᐊᒋᑦ ᑲᔪᓱᖓᓇᖅᑲᓱᖓᓄᑦ ᓄᐅᓪᓂᔭ ᑎᐱᓪᓂᔭ ᑐᓂᓯᒪᔭᖅ 64°4'-ᒍᑦ. ᑕᒪᖓᑦ, ᓴᓇᓕᒍᑦ ᓂᑎᐊᒋᑦ ᑲᔪᓱᖓᓇᖅᑲᓱᖓᓄᑦ ᓄᐅᓪᓂᔭ ᑎᐱᓪᓂᔭ 63°50'-ᓄᑦ ᐊᓚᒐᓂ ᓴᓇᓕᒍᑦ ᑲᔪᓱᑎᓪᓂᔭ 82°10'-ᒍᑦ; ᑕᒪᖓᑦ, ᐃᓄᓕᒍᑦ ᐅᐊᖓᓇᖓᓚᐅᒥᒃ ᑲᔪᓱᖓᓇᖅᑲᓱᖓᓄᑦ ᐱᖏᓂᖏᓕᒃ ᐅᖃᐅᓯᐅᒪᔪᓕᑦ ᑕᒪᖓᒥᒃ ᑲᑎᓪᓂᒃᑯᑦ ᑕᐃᔪᑎᖃᖓᓕᒫᑦ ᐅᖃᐅᓯᐅᒪᔪᓕᑦ ᑕᒪᖓᒥᒃ ᑲᑎᓪᓂᒃᑯ

Ↄϼᓯᑎᕋᑎᐊᑭᕋᑎᐊᖅᐅᐅᔪᓂ<>ᕋ ᐊᓗᒪᓗ ᑕᑯᔭᕆᐅᓪᓗᑎᒃ. ᑕᒪᕋᒥᒃ ᐊᓯᓪᒥᒪᖏᕋ ᑕᑯᔭᕆᕕᐃᕋ ᑦᒃᑯᓇᓂ ᐊᐅᓪᓚᕋᑎᓂᕐᒧᑦ <ᕋᓇᐅᑎᓂᒃ ᓴᕋᑭᐅᒪᑎᕋᑎᐅᖅᓴᖅ ᓯᓪᓚᔪᓂᒃ ᑕᒪᒪᓯᒪᔪᓂᖅᑐᓂᒃ ᑎᖏᒥᐊᖅᑲᕋᐅᒃ ᑭᖅᓴᖅᓯᒪᐅᓂᒃ.

Environment and Climate Change Canada
Environnement et Changement climatique Canada



2020

ᐃᑲᕐᑐᐊᖏᑦ ᑎᓴᒥᐊᖅᑲᖏᓐᓇᖅ ᐊᐅᓚᕆᑎᓂᖓᔪᓂ ᐸᖏᓐᓄᐅᑎᓪ

[ᓴᓇᐊᓗ 2020]

ᐃᓕᓴᕆᔨᖅ:

Ì°ᓇᐅᓕᖅᑐᑦᖅ ᓯᕗᓂᐊᒍᓪᓗ ᑲᑎᒪᔨᐅᓕᐅᖅᑐᑦ, ᑲᑎᒪᔨᐅᓕᖅᑐᑦᓪᓗ ᐃᔅᓂᐅᕐᖃᐃᖅᑯᑦ ᐊᕐᓇᖅᑐᑦᓯᓕᕐᓂᖅᓂᕆᔨᐅᔅᓴᖅᑕᐅᓂᔅᓯᒪᑦ ᑲᑎᒪᔨᕋᓱᒃᑐᑦ (ᐃᔅᓂᐅᕐᖃᐃᖅᑯᑦ ᐊᐅᓕᕐᑎᐅᔨᐅᖅᑲᑎᑦ) ᓴᕐᑭᑎᕐᓕᐅᖅᓴᓕᕐᓂᖅ ᐅᑯᓂᖅᑐ. ᑲᑎᒪᔨᐅᓕᖅᑎᓪᓗᒍᑦ ᐅᑯᐊ, ᓄᐊ ᑲᑦᓕᒃ, ᖃᓄᖅ-ᑭᕋᓂᓯᐊ ᑐᑭᐊᔅ, ᐅᐃᓕ ᐊᑕᓴᔅ, ᐊᑯᓂ ᓂᖅᑎᐅᖅᓱᓂᖅ. ᐊᑕᓴᐅᖅᑎᓪᓗᒍᑦ ᐅᑯᐊ, ᓄᐊ ᑲᑦᓕᒃ, ᐱᓕᕆ-ᑭᕋᓂᓯᐊ ᑐᑭᐊᔅ, ᐅᐃᓕ ᐊᑕᓴᔅ, ᐱᐊᕐᓕ ᐊᖓᑎᐊᓗᒃ, ᐊᖓᒪ ᐃᖓᑎᐊᓗᒃ, ᐊᖓᒪᑦ ᐊᖓᑎᐊᓗᒃ, ᑎᑎ

Environment and Climate Change Canada
Public Inquiries Centre
Fontaine Building 12th floor
200 Sacré-Coeur Blvd
Gatineau, QC. K1A 0H3
ᐅᖅᑲᓕᐅᖅ: 819-938-3860
ᐅᖓᓱᓴᑎᖅ ᐊᑭᖏᕐᑐᖅ: 1-800-668-6767 (ᑲᓇᑕᒥ ᑭᓱᐊᓂ)
ᖃᓄᑕᐅᔪᒃᑯᑦ ᑎᑎᖃᖅᑕᕐᕕᖏᓐᓄᑦ: ec.enviroinfo.ec@Canada.ca

ᖃᖅᖏᓂᑐᖅ ᐊᐅᓚᑦᑎᐊᕐᒃ: © ᐊᐲᑎᓕᓗᓴᕐᔪᐊᒃᑯᑦ ᓱᓕᐅᖅᓯ ᐊᓱᐅᖅᓯᒪᔪᓕᐊᓂᖓ ᑲᓇᑕᒥ, ᖁᔭᓂ-ᑭᑦᓂᓱᐊ ᑐᑭᐊᑦ
© ᑯᐃᓐᒋᑦᖅ, ᐱᔪᓐᓇᐅᑎᖃᕐᓂᖏᓐᓄᑦ ᑲᓇᑕᒥ, ᑭᖏᖅᑐᖅᑕᐅᖅᓯᓗᓂ ᒥᓂᔅᖏᓐᓄᑦ ᐊᐲᑎᓕᓗᓴᕐᔪᐊᒃᑯᑦ ᓱᓕᐅᖅᓯᓗ ᐊᓱᐅᖅᓯᒪᔪᓂᖓᓄᑦ, 2020

Aussi disponible en français
ᐊᑐᐃᓐᓇᐅᖏᓪᓗᒍ ᑕᒃᑯᐊ ᐅᐃᓴᖅᑎᑐᑦ

ᐱᓴᓯᑎᑭᓪᓗᑭᑦ ᐊᕐᕋᑎᓕᓚᕆᕐᓴᐅᑲᓂᕐᑐᑦ ᓱᓕᐅᕐᓪᓗ ᐊᓱᓴᖅᓯᕐᓕᐊᓂᖏᓐᓄᓚᓂᑦ ᑲᓇᑕᐅᕐ ᓴᓐᐊᖅᑕᐅᓱᓕᓂᖏᑐᓚᓂᑦ ᓄᓇᐃᑦ ᐊᓚᒪᓗ ᐊᐅᓪᓕᑕᐅᓂᖏᑐᓚᓂᑦ ᕿᖅᓇᐅᑎᑦ

ᓱᓇᐅᖏᓪᓕ ᐊᕐᕋᑎᓕᓚᕆᕐᓴᐅᑲᓂᕐᑐᑦ ᓱᓕᐅᕐᓪᓗ ᐊᓱᓴᖅᓯᕐᓕᐊᓂᖏᓐᓄᓂᑦ ᑲᑎᑕᐅᕐ ᓄᓇᕗᑦ ᓄᓇᕗᑦ ᓴᓐᐊᖅᑕᐅᓂᕐᒪᓱᓕᕐᓂᑦ?

ᑎᓐᓕᐊᖅᑲᕃᐅᓪᓗᖅᑐᑦ ᓴᓐᐅᓂᐊᖅᑕᐅᓱᓕᕐᓂᑦ ᓴᖅᑭᑎᑕᐅᓕᐅᖅᓱᓕᔪᑦ ᒪᓕᖅᑕᐅᕐᓕᓗᑎᑦ ᒪᓕᑐᖃᔪᐊᑦ ᑕᐃᓴᐅᔪᑦ ᑎᓐᓕᐊᑦ ᓂᕿᐊᓚᓴᐅᕐᐅᖅᑐᑦ *ᑲᑎᒪᖃᔪᐊᕐᑕᐅᓂᖏᑐᓄᑦ ᒪᓕᑐᓕᐊᑦ, 1994-ᒥ*, ᐃᓂᐊᓴᖅᑭᑕᐅᕐᓕᓗᑎ ᑎᓐᓕᐊᓄᑦ ᐅᑎᖅᕿᖅᑐᑦ ᐊᕐᕌᒍᑦᓭᒃᖅ ᐃᒪᓕᖃᑦᓄᑦ ᓄᓇᕗᑦᑐᓚᓂᑦ ᓇᔪᒍᕐᓕᓗ ᓀᔪᖅᕿᖅᑕᑐᖅᓴᖅᑎᓚᖏᓐᓄᓂᑦ. ᐊᕐᕋᑎᓕᓚᕆᕐᓴᐅᑲᓂᕐᑐᑦ ᓱᓕᐅᕐᓪᓗᖅ ᑕᓚᐅᕐᒥ ᐃᒪᓕᖅᓂ ᐊᒪᓚᓗ ᓄᓇᕗᑦ ᑲᑎᑕᐅᕐᓕᓗᑎᑦ ᐃᒪᓕᖅᑐᖅᓴᖅᑎᓚᐅᖅᓯᒪᓕᖅᑐᑦ ᑲᑎᑕᐅᕐᓕᓗᑎᑦ ᐃᒪᓕᖅᑐᖅᓴᖅᑎᓚᐅᖅᓯᒪᓕᖅᑐᑦ ᑲᑎᑕᐅᕐᓕᓗᑎᑦ ᐃᒪᓕᖅᑐᖅᓴᖅᑎᓚᐅᖅᓯᒪᓕᖅᑐ

ᓱᓇᐅᖏᒪᑕ ᐊᐅᓪᓚᕆᑎᑲᓕᐅᑦ <ᕀᓇᐅᑏᑦ?

ᐊᐅᓪᓚᕆᑎᓚᒋᔪᑦ <ᕀᓇᐅᑏᑦ ᐱᑕᖏᕀᓕᐅᖅᑐᑐᑦ ᐱᓪᓗᓐᑕᐅᔨᓴᓇᓂᒋᔪᑦ, ᐊᐅᓪᓚᕆᑎᐅᐅᔨᓇᓂᑦ ᐃᓱᒪᓪᐅᕈᑕᐅᔭᓘᐊᓪᖏᓇᓂᑦ. ᑐᑦᖏᒪᐅᑦ ᐊᑐᖏᑕᐅᓂᒋᔪᑦ ᐊᐅᑎᓪᓗᓂᔭᕐᔨᐊᖏᑦ ᓱᓪᒃᐅᓂᑦ ᐊᓱᓴᖏᑦᓯᓪᓗᓂᑦ ᑲᑎᒪᓂᐅᕈᑕᐅᔭᓘᐊᓪᖏᓇᓂᑦ, ᐱᓐᐊᖏᑐᒃ ᐱᔨᐊᑎᖏᖏᑐᓂᑦ ᐱᔨᓂᐊᕐᑎᓂᐅᕈᑕᐅᔭᓘᐊᓪᖏᓇᓂᑦ ᑲᑎᒪᓂᐅᕈᑕᐅᔭᓘᐊᓪᖏᓇᓂᑦ. ᐊᐅᑎᓪᓗᓂᔭᕐᔨᐊᖏᑦ ᓱᓪᒃᐅᓂᑦ ᐊᓱᓴᖏᑦᓯᓪᓗᓂᑦ ᑲᑎᒪᓂᐅᕈᑕᐅᔭᓘᐊᓪᖏᓇᓂᑦ ᐊᐅᓪᓚᕆᑎᒋᔪᑦ <ᕀᓇᐅᑏᑦ ᑕᒪᐃᓐᓂᑦ ᑲᑎᒪᓂᐅᕈᑕᐅᔭᓘᐊᓪᖏᓇᓂᑦ, ᑐᑦᖏᒪᐅᑦ ᓄᓇᕗᒻᒥᓂᑦ ᐊᐅᓪᓚᕆᑎᒋᔪᑦ ᐃᓐᓇᐃᓐᓇᓂᑦ ᐊᐅᓪᓚᕆᑎᒋᔪᑦ ᐱᐅᔪᓐᓇᓂᑦ ᐊᐅᓪᓚᕆᑎᒋᔪᑦ ᐃᓐᓇᐃᓐᓇᓂ

ᑎᑎᕀᖅᑕᐅᓯᒪᓕᖅᓂᖅᓯᕐᓂ ᑭᖓᓪᓗᓴᒃᑯᖓᑦ

ᐊᕐᑎᑎᓪᓗᓴᐅᕐᒪᑦ ᓯᓚᐅᑦᓗ ᐊᓯᔪᖅᑯᑦᓪᓗᐊᓂᖓ ᑲᓇᑕᒥ ᑎᑎᕀᖅᑎᑕᖅᓴᑦ ᐊᐅᓚᕆᑎᓂᕐᒧᑦ ᕿᖏᐅᑎᒃᑲᓂᒃ ᑕᒪᐃᓐᓄᑦ ᑎᓐᓕᐊᖅᑲᖏᖅᓄᑦ, ᐊᐅᓚᑕᐅᔪᓂᒃ. ᑕᒪᒃᑯᐊ ᐃᓴᑭᐅᒪᓂᖓᓂ ᐊᑐᖅᑕᐅᖅᑲᕐᑎᑕᑭᓐᓇᕐᓂᕐᒧᑦ ᐊᓯᖏᓐᓄᑦ ᐱᓕᓚᖏᐅᔪᓄᑦ ᑐᕌᖏᐊᕐᓂᑯᓐᓄᑦ ᑎᑎᕀᖅᑎᑕᖅᓴᑦᑕᓐᓇᖅᑐᓂᖅᓄᖅ ᐊᔪᕐᓗᓂᖅᑐᓂᖅᓄᖅ ᐊᐅᓚᕆᑎᓂᕐᒧᑦ ᕿᖏᐅᑎᓂᒃᑲᓂᒃ, ᓯᕗᓕᓕᖅᒥ ᒪᓕᑐᖅᑎᒍᑦ ᖃᐅᔨᖅᑕᐊᖅᑕᐅᖅᑎᑕᖅᑕᖅᓯᓗᑎᒃ ᑕᓕᓚᒪᑕᒧᑦ ᐊᕐᕌᒍᐃᓄᑦ ᓇᕐᓂᓚᖓᔪᖅ, ᐊᖏᕈᑕᐅᓚᐅᖅᑎᓕᓗᑎᑦ ᓯᕗᓕᓚᓇᔨᐅᓪᓚᑦ ᕿᖏᐅᑎᓴᓐᓂ, ᑭᖓᓄᓕᐊᒍᑦ ᖃᐅᔨᖅᑕᐊᖅᑕᐅᑲᖏᓐᓄᕐᓇᖅᓯᓂᑦ ᖃᐅᓕᓂ 10 ᐊᕐᕌᒍᐃᓄᑦ ᓇᓴᕐᖏᓚᒥᒃ.

ᖃᐅᔨᓴᕆᑎᐊᖅᑲᓂᕐᒪᓵᓄᑦ

ᖃᐅᔨᓴᕐᑎᐊᖅᑲᓂᕐᒪᔪᐃᓕᑦ ᐱᔾᔨᑎᕈᖃᖅᑕᑐᓂᒃ ᐊᕐᑎᑎᓪᓗᓴᐅᕐᒪᑦ ᓯᓚᐅᑦᓗ ᐊᓯᔪᖅᑯᑦᓪᓗᐊᓂᖓᓄᓂᒃ ᑲᓇᑕᒥ ᓴᓐᓂᓂᓕᐅᖅᑐᓂᖅᓄᖅ, ᐊᑎᑐᖅ ᑕᐃᑎᐊᖅᓂᐊᖅᑯᐃᑦ ᖃᐅᔨᖅᑕᐊᖅᑕᐅᖅᕿᓚᓂᑦ ᖃᐃᓂᐊᖅᑕᐅᖅᕿᓚᓂᑦ ᐅᕐᓂ https://www.Canada.ca/en/environment-climate-change/services/ɹonational-wildlife-areas.html ᐅᕐᓂᓚᓂᓂᑦ ᐅᖅᑲᖏᐱᓚᑐᑦ ᑲᓇᑕᒥ ᐅᒪᔨᓚᓂᓚᐅᓂᒃᑯᑦ ᐱᓕᑎᑎᐃᓕᓴᕐᑎᑦ.

ᐃᑲᑐᐊᖅᑎ ᑎᖏᑕᐊᖅᑲᕐᓚᖅ

ᐃᑲᑐᐊᖅᑎ ᑎᖏᑕᐊᖅᑲᕐᓚᖅ (ᐃᑲᑐᐊᖅ) ᐃᓂᓕᖅ ᑕᐅᖅᓂ ᐅᐊᖏᓐᓇᒪᓇᖏᓂ ᕼᐊᑦᔭᐧᐋᒃ, ᑕᒧᐅᕗᖅᓯᐊᕐᓗᐊᓂᖅ, ᑕᒪᓐᓂ ᓴᓐᓂᐅᑎᓂ ᐊᖏᖏᖅᓯᒪᓂᖏᓂ ᓄᓇᕗᑦ, ᔭᐅᔭᖅᑕᖏᓂ. ᐃᓂᓕᖅ 110 ᑭᓪᒥᑕᓂᖅ ᓂᑎᐊᑕ ᐱᖏᓐᓇᖅᑲᓯᖏᓂ, ᐅᖏᓯᖏᓂᓕᒥ ᔭᑦᑲᖅᓂᖏᓐ, ᑕᐃᖅᓇ ᐃᑲᑐᐊᖅᑎ ᑎᖏᑕᐊᖅᑲᕐᓚᐅᑦ ᑯᖏᑦᑕ ᐃᒪᓐᓇᖅᑐᑦ (ᐳᐊᓯᐅᑦ ᑯᖏᓂᖅ ᑕᐃᔨᐅᖅᑐᑎᒃ ᖃᐅᔨᖅᑲᐃᑎᑐᑦ). ᑕᒪᖏᓇ ᑎᖏᑕᐊᖅᑲᕐᓚᖅ ᐃᓕᖅᖃᓇᖅᖢᖅ ᓄᓇᒥᖅ ᐃᓗᑐᓂᐊᓗᖏᒥᖅ ᑯᐃᓚᐅᓕᖅᓯᓂᖏᓐᓄᓕᓂ ᓯᐅᑦᖁᑦᓯᓕᐊᓯᒪᓂᑯᖅᑎᒃ ᔭᖏᓂᖅ ᐊᒪᒪᓗ ᐊᑯᖅᑐᑦ ᑕᐅᖅᓂ ᐊᑭᐊᓂᖅ ᐊᓯᖅᑐᐃᑎᖅᑐᓂᖅᑐᑦ ᑕᒪᖏᓇ ᑎᖏᑕᐊᖅᑲᕐᓚᖅ ᑐᖅᑲᐃᖅᑲᖅᓂᖏᓐᓂᖅᑐᑦ. ᑕᒪᖏᓇ ᑎᖏᑕᐊᖅᑲᕐᓚᖅ ᑐᓴᖅᑲᖅᑐᑦ ᐅᐱᖅᑎᑦ ᐅᐱᖅᑲᐃᖅᑐᑦ ᓯᑎᐊᓚᖅᑲᖅᑐᑦ ᐱᖏᓐᓇᖅᑐᑦ ᓯᑎᐊᓚᖅᑲᖅᑐᑦ ᓯᓂᓗᐊᓂᖅᑐᑦ ᐃᓚᖅᑲᐃᓐᓇᖅᑐᑦ ᓯᓂᓗᐊᓂᖅᑐᑦ ᑕᒪᖏᓇ ᑎᖏᑕᐊᖅᑲᕐᓚᖅ ᑐᖅᑲᐃᖅᑲᖅᓂᖏᓐᓂᖅᑐᑦ. ᑕᒪᖏᓇ ᑎᖏᑕᐊᖅᑲᕐᓚᖅ ᑐᖅᑲᐃᖅᑲᖅᓂᖏᓐᓂᖅᑐᑦ

ᐊᒥᓱᓂᕆᔨᓂᖅ ᑕᒪᓂ 690,000-ᓂᒃ ᑲᖓᔪᓂᒃ ᐊᐸᑭᖏᑦᑐᑎᑕᐅᔪᓂᒃ (ᐅᖏᓪᓗᓂᑦᑐᑦ >5% >ᓴᖅᑎᖏᑭᓂᖅᑕᐅᕐᓗᑎᒃ ᑖᒃᑯᓇᓂ ᒥᑭᓂᖅᓴᐅᓂᖅᑕᐅᕐᖓᑐᑦ ᑲᔪᓯᖏᓪᓗᓂ, ᓇᓂᑐᐃᓇᖏᑎᐊᖅ ᐱᑕᖅᑯᓱᓂᖏᑭᓂ, ᑕᐃᒃᑯᓇᓂᓗ ᑲᓇᑕᐅᕐᔨ ᓄᓇᕗᑦᑕ ᖃᑎᖅᑳᓱᐊᓄᐊᖅᑕᕐᒥᔪᓂᒃ ᑎᖅᒥᐊᓚᕆᐅᖓᖓᑐᑐᒃ). ᑲᖓᐃᑦ ᐱᑕᖅᑲᓂᖅᑭᓂ ᐅᐊᒥᓱᓂᖅᐸᑎᐊᖓᕐᑐᑦ ᑕᐅᖓᓂ ᐃᑲᑕᑐᐊᑦ ᑯᓴᖅᑲᐃᓕᓂᑦ. ᐱᑐᖅᑐᖅᑲᓂᖅᑎᑦ ᐃᓚᖅᓴᐃᓇᐅᓂᖅᑭᓂᓂ ᐊᒥᓱᓕᑕᐅᖓᑐᑦ ᐅᓪᓗᖅᑲᖓᓚᐅᓇᓂᕐᒪᑦ. ᐱᑐᖅᑐᑦ ᑭᓪᓚᓇᐃᓇᐅᓂᖅᑭᓂᓂ, ᖃᖅᑲᑦᖅᑲᖅᑭᓂᓂᐅᔪᓂ ᐊᑎᑕᑐᑎᑕᐅᔪᓂᒃ ᓇᓚᒪᓗ ᐃᓚᖃᖅᑲᓂᖅᑭᓂᓂᓗᓐᖓᓂᒃ ᐊᑎᒪᓗ ᐃᓚᖃᖅᑲᓂᖅᑭᓂᓂᓗᓐᖓᓂᒃ ᐊᒥᓱᓕᑕᐅᖓᑐᑦ. ᐃᓕᓴᐅᕐᑕᖅᑕᐅᔪᓂᒃ ᐊᒥᓱᓕᑕᐅᖓᑐᑦ ᐊᑎᒪᓗ ᐃᓚᖃᖅᑲᓂᖅᑭᓂᓂᓗᓐᖓᓂᒃ ᐊᒥᓱᓕᑕᐅᖓᑐᑦ. ᑕᐃᒃᑯᓇᓂ ᑭᓪᓚᓇᐃᓇᐅᓂᖅᑭᓂᓂᓗᓐᖓᓂᒃ ᐊᒥᓱᓕᑕᐅᖓᑐᑦ ᐃᓚᖃᖅᑲᓂᖅᑭᓂᓂᓗᓐᖓᓂᒃ ᐊᒥᓱᓕᑕᐅᖓᑐᑦ ᑕᒪᓐᓇᖅᑐᒥ

ᐊᐅᓪᓗᕆᑐᐅᖅᑕᐅᓂᖅᓯᓂ ᐊᓚᓗᓐ ᐊᖏᕈᑕᐅᓴᓚᐊᓪᓕᓂ ᑲᒪᕈᓴᐅᕙᖏᖅᓯᓂ

ᒪᓪᓕᑕᐅᓴᓚᐊᕐᑲᖅᓂᖅᓯᒪᓂᓄᓐᓂ ᓄᓇᕗᓴᒥ ᐊᖏᕈᕈᑎᑦ, ᐃᓄᐃᑦ ᐊᐅᑐᖅᑕᐅᓂᓴᓕᓚᖏᖅᓯᒪᓂᓄᓐᓂ ᐃᑲᔪᓴᐊᖅᑕᕈᑎᑎᐅᓴᖅᑐᓂᓪᓗ ᐊᖏᕈᕈᑎᓂ ᑲᓇᑕᓕᒫᒃ ᐅᒪᔨᕐᑲᖅᓇᖅᓄᓐᓂ ᐊᓚᓗᓐ ᑎᖏᐅᐊᖅᑲᖅᓇᓕᓪᓕᖅᑐᒥ, (ᐃᓄᐃᑦ ᐃᑲᔪᓴᐊᖅᓴᓚᐊᕐᓄᓐᓂ ᐊᖏᕈᕈᑎᑎᑦ) ᔨᓐᓕᓕᑦᑐ ᐱᓴᓱᖅᑕᐅᓴᖅᑲᒪᒃᑐᖅᑐᓂ 2006-ᖓᑎᓕᓪᔪᒍ, ᐊᑐᖅᑕᐅᓕᒃᐅᖅᑐᓂᑐᒃ 7-ᓄᓂ ᐊᐅᖏᒍᓄᓂ, ᐊᖏᕐᕋᑲᑎᑦᑕᐊᖅᑲᓂᕐᓂᓕᕆᓕᓪᓕᒫᓕᓪᓗᓂᑐᒃ 7-ᑲᓐᓂᐅᓂᐊᓕᕐᑎᔪᓄᓂ ᐊᐅᖏᒍᓄᓂ. ᒪᓪᓕᑕᑐᓄᓂ ᐊᐃᓴᑐᖅᓯᒪᓂᖅᓯᓂ *3*-ᖓᓚᓪᓕᑐᑐᓂ (ᐱᔪᔨᑎᕋᖅᑐᑐᓂ ᐊᐅᓪᓗᕆᑐᐅᖅᑕᐅᓴᖅᑕᐅᔪᓂᖅᑐᒃ) ᑕᒪᖅᑯᓂᖓᓪᓚᕆ *ᐃᓄᐃᑦ ᐃᑲᔪᓴᐊᖅᓴᓚᐊᕐᓄᓐᓂ ᐊᖏᕈᕈᑎᓂ ᑎᑎᕋᖅᑲᐃᐧ ᐅᕋᖅᑕᐅᓴᖅᑲᒪᒃᑐᖅᑐᓂᑐᒃ ᐅᕋᖅᑕᐅᓴᖅᑲᒪᒃᑐᖅᑐᓂᓪᓗ ᐃᒪᓕᒫᐅᔪᓂᖅᑐᒃ ᓄᓇᕗᓴᒥ ᐊᖏᕈᕈᑎᓂ ᐊᓚᓗᓐ ᐊᐃᓴᑐᖅᓯᒪᓂᖅᓯᓂ ᐊᖏᕈᕈᑎᓂ ᐊᓚᓗᓐ ᐱᓕᕆᔨᒃᑯᓂᖅᓯᓂ ᐊᐅᓪᓗᕆᑐᐅᖅᑕᐅᓴᖅᑕᐅᔪᓂᖅᑐᒃ ᓄᓇᕗᓴᒥ ᐅᒪᔨᕐᑲᖅᓇᖅᓄᓐ

ᐊᐅᓪᓚᕐᓂᖅᓯᒪᔪᓂ ᐸᖅᓇᐅᑎᓂᒃ ᑕᐃᒃᑯᓇᖅᑐᒃ ᐃᓚᔨᓪᓚᕆᔨᖅᑕᐃᒃᑯᓂ ᑲᑎᒪᔨᖅᑭᒃᓇᓂ (ᐃᓚᔨᓪᓚᕆᔨᖅᑕᐃᒃᑯᓂ) ᐊᖏᕐᕋᐅᔾᔨᑯᓗᑎᓂ ᐊᑐᖅᓯᒪᔭᓂᑐᓂᒃ *5.2.34(c)*-ᒥ ᐊᓚᒪᓗ *5.3.16*-ᖅᓱᓂ ᓇᓇᕐᓚᒥ ᓇᓇᑕᐳᑎᓂᒃ ᐊᖏᕐᕈᑎᓂᒃ. ᐊᐅᓪᓚᕆᑎᐅᔾᕋᕐᕋᐅᓂᖅᓯᒪᔪᓂ ᑲᑎᒪᔨᕿᓕᓂ ᑐᓂᓯᔪᐊᖅᑐᑦ ᑭᓐᓚᕐᓂᒥ ᐃᓇᐃᓂ ᑲᑐᔨᖅᑲᑎᕐᑎᓂᒃ, ᑐᓴᐅᖏᓐᓂᒃ ᐱᓕᕆᖅᑕᐅᓯᒪᔪᓂᒃ ᐊᐅᓪᓚᕆᓂᖅᓯᒪᔪᓂ ᐸᖅᓇᐅᑎᓂᒃ, ᑐᔪᐃᓂᖏᓐᓂᒃ ᐸᖅᓇᐅᑎᓂᒃ ᐊᐅᓪᓚᕆᓂᖅᓯᒪᔪᓂ ᐸᖅᓇᐅᑎᓂᒃᑲᓐᓂᕐᒪᑐᐃᓐᓇᖅᑐᒃ ᓇᓇᕐᓚᒥ ᐃᓚᔨᓪᓚᕆᔨᖅᑕᐃᒃᑯᓂ (ᐊᑐᖅᓯᒪᔭᓂᒃ ᐃᓇᐃᓂ ᐃᑲᔪᓯᐊᖅᑎᖏᓐᓂᒃᓂ ᐊᖏᕐᕈᑎᓂᒃ *3.6.2*). ᒪᓕᓐᓯᒪᔭᒃ ᐃᓯᒪᓕᐅᕗᖅᑎᐅᔨᓴᖅᓂᖏᓐᓂᒃ ᐱᔨᓴᕆᓂᖏᓕ ᐊᑕᒧᓂ ᓇᓇᕐᓚᒥ ᓇᓇᑕᐳᑎᑎᒧᓂ ᐊᖏᕐᕈᑎᓂᒃᑯᓂ, ᐃᓚᔨᓪᓚᕆᔨᖅᑕᐃᒃᑯᓂ ᐅᕙᓚᐅᔾᔨᖅᑕᐃᒃᑯᓂ ᐅᓂᒃᑳᖅᑐᓂ ᐊᖏᕐᕈᑎᓂᒃ, ᐊᒻᒪᓗ ᐅᑎᖅᑎᑕᐅᔪᓐᓇᖅᑐᒃ ᐊᐅᓪᓚᕆᑎᐅᔾᕋᕐᕋᐅᓂᖅᓯᒪᔪᓂ ᐅᖃᓕᕆᔨᒃᑯᓂ ᐊᐅᓪᓚᕆᓂᖅᓯᒪᔪᓂ ᐸᖅᓇᐅᑎᓂᒃ ᐃᓚᔨᓪᓚᕆᔨᖅᑕᐃᒃᑯᓂ ᐅᖃᓕᕆᔨᒃᑯᓂ ᐃᓕᓴᕐᕋᐅᓯᒪᔪᓂ ᐊᖏᕐᕈᑎᓂᒃᑲᓐᓂᕐᒪᑐᐃᓐᓇᖅᑐᒃ ᐃᓚᔨᓪᓚᕆᔨᖅᑕᐃᒃᑯᓂ ᐊᖏᕐᕈᑎᓂᒃᑲᓐᓂᕐᒪᑐᐃᓐᓇᖅᑐᒃ

ᒪᑲᐱᒪᐃᑦ ᖃᑲᓂᐃᓕᖅᓯᖏᑦ

ᑎᑎᕃᓕᓂᖏᑦ ᐊᐃᑲᑐᓕᕆᓂᑲ ᑕᔪᑎᑦᑎᐊᑦᑕ.....x
ᑎᑎᕃᓕᓂᖏᑦ ᑕᔪᐅᓴᐃᐅᑦ.....xi
ᑎᑎᕃᓕᓂᖏᑦ ᓇᐃᒡᓗᑭᐊᖅᑕᐅᓕᕆᑦ ᑎᑎᖅᐃᑦ ᖅᓪᓗᓂᔪᓚᑐᑐᑦ.....
1 ᐅᖅᐅᓕᐅᑦᑎᐊᕐᓂᖏᑦ ᓴᐳᔨᓂᐊᖅᑕᐅᓕᕆᑦ ᓄᓇᐃᑦ.....1
1.1 ᐊᐃᑲᑐᓕᕆᓂᖓᑕ ᖃᓄᐃᓕᓂᖓᓂᖓ.....8
1.2 ᖃᓄᖅ ᐱᓇᓪᓕᐊᓕᐅᖏᓐᓂᑦ.....9
1.3 ᓄᓇᕐᔪᑎᑎᓴᐅᓂᖏᑦ ᐊᒻᒪᓗ ᐊᑐᖅᑕᐅᓕᔨᓇᕐᓂᖏᑦ.....10
1.4 ᐃᓱᓗᕐᔪᑎᑎᓴᐅᔨᑦ ᐱᕐᔪᐊᓪᓗ.....11
2 ᐊᓯᖏᓂᖏᑦ ᐊᓯᖏᑎᐅᑕᖃᕐᓂᖏᓪᓗ.....13
2.1 ᓄᓇᒥᐅᑕᐃᓪᓗ ᐃᒪᕐᒥᐅᑕᐃᓪᓗ ᐅᒪᔨᖃᕐᐸᑦ.....13
2.2 ᑎᓕᒥᐊᑦ.....15
2.2.1 ᑲᔪᐃᐃᑦ.....15
2.2.2 ᓱᓴᖅᓯᓕᐅᓂᖅᑭᓱᑦ ᑎᓕᒥᐊᑦ.....20
2.2.3 ᐊᓱᓕᑦ ᑎᓕᒥᐊᑦ.....20
2.3 ᐊᓱᓕᑦ ᐅᒪᔨᐃᑦ.....21
2.3.1 ᓄᓇᒥᐅᑕᐃᑦ ᐅᒪᔨᐃᑦ ᐊᐱᒪᑲᑎᑦᑎᓱᑦ ᐱᒥᓂᑲ.....21
2.3.2 ᑕᕆᐅᖅᑎᐅᑕᐃᑦ ᐅᒪᔨᐃᑦ.....22
2.3.3 ᑕᕆᐅᖅᔪᕐᓂᓕᒥ ᑕᓱᖅᑎᐅᑕᐃᑦ ᑕᕆᐅᖅᑎᐅᑕᐃᓪᓗ ᐃᖃᓄᐊᓕᕐᐃᑦ.....22
2.4 ᐱᑭᖅᑐᖃᖅᓂᖓ ᓄᓇ.....23
2.5 ᐅᒪᔨᐃᑦ ᐊᒥᓱᓕᓪᓚᐊᓕᓂᖅᑐᑦ.....23
2.6 ᑕᐃᓂᖅᑎᐅᑕᑐᖃᖅᓂᖅᑐᑦ ᐅᒪᔨᐃᑦ/ᖃᐅᔨᓴᕐᓗᓂᓪᓚᕐᓂ, ᐱᑭᖅᑐᓪᓗᓂᓪᓚᕐᓂ ᐱᑕᖃᖅᓂᖓᓂᑦ.....25
3 ᐃᓕᓂᐅᔪᑐᖃᖅᑎᒍᑦ ᐊᑐᓚᓗᐅᐊᓂᖓᓂᑦ.....26
3.1 ᐃᓕᓂᐅᔪᑐᖃᖅᑎᒍᑦ ᓄᓇᒥ ᐱᖃᑎᖅᐃᓚᓂᑦ ᐊᑐᖅᑕᐅᐊᓕᐅᖅᓂᖓᓂᑦ ᑐᑭᖃᖅᓂᖓᓕᓂᓪᓗ ᖃᐅᔨᓴᖅᑕᐅᓕᕆᑦ.....26
3.2 ᑕᐃᒪᓐᓇᓂᑲ ᐃᓄᐃᑦ ᐊᑐᖃᖅᑕᖅᓂᖓᓂᑦ ᓄᓇᒥᑲ.....27
3.3 ᐊᓪᓚᐅᓕᖅᑐᖅ ᐃᓄᐃᑦ ᐊᑐᖃᖅᑕᖅᓂᖓᓂᑦ ᓄᓇᒥᑲ.....32
4 ᑐᖏᓚᓗᐅᔨᑦ ᐊᒻᒪᓗ ᐱᔨᓱᖅᑕᐅᔨᒪᔨᑦ.....33
4.1 ᑕᐅᑐᓚᓗᐅᔨᑦ.....33
4.2 ᑐᖏᓚᓗᐅᔨᑦ ᐊᒻᒪᓗ ᐱᔨᓱᖅᑕᐅᔨᒪᔨᑦ.....33
4.3 ᖃᐅᔨᓕᐊᖃᑕᖅᓂᖅ.....35
5 ᐊᐅᓪᓕᑦᑎᓂᖅᒥᑦ ᐃᓕᒪᔨᖏᖅᓴᐅᑎᑦᑎᐃᓕᑦ.....36
6 ᐊᐅᓪᓕᑦᑎᓵᐅᓂᖅᒍᑦ ᖃᓄᐃᓕᓕᐅᕐᑐᑎᑦᑎᐃᓕᑦ.....42
6.1 ᐱᐅᓴᑐᖃᖅᑎᒍᑦ ᓄᓇᒥᐅᑕᓂᑲ ᐊᐅᓪᓕᑦᑎᓂᖅ.....42
6.1.1 ᐱᐅᓴᓗᒥᒍᑦ ᐊᒻᒪᓗ ᐃᓄᓕᕐᔪᑎᑎᓂᓄᑦ ᐊᑐᖅᑕᐅᐅᖃᐃᓄᐊᓕᓂᖓᓂᑦ.....42
6.1.2 ᐅᒪᔨᖃᖅᐃᑦ ᐱᓪᓚᓗᐅᔨᑦ ᐃᓄᖅᓄᑦ.....43
6.1.3 ᐃᓄᐃᑦ ᑕᐅᔭᓂ ᑕᐃᔪᓴᑐᖃᖅᓂᑦ ᓄᓇᒧᑦ ᐃᒪᓕᓄᓪᓚᓪᓗ.....44
6.2 ᐅᒪᔨᓪᓗᓂᒥᑲ ᐊᒻᒪᓗ ᐅᒪᔨᖃᖅᓄᐅᔭᖅᓂᖓᓂᓄᑲ.....44
6.2.1 ᐊᓱᓴᖅᑕᐅᓂᖅᓯᒪ ᐃᓕᓐᓂᓗᐅᑎᓕᔨᖅ ᖃᓄᐃᓕᑕᑐᓂᖅᓯᒪ ᓴᐳᔨᓂᐊᖅᓴᖏᓐᓄᐅᓂᖅ.....44
6.2.2 ᐊᒥᓴᓪᓚᐊᓕᕐᓂᖓᓂᑦ ᑲᖏᐃᑦ.....45
6.2.3 ᐅᒪᔨᐃᑦ ᐊᒥᓱᓕᓪᓚᐊᓕᖅᑐᑦ.....45
6.2.4 ᑕᐃᓂᖅᑎᐅᑕᖃᐅᔨᖅᑐᑦᑕᑦ ᐅᒪᔨᐃᑦ/ᖃᐅᔨᓴᕐᓗᓂᓪᓚᕐᓂ ᐱᑭᖅᑐᓪᓗᓂᓪᓚᕐᓂ ᐱᑕᖃᖅᓂᖓᓂᑦ.....46
6.2.5 ᐊᖏᓇᓄᒃᓴᖓᓂᖅ.....46
6.2.6 ᑎᓕᓴᓂ ᐃᖏᕐᕋᓭᓂᖓᓂᑦ ᑕᐅᔭᓂ.....47
6.2.7 ᑕᕆᐅᖅᑎ ᐅᒥᐊᖅᑐᕐᓂᖅ/ᐅᒥᐊᕐᔪᐊᖃᖅᓂᖅ.....47
6.3 ᓇᐅᑦᑎᖅᑐᐃᓕᖅᓴᖃᖅᑐᑦᑕᓂᖓᓂᑦ ᖅᐅᓵᕐᓄᓇᖅᓭᓭᓂᖓᓂᓪᓗ.....48

6.4 ᐃᓄᖅᑎᑎᕆᐅᔪᓂᑎᓂᖅᓴᐅᕆᕐᓂᖅ ᐊᒻᒪᓗ ᖃᐅᔨᓴᕆᕐᓂᖅ ᐊᐅᓚᑦᑎᐊᓂᖅ 49
6.4.1 ᓄᓇᓕᒃ ᐅᓴᐅᓯᖅᓂᖅᓴᐅᔪᓂᖅᕈᑦ ᑐᐱᓯᐅᒪᕆᕐᓂᖅᕈᑦᓯᓗ 50
6.4.2 ᐃᓄᖅᑎᑎᕆᐅᔪᑦ ᐅᓴᐅᓯᖅᓂᖅᓴᐅᖃᐅᔪᓚᐅᔪᓂᖅᕈᑦ 50
6.4.3 ᐃᓕᓴᓂᐊᖅᑐᓄᓕ ᑐᕐᖏᓚᔪᑦ ᐱᓕᕆᐊᓴᓴᐃᑦ 51
6.4.4 ᓂᐅᕐᑭᐊᓂᖅ ᐳᓪᓕᕐᑐᓕᕆᓂᑦᑎᒃ ᐱᓕᕆᐊᖅᑎᓂᖅ 52
6.4.5 ᓄᓇᒋᕐᕕᐅᓂᖅᓄᑦ ᑕᖏᒧᖅᕕᐅᓂᖅᑐᓗ 53
6.4.6 ᐃᓄᖓᓄᑦ ᐱᖅᑎᑎᕆᐅᔪᑦ ᓄᓇᐃᑦ 54
6.4.7 ᒪᓕᕆᓚᐅᔪᓂᖅ ᒪᓕᒃᑕᐅᖅᐃᓇᓕᔪᓂᖅ 54
6.4.8 ᐊᕕᖅᑐᖅᓯᒪᓂᕐᓂᓂᓂ ᖃᓄᐃᓇᓕᐅᕈᑕᐅᕕᖅᑐᑦ ᐱᑎᐊᖅᑎᑕᐅᓯᒪᓕᕐᖃᑐᓗ 55
7 ᐊᖏᕐᕈᑕᐅᓯᒪᓚᓚᐊᓕᑦ ᐱᓕᕆᐊᒃᓴᐃᑦ, ᐊᔪᖅᓴᖅᑎᑕᐅᔪᑦ ᐱᓕᕆᐊᕕᐅᔪᓴᐅᓴᖅᓯᒪᔭᑦ ᐊᒻᒪᓗ ᐊᔪᖅᑎᑕᐅᔪᑦ ᑎᑭᓯᒪᕕᐅᔪᓴᐅᓴᖅᓯᒪᔭᓂᖅᑐᓴᓂᑦ 57
7.1 ᐊᐅᓚᑦᑎᐊᓄᑦ ᐊᖏᕐᖃᑕᐅᒃᔪᓯᑦ 58
7.1.1 ᐊᖏᕐᖃᑕᐅᖃᑕᖅᓂᖅᕈᑦ ᐱᔪᓴᓚᐅᑦ 58
7.1.2 ᓄᓇᕗᒻᒥ ᐃᓄᐃᑦ ᐱᓕᕆᔪᓴᓴᖅᓂᖅᕈᑦ ᐱᔪᓴᓚᐅᑎᖃᖅᕈᓐᓃᑎᒃ 59
7.1.3 ᓄᓇᕗᒻᒥᐅᑦ ᐃᓄᐃᑦ ᐱᓕᕆᐊᕆᔪᓴᓴᖃᑕᖅᕈᑦ ᑭᓯᐊᓂ ᐱᔪᓴᓚᐅᑎᖃᖅᑕᑎᒃ 60
7.1.4 ᐃᓄᕐᓯᓚᐅᕖᕈᑦᑐᓄᓕ ᐱᓕᕆᐊᖃᕈᓴᓴᖅᓂᖅᕈᑦ ᐱᔪᓴᓚᐅᑎᖃᖅᕈᓐᓃᑎᒃ 60
7.1.5 ᐃᓄᕐᓯᓚᐅᖅᕈᑦᑐᑦ ᐱᓕᕆᔪᓴᓂᖅᕈᑦ ᐊᖏᕐᖃᑕᐅᔪᓴᖅᑐᑦ ᑭᓯᐊᓂ ᐱᔪᓴᓚᐅᑎᖃᖅᓯᓗᑎᒃ 60
7.1.6 ᐱᓕᕆᔪᓴᓂᖅ ᐊᖏᕐᖃᑕᐅᓯᓚᔪᑦ ᐱᔪᓴᓚᐅᑎᖃᖅᓂᓯᒧᑦ ᐃᓄᖓᓄᑦ ᐱᖅᑎᑎᕆᐅᔪᓴᐅᓴᖅᓯᒪᔭᓂᖅᑐᓴᓂᑦ 61
7.1.7 ᑲᓇᑕᒥ ᐱᔪᓴᓚᐅᑎᓂᑦ ᐱᓚᓯᐊᖃᖅᑕᑕᒪᓴᓴᖅᓂᓯᒧᑦ ᐱᐅᓯᓚᑦᑎᐅᓴᖅᓴᐅᑐᑦ ᐊᑐᖅᑕᐅᔪᓚᐊᓕᓕᑦ 61
7.2 ᐊᓯᖏᑦ ᑲᓇᑕᐅᑦ ᓴᖅᒪᑦᑯᖏᓴᓄᑦ ᐊᒻᒪᓗ ᓄᓇᕗᑦ ᓴᖅᒪᓇᖏᓴᓄᑦ ᐊᖏᕐᕈᑕᐅᖅᓴᐅᑐᑦ ᐊᒻᒪᓗ ᐊᒻᒪ ᐱᔪᓴᓚᐅᑎᐅᕖᖅᑎᓗᔪᑦ 63
8 ᓄᓇᖏᑦ ᑎᖅᑎᐊᖅᓴᐃᕐᓂᖅᓴᐅᓂᖏᓴᓄᑦ ᑕᐃᔨᐅᓂᖏᑦ ᐃᓕᑕᒪᔨᐅᓯᒪᓂᖏᓴᓗ 64
9 ᐋᓴᓂᖅᑕᐅᑦᑕᐃᓕᒪᑎᑦᑎᓂᖅ, ᐋᓴᓂᐊᖅᑐᑕᒪᓂᖅ ᐊᒻᒪᓗ ᐅᓗᓪᓚᐊᖅᑐᑕᒪᓂᖅ 70
9.1 ᓇᓄᐃᑦ ᑐᖅᓄᑕᐅᕙᓂᓂᖏᑦ 71
10 ᐊᑐᕐᖅᑎᑦᑎᓕᓪᓕᐊᓂᖅ 72
11 ᐸᖃᓇᐅᑦ ᐊᑐᕐᖅᑎᑕᐅᓂᖏᑦ 74
11.1 ᐊᐅᓚᑦᑎᓂᔪᓗ ᐸᖃᓇᐅᑦ ᐊᐠᑭᓚᐊᖅᑕᐅᔪᓚᐅᖃᖅᕈᓐᓃᑎᒃ 77
11.2 ᐊᐅᓚᑦᑎᐅᐳᐅᔪᓚᖅᑐᑦ ᐊᒻᒪᓗ ᑎᓪᓗᐅᓄᓓᔪᓂᖏᑦ 77
12 ᐱᓕᕆᖅᑎᑦᑎᐊᕐᕕᓚᖃᓂᖅ 78
12.1 ᐃᓄᐃᑦ ᐊᒻᒪᓗ ᑲᙰᒪᐃᑦ ᐱᓕᕆᖅᑎᑎᒍᓚᖃᓂᖏᑦ 78
12.2 ᓄᓇᕗᑦ ᑲᙰᒪᐃᑦ 78
13 ᐅᖅᓀᒪᓕᖅᓂᒃ ᑐᐱᓄᓂᐊᖅᐳᓄᖏᑦ 80
ᑎᑎᖅᐃᑦ ᐃᓕᓴᖅᓂᖏᑦ 84
ᑎᑎᖅᐃᑦ ᐃᓕᓴᖅᓂᖏᑦ A: ᒪᓕᒪᖅᑎᔪᑦ ᐅᓂᒃᖅᑕᐅᑦᑎᐊᖅᓂᖏᑦ ᓄᓇᐃᑦ ᓇᓗᓂᓂᖏᑦ 85

ᑎᑎᕋᖅᓯᒪᓂᖅᓄᑦ ᐊᕐᕌᒍᖅᓯᒪᔪᓂᒃ ᑕᔪᑎᑕᑎᑦᔪᓱᑦ

ᐊᕐᕌᒍᖅᓯᒪᔪᓂᒃ ᑕᔪᑎᑕᑎᑦᔪᓄᑦ 1: ᐃᒃᑐᑐᐊᖅᑐᑐᒃ ᑎᖅᑎᐊᕋᖅᑕᕐᕕᖅᑎᒃ ᓇᐃᓐᓇᖅᓯᒪᔪᓂᒃ ᖃᒃᐅᓕᒪᑦᑎᑎᐅᑕᐅᑎᑦ.....3

ᐊᕐᕌᒍᖅᓯᒪᔪᓂᒃ ᑕᔪᑎᑕᑎᑦᔪᓄᑦ 2: ᐊᕐᕌᒍᖅᓯᒪᓂᖅᓴᑕ ᖃᓄᐃᔨᓂᖅᓴᓂᑦᑎᑦ ᐊᓘᒪᓗ ᐅᒪᔪᕋᖃᕋᓂᖅᑎᑦ ᐊᐅᓚᐊᓂ ᓇᓗᓇᐃᒃᑯᑎᖅᑕᐅᓯᒪᓂᖅᑎᑦ ᑎᖅᑎᐊᕋᖅᑕᕐᕕᖅᑎᒃ.....9

ᐊᕐᕌᒍᖅᓯᒪᔪᓂᒃ ᑕᔪᑎᑕᑎᑦᔪᓄᑦ 3: ᐃᓗᓪᓕᔪᑎᑎᕆᐅᔨᓄᑦ ᐱᓕᔪᑏᑦᓗ ᑎᖅᑎᐊᕋᖅᑕᕐᕕᖅᑎᒃ.....11

ᐊᕐᕌᒍᖅᓯᒪᔪᓂᒃ ᑕᔪᑎᑕᑎᑦᔪᓄᑦ 4: ᐅᒪᔪᐃᐃᒃᑐᓗᐊᓕᕐᓂᖅᓴᑕ ᑎᖅᑎᐊᕋᖅᑕᕐᕕᖅᑎᒃ.....24

ᐊᕐᕌᒍᖅᓯᒪᔪᓂᒃ ᑕᔪᑎᑕᑎᑦᔪᓄᑦ 5: ᐃᓄᐃᓪ ᐃᓕᓴᔪᔪᑐᖅᑐᒃ ᑕᐃᔪᒍᑦ ᓄᓇᖅᑐᒃ ᐃᒪᓕᕐᓂᓗᐖᓂᑦ ᑎᖅᑎᐊᕋᖅᑕᕐᕕᖅᑎᒃ (ᐊᕐᕈᑎᐊᕋᖅᑕᐅᓕᑎᐊᕋᒃᖅᑐᓂᕋᖅᑐᑦ ᐃᓄᐃᓪ ᐱᐅᔪᑐᒃᖅᑐᖅᑐᒃ <<ᕕᑎᐅᓴᒃᔪᑦ ᑎᑎᕋᖅᑕᔪᑎᑎᖅᑐᖅᑐᒃ).....30

ᐊᕐᕌᒍᖅᓯᒪᔪᓂᒃ ᑕᔪᑎᑕᑎᑦᔪᓄᑦ 6: ᐊᐅᓪᓚᕕᑎᓕᖅᑐᑦ ᑐᖏᓚᓱᐅᐅᔨᓄᑦ ᐊᓘᒪᓗ ᐱᓴᓱᖅᑕᐅᔨᓚᔨᓄᑦ.....34

ᐊᕐᕌᒍᖅᓯᒪᔪᓂᒃ ᑕᔪᑎᑕᑎᑦᔪᓄᑦ 7: ᓇᐃᓐᓇᖅᑕᐅᔪᓚᔨᓄᑦ ᐊᐅᓪᓚᕕᑎᓕᖅᑐᖅᑐᒃ ᐃᔪᓚᐅᖏᖅᑐᐅᔪᕕᑎᑐᐅᑎᐅᖏᓐᓄᑦ ᐊᓘᒪᓗ ᖃᒃᓄᐃᓪᓚᐅᕕᑎᑐᐅᑎᐅᖏᓐᓄᑦ.....39

ᐊᕐᕌᒍᖅᓯᒪᔪᓂᒃ ᑕᔪᑎᑕᑎᑦᔪᓄᑦ 8: ᒪᓪᓯᓚᐅᐅᔨᓄᑦ ᖃᒃᓄᐃᓪᓚᕕᑎᖅᑎᓕᖅᑐᖅᑐᒃᓄᑦ ᑎᑯᔪᐊᖅᑕᐅᔨᓴᓇᖅᑎᓂᐊᕐᓂᖅ ᐃᒃᑐᑐᐊᖅ ᑲᓇᑕᒥᒪᒥ ᑎᖅᑎᐊᕋᖅᑕᕐᕕᖅᑎᒃᐅᓕᖅᑎᓄᑦ.....65

ᐊᕐᕌᒍᖅᓯᒪᔪᓂᒃ ᑕᔪᑎᑕᑎᑦᔪᓄᑦ 9: ᓇᐃᓐᓇᖅᑕᐅᔪᓚᔨᓄᑦ ᖃᒃᐅᓕᓴᖅᑎᑎᐅᓪᓚᕕᐅᕕᖅᑐᑦ ᓇᐅᔨᓕᖅᑐᐃᐅᕐᖅᑐᑦ ᖃᒃᐅᓕᓴᖅᑐᑦᑕᕐᕕᖅᑎᒃᓯᒪᔪᓂᓗ ᑎᖅᑎᐊᕋᖅᑕᕐᕕᖅᑎᒃ.....67

ᐊᕐᕌᒍᖅᓯᒪᔪᓂᒃ ᑕᔪᑎᑕᑎᑦᔪᓄᑦ 10: ᑕᒡᓕᒪᓂᑦ ᐊᓕᖅᓴᒍᓂᑦ ᐊᐅᓪᓚᕕᑎᓕᖅᑐᖅᑐᒃ <ᖃᓇᐅᑎᖅᓴᓂᒃ ᐊᑐᓕᖅᑎᑎᑎᑦᒡᓕᐊᓕᕐᓂᖅ.75

ᑎᑎᕃᓱᒪᓕᕆᔨᑦ ᑕᐃᔭᕐᓴᐃᑦ

ᑕᐃᔭᕐᓴᖅ 1: ᐃᓂ ᑎᖏᑕᐅᖅᓯᐅᑦ ᐊᓪᓚᓗ ᐃᓄᐃᑦ ᐅᖃᐅᓱᑐᒃᔪᑦ ᓄᓇᐃᑦ ᑕᐃᔭᐅᓂᖏᑦ (ᐱᔭᐅᓱᒪᔾᑦ ᐃᓄᐃᑦ ᐱᐅᓱᑐᖃᖅᓯᓂᖅ ᐸᐸᕋᑎᓕᑦ ᑎᑎᕃᖅᑕᐅᓂᖏᓐᓂᖅ). 7

ᑕᐃᔭᕐᓴᖅ 2: ᐃᓄᖓᓄᑦ ᓄᓇᕋᐃᑎᑭᐅᔾᔨᑦ ᓄᓇᖏᑦ ᑎᖏᑕᐊᕐᓴᖅᓯᒪᓕ. 12

ᑕᐃᔭᕐᓴᖅ 3: ᓇᔨᓚᓴᐅᑦᑐᐊᑕᕐᓴᕆᑕᕐᓂᖏᓐᓂᖅ ᐊᕐᕕᖅᑕᐅᓱᒪᓕᕆᔨᑦ ᑎᖏᑕᐊᕐᓴᖅᓯᐅᑦ (ᒪᑕᒃᑕᐅᓱᒪᑦᑐᑎᒃ ᐱᑕᓚᑕᐅᕐᖄᑦ Fᐊᖅᑎᓵ ᐊᓪᓚᓗ ᐃᓗᓚ-Fontaine and Mallory, 2011). 14

ᑕᐃᔭᕐᓴᖅ 4: ᐃᓄᐃᑦ ᕿᐅᐸᒪᔭᑐᖅᓯᒪᔨᑦ ᓇᓂᓱᐊᓂᖏᑦ ᑲᖓᐃᑦ ᑎᖏᑕᐊᕐᓴᖅᓯᐅᓂᖏᑲᒃ (ᒪᑕᒃᑐᑦ ᑲᑕᑦ, ᐱᑕᓚᕿᑎᖏᑦᑕᓗ-Carter et al. 2018-ᒥ ᕿᐅᐸᔪᓂᓂᕐᕕᓂᖏᓐᓂᒃ. ᐱᑕᕿᓴᕐᕕᓂᖏᑦ ᓴᓇᐅᒪᔾᔨᑦ ᑕᒪᐃᓐᓇᑦ ᐃᓄᐃᑦ ᕿᐅᐸᒪᔭᑐᖅᓯᒪᓕᕆᓂᖏᓐᓂᒃ ᒪᑕᒃᓱᑎᑎᒃ 21-ᓂᒃ ᐃᓄᖓᒃ ᐃᑕᕿᑲᑕᐅᖏᒪᑕ. 18

ᑕᐃᔭᕐᓴᖅ 5: ᑲᖓᐃᑦ ᐅᓪᓗᕿᑎᕐᓂᑕ ᓇᓂᓂᖏᓂ ᑐᑭᓕᐅᖅᑕᐅᔪᑎᒃ 2014-ᒥ ᓴᓪᓕᐅᑦ ᕿᑭᖅᑕᒪᔾᔪᐊᓂᒃ ᕿᑯᓕᐅᑎᓪᓗᑎᒃ ᕿᐅᐸᔪᓴᖅᑕᐅᓂᖏᓐᓂᒃ ᐊᐳᔪᓕᐅᖅᑕᐅᓕᐅᖅᑐᓂᒃ. 19

ᑕᐃᔭᕐᓴᖅ 6: ᑕᐃᒪᓐᓇᓂᒃ ᐃᓄᐃᑦ ᐊᑐᖅᑕᑦᑎᓂᖏᓂ ᓄᓇᒥᒃ ᓴᓪᓕᐅᑦ ᕿᑭᖅᑕᒪᔾᔪᐊᓂᒃ. 31

ᑕᐃᔭᕐᓴᖅ 7: ᑕᐃᒪᓐᓇᓂᒃ ᐃᓐᓇᐅᔪᓕᐅᑐᑎᓪᓗ ᐃᓅᓴᓪᑎᐱᓴᓂᒃ ᐊᖏᔪᓇᓱᒃᓯᓂᖏᓂ ᓴᓪᓕᓭᐅᑦ ᕿᑭᖅᑕᒪᔾᔪᐊᓂ. 32

ᑕᐃᔭᕐᓴᖅ 8: ᑎᖏᑕᐊᕐᓴᖅᓯᒪᓕ ᐱᔨᑲᓇᐅᑎᑕᓴᓪᔨᐊᓂᔭᓕᑦ ᐅᓄᑐᓗᑦ ᕿᓄᐃᑕᐅᑐᑕᐅᕙᖏᓂᖏᓂ 63

ᑎᑎᕎᖅᓯᒪᓂᖅᓯᓂ ᓇᐃᓴᓕᕆᐊᖅᑕᐅᓯᒪᔪᓂ ᑎᑎᖅᑐᐃᓂ ᖃᖅᓴᓄᓇᑎᑐᓂ

ACMC	ᐊᕐᓇᓵᖅᓯᒪᓂᖅᓄᓂᒃ ᐊᐅᓪᓚᕆᑎᓕᐅᖅᑕᐅᓕᕆᓂᖅ ᑲᑎᒪᓕᕎᓪᓂ
COSEWIC	ᑲᓇᑕᒥ ᑲᑎᒪᓕᕎᓪᓂᖅᓄᓂᒃ ᖃᓄᓇᐃᓕᓴᖅᑎᑐᖅᓄᓂᒃ ᐊᒥᓱᔪᓄᑦᓴᖅᖃᓪᓚᐊᓕᕎᖅᑕᐅᓯᒪᓂᖅᓄᓂᒃ ᐅᒪᔪᐃᓂᒃ
CWA	*ᑲᓇᑕᒥ ᐅᐃᒪᔪᓕᓚᓕᕐᓕᒃ ᒪᓕᓴᐃᓂᒃ*
CWS	ᑲᓇᑕᒥ ᐅᒪᔪᓕᓚᓴᓂᒃ ᐱᓕᕆᑎᕎᐃᓕᓴᖅᓯᒪᓂᒃ
IIBA	*ᐃᓄᐃᓂᒃ ᐊᖏᑐᖅᑕᐅᓯᒪᓂᖅᓄᓂᒃ ᐃᑲᔪᓱᐊᖅᑕᒋᑐᑎᖏᖅᓯᒪᓂᓄᓂᒃ ᐊᖅᑭᑐᑏᑦ ᑲᓇᑕᓕᒥᒃ ᐅᒪᒻᒪᓕᖅᓯᒪᓂᖅᓄᓂᒃ ᐊᖅᑭᑐᑏᑦ ᓄᓇᕗᑦ ᓄᓇᕈᕐᓇᐃᐅᓯᒪᔪᓄᑦ ᓄᓇᕐᔪᑎᖅᓯᒪᓂᒃ (ᐃᓄᐃᓂᒃ ᐊᖏᑐᖅᑕᐅᓯᒪᓂᖅᓄᓂᒃ ᐊᖅᑭᑐᑏᑦ)*
CIRNAC	ᑲᓇᑕᐅᕙ-ᓄᓇᖅᑐᑐᖅᑐᐃᓕᓴᒃ ᐱᓕᓚᖅᑎᑎᑦᓯᖅᓯᒪᓂᒃ ᐅᑭᐅᖅᑕᖅᑐᒥᓴ ᐱᓪᓕᓕᐊᔪᓕᓚᔪᓂᒃ ᑲᓇᑕᒥ
DFO	ᐃᖅᑲᓴᓕᓚᓵᖅᓴᔪᐊᖅᑯᓂᒃ ᑕᓚᐅᖅᑎᐅᑕᓕᓕᓚᓵᕙᓴᓴ ᑲᓇᑕᒥ
ECCC	ᐊᖅᑭᑎᓕᓕᓚᓵᖅᑎᐊᖅᑯᓂᒃ ᓱᓕᐅᕙᓴ ᐊᓱᓵᖅᐸᕙᓕᓂᔪᓄᑦᓯᓄᓂᒃ ᑲᓇᑕᒥ
GN	ᓄᓇᕗᓂᒃ ᓴᖅᓚᐃᓂᒃ
HTO	ᐊᖏᕐᓇᓱᑉᑎᓂᒃ/ᒪᖅᑲᐃᑎᓂᒃ ᒥᑭᑎᐊᖅᓂᐊᖅᑎᕙᓴ ᑲᑐᓵᖅᑲᑎᑎᑦᓯᓂᒃ
IHT	ᐃᓄᐃᓂᒃ ᐱᐅᓱᑐᖅᖃᓂᓪᓂᑲᓴ ᐸᐸᕈᑎᓴᓂᒃ
IOL	ᐃᓂᓄᓂᒃ ᐱᖅᑯᑎᕐᓴᐅᔪᓂᒃ ᓄᓇᐃᓂᒃ
IQ	ᐃᓄᐃᓂᒃ ᖃᑲᓵᒪᔪᑐᖅᓯᒪᓂᒃ
KivIA	ᑭᓚᓕᓕᒥ ᐃᓄᐃᓂᒃ ᑲᑐᓵᖅᑲᑎᑎᑦᓯᓂᒃ
MBCA	*ᑎᓪᓗᐊᖅᓂᓴ ᐅᑎᖅᐸᓴᑐᓂᓴ ᑲᑎᓪᓚᒪᐅᑕᐅᓗᑎᓴ ᒪᓕᓚᓕᐊᕐᓯᓚᒪᔪᓂᒃ*
MBS	ᑎᓪᓗᐊᖅᑲᖏᐃᓂᒃ
MBSR	*ᑎᓪᓗᐊᖅᑲᖏᓕᓂᒃ ᒪᓕᓚᓪᓗᑐᐃᓂᒃ*
NIRB	ᓄᓇᕗᒻᒥ ᐊᖏᑎᐸ ᐊᖅᑐᖅᑕᐅᔪᓐᓇᖅᓂᖅᓯᒪᓂᒃ ᑲᑎᒪᓵᓂᒃ
NA	ᐊᖅᑭᑐᑕᐅᓚᒪᔪᓂᒃ ᐃᓄᓂᒃ ᓄᓇᕈᖅᑕᐅᓚᒪᔪᓚᓄᓂᒃ ᓄᓇᕗᒻᒥ ᑯᐃᓐᒍᓪᓴᓪᓚᖅᑐᖅᓂᖅᓯᒪᓄᓂᒃ ᑲᓇᑕᒥ, ᑕᐃᔨᐅᖏᖅᓯᒪᔪᖅᓱ ᓄᓇᕗᒻᒥ ᐊᖅᑭᑐᑕᑎᓂᒃ
NPC	ᓄᓇᕗᒻᒥ ᓄᓇᐃᓂᒃ ᐊᑐᑎᑦᓚᖅᑭᑐᓂᒃ ᐸᖅᓇᐃᓴᓂᒃ ᑲᑎᓯᓇᐃᓂᒃ
NTI	ᓄᓇᕗᓂᒃ ᑐᓂᖏᓚᐅᑯᓂᒃ ᑎᑎᖅᑯᑎᖏᓕᓂᒃ
NuPPAA	*ᓄᓇᕗᒻᒥ ᐸᖅᓇᐃᓂᕐᓕᒃ ᐱᓕᓚᐊᐅᓚᓂᑐᓴ ᖃᐅᓴᓚᑐᓇᖅᓂᕐᓕᒃ ᒪᓕᓚᓕᐃᓂᒃ*
NWA	ᑲᓇᑕᓕᒧᒥ ᐅᒪᔪᖅᖃᓚᐅᔪᖅ
NWMB	ᓄᓇᕗᒻᒥ ᐅᒪᔪᓕᓚᓵᖅᓴᔪᐊᖅᑯᓂᒃ ᑲᑎᒪᓴᖏᓂᒃ
SARA	*ᐅᒪᔪᐃᓂᒃ ᐊᒥᓱᓕᑐᓕᓂᖅᓯᒪᓂᖅᓄᓂᒃ ᒪᓕᓚᓕᐃᓂᒃ*

1 ᐅᖃᐅᓯᐅᑎᐊᖃᓂᖅᓱᓕᕆᓂ ᓴᓇᓂᐊᖅᑕᐅᓯᒪᔪᓂ ᓄᓇᕗᑦ

ᑕᒪᓐᓇ ᐃᑲᑦᑐᐊᖅᑎ ᑎᓯᒥᐊᖃᖅᖃᐃᓐ ᐱᓐᓴᖅᑎᑕᐅᓕᐅᖅᓯᒪᔪᖅ 1959-ᓴᑎᑦᓗᐃ, ᐊᑐᖅᑕᐅᓗᑎᒃ ᑲᓇᑕᓕᒫᒪᑦ *ᑎᑭᒃᖃᓗᓴᕐᕕᒃᑐᑦ ᑲᑎᒪᖃᖅᑕᐅᕆᐅᓂᖅᓄᓐᓄᑦ ᒪᓕᑕᓕᐊᑦ, 1994-ᒥ (MBCA-ᒪᓕᑕᓕᐊᖅᑎᐃᔪᑦ)* ᓴᐳᓂᐊᖅᑕᐅᑭᐅᐊᖅᓯᑎᒃ ᐊᒥᓯᑭᖅᓯᐊᓴᐅᑦᓯᓗᑎᓗ ᑎᑭᕐᕕᑦ ᑎᓯᒥᐊᑦ. ᑎᓯᒥᐊᖅᑲᖃᒃ ᑕᒧᓂ ᑭᐸᓵᓴᑐᑐᒃ 1,433 km² ᑭᓚᒥᑕᓂᒃ ᐊᓯᑎᓂᓕᒃ, ᐃᓂᖅᑲᖅᓯᓂ ᐅᐊᓂᓇᖅᑕᓯᓂᒪᑕ ᑕᓓᐅᓂᒪᓂ ᕼᐊᑦᓴᐳᐧᐃᐸ, ᓴᑦᓂᒃᐅᐸ ᖁᑭᖅᑕᓵᐊᓂ, ᓇᓐᒃᒪᒥ. ᑎᓯᒥᐊᖅᑲᖃᐃᐅᔪᖅ ᑐᑭᒧᐊᒃᑐᑦ ᐅᓂᒪᑦᓇᓂ 50 km ᑭᓚᒥᑕᓂᒃ ᑕᑭᓂᓂᒥᒃ ᐅᒃᑐᑎᐅᑎᒃᑯᑦ ᐊᓚᒪᓗ 40 km ᑭᓚᒥᑕᓂᒃ ᓂᑕᑐᓂᖅᑭᖓᓂᖅᓱᓂ (ᑕᑯᔭᒃᓴᖅ 1).

ᐃᑲᑦᑐᐊᖅᑎ ᑎᓯᒥᐊᖅᑲᖃᐃᖅ (ᐃᑲᑦᑐᐊᖅ) ᐃᓂᓂᒃ ᐃᓗᐊᓂ ᑯᓘᐃᐅᕐᓂᖅᓯᑎᓂᓄᑦ ᐃᑲᑐᐊᑦ ᑯᓘᒪᓗ ᐃᓕᓵᖅᑯᓂᒃ ᐊᓚᒪᓗ ᐃᓕᖅᑲᖅᓯᑎᒃ ᓇᓇᒪᓐ ᐃᓗᑐᓂᐊᓗᓵᖅᒥᒃ ᑯᓘᐃᐅᖅᑎᓂᓗᓪᓕᒫᓂᒃ ᐃᓕᑐᐊᑦ ᑯᓘᒪᓗ ᐊᑯᖅᑐᓂᒃ ᐅᓪᓗᑎᒃ ᑕᐅᐅᓂᒃ ᐊᐅᓚᐅᑎᓂᓗᓪᓕᒫᓂᒃ ᓴᖅᑭᑦᓯᐊᕐᓂᖅᑐᓂᒃ. ᐱᐸᖅᑐᖅᑲᖅᑐᖅ ᐃᐊᖏᓴᓂᓂᒃ ᓇᓇᒪᓐ ᐃᓗᑐᓂᐊᓗᐊᑦᓂᒃ ᐅᓪᓗᖅᑕᕐᓇᓂᓂᓕᒫᓂ ᐊᒥᓱᓂ ᐊᓱᐅᑦᓯᑐᓄᐅᓂᒃᑕᕐᓇᓂᓂᓕᒫᓂ ᑕᐅᐅᓂ, ᐃᒪᓕᒫᓂᓱᐊᓂᒃ ᐅᓪᓗᖅᑕᕐᓇᓂᓂᓕᒫᓂ ᐊᒥᓱᓂᓪᓗ ᐊᓱᐅᑦᓯᐊᓂᒃ ᓇᓇᖅᐱᐅᓂᓂᑎᓱᐊᓂᒃ. ᑕᒪᐃᓂᐊ ᐱᐅᖅᐃᐊᐅᖅᑕᓂ ᑭᓚᓇᖅᓂ, ᒪᓴᓴᐅᓂᖅᓯᒪᔪᓂ ᐊᓯᖏᓐᓄᑦ ᓇᓇᖅᑐᐃᓂᖅᓱᓂ ᑕᐅᐅᓂ ᐃᒪᓕᓴᖅᑭᑦᓯᐊᓂᖅᓱᓂ ᐊᒥᓱᓂᓪᓗ ᐊᓱᐅᑦᓯᐊᓂᒃ ᐊᓯᖏᓐᓄᑦ ᓇᓇᖅᑐᐃᓂᖅᓱᓂ ᒪᓕᒫᓂᒃ ᐃᓕᓴᕐᓂᖅᓱᓂ ᐊᒥᓱᓂᓪᓗ ᐊᓱᐅᑦᓯᐊᓂᒃ ᓇᓇᖅᑐᐃᓂᖅᓱᓂ ᓴᓇᓕᒫᓂᒃ ᐊᒥᓱᓂᓪᓗ ᐊᓱᐅᑦᓯᐊᓂᒃ ᓇᓇᖅᑐᐃᓂᖅᓱᓂ

ᑎᖏᑕᐊᖅᖃᐅᔨᑦ ᐊᑐᑎᖅᑲᖅᓯᒪᓂᖅᓱ ᓴᓗᓂᓄᖅᑕᐅᖅᖁᐃᐅᓂᖅᓱᓇᓐᑦ, ᖃᐅᐅᓰᒪᔨᐅᓱᓇᓐᖅᓯᑎᖅ ᐱᓪᒪᓇᐅᓂᖅᑭᑦ ᑲᖏᖅᓴᓐᑦ ᑎᖏᑕᐊᖅᓗᓐᑦ.

ᑕᐃᖅᑯᐊ *ᑎᖏᑕᐊᖅᓂᖅ ᐅᑎᖅᐸᖅᑐᓂᖅ ᑲᑎᓪᒪᐅᑕᐅᓗᑎᖅ ᒪᑕᓚᑕᐊᖏᓯᒪᔪᑦ MBCA* ᐊᓪᒪᓗ *ᑎᖏᑕᐊᖅᑲᖅᓴᓐᓂᖅ ᒪᑦᓗᓚᑦᓗᑐᖏᑦ (MBSR-ᑯᑦ)* ᐱᔪᓐᓇᖅᑎᑦᑎᐅᖅᑦ ᑎᐱᓯᒪᖅᓴᐅᓇᖅᑕᖅᑕᑭᓐᓇᖅᓂᖅᓴᓂᖅᑐ ᐃᑲᑦᑐᐊᐸᑦ ᑎᖏᑕᐊᖅᑲᖅᓴᓐᓂᖅᓱᓇᓐᑦ. ᓄᓇᕗᓪᒥᐅᑕᐃᑦ ᐃᓄᐃᑦ ᐱᔪᓐᓇᖅᓂᖅᑲᖅᑐᑐᕗᑦ ᐊᖏᔪᓯᑎᐊᖅᑲᑦᑕᑭᓐᓇᖅᓂᖅᖃᖅ, ᐊᓪᒪᓗ ᐱᔪᓐᓇᐅᑎᖅᖁᑎᖅᑲᐅᑎᖅ ᐱᑕᖃᔪᓐᓇᖅᑐᖅᑕᑦ ᐊᖏᔪᓯᒪᓕᖅᑐᑦ. ᑕᒪᖁᖅ ᐊᓯᖏᑦ (ᓯᑦᓗ ᐃᓄᑦᑕᓪᒪᐅᖏᑦᑐᑦ), ᒪᑕᑐᖅᑎᔪᑦ ᐊᔪᖅᑎᑦᑕᐅᔪᑦ ᑎᐅᔪᒪᔪᓐᓇᖅᓂᖅ ᑕᐅᕐᖏᖅᑐ, ᐱᓯᐊᓂ ᐱᔪᓐᓇᐅᑎᖅᖁᖅᑎᑕᐅᔪᑎᑎᖅ ᐊᐋᑎᑕᓪᒪᓇᔨᔪᐊᖅᑯᑦ ᐅᒪᔪᑕᓪᒪᐅᖏᑦᓄᓐᓗ. ᒪᑯᓂᖅᑐᑦ ᐊᑐᑭᓐᓇᑦᑎᔪᑦ ᒪᑕᑐᖅᑎᔪᑦ 1) ᑎᖏᑕᐊᖅᓯᐅᑭᓐᓇᖅᓂᖅᖁᖅ, 2) ᔪᖅᓚᕆᐃᔪᓐᓇᖅᓂᖅᖁᖅ, ᓯᑎᖅᑎᓪᒪᔪᓐᓇᖅᓂᔨᓐᓂᑦ ᑎᖏᑕᐊᑦ ᒪᓐᖏᓐᓂᖅ, 3) ᐱᓯᒪᔪᓐᓇᖅᓂᖅᖁᖅ ᐅᒪᔪᓂᖅ ᑎᖏᑕᐊᖅᓂᖅ, ᑎᒥᖏᓐᓂᔨᓐᓂᑦ, ᐊᒥᖏᓐᓂᖅ, ᐅᑦᓗᖏᓐᓂᖅ, ᒪᓐᓇᖅᑎᓪᒪᔨᓐᓂᑦ ᑎᖏᑕᐊᑦ, 4) ᖃᐅᑭᐅᓯᕈᓐᓇᖅᓂᖅᖁᖅ ᐊᓯᖏᓐᓂᖅᓱᔨᓐᓂᑦ ᐊᖏᔪᓯᐳᑐᖅᓯᐅᑎᓂᖅ ᐊᑐᑕᓐᓇᖅᓂᖅᖁᖅ.


ᐊᕐᓇᖅᑐᓂᓗᐊᓂᖅ ᑕᑯᑎᑎᑦᓴᐊᑦ 1: ᐃᒃᑕᑐᐊᖅᑐᒥ ᑎᖏᒥᐊᖅᑎᓇᖏᑦᑐᒃ ᓀᐃᐹᓕᓴᐅᓂᖅᑐᑎᒥᐊᓂᖅ ᖃᐅᔨᓴᖅᑎᑦᑎᐊᕈᑎᓯᑦ

ᓴᓗᒥᐊᖅᑕᐅᓂᒪᖕᓇᖅᓯᓂᒃ	ᑎᖏᒥᐊᖅᑎᓇᖅ ᓴᓗᒧᐊᖅᑕᐅᖕᖏᓇᖅ
ᒪᑕᑦᓂᖅ ᒪᑕᔪᑕᐅᔨᑦ ᓴᓗᒧᐊᖅᑕᐅᔨᒧᓐᓇᖅᓂᖅᓯᒪᑦ ᑎᒃᓀᐊᖅᑕᐅᓂᒪᖕᓇᖅᓯᓂᒃ	ᐃᒧᑦᑎᓇᐅᔨᑦ ᐃᓂᒪᑎᓴᐅᔨᒧᓐᓇᖅᑕᑦ ᐅᓪᒪᖏᑕᑎᖏᓇᐅᔨᒧᓐᓇᖅᓂᖅᓴᓗᒧᐊᖅᑕᐅᓂᒪᖕᓇᖅᓯᓂᒃ ᓀᐃᐹᓕᓴᐅᓂᒪᖕᓇᖅᑎᑦ ᒪᓀᓂᓴ ᒪᑕᓴᓂᒪᓕᕐᓗᑎᒃ ᐊᑕᐅᓂᔪᑦᑐᒃ ᐊᒥᓂᔪᓂᒃᓴᐅᓂᓚᐅᖅᓂᑦ: 1. ᐱᑕᖅᑎᓇᐅᔪᑎᒃ ᐊᒥᓂᓂᒃ, ᖃᖏᓴᑐᐃᓐᓇᐅᑎᓂᖅᓴᓂ ᐊᕐᕿᔪᒥ, ᒪᑕᔪᒪᓂᕐᒧᑦ ᐊᑕᐅᓂᔪᑦᑐᒃ ᐊᒥᓂᓂᒃᓴᐅᔨᓂᓚᐅᖅᓂᑦ ᐱᑕᖅᑎᒪᐊᖅᑎᒃᖅᑐᓂᒃ; ᑕᐃᒪᐃᐅᓂᖏᑦᓂᓗ, ᓄᓇᐃᑦ ᐊᑐᑎᑕᒪᒪᓕᐅᔪᑎᒃ ᐱᑕᖅᑎᒪᐊᖅᑎᒃᖅᑐᓂᒃ ᐊᐅᑕᓪᓕᐅᓇᓂᖅᓴᖅ ᐊᕐᓇᖅᑐᖃᓕᖏᑐᓂ ᐱᒋᔨᓂᒃ ᐱᑕᖅᑎᓇᐅᑦᑐᒃᓴᓂ ᑎᒥᖅᑐᓂᒃ ᐅᑎᖅᑕᒃᑐᓂᒃ. 2. ᓄᓇᒥ ᐃᓂᖏᑦᑕ ᐊᒃᑐᖅᑕᐅᕿᑎᐃᓂᐅᓂᖅᑎᒃᑯᑎᑎᒃ. ᐱᓕᒪᓕᐅᔪᒥᒃ ᐅᒪᔨᖏᑦᑕ ᐊᒃᑐᖅᑕᐅᔨᒪᔨᓇᕈᑎᒃ, ᐅᓗᓕᐊᓇᖅᑐᖅᑭᓇᖅᑐᓂᒃ ᐊᒥᔨᓗᐊᖅᑐᓄᓪᓕ ᐊᖏᔪᓇᙵᓚᐅᖅᓯᒪᓇᖅᓂᖅᓴᖅ, ᖃᓄᖅᓚᐅᔪᓇᓇᖅᓂᖅᓴᖅ ᐱᓯᒪᓕᐊᔪᑕᐅᔪᒪᔨᓂᒃ, ᐊᔨᖏᑐᓂᑐᓗ. ᑕᒪᑯᐊᐊ ᐱᓕᒪᓕᐅᔪᑦ ᓇᔪᖅᑕᐅᓯᒃᑐᑦ ᐃᑕᖅᑭᓇᓕᖅᑎᑦ ᐃᓚᐅᓯᒃᑐᓂᖅᓴᖅ ᐅᑦᓗᖅᓴᖏᓚᐅᖅᓯᒪᓂᖅᓴᖅ, ᑎᖏᑦᓇᓇᐃᑕᔨᒪᓚᐅᖅᓴᖅᑎᖅᑕᑦᓂᖅᓴᖅ ᐃᓕᓴᖏᓚᐅᖅᓯᒪᓂᖅᖏᑦᑐᓂᒃ, ᐅᑭᐃᓚᐅᖅᓯᒪᓂᖅᖏᑦᑐᓂᒃ, ᐅᖃᓚᐅᖅᓂᒃ ᐅᒪᔨᖏᑦᓄᓪᓕ ᓄᓇᒥᖅᓯᒪᓂᖅᖏᑦᑐᓂᒃ ᐊᑐᐊᖅᑐᓂᖅᓴᖅ ᐃᓄᖅᓯᒪᓂᖅᖏᑦᑐᓂᒃ. 3. ᐃᓂᖅᑎᓇᐅᓯᒃᑯᑎᒃ ᐅᒪᔨᖏᑦᓄᓪᓕ ᐃᓕᐃᓐᓇᖅᑎᓂᖅᑕᑭᓐᓇᖅᑐᓂᒃ ᓄᓇᓂᒃ ᐊᓪᓚᓗ ᐊᒃᑐᖅᑕᐅᓕᕐᓗᐊᑦᑎᓐᓇᖅᑐᒃ ᐃᓄᖅᓄᓪᓕ ᕿᓚᕕᖅᑕᐅᔨᒪᓚᐅᖅᓯᒪᓂᖅᖏᑦᑐᓂᒃ ᐃᓄᖅᓄᓪᓕ ᐅᖅᑲᐅᓯᒃᑯᑎᒃ ᐃᓄᖅᓄᓪᓕ ᐱᑕᖅᑭᓇᓕᖅᑎᓂᖅᓴᖅ ᐅᑦᓗᖅᓴᖏᓚᐅᖅᓯᒪᓂᖅᖏᑦᑐᓂᒃ ᐃᓕᓴᖏᓚᐅᖅᓯᒪᓂᖅᖏᑦᑐᓂᒃ ᐅᑭᐅᓂᖅᓴᖅ ᐃᓕᓴᖏᓚᐅᖅᓯᒪᓂᖅᖏᑦᑐᓂᒃ

ᑲᓇᑕᐅᑉ ᐊᕐᓚᑲᖏᑦᑐᒍᓐᓇ ᐅᖃᖅᓯᓂᒃ ᑲᓇᑕᐅᑉ ᐅᑭᐅᖏᑦᑕᖏᑦᑐᖅᓯᐅᒍᓐᓇᑕ ᐊᕐᓚᑲᖏᑦᑐᒍᓐᓇ	ᓄᓇᕗᑦ
ᐊᕐᓚᑲᖏᑦᑐᒍᓐᓇᕐᓂᕐᒥᑦ	ᑭᖅᑎᓪᓗᖅ
ᐃᓕᐅᕈᕆᐅᒍᓐᓇᕐᓂᒃ ᖃᐅᔨᒪᓂᒃᑯᑦ ᐊᑲᖏᑦᑕᐅᓯᒪᔪᑦ ᓄᓇᓕᒫᑦ	ᓴᖅᑎᓪᓗᖅ
ᓄᓇᖓᐊᑎᒍ ᑎᑎᖏᐅᖅᓂᒃ ᓴᓂᒍᑦ ᑲᓇᖅᑐᒥᐅᑦ/ᐱᖓᓱᓂᑎᒍᑦ ᐊᓚᒪᓗ ᑐᑭᒍᑦ ᐅᐊᖓᓇᒥᐅᑦ ᓂᕈᕐᒍᑦ	63°45’ N ᐅᐊᖓᓇᒍᑦ/85°40’ W ᐱᖓᓱᓇᒥ
ᐊᖓᕈᓂᖅ	1,433 km² ᑭᐸᓰᓴᐅᑐᓂ ᑭᓯᒥᑕᓂᒃ, ᐃᓕᖏᑲᖏᓂᐅᒃ 190 km² ᑭᐸᓰᓴᐅᑐᓂᒃ ᑭᓯᒥᑕᓂᒃ ᑕᓚᐅᖏᒥ ᐃᒪᖏᒃ ᐅᒪᔨᖏᑲᐃᐅᓂᖓᕈᓐᓃᓂᒃ
ᐳᖏᑐᓂᖏᑲᖏᑎᑕᓂᖓᓂ (m ᒧᑕᒃᑯᓂ)	ᑕᓚᐅᐸ ᓇᑎᐊᓂᒃ ᑕᒪᐅᖓᓚ ᖏᑯᒪᒧᒃ ᐃᒪᖓᓕ ᐳᖏᑐᓂᖏᑲᓰᓚᖃᖓᓚᓄᓕᓂ ᑎᑭᐅᒪᓂᖏᑲ ᖏᓂᕐᖓᓚᓄᓕᓂ 60 m ᒧᑕᓕᓐᓄᓂ
ᐊᖏᖃᔪᒥ ᓴᖏᑭᑎᑕᐅᓯᒪᔪᑦ ᒪᓕᖅᓕᐊᑦ ᐅᖏᓀᒪᒪᓚᖏᑎᔪᑦ ᑲᓯᓕᑯᑦ-Gazette-ᑯᑦ)	1959
ᓴᐳᓂᐊᖏᑕᐅᓯᒪᕐᐃᐅᓂᖏᒥᑦ ᖏᓀᓄᐃᓕᓐᓂᖓᓕᒫᓂᒃ ᑎᑭᐅᑎᑎᑦᑕᐅᓂᒃ	ᑕᐃᒪᓐᓂᓂᒃ: ᓴᓴᑭᑎᑕᐅᓕᐅᖏᓯᒪᔪᑦ ᓴᐳᓂᓂᐊᖏᑕᐅᖏᖏᑦᑐᓗᑎᑦ ᐃᕐᕐᐃᐅᑕᖓᓂᒃ ᐅᓪᓗᓕᐅᕐᓂᖓᓂᒃ ᓄᓇᐃᑦ ᑲᖓᓯᑲᓂᒃ (10% ᐳᓴᓂᑎᖓᓂᑐᑎᒃ ᓄᓇᖏᔪᐊᓕᒫᒥ ᑲᖓᓯᓄᑦ ᐊᑐᖏᑕᐅᕐᓂᖓᕈᓐᓃᑦ, ᑕᒪᑯᔨᓂᖓᓚᓗ (Atlantic-ᓂᒃ) ᓂᕆᓂᖓᓇᖏᑦ (4% ᐳᓴᓂᑎᓂᐅᑐᑎᒃ ᓄᓇᖏᔪᐊᓕᒫᒥ). ᐃᓂᓈᐅᓕᓂᖏᑐᓂᒃ: ᓄᓇᖓᓚ ᐱᑕᖏᑲᖏᐅᔪᖏᒃ 5% ᐳᓴᓂᑎᖓᕐᓂᑐᑎᒃ ᑲᓇᑕᐅᑉ ᓄᓇᖏᔪᐊᖓᓚᑕ ᖃᑎᖏᐸᓯᐊᓂᓐᓇᓴᑲᑦᑕᖏᑐᑎᒃ ᒥᑭᓈᓂᖏᖏᑲᐅᓂᖏᑕᐅᔪᓂᒃ ᑲᖓᓯᖏᑦ ᐊᑐᖏᑕᐅᕐᓂᖓᕈᓐᓃᑦ, 1% ᐳᓴᓂᑎᖓᓂᓐᓂᑦᑐᓗ (Atlantic brant-ᓂᒃ) ᓂᕆᓂᖏᓇᑲᓰᓴᔪᓐᓇᑐᓂ (ᑕᐃᒪᐃᓇᓯᒪᔨᐅᑐᐃᓇᖏᑦᑐᑎᒃ).
ᓴᐳᓂᐊᖏᑕᐅᓯᒪᓕᖓᓂᓈᓂᒃ ᓇᓐᓇᓇᐃᔨᖏᑲᐅᓯᒪᔭᑎᖓᓂᒃ	ᐊᕐᓚᑲᖏᑦᑐᒍᓐᓇᑕᒃ A-ᒥᒃ, ᖏᑯᑎᑲᖏᑦᑐᒃ ᐱᑕᖏᑲᓚᐊᖏᑲᓕᓚᓂᖓᓂᓈᓂᒃ ᐃᓕᑕᓚᔨᐅᔨᔪᑎᑦᓂ ᐅᒪᔨᖏᑲᓂᖓᓂᓈᓂ ᓇᔪᖏᓚᓕᑲᑲᑎᑲᑎᐊᓚᖏᑲᓂᖓᓂᓈᓕᓐᓂᓗᓂᑦᓂ ᓴᐳᓂᐊᖏᑕᐅᓯᒪᓗᑎᒃ
ᓄᓇᖏᔪᐊᖏᒥ ᓄᓇᕗᓕᔪᐊᑦ ᑕᐱᓱᓂᓂᑦ ᓴᐳᓂᐊᖏᓂᖏᒥᑦ ᐱᐅᓯᐅᑦᑕᐅᓯᒪᔨᐅᓂ IUCN-ᑯᑦ	ᐊᕐᓚᑲᖏᑦᑐᒍᓐᓇᑕᒃ 1b (ᐅᒪᔨᖏᑲᖏᐃᑐᑲᐅᖓᓂᓈᓕᓐᓂᖓᓚᓄᓕᓂ)
ᑲᖇᒪᐃᑦ ᒥᓂᑕᐃᑦ ᑲᑎᒪᔨᒥᑦ ᓴᓇᓕᒫᑦ	P.C. 1959-629 (SOR/74-514)

ᓇᓗᓇᐃᖅᑯᓂᖅ ᑲᓇᑕᐅᕐ ᓄᓇᕗᑎᖓᓂᒃᑐᓄᖅ ᐱᕐᔪᐊᑎᖓᓂᒃᓴᓂ ᓱᕐᔪᐊᑦ	ᐊᑐᐃᓂᖅᑐᑦ
ᐃᓄᖃᓂ ᐱᕐᔪᐊᑎᖓᔪᐊᕐᓂ ᓄᓇᐃᓂ (ᓄᓇᕗᑦᑎᓂᒃᑯᓂᖅ ᐊᐃᓕᖅᑐᓂ)	**ᐊᐃᒃᑐᑐᖅᓯᒪᔪᓂ ᓇᓗᓇᐃᓄᑦ**	**ᐱᕐᔪᐊᑎᖓᓂᒃᓴᓂᖅ ᐊᓂᖅᓂᖅᓂ**	**ᑲᑎᒪᔨᖅᑕᐃᓕᓂᖅ (ᐄᐃᐊᒃᑕᐃᓕᓂᑦ)**
	CH-06	ᓄᖅᑎᓂᑐᑦ ᓄᓇᐃᓂᑦ	714km^2 ᑭᒃᑯᓕᖅᑐᑦ ᑭᓪᓗᒍᑦ
ᐊᓯᖅᓂᖅᓯᒪᓂᖅᑕᐅᓄᖅ ᑎᖅᑯᐊᓄᖅᑕᐅᓯᒪᓕᖅᓯᓂᖅ	▪ ᐃᖅᑐᐊᑦ ᑯᖓᒥᓂ (ᓄᓕᑦᓴᓇᑎᑐᑦ ᑕᐃᔭᐅᔪᒥᐅᖅ ᐊᐊᓯᑦ ᑯᖓᒥᓂᖅ) ᓄᓇᐃᑦᓴ ᐊᕐᓚᐊᓂ ᒪᓕᓴᐃᓂᖅᓯᖅ ᑎᓯᒥᐊᓄᖅᑐᑎᑦ ᐅᓇᓱᐊᕐᓂᖅ (NU022) ▪ ᐱᓕᓕᐅᓂᖅᓯᖅ ᓄᓇᐃᓂ ᑎᓯᒥᐊᓄᑦ ᓄᓇᑭᒪᒥ (ᓄᓇᖓᓂ 13) ▪ ᐃᖅᑕᐊᑦ ᑯᖓᒥᓂ (ᑕᐃᔭᐅᓂᖅᓂ ᓴᓪᓕᐊᑎᑦ ᐊᐊᓯᑦ ᑯᖓᒥᓂᖅ) ᓄᓇᕐᔪᐊᑕᒧᒥ ᐅᓪᓚᐊᓴᓄᖅᑐᓂᖅᓂᓄᑦ ᐱᑕᓕᐊᐅᓂᖅᓂᖓ (ᓄᓇᖅᑐᓂ 6-5) ▪ ᐱᓕᓕᐅᔪᓴᖅ ᓄᓇᒥ ᓇᔪᓴᑕᐅᖅᓂᐊᖅ ᑎᓯᒥᐊᓄᑦ (NU ᓄᓇᖓᓂ 43)	
ᓇᓂᓂᖅᓯᑦ ᐅᓚᔪᓴᓂᓪᓂ ᓄᓇᐃᑦ ᐊᓚᒪᓗ ᐱᑭᓴᖅᐃᐊᔪᓴᓴᖅᑐᑦ ᐱᓕᓕᐅᓂᖅᓯᑦ	• ᐱᓕᓕᐅᔪᓂᒃ ᑎᓯᒥᐊᓴᓂᓪᓚᖅᓄᓕᑦ ᒪᓕᐅᖅᑐᑦ ᒪᓕᓕᐅᓂᓴᓂᖅᓴᖅᓯᓂᕐᓂᐊᔪᓄᖅᓯᒥᑲ: ᑕᒪᓂ ᐊᑭᐅᓄᖅᓯᑐᑎᓂᒃ ᐊᒥᓱᓂᒃ ᐅᓚᔪᓴᓴᓂᓪᓚᖅᓄᓕᑦ, ᐃᒪᖏᒥᑕᓂᒃ, ᑎᓯᒥᐊᓄᖅᑐᑦ ᐊᒥᓱᓂᒃ ᐊᑭᐅᓄᖅᓯᑐᑎᑦᑐᐅᔪᓂᒃ • 5% ᐳᓴᓄᖅᓯᐅᓂᑦ ᑲᓇᑕᐅᔪᓗ ᐊᒥᐊᓕᑲᐅᔪᓗ ᓄᓐᓗᐊᓂᓱᓂ ᑲᖏᐃᑦ ᓄᓕᐊᓴᖅᑕᖏᒪᑕ ᑕᒪᓂ • 4% ᐳᓴᓄᖅᓯᐅᓂᑦ ᓂᖏᓕᓴᐃᑦ (Atlantic brant) ᑕᒪᓂ ᓄᓕᐊᓴᖅᐸᓪᐅᔪᖏᑦᔪᓂ ᓱᐅᓂᐊᒍᑦ • ᐱᑕᓴᖅᓇᐅᐅᖅᑕᖅ ᑕᒪᓐᓇ ᑎᑭᓱᒪᖏᓇᐅᖅᑲᐃᓐᓇᓴᑕᖅᑕᓂᓴᖅ ᑲᔪᓱᐅᓕᓪᓕᐊᓂᖅᓄᓂ ᓂᑭᖅᐸᓱᓚᑦ ᓴᓄᖅᑕᓪᓕᐊᓪᓕᖅᑎᓪᓗᑎᑦ ᓴᓪᓚᖅᑕᐃᓕᐅᖅᑐᑦ ᐅᑎᖅᐸᓕᓪᓕᐊᓪᓕᖅᑎᓪᓗᑎᑦ ᐅᑭᐅᖅᑕᖅᑐᑦᓯᐊᖅᑐᑦ ᐅᓕᓴᖅᓯᐅᓕᐅᖅᑐᑦ ᓱᓴᖅᐸᓱᓕᕐᓂᖅ ᑎᓯᒥᐊᑦ ᓱᓴᓗᐊᓕᔪᓚᐅᑦ • ᐱᓕᓚᐊᔭᔪᖅ ᐊᐅᓵᖅᑯᑦ ᐃᓄᖅᓴᖅᐊᐅᖅᓯᓂᖓᓂᓄᑦ • ᐃᓵᖅᑐᓴᖅᓯᖅᑎᑦᑎᐅᑦ ᑐᖅᑐᐃᑦ ᐃᓚᖏᓂᓄᖅ ᓄᓇᓂᖅ ᑕᒪᓂ ᐊᑐᖅᐸᓚᐅᖅᑐᑦ ᓄᓕᐅᑦᓴᓪᓚᐅᑦ	
ᑕᒪᓂᖅᑎᐅᑕᑐᖅᑲᐅᖏᓪᓕᖅᑐᓄᓇᓂᑦ ᐅᓚᔪᓄᑦ/ᖃᐅᔨᐱᖓᓄᑦ ᐱᑕᖅᓴᐊᐅᓕᕐᓂᖅᓂᖓ	ᖅᑲᐅᔭᒪᔨᐅᔪᖅᓯᖅᑐᑦ ᓂᓚ

ᐅᒪᓪᐃᐧᐃᑦ ᐊᒥᓯᖏᑕᐊᓕᕐᓂᖏᑦ	**ᒪᓕᒐᖅᑎᔪᑦ ᐃᓕᓴᕆᐅᓯᒪᓂᖏᑦ ᑲᓇᑕᒥ *ᐅᒪᓪᐃᐧᐃᑦ*** ***ᐊᒥᓯᖏᑕᐊᓕᕐᓂᖏᖅᓄᓂᑦ ᐃᓕᓴᕐᑎᑎᔪᑦ*** ᓄᖓᒍᓕᑐᕐᖏᑐᖅ: ᑕᐃᓪᓕᑕᐃᓪᓕᓯᖅ (ᓴᐅᑦᕋᐃᖅ) ᐊᐸᐅᖏᖅᑕᑐᒃ ᐃᓯᒪᓕᑕᐅᓂᖏᑦ: ᐊᐸᐅᖏᖅᑕᑐᒃ ᐃᓯᒪᓕᑕᐅᓂᖏᑦ: ᓇᓄᖅ ᓴᐅᑦᕋᖅ ᑲᔪᓂᖅᓴᖅ ᒥᑭᓂᖅᓴᖅ ᑲ ᖁᖏᕐᐃᖅ **ᖁᑲᐅᓴᖅᑕᐅᓯᒪᔪᑦ COSEWIC-ᑯᓄᓂᑦ ᑲᓇᑕᒥ ᑲᑎᒪᓴᕋᓯᖏᓐᓂᑦ** **ᖁᓄᐃᑕᖏᓚᓂᖏᑭᓄᓂᑦ ᐊᒥᓯᔪᓐᓇᖅᐸᑦᓯᓕᐊᓂᕋᖅᑕᐅᓯᒪᔪᓄᑦ ᐅᒪᓕᔪᐃᑦ** ᐊᒥᓯᖏᑕᐊᓕᕐᓂᕐᒧᑦ ᐅᓗᓚᐊᓇᓕᓂᒥᒋᑦᑕᖅᑐᑦ: ᓇᓵᐸᖅᑐᓕᖏᑕᕐᒧᑦ ᑐᐃᑐᐃᑦ ᓂᐱᐃᑦ ᖁᑯᓂᖅᑐᑦ ᐸᐃᐃᑦ ᐊᐸᐅᖏᖅᑕᑐᒃ ᐃᓯᒪᓕᑕᐅᓂᖏᑦ: ᐊᐃᐃᑦ ᖁᑭᓇᓗᒪᐃᑦ ᐊᖁᐃᑦ ᑐᐄᓂᑦ ᖁᑭᓇᓗᒪᐃᑦ ᓇᑦᑏᑦ
ᐊᐅᓪᓚᕆᑎᔪᐅᓂᒪᓕᑦ **ᑎᒥᖁᑎᑎᒋᔭᐃᑦ**	ᐊᐳᑎᓕᓚᔅᓴᕐᐸᐅᑯᑦ ᓯᓪᐅᑦᓪᓗ ᐊᓯᔪᖅᐸᑦᓪᓕᐊᓯᓕᓚᔅᓴᖏᑦ ᑲᓇᑕᒥ (ᑲᓇᑕᒥ ᐅᒪᓕᔪᓕᓚᔅᓴᓂᑦ ᐱᔪᑎᕆᐃᐅᔅᖏᑦ) ᐃᓪᒥᑦᓪᓗᓂᑦ ᐃᖅᓂᐅᖅᑐᑦ ᐊᖁᔅᑐᐃᓐᓇᖅᑐᑦ ᐊᐅᓪᓚᕆᑎᔪᐅᓂᒪᓕᑦ ᐊᐳᖓᓕᖏᓐᓂᑦ ᐊᐅᓪᓚᕆᑎᔪᐅᓂᒪᓕᑦ ᑲᑎᒪᓴᕋᓯᖏᓐᓂᑦ
ᐃᓄᖅᓄᑦ **ᑎᑭᓯᒪᕐᕕᐅᔪᓐᓇᖅᑐᑦ** **ᐊᓚᒪᓗ ᐊᑐᖅᑕᐅᔪᓐᓇᖅᑐᑦ**	ᐃᓄᐃᑦ ᓄᓇᕗᓚᒥᐅᑦ ᐊᑭᖅᖏᑦᑐᑐᒃ ᐊᓚᒪᓗ ᐊᔪᖅᓇᖅᑎᑕᐅᓕᖅᖏᑦᑎᐊᖅᑐᒃ ᑎᑭᓯᒪᔪᓕᓇᖅᑐᓕᖅᑐᒃ ᑕᒪᐃᓕᖅᑐᒃ ᓄᓇᕗᓕᖅᑐᒃ, ᐃᒪᖅᑐᒃ ᐊᓚᒪᓗ ᑕᓪᐅᐅᖅᑐᒃ ᐃᒪᖅᑐᒃ ᐃᑐᐊᓂᑐᑐᒃ ᑎᖅᑕᐊᖅᖏᓐᓇᖅᑐᒃ (ᒪᓕᒃᑕᐅᓕᓕᑐᒃ ᐃᑐᓕᖅᑐᑦ *5-ᖅᑲᓂᑐᑦ* ᐃᓄᐃᑦ ᐃᑲᔪᓯᐊᖅᓴᖅᑐᓄᓂᑦ ᐊᖅᑭᑕᑎᓂ) ᐊᓚᒪᓗ ᒪᓕᒃᑕᐅᓕᕆᓚᐊᖅᑲᓚᑎᒃ ᐃᑐᓕᖅᑐᑦ *5.7.18-ᒋᑦᑐᑦ* ᓄᓇᕗᓚᒥ ᐊᖅᑭᑕᑎᓂ). ᐃᓄᓕᓚᐅᓴᖅᑭᑐᑦ ᐊᑐᕈᓐᓇᖅᑐᑦ ᑎᖅᒥᐊᖅᑎᖅᓇᖅᓂ ᐃᖅᑭᐊᖅᐳᐃᐅᓕᕐᕕᐅᓂᔪᑦ ᐊᔪᖅᑎᓄᓕᓂᑐ, ᐅᓚᒪᔪᓂᒃ ᐱᔪᓴᐅᐃᑎᖅᑐᐊᕈᑎᒃ, ᒪᓕᒃᑕᐅᓕᓕᑐᒃ ᑎᖅᒥᐊᖅᓂᒃ *ᐅᑎᖅᑕᐅᑐᒃ ᑲᑎᒪᓴᕋᓂᒃ ᑲᑎᒪᔨᓕᕐᕕᐅᑎᒃ ᒪᓕᑕᒪᐊᖅᔪᓕᓂᖏᑦ.*



ᑕᔪᒃᓴᖅ 1: ᐃᓂ ᑎᖏᑕᐅᖅᑲᐃᓕᓴᐅᑦ ᐊᒻᒪᓗ ᐃᓄᐃᑦ ᐅᖅᑲᐅᓯᑐᖅᑐᒍᑦ ᓄᓇᐃᑦ ᑕᐃᔨᐅᓂᖏᑦ (ᐱᔨᐅᓯᒪᔪᑦ ᐃᓄᐃᑦ ᐱᐅᓯᑐᖅᑲᖏᓐᓂᖅ <<ᕐᓂᐊᕐᑦ ᑎᑎᖅᑎᔪᖏᓐᓂᓂᖅ).

1.1 ᐊᕕᑦᑐᓂᓯᓚᑦᓂᑕ ᖃᓄᐃᒋᓚᑦᓂᓚᑦ

ᐃᑲᑐᐊᖏᑎ ᑎᖅᒥᐊᖏᑲᕕᐃᑲ (ᐃᑲᑐᐊᖏᓐ) ᐃᓕᖏᑲᖅᓴᖅ ᑕᐅᔭᓂ ᓕᑭᖅᑲᓂᐸᓂᐊᑕ ᐱᖅᑐᓇᖅᑲᓂᓚᓂ ᓴᑦᒥᐅᓕ ᕿᑭᖅᑲᓂᐸᓂᐊᑕ ᐃᒋᖅᓯᓗᑕ, ᐊᖅᑭᓵᐊᓗ ᕿᑭᖅᑲᓂᓗᐊ ᑕᐅᔭᓂᒪᑦ ᕼᐊᑦᓴᓪᐹᐸ ᑕᓚᐅᖅᑕ ᐅᐊᓪᓇᐅᓕᖅᑐᖅᓚᓂ, ᑭᔭᑦᒥᐅᐸ ᐊᕕᑲᖅᑲᓂᓚᓂ, ᐅᓇᐅᓴᒥ. ᑕᒪᓪᓇ ᑎᖅᒥᐊᖏᑲᕕᐃᑲ ᐅᖅᓱᓂᓚᒥᓄ 110 km ᑭᓯᒥᑕᓄ ᐅᓇᒥᑐᐊᓚᓴᖅᓱᓪᓂᓄ ᕿᑭᖅᑕᖅᓱᓂ, ᐅᓇᒥᖅᓄ ᓴᑦᒥᓄ, ᐃᓄᖏᑲᖅᑐᓄ 891-ᓄᑦ. (ᐅᓴᓴᐃᓯᑦ ᑲᓇᑕᒥ, 2016-ᒥ).

ᐱᓗᐊᖏᑐᒥᑲ ᐅᓇᖅᓱᓂ ᐱᑕᖏᑲᖅᓴᖅᑐᓂ ᑎᖅᒥᐊᖏᑲᕕᐃᑲᐅᑦ ᐅᓇᖅᓱᓂ ᑯᖅᑦ (ᖏᑦᑕᓗᓇᑎᑐᑦ ᑕᐃᔨᐅᔪᒥᑲ ᐳᐊᓯᑦ-Boas ᑯᖅᓱᓄᑦ) ᑯᑦᐸᑲᑐᑦ ᑕᒪᒪᐅᖅ ᐃᑲᑐᐊᖏᑎᓕᕕᐃᑲᐅᑦ. ᐊᒥᓯᑐᓴᓵᓄᑦ ᖃᖏᖃᓴᖏᑲᐅᖅᓴᖅ ᑎᖅᒥᐊᖏᑲᕕᐃᑲᐅᑦ ᑲᓇᖅᓇᖅᐸᓯᖅᓱᓂ, ᐊᖏᒪᓗ ᒪᔪᖅᑐᓄᑦ ᖏᑯᓕᖅᑐᓄᑦ ᓯᓴᖏᖅᑲᓯᓂ ᐱᖅᖅᓱᓂ. ᑎᖅᒥᐊᖏᑲᕕᐃᑲ ᐱᑕᖏᖅᒥᔪᖅ, ᑕᑭᓂᓕᒥᑦ 45 km ᑭᓯᒥᑕᓄ ᓯᓴᒥᑦ ᐊᖏᒪᓗ ᐃᑲᑐᐊᖏᑦ ᒪᕐᕕᐊᑲᐅᓂᒥᑦ ᐊᖏᒪᓗ ᐱᑐᖅᑐᑐᖅᓱᓂ ᑭᓚᓇᖅᓂᓂᐅᓂᓕᖅᑐᑦ ᐃᑲᑐᐊᖏᑦ ᐅᓇᖅᑐᓂᓄ ᐊᑎᑕᐅᓂᖅᑐᑦ ᐃᑲᑐᐊᖏᑦ ᑯᖅᑦ ᐅᑐᓂᓯᐅᑦ ᐃᑲᑐᐊᖏᑦ ᐅᓇᖅᓱᓂ ᑕᒥᓄ ᐅᑐᑐᓂᐅᓂᓚᖅᓄᑦ 60 m ᑭᑕᓚᖅᓄᑦ.

ᓴᓪᓚᓄ, ᑎᖅᒥᐊᖏᑲᕕᖅᓗᓚ, ᓄᓇᖅᓱᒪ ᓴᓇᕐᕕᐅᓯᒪᓴᐊᖅᑭᑐᑐᖅᑐ ᐊᒪᒪᓗ ᐊᐅᑐᖅᑕᐅᓯᒪᓗᐊᖅᑭᑐᓴᓯᓄ. ᐃᓅᔨᔪᑎᐅᓴᖅᑭᑐᓄᑦ ᐊᖅᐳᓇᓯᑎᐊᖏᖅᐸᐅᑐᑦ ᐱᒪᒪᓇᐊᓗᖅᑎᑦ ᐃᓕᓴᐅᓂᖅᑐᓄᑦ, ᐃᓅᔨᔪᑎᖅᑭᑐᓄᑦ ᐊᒪᒪᓗ ᐃᓅᔨᔪᑎᐅᓴᖅᑲᓇᕐᑕᖅᑭᑐᓄᑦᓗᓗ ᓴᓪᓚᓄᐅᓯᒪᓗᐊᖅᑭᑐᓴᓯᓄ (ᐃᓗᓕᖅᑭᑐ 3.2 ᐊᒪᒪᓗ 3.3-ᒥ ᐅᓴᐅᑲᓯᐅᔪᑦ). ᐃᓅᐃᑦ ᓅᐸᐅᑐᑦ ᕿᑭᖅᑕᖅᑲᓂᓄᑦ ᐱᓗᐊᖏᑐᒥᑲ ᓯᑭᑐᐅᔪᑦ, ᕿᑭᓯᓯᔪᓗᓄᓅᖅᑐᑦ ᐅᑭᐅᖅᔪᑦ ᐊᒪᒪᓗ ᐊᐅᖅᔪᑦ ᖃᐅᔨᓴᖅᑭᑐᓄᑦ ᐱᑕᖏᑲᖅᓴᖅᑐᑦ ᐱᖅᑐᓇᖅᑲᓂᓚᓂᓄᑦ ᐱᓕᕆᔨᒪᓂ

ᐊᕕᒃᑐᖅᓯᒪᔨᓂᒃ ᑕᑯᑎᑎᓴᔨᑦ 2: ᐊᕕᒃᑐᖅᓯᒪᓂᖅᓗᑕ ᖃᓄᐃᓕᓕᖅᓂᖅᑎᑦ ᐊᓚᒪᓗ ᐅᒪᔨᖃᖅᖢᓂᖅᑎᑦ ᐊᐅᓗᐊᓂ ᓄᓇᕗᑦ ᐅᓇᓗᐃᒃᑯᑎᖅᑕᐅᓯᒪᓂᖅᑎᑦ ᑎᖅᑕᐊᖃᖅᖢᓕᖅᓄᖅ

ᐊᕕᒃᑐᖅᓯᒪᓂᖅᓗᑕ ᖃᓄᐃᓕᓕᖅᓂᖅᑎᑦ	ᑲᓇᑕᐅᑦ ᓄᓐᓗᓗᐊᓂᒃ ᑕᐃᔨᓯᒪᓂᑐᖃᖅᓚᓗᒃ ᓴᕐᑭᖅᐸᑦᓕᐊᑕᓕᐅᖅᓂᖅᑎᑦ
ᓄᓇᖅᓗᑕ ᑲᓇᑕᒥ ᐊᕕᒃᑐᖅᓯᒪᓂᖅᓗ	ᕿᐊᑦᓴᐟᐸᑦ ᐊᕕᒃᑐᖅᓯᒪᓂᖅᓗᓄᑦ ᐃᑕᑭᐅᐅᔨᖅ
ᐊᐢᑎᑎᐅᑕᖅᑎᑦ ᖃᓄᐃᐃᑐᖃᖅᓂᖅᑎᑦ	3 ᓂᑭᖅᐸᓯᐊᓂ ᐅᑭᐅᖅᑕᖅᑐᑦ
ᓄᓇᖅᑎᑦ ᐅᒪᔨᖃᖅᓂᖅᓱᓄᓪᓗ ᐅᓴᐹᖅᓂᖏᓐᓂᖅᑎᑦ	3.2 ᑭᐢᓪᓕᖅᒥ ᐊᑎᑎᒃᑐᑎᐅᓂᖅᑎᑦ ᓄᓇᖅᑎᑦ
ᓄᓇᖅᑎᑦ ᐅᒪᔨᖃᖅᓂᖅᓱᓄᓪᓗ ᐅᓴᐹᖅᓂᖏᓐᓂᖅᑎᑦ	3.2 ᑭᐢᓪᓕᖅᒥ ᐊᑎᑎᒃᑐᑎᐅᓂᖅᑎᑦ ᓄᓇᖅᑎᑦ
ᓄᓇᖅᑎᑦ ᐅᒪᔨᖃᖅᓂᖅᓱᓄᓪᓗ ᐳᑐᒃᑕᐅᓗᑐᓕᐊᖃᖅᑕᐃᓕᒪᔨᑦ (ᓄᓇᐃᑦ ᓇᒥᐅᑕᖅᑭᑦᓗ)	3.2.46.185 Boas ᖃᓪᓗᓈᑎᑐᑦ ᑕᐃᔨᐅᔨᑦ ᐳᐊᓯᑦ ᑯᖏᒐ
ᑕᓚᐅᖏᒥᐅᑕᐃᑦ ᐅᒪᔨᖃᖅᓴᓚᓕᕆᐅᓂᖅᑎᑦ	ᐊᔨᓴᐹᖅᑭᑐᑎᓂᒃ ᕿᐊᑦᓴᐟᐱᖓᓂᖏᓐᓄᑦᓯᑦ

1.2 ᖃᓄᖅ ᐱᓕᓪᓕᐊᓚᐅᖏᓪᓚᓂᑦ

ᑕᒪᓐᓇ ᑲᖓᖃᖅᓴᒃ ᐃᑲᑕᑐᐊᑦ ᑯᖏᒍᓂ ᖃᑲᐅᓕᖅᑕᐅᓯᒪᓂᖅᐸᖓᔪᖅ ᓄᓇᖅᑭᓂ ᑲᓇᑕᐅᑦ ᐅᑭᐅᖅᑕᖅᑕᖅᑐᑦ ᐊᑯᓂᓂᑭᓂᒃ ᐊᕐᕌᒍᐃᑦ 1930ᑦ ᐊᒻᒪᓗ ᐃᓱᖅᐸᓱᑭᓂᓂ ᐊᕐᕌᒍᐃᐃ 1950ᑦ. ᖃᑲᓂᑯᑭᓂᓂᒃ 20,000 ᑲᖓᐃᑦ ᓂᐅᖏᒥᐅᑕᖅᑕᐅᓯᒪᔨᖅᑕᐃᓕᒪᔨᑦ ᐊᕐᕌᒍᐃᑦ ᐊᑯᓂᓂᑭᓂᓂ 1952 ᐊᒻᒪᓗ 1958. ᑎᖅᑕᐊᖃᖅᓴᒃ ᑲᒪᑭᐅᔨᒪᓚᐊᖏᓕᐅᖅᑕᖅᑐᖅ ᐅᒪᔨᖃᓕᓚᓄᓪᓗ ᐱᑕᖃᖅᓂᕐᓗᑕᐅᓂᖅᓂᑭᓂᓂᓪᓗ “ᖃᑲᖏᐃᑦ” “ᑲᖓᐃᐅᑦᓗ” ᓄᓕᐊᖃᑕᐅᑎᓱᖓᓂᖅᑕᐅᓂᖅᓂᑭᓂᓂᓪᓗ ᑕᐅᖏᓂ (Cooch & Barry, 1957). ᑕᐃᔨᓱᒪᓂ, ᑕᐃᔨᓱᒪᓂᓕ, ᒪᕐᑕᖓᓂᒃ ᑕᖃᒪᖃᖅᑎᑎᑎᑦᑎᐅᓱᑎᓂᖅᓂᑭᓂᓂᓪᓗ ᐊᔨᓴᑦᑐᐃᑕᐅᓇᓱᑎᐅᖏᓕᐅᖅᑐᑦ ᑎᖅᑕᐊᒥ. 1957-ᒥ, ᐅᒪᔨᓕᓚᓄᔪᑦ ᐱᑎᐊᖃᖅᑎᑎᐅᔨᒪᓕᓚᐅᖅᑐᑦ ᑎᖅᑕᐊᖃᖅᓴᖅᑭᒃ ᔪᑐᓂᓂᐊᖃᖅᑕᐅᔪᑦᓗᑦᑎᑦ ᑎᖅᑕᐊᑦ ᐃᑲᑕᑐᐊᑦ ᑯᖏᒍᓂ, ᓄᓕᐊᖅᓴᐅᓂᖅᓂᑦᓗ

ᓂᕆᓪᓗᓐᓄᑦ (Atlantic brant-ᓂᒃ) ᑕᐃᓵᓕᒐ ᒪᓕᑦᑲᒪᐅᓂᖅᓯᖏᓐᓄᑦ ᐊᒥᓱᓐᓂᒍᐊᓕᕆᓂᖅᑕᐅᓂᖏᓐᓄᑦ, ᖃᐅᔨᒪᔭᐅᓂᖏᓐᓄᑦᓗ ᑐᓴᕆᓚᔨᐅᔪᑎᑐᓂᖅ ᓄᓕᐊᖅᐱᐅᒪᖅᓂᖅ ᑲᖏᒃᐃᑦ (Ross-ᒋᓴ), ᑲᓇᑕᐅᕐ ᐅᑭᐅᖅᑕᖅᑐᖅᓯᓱᐊᓂᖏᓐᓄᒐᓂ.

ᑕᒪᓂ ᒪᐃ 21, 1959-ᖏᑎᓪᓗᒍ ᐃᑲᓂᑐᐊᕐ ᑯᖏᓂᒃ ᐱᓂᖏᖅᑎᓂᑎᓕᐅᖅᓴᓂ ᕿᐊᐅᓕ ᑎᓯᓱᐊᕐ ᑎᖏᒥᐊᖃᖅᑲᖏᓐᓄᒃ, ᒪᓕᑎᓕᐅᖅᑎᖏᓂᓄᑦ (ᑕᖓᒪᑎᔾᔨᐅᓂᖏᓂᒃ P.C-ᑯᓕ. ᐊᖅᖁᒍᒥ 1959-629). ᑎᖏᒥᐊᖃᖅᑲᖏᕐᓄᑎᑕᐅᓯᖅᓄᑦ ᑕᐃᔾᔨᐅᓕᐅᖅᑐᖅ ᐅᐱᒋᔾᔨᐅᓂᖏᓐᓄᓕᓂ ᕿᐊᐅᓕ ᑎᓯᓱᐊᕐ ᐅᓇᐃᓂᓄᑦ (ᐃᓄᓕᐅᖅᑐᖅ ᑕᒪᓐᓇ. 1900-1954). ᐅᓇᐃᓂᓄᑦ ᐃᓄᓕᐅᖅᑐᖅ ᐅᓄᑯᓕᓂᕐᒥᓗ, ᐱᓕᓚᐅᒋᔾᔨᐅᓕᐅᖅᑐᖅ ᕿᐊᑦᓴᓂᕐᐳᓗ ᑲᒪᓕᑐᒃᑐᐅᓂᖅᓗ ᐊᒥᐊᓕᓚᐅᐅᓂᖅ ᐅᓇᑕᖅᑐᓴᖏᓐᓄᑦ, ᐊᒥᓱᓕᕆᓂᖅᓗ ᑲᑎᒪᓂᖅᓄᑦ ᐅᓇᑕᖅᑐᓴᖏᓐᓄᑦ ᐊᒥᓱᓂ ᑲᖏᖅᑐᑦ 1933-ᖏᑎᓪᓗᒍ, ᐱᔨᑎᑎᐃᔨᐅᒋᔾᔨᐅᓂᖏᓐᓄᓕᓂ ᐊᐅᓪᓚᒍᔨᐅᓂᖏᓐᓄᑦ ᐊᒥᓱᓕᕆᓂᖅᓗ ᐊᒥᓱᓂ ᖃᐅᔨᒪᔭᐅᓂᖏᓐᓄᑦ (Cooch, 1957).

ᑎᑎᕿᖅᑕᐅᓕᓕᐅᖅᑐᓂ ᐊᐅᓕᓴᑎᓂᖅᓗᑦ ᕿᖃᓇᐅᑦᑐᓂ 1986-ᖏᑎᓪᓗᒍ, ᓴᐅᓂᐊᔪᓂ ᓄᓇᕐᒥᓇᐅᖅᑐᒃ ᓄᓇᑦᓇᐅᖅᑎᑦᑎᓂᖅᑐᓂ ᓴᓕ ᐊᒪᓚᓗ ᐊᑎᓕᐅᖅᑕᐅᔨᕆᖃᓇᐅᖅᑎᓂᖅᓄᑦ ᓴᓕ *ᐃᓄᓐᓄᑦ ᐃᑲᔪᓴᐊᓵᕆᓂᖅᓄᓂ ᐊᖏᕐᕉᑦ.* (Stephenson & McCormick, 1986).

1.3 ᓄᓇᕆᐊᑎᕈᐅᓂᖅᑎᓂᑦ ᐊᒪᓚᓗ ᐊᑐᖅᑕᐅᔨᓴᓂᖅᑎᓂᑦ

ᑕᒪᓐᓇ ᓄᓇᖅᑎᓂᒃ ᐊᖅᑎᓂᖅᓴᓕᐅᕐᓂᑦ ᑲᓇᑕᐅᕐ ᑲᕿᒪᓂᖅᓄᑦ ᓄᓇᕆᐊᑎᕈᐅᔨᓂᑦ ᑕᐃᓂᓇ ᐃᓄᖅᓄᑦ ᐱᑎᕈᐅᔨᑐᑎᓪᓗᒍ ᑎᓪᓗᒍᓴᐅᓂᑦ (CH-06, ᓄᓇᐅᕐ ᖃᖅᖢᓕᒫᓂᒃ ᐱᔨᓇᐅᕈᑎᖃᖅᖢᓕᒃ; ᑕᑯᐅᒃᓴᖅ 2). ᐃᓗᓐᓄᖅᓄᑦ ᐱᕆᑯᑎᕈᐅᔨᔪᖅ ᓄᓇᖅᓱ ᐊᐅᓪᓚᑕᐅᕐᓂᖅ ᑭᓪᓕᓪᓂᕆᑦ ᐃᓄᐃᓂᓕᒫᑎᓂᒃ ᐅᖃᓕᓂᐅᔪᓂᒃ ᑭᒪᔭᐃᓂᓕᒫᑎᓂᒃ ᑕᒪᐃᓐᓂᒃ ᐃᓄᐃᓂᖅ. ᑲᓇᑕᐅᕐ ᒥᓂᔅᓴ ᑲᓇᑕᐅᕐ ᑲᕿᒪᓂᖅᓄᑦ-ᓄᓇᖃᖅᑐᑐᖃᖅᑲᐃᓪᓗᓕ ᐱᓪᓚᖃᖅᑎᑦᑕᑎᐊᕈᓐᓇᖅᑐᖅᓄᑦ ᐅᑭᐅᖅᑕᖅᑐᒥ ᐱᓪᓕᓪᓚᓕᓚᓂᓪᓕᒍ ᑲᓇᑕᒥ, ᐊᑐᖅᓴᓂ *ᑲᓇᑕᐅᕐ ᐅᑭᐅᖅᑕᖅᑐᖅᓯᓱᐊᓂᖅᓱᓂᓗ ᓄᓇᖅᑎᓂᒃ ᒪᓕᑦᑲᐃᓪᓗ*, ᑎᐃᒥᐊᖅᑎᐅᐅᓂᖅ ᑕᒪᐃᓐᓂᒃ ᓄᓇᕆᐊᑎᕈᐅᔨᑐᑎᓪᓗᒍ ᑲᑎᒪᓂᖅᓄᑦ ᐃᓄᐃᓂᖅ ᑲᕿᒪᓂᖅᓄᑦ ᐱᔨᑎᑎᐃᔨᐅᓂᖅᓱᓂᓗ ᓄᓇᓂ ᑎᓪᓗᒍᐊᖅᑐᖅᓄᑦ ᐱᔨᑎᑎᐃᔨᐅᓂᖅᓱᓂᓗ ᓄᓇᓂ ᐃᓄ


ᓴᐳᵒᓂᐊᖅᑕᐅᓯᒪᖏᖅᓴᐅᓂᖏᑦ ᐃᵒᓇᐅᔨᖅ ᐃᓕᖅᖃᖅᑐᑦ ᖃᖅᖃᓴᐅᖅᑐᑉᐸ ᑕᓇᐅᖏᑕ ᐃᒪᖅᔪᑎᑦᑐᵒᓂᒃ ᑎᖏᑦᖃᐊᖅᖁᖏᑦᒃ, ᐱᓯᐊᓂ ᐃᖅᑲᓗᓪᓚᕆᔨᐊᕆᔪᑦ ᐊᒻᒪᓗ ᐃᓕᑭᖅᑕᖏᑦ ᑐᑭᓯᐅᕐᓇᑦᑐᑦ ᐱᖓᓱᓕᖅᖁᒪᖏᖅᖁᖏᑦ ᑕᓇᐅᑉ ᓴᐳᵒᓂᐊᖅᑕᐅᓯᒪᖏᖅᓴᖅᑐᓂᒃ, ᑐᑭᓯᑎᐊᕈᑕᐅᑎᐊᓕᐅᖅᓵᑐᒃ ᐊᒻᒪᓗ ᐃᓄᖅᓂᒃ ᑭᓇᐅᔪᑦᑐᐃᓐᓇᖅᑐᒃ ᑎᑎᕋᔪᑎᐊᖅᑐᓂᒃᓴᖅᑐᒃ ᖃᐅᔨᓴᖅᑎᑦᓴᖅᑕᐅᖅᓵᑐᒃ.

1.4 ᐃᓗᓕᔪᑎᑎᑦᓴᐅᔪᑦ ᐱᕐᔪᐊᓱᓪᓗ

ᐊᑕᐅᓯᕐᒥᒃ ᐃᓗᓪᓕᕆᓪᓕᖅᑕᓪᓕᒃ ᑎᖏᑦᐊᖅᑲᖅᖁᖏᑦ ᓇᖏᒥᓂᓴᒥᒃ ᐊᐃᐃᒐᑦ ᐊᖏᔪᓇᓴᐅᑦ ᒥᑭᑎᕐᓂᐊᖅᑲᖅᓱᓪᓗ ᑲᑭᑎᑦᓂᐊᕐᓂᖅ (ᑕᔪᓴᒃᓴᖅ 1; ᐊᖁᒃᑐᖅᓯᒪᔪᓂᒃ ᑕᔪᑎᑦᑎᓴᔪᑦ 3). ᑕᐃᵒᓇ ᐃᓗᓪᓕᔪᐊᖅ ᖃᔪᓕᒃᓴᐅᓕᖅᑐᑐᓂ 10 ᐊᑦᕿᔭᓂ ᑕᐅᖅᓀᓕᖅᑐᖅ ᐊᑐᖅᑕᐅᖅᑎᑦᑕᖅᓯᓂᓗ ᔪᔪᓯᒪᓂᖅᓴᓂ ᐊᖏᔪᓇᔭᒪᒃᓴᓄᑦ ᑐᑦᖃᖓᓴᓂᒃ, ᐱᓗᐊᖅᑐᒥᒃ ᓇᵒᓂᐊᖅᑎᑦᓴᖅᑐᒃ.

ᐊᖁᒃᑐᖅᓯᒪᔪᓂᒃ ᑕᔪᑎᑦᑎᓴᔪᑦ 3: ᐃᓗᓪᓕᔪᑎᑎᑦᓴᐅᔪᑦ ᐱᕐᔪᐊᓱᓪᓗ ᑎᖏᑦᐊᖅᑲᖅᖁᖏᑦ

ᓯᓇᐅᓂᖏᑦ (ᐱᕐᔪᐊᓱᑦ)	ᖃᐅᔨᐃᓂᖏᑦ	ᐊᖏᓂᖏᑦ ᐅᒃᑐᑐᐅᑎᒃᔪᑦ Fᐃᑦᔪᑦ	ᖃᐅᖓ ᓴᓇᔨᐅᓂᔪᑦ	ᐅᒪᔨᐅᔪᑦ ᐱᕐᔪᐊᑎᓂᒃ	ᓇᓂ
ᐃᓗᓪᓕᕆᓪᓕᖅ	ᖃᐅᔨᓴᒪᔨᐅᓐᓂ	10 x 10	2008	ᐊᐃᐃᒐᑦ ᐊᖏᔪᓇᓴᐅᑦ	63° 39’ 17”, -86 ° 09’ 42”




ᑕᘻᒃᓴᖅ 2: ᐃᓄᖕᓄᑦ ᓄᓇᕗᑎᑎᕆᐅᔪᑦ ᓄᓇᖏᑦ ᑎᖅᒥᐊᖅᑲᓇᖕᒥ.

2 ᐊᕐᑎᕐᓗᒃ ᐊᕐᑎᑎᐅᑕᖅᑲᓂᕐᓗᒃ

2.1 ᓄᓇᒥᐅᑕᐃᓪᓗ ᐃᒪᖅᒥᐅᑕᐃᓪᓗ ᐅᒪᔨᖅᑲᐃᓕᖅ

ᑕᓚᐅᐸ ᐸᖏᓚ ᑯᐃᕐᐃᐅᕐᓂᖏᓚ ᖃᑎᖅᐸᓯᐊᑯᓗᒪᓕᑦ ᓴᓪᓕᐅᐸ ᖃᑭᖅᑕᒡᓗᑕ, ᐃᑲᖅᑐᐊᐸ ᑯᖏᓚ ᐊᖏᑕᓂᖅᐸᖅᑎᐊᖓᔪᖅ ᑯᖓᓂ ᑕᒪᓐᓂ ᑯᐃᓴᐅᕐᖏᑐᑐᓂ. ᑕᒪᖏᓐᓇ ᑯᕆᑐᐊ ᓂᕈᑐᐊᓚᑦᑐᔪᖅ, ᐃᑲᖏᑕᐊᓚᑦᓗᓂᓗ, ᑕᓚᐅᕆᑐᐊᑕ ᓇᑎᖅᐸᓯᖓᓂᑲ ᖃᑯᑎᖏᓂᖅᓴᐅᑐᐃᓇᕆᑕ ᐊᒥᓯᖏᑕᑐᓂᑲᑦ ᐅᑲᑐᐅᑎᓂᑲ ᑭᓯᐅᑕᑐᖅ. ᐃᑲᖅᑐᐊᐸ ᑯᖓᓚ ᑲᔨᓯᔪᖅ ᓄᓇᑲᑯᓂ ᐱᓕᕆᑕᓂᖓᓚᒍᓂ, ᕈᑕᖅᓇᐃᓇᐅᕐᓇᒪᒍᓂ ᐊᒻᒪᓗ ᓄᓇᖏᓂ ᐃᓗᑐᓂᑕᐅᑕᓂᖓᓚᒍᓂ, ᑕᑕᒪᓂᑲ 5 ᕿᒥᒥᓂᑲ ᓂᕈᑐᓂᖅᑲᖅᑎᑎᔨᒃ ᐊᒻᒪᓗ 13 ᕿᒥᒥᓂᑲ ᑕᕿᓂᑕᖅᑲᒃ ᑯᖓᓂᖅᓴᐅᑐᐃᓐᓇᓂ ᐃᑲᖅᑐᐊᖅᓐᓇᓂ. ᐃᑲᖅᑐᓂᖅᓴᓂᑲ ᒪᕐᕉᐃᓇᓗᐊᖅᓴᖓ ᑕᒪᖏᓐᓇ ᓂᕈᑐᐊᓗᓚᖏᓚ ᑕᕿᓂᑕᖅᑲᒃ 13-ᕿᒥᒥᑕᓂᑲ ᐅᖏᓕᒥᓂᓗᖏᓂ ᓱᓴᕐᖅᐸᓯᐊᔪᖅᑐᖅ (Latour, 2008). ᐊᖅᑎᐅᑐᑎᓂ ᓄᓇᕈᓂ ᐊᑕᓂ 30 m ᒥᑕᓂᐳ >ᖅᑐᓂᓕᖅᒥᐳ, ᐊᒥᓯᑦᓗᑎᓚᓗ ᑕᓯᓚᕆᐃᑦ ᐊᓯᖏᓂ.

Fontaine et al-Fᐊᐟᖏᑎᖏ ᐊᒻᒪᓗ ᐱᑕᓇᖅᑎᑎᕈᓂ (2011) ᓄᓇᖏᓂ ᓯᓇᑕᖅᑯᓂᖓᕈᓂᐳ ᖃᖏᓚᓂᑐᑐᓂᐳ ᓇᓗᓇᐃᐳᔪᓂᑐᑐᓂᖅᑐᑕᐅᓂᑕᖅᑐᑦ ᑕᒪᐃᓂᖏᓚ ᑭᑦᑕᐸ ᖃᑭᑦᑕᖅᑐᐳᓂᖓᓂᖏᓚᓂ ᓯᓗ ᓄᓇᖏᓂ ᐊᑐᑎᓕᒪᓇᖏᓕᐳ ᑐᖏᓚᐅᓂᖅᓴᖓ ᐅᓇᓂᐊᐳᓇᖅᑐᖅᓴᓂᖅ ᐅᒪᔨᖅᑲᓐᓂᐊᕈᓂᖓᓂᕈᓂᐳ ᐊᓯᔪᖅᑲᖅᐸᓕᕆᔨᐳᖅᑐᐳ ᖃᐅᔨᖅᑲᓐᓂᐊᕈᓂᖓᓂᕈᓂᐳ. 20-ᖏᓂᓕᒪᒃᑕ ᐊᔪᕐᓂᖏᑐᑎᑎᓂ ᐅᔨᒪᓇᓴᐅᓂᕈᑐᑦ ᐃᓕᑕᒪᓇᓴᐅᓂᔨᔨᓂ ᐃᑐᐊᓂ ᐃᑲᖅᑐᐊᐸ ᑎᓐᒥᐊᖅᑲᖅᓇᖓᓂ. ᕈᓯᐊᓂᓕᑕᐅᖅᑲ, ᐱᔨᔨᑎᑎᓕᓕᑦᑐᑎᑐᑦ ᑭᓴᐳᓇᖅᑐᖅ ᐊᖓᔪᖅᑐᑎᑦ ᐃᑐᐊᓂᖅᑐᑕᐅᑎᐊᖅᓯᒪᔨᓂᓂ ᓄᓇᑕᓐᓄᑦᑐᑎᑦ ᐅᓇᓂᐊᓴᐅᓂᕈᓂᖓᓂᖅᑲᖅ. ᐊᖏᓕᒃᑎᑎᐅᓂᖅᑲᓇᑦ ᓄᓇᓕᑦ ᐅᓇᖅᑕᐅᓂᕈᓂᖓᓂᑕ ᐱᔨᔨᑎᑎᓕᓕᑦᑐᑎᑦ (35% >ᓴᖏᑎᓂᖅ) ᐊᒻᒪᓗ ᖃᐅᔨᓴᓂᐅᓂᕈᑐᑎᓂᖅ (28% >ᓴᖏᑎᓂᖅ), ᑕᐃᔪᓂᖓᓚᓗ 20% >ᓴᖏᑎᖏᓕᓂᑐᑎᓂᖅ <ᓂᓂᓂᖅᑲᐅᔨᓂᖅ, ᐊᒻᒪᓗ ᐱᔨᖅᓄᐅᓂᕈᓂᑐᑎᓂᖅ ᓄᓇᖏᓂᖅᓂᖅ (ᓯᑦᓗ ᐅᖅᑐᑎᑎᓗᔪ ᓯᐅᕆᐃᑦ, ᒪᕆᐃᑦ, ᑐᐊᐸᐃᑦ, ᐅᓴᖏᑦᓭᓂᐊᓗᐃᑦᓗᓂᖏᓕᓂ) ᐊᒻᒪᓗ 18% >ᓴᖏᑎᖏᓂᖅ ᐃᒪᐃᑦ (ᑕᔪᓴᖅᑲᖅ 3).



ᑕᘻᖅᓴᖅ 3: ᓇᓱᒪᓗᐅᔪᓪᓗᐊᑕᖅᑕᕐᓂᖅᑭᓂᒃ ᐊᐃᖅᑐᖅᑕᐅᓯᒪᓂᖅᑭᓂ ᑎᖏᒥᐊᖅᕿᕐᐃᐅᔪᓂ (ᒪᓕᖅᑕᐅᓯᒪᔪᓪᓗᑎᒃ ᐱᓕᓚᓪᒋᐅᖅᑐᑦ Fᐃᐢᑎᐢ ᐊᒻᒪᓗ Ỉᓗᓚ-Fontaine and Mallory, 2011).

2.2 ᑎᖅᑕᐊᓕ

ᐃᑲᓂᐃᔪᖅᑎ ᑎᖅᑕᐊᖅᑲᓇᑦ ᐃᓕᓴᖅ ᑕᒪᓐᓇ ᓴᓪᓂᐊᖅᑕᐅᓯᒪᖅᓯᓚᖅᓴᓂ ᑲᑎᒪᔨᖅᓯᓚᖅᓴᓂ 3-ᑕ ᐅᑭᐅᖅᑕᖅᑐᑦ ᐊᕐᓇᖅᑐᓯᒪᓕᖅᓴᓂ ᓇᒻᒥᖅᑐᖅᓯᑕᖅᑐᑦ ᐊᒻᒪᓗ ᑭᖅᑕᓴᓪᓂᖅ (ᐊᐅᑎᓕᓐᓄᓴᖅᐊᐅᑉᑐᑦ ᑲᑎᒪᔨᖅᑕᐃᓐ, 2013). ᑕᒪᖅᑕᖅ ᓄᓇᒥᐅᑕᓂᖅ ᓯᓚᖃᖅᑕᓯᐅᒥᐅᑕᓂᖅᓗᓐ ᓇᓯᖅᑕᐅᐊᖅᓂᖅᑎᑦ ᐊᒥᓯᐊᓗᖅᓂᖅ ᑎᖅᑕᐊᖅᑲᓇᐅᐊᖅᑐᖅᑐᑦ. ᐅᖅᓚᑕᓂᑦᑐᑦ 50 ᑎᖅᑕᐊᑦ ᐊᔪᐃᖅᑎᑦᑐᑲᓱᑦ ᑕᐅᐊᓂᖅᑕᑦᑕᖅᑐᑦ ᑎᖅᑕᐊᖅᑲᓇᖅᓚᓂᓂᖅ, ᓇᑦᑕᑦᓗᐊᖅᑎᑦᓗ ᓄᓕᐊᖅᑕᓯᓯᑎᖅ ᑕᐅᐊᓂᖅ ᖃᓄᑎᓴᖅᑎᓂᖅᓗᓂᓂᓂᑦ ᐊᐊᓂᖅᑎᓂᖅ ᓄᓇᒥᖅ. ᖃᐅᔨᓴᖅᑕᐅᓯᒪᕐᓇᐊᖅᑎᑦᑕᑦ ᑕᐃᒪᖅᓯᓚᖅᓴᓂ 1950-ᖅᑎᓂᖅᓴᓂ, ᐊᒥᓯᔪᖅᑎᑕᕐᓂᖅᓯᓚᑎᓗ ᑎᖅᑕᐊᖅᑲᓇᖅᑎᑦᑕᐊᖅᑐᑦ ᐃᓄᐃᑦᓗᑦ/ᓂᐅᖅᑭᐊᑦ, ᑕᒪᖅᑯᐊ ᓄᓕᐊᖅᑎᐊᖅᑎᑦᑕᖅᑐᖅᓂᖅ, ᑕᐃᒪᐃᓂᖅᓱᓐᓗ ᐊᒥᓯᓂᖅᓴᐅᖅᑲᑎᑕᖅᑐᖅᑕᖅᓴᐅᓕᖅᑕᖅᑐᑦ ᑕᐅᐊᓂᖅ ᑎᖅᑕᐊᖅᓴᓴᑦ ᐱᑕᖅᑐᖅᑕᖅᑐᑦ. ᑎᖅᑕᐊᖅᑲᓇᖅᓯᓚᖅᓴᓂᖅ ᐊᔪᐃᖅᑎᑦᑐᑲᓱᑎᐅᖅᑐᑦ ᑕᐃᖅᑯᑎᑐᓐᓇ ᐊᓯᖅᑐᓂᖅ ᖃᐅᔨᓴᖅᑕᐃᓕᖅᑐᑦᑕᖅᓯᒪᕐᓇᐊᓂ ᓇᓯᓚᓴᐅᐊᖅᓯᖅᑐᓂᖅ ᓄᓇᖅ ᓴᓕᑦᑐᐅᐊᖅ ᖃᐅᖅᑎᑦᑐᓐᓇᐅᑦ ᑎᑎᖅᑕᖅᑲᓇᐅᖅᑐᖅᓂᖅᓗᓂᓂᑦ.

2.2.1 ᑲᖓᐃᐃᑦ

ᑎᖅᑕᐊᖅᑲᓇᖅ ᐱᐅᓇᐊᓗᑐᐃᖅᓴ ᓇᓯᓚᓴᐅᐅᓇᖅᓂᖅᓕᑦ ᑲᖓᓴᓄᑦ. ᐊᒥᓯᐊᓗᐃᐃᑦ ᑲᖓᐃᐃᑦ (ᒥᑭᓂᖅᓴᐅᓂᕆᖅᑕᐅᖅᑐᑦ) ᓄᓕᐊᖅᑐᑦ ᐃᑲᓂᐊᖅᑐᑦ ᑯᓐᓴᑦ, ᐊᒥᓯᓴᓴᐅᓕᓗᑎᖅ ᐅᑦᓗᖅᑲᓇᐅᖅᑐᖅᑕᖅᑐᑦ ᑲᖓᓴᓄᑦ ᓴᓕᑦᑐᓄᑦ. 2014-ᒥ ᑲᖓᓴᓴᐃᑦ (ᓯᓴᓗ Lesser snow geese ᒥᑭᓂᖅᓴᐅᓂᕆᖅᑕᐅᓛᑦ) ᐊᒻᒪᓗ (Ross-ᑲᓴ) ᑲᖓᐃᐃᑦ) ᐊᒥᓯᓂᖅᑲᓇᓯᑎᐅᓕᐅᖅᑕᖅᑐᑦ ᖃᐅᔨᓴᖅᑎᑎᓂᖅ 689,700-ᖅᑎᓂᑎᑕᓂᖅ ᐅᑦᓗᖅᑲᖅᓯᖅᑐᑦ, ᑲᖓᐃᐃᑦ (ᐅ. ᑕᐢᓗᐊᖅᓴ-J. Leafloor, ᐅᒪᓛᓕᓐᓄᓂ ᑐᓴᐅᒪᖅᑕᑦᑕᐅᑎᓕᓐᓄᑦ). ᑕᐃᒪᓂᐊ ᑲᖓᐃᐃᑦ (Ross-ᑲᓴ) ᐊᒥᓯᔨᓐᐊᑲᓂᖅᑎᓂᕆᖅᑐᑦ ᐱᒻᒪᓐᐅᓛᕐᓂᖅᑐᑦ ᕿᑎᖅᑕᐊᖅᑲᓇᖅᑐᖅ ᓴᓕᑦᑐᓄᑦ ᑕᐃᒪᓕᖅᑐᑦ ᐊᖏᕐᓇᖅᑐᖅᑐᑦ 1970-ᓂᖅ (ᑯᐱᓴ, ᐱᓕᓐᓄᖅᑎᓗ-Kerbes, et al., 2014), ᐱᓴᓇᑕᐅᓗᑎᖅᓗᓐ ᐊᒥᓯᔪᖅᑐᖅᑲᓇᓯᑎᓕᖅᑎᓯᖅᑐᓂᖅ ᐱᑕᖅᑲᓇᐅᓯᒪᓕᖅᓴᖅᑐᓂᖅ ᑲᖓᐃᐃᑦ ᖃᐅᔨᓴᖅᑕᐃᓕᖅᑐᑦᑕᖅᑐᖅᑕᖅᓴᐅᓕᖅᑕᖅᑐᑦ ᑎᑎᖅᑕᖅᑲᓇᐅᖅᑐᖅᓂᖅᓗᓂᓂᑦ ᐅᑦᓗᖅᑲᖅᓯᖅᑐᑦᑕᖅᑐᖅᑕᖅᓴᐅᓕᖅᑕᖅᑐᑦ ᐊᖏᕐᓇᖅᑐᖅᑐᑦ ᐱᑕᖅᑲᓇᐅᓯᒪᓕᖅᓴᖅᑐᓂᖅ ᑲᖓᐃᐃᑦ ᖃᐅᔨᓴᖅᑕᐃᓕᖅᑐᑦᑕᖅᑐᖅᑕᖅᓴᐅᓕᖅᑕᖅᑐᑦ. (Ross-ᑲᓴ) ᑲᖓᐃᐃᑦ ᐊᒥᓯᔪᑎᑎᓕᖅᑕᖅᑐᑦ ᐅᖅᓚᑕᓂᑦ <1% ᐳᓴᐅᑎᓂᖅ ᖃᐅᔨᓴᓪᓚᖅᑕᓂᓂᑦ ᓂᐅᖅᑎᐅᑕᓂᖅᑕᐅᓯᒪᓕᕐᓇᐃᓂᐅᓂᖅᑎᓂᑦ, ᐊᖏᕐᓇᑕᒪᓴᖅ ᓴᓕᑦᑐᓄᑦ, ᓯᖃᓂᐊᓕᑦ ᐊᖏᕐᓇᔪᑦ 1980-ᓂᖅ (ᑯᐱᓴ ᐱᓕᓐᓄᖅᑎᓗ-Kerbes, et al., 2014), ᑭᓯᐊᓂᓕᑎᖅᓴᖅ ᐱᑕᖅᑲᓇᖅᑎᑦ ᖃᐅᔨᓴᖅᑕᐃᓕᖅᑐᑦ 12.5% ᐳᓴᐅᑎᖅᓂᖅᓴᓂᖅ ᑕᒪᐃᓂᖅᑐᖅ ᑲᖓᐃᐃᑦ ᓂᐅᖅᑎᑎᐅᑎᖅᓴᖅᑎᑎᓕᖅᓴᖅᑐᖅ ᐊᖏᕐᓇᔪᓴᐅᖅᑐᖅ ᑕᐃᒪᓐᓇ 2009-2018-ᒥ (ᐅ. ᑕᐢᓗᐊᖅᓴ-J. Leafloor, ᐅᒪᓛᓕᓐᓄᓂ ᑐᓴᐅᒪᖅᑕᑦᑕᐅᑎᓕᓐᓄᑦ).


ᑲᖓᐃᐃᑦ (ᒥᑭᔨᑎᐅᓂᖅᓴᐅᓂᕆᖅᑕᐅᖅᑐᑦ) ᑕᐅᐊᓂᖅᑕᑦᑕᖅᑐᑦ ᐃᑲᓂᐊᖅᑐᑦ ᑯᓐᓴᑦ ᐊᒥᓯᓂᖅᑲᖅᓯᐳᑦ ᐅᖅᓚᑕᓂᑦ 5% ᐳᓴᐅᑎᓂᖅ ᑲᑎᒪᔨᑦ ᐊᒥᐊᓕᑲᐅᑦᓗᓐ ᓄᓇᑕᐅᖅᑐᑦ ᐊᒻᒪᓗ ᐊᒥᓯᓂᖅᑲᖅᓯᐳᖅᑐᖅ. ᑕᒪᓐᓇ ᐅᑦᓗᖅᑲᖅᓯᐅᖅᓂᖅᓴ ᐊᔪᓴᖅᓯᒪᓕᓐᓄᑦ ᐊᑯᓂᐅᔨᖅᑐᑦ, ᐃᓐᓇᐅᓕᖅᑐᑦᓗᓐ ᓯᓚᑦᑕᓂᖅᑐᓐᓇᑦᓗᓐ ᓄᓕᐊᖅᑕᓯᓯᑎᓕᖅᑐᑦᓗᓐ ᑭᓴᖅᑲᓇᐅᓂᖅᑎᓂᖅᓴ ᐃᑲᓂᐊᖅᑐᑦ ᑎᖅᑕᐊᖅᑲᓇᐅᖅᑐᖅᑕᖅᓴᐅᓕᖅᑕᖅᑐᑦ (ᑕᑯᔭᖅᓴᐃᑦ 4 ᐊᒻᒪ 5). ᐃᓕᓴᖅᓂᖅ ᐊᖏᕐᓇᔪᓂ (ᓯᓴᓗ ᐅᖅᑐᑎᑎᓕᓗᔪ 1997-ᒥᖅ 2004-ᒧᑦ),


ᑎᖅᑕᐊᖅᑲᕆᐅᕐᓂᕆᐅᖅᑐᑦ ᐱᑕᖅᑲᖅᐸᓂᐅᖅᑐᑦ ᐊᖅᑭᓯᐊᓗᖅᑕᒃ ᐅᑦᓗᓯᖅᑲᕆᐅᕐᓄᖅᑐᒃ ᑕᐅᕐᓂ ᖃᖅᓱᐊᖅᑕᑦᑕᖅᑕ (ᐊᐅᓪᓗᓈᓂᒃᑯᑦ ᓄᓇᕆᔨᓕᒃ), ᐱᑕᖅᑳᐊᓕᓪᓂᐅᖅᑐᒃ ᑕᐅᕐᓂ 1970-ᖅᑭᓐᓂ (ᑯᐱᔭ-Kerbes, 1975).

ᑲᖓᐃᑦ ᒥᑭᔨᑎᐅᓂᖅᓴᐅᓂᖅᑕᐅᔨᑦ ᓯᓗ ᓂᕿᑲᖅᓴᓕᑳᐅᑦᓪᓚᐅᖅᑐᑦ ᓴᑦᓕᒥᐅᑕᓄᑦ, ᐱᓗᐊᖅᑐᒥᒃ ᐊᑎᖃᔪᓂ ᑐᐅᖅᑐᑕᐊᕆᐊᓂᖅᑎᓗᒍ ᑕᐅᕐᓂ (Carter, et al., 2018). ᑲᖓᐃᑦ ᑎᑭᐸᖅᑐᑦ ᑕᒪᐅᖅᓱ ᑕᓂᕿᑎᖅᑐᓂ ᓄᑭᑐᐊᓂ ᓄᓇᓕᒻᒥᐅᑦ ᐅᑦᓗᓕᕆᓂᖅᐸᑦᓕᓕᐊᓴᖅᑲᕆᖅᓯᑎᒃ ᕿᑭᖅᑕᖅᓱᓂ ᐸᓂᖅᑐᑎᖅᑭᓐᓂᐊᖅ ᓄᓇᓂᖅ ᖃᐅᔨᑎᖅᓯᖅᑐᓗᓗ ᐃᒃᑐᑎᐅᔨᓂ ᐃᒪᐅᑎᖅᑭᓐᓂᐊᖅ (Sutton, 1931; Bray, 1943). ᒪᓐᓂᖃᑦᖅᑎᑦᓗᑎᑦ ᕿᑎᖅᐸᓯᐊᓂ ᑕᖅᑭᐅᐸ ᔨᓕᐃᐸ ᐊᖏᓗ ᖃᖅᖓᑕᕿᓐᓂᕆᒪᓂᖅᑎᑦ, ᑲᔪᖃᑦᑕᖅᐳᑦ ᓄᓇᖅᐱᐅᓂᔪᑦ ᐃᒃᑕᑐᐊᔨᓂ ᐊᖏᓂᖅᐸᑦ ᑯᖓᓕᒃ (Cooch, 1957; Bray, 1943; Parker & Ross, 1973). ᑲᖓᐃᑦ ᓄᑦᐊᖅᓯᒪᓕᕐᓂᑦ ᐊᐅᑦᓕᖅᐸᑦᓕᓕᐊᑕᖅᑦᑕᖅᑐᑦ ᓴᑦᓕᓂᑦ ᐊᖏᓯᐅᐸ ᕿᑎᖅᐸᓯᐊᓂ, ᓄᑦᐊᓕᐅᖅᑐᑎᓗ ᐃᒃᑐᑎᐅᔨᓂ ᐊᐅᑦᓕᖅᐸᑦᓕᓕᐊᑕᖅᑦᑕᖅᑐᑦ ᓴᑦᓕᓂᑦ ᐊᐅᑦᓕᖅᐸᑦᓕᓕᐊᑕᖅᑦᑕᖅᑐᑦ ᓴᑦᓕᓂᑦ ᐊᖏᓂ

ᑕᓗᐅᖓᕐᕉᑐᓂᓕᕆᑎ (Bray, 1943). ᒪᵒᓂᖓᕐᑎ ᐃᵒᓄᒃᓯᓕᕆᓕᕆᖅᑕᕐᑐᑦ ᖁᑎᖅᓯᓴᐅᓂᒃ ᐃᓴᖅᓯᓴᐅᓄᑦ ᑕᖁᐅᓯᓕᐅᑦ ᐊᓴᖅᓯᓴᐅᓄᑦ ᐊᓯᖅᓯᓴᐅᓄᑦ (Atlantic brant) ᓂᕆᓕᕆᓄᑦ ᑎᓂᖓᓂᕆᔭᐅᖏᓐᓄᖅᓯᓴᐅᓕᖅᓴᓴᑦ (Cooch & Barry, 1957).

ᑕᒪᒃᑯᐊᕐᑕᐅᖅ ᓂᕆᓕᕆᓄᐃᑦ (ᖃᒃᓕᓄᐹᑎᑐᑦ Cackling geese) ᐱᑕᖃᕆᔨᖓᒥᔨᑦ ᑎᖓᒥᐊᖃᖅᖁᖓᒥ. ᓂᕆᓕᕆᓄᐃᑦ ᑕᒪᒃᑯᐊ ᑎᑭᖃᑕᕆᓂᖓᓂᑦ ᐅᓪᓗᕆᖃᓄᖓᓂᓗ ᐱᐅᓴᖃᑐᐃᓄᐃᓇᑲᑎᑦᑎᐊᕆᐅᑦ ᑲᖓᖅᑎᑐᑐ. ᐊᒥᓴᓂᖓᓂᑦ ᑕᒪᒃᑯᐊ ᓂᕆᓕᕆᓄᐃᑦ ᐱᒻᒪᓕᖓᒥᒃ ᐊᒥᓴᑕᓕᐊᖅᓴᒪᔨᑦ ᐊᓴᖓᓐᓂᓂᒃ ᓄᐊᖓᓐᓂᓂ ᓴᓪᓕᐅᑦ ᕿᑭᖅᑕᑐᐊᓂ ᑕᐃᒪᓐᓇᓂᒃ ᐊᖏᕆᔪᖓᓐᓂᒃ 1979–1980 (Nissley, 2016; Sharp & Abraham, 2010). ᖃᓄᐃᑕᕆᓂᖓᓇᓂᖓᓂᑦ ᑕᒪᒃᑯᐊ ᓂᕆᓕᕆᓄᐃᑦ (Cackling) ᐃᑲᑕᐊᖏᑦ ᑎᖓᒥᐊᖃᖅᖁᖓᒥ ᖃᑲᐅᒪᔭᐅᖓᑕᓪᓕᐅᑦ.


ᐊᐢᑎᓕᓪᓚᕆᐊᒃᑯᑦ ᓱᓕᐅᔪᓐᓇ ᐊᒻᒪᓗᓂᖅᓯᓕᓪᓚᕆᐊᓕᓪᓚᕆᐅᓪᓚᕆᐅᔪᓐ
ᐃᑲᑕᐅᐊᖅ ᑎᖏᒥᐊᖅᖃᖅ
0 2.5 5 10 15 km
N
ᓄᓇᓕᒻᒥᐅ ᓄᓇᓕᖕᒥᓄᑦ ᖃᐅᔨᓴᖃᖅᑐᑦ ᐃᓄᐃᑦ ᐊᑐᓂᖅᓯᒪᔨᖅᑕᐅᓂ



ᑕᐻᓴᓴᖅ 5: ᑲᖓᐃᐱ ᐅᓪᓗᖅᕿᖏᑦᑕ ᓇᓂᓂᐅᓕᖏᑦ ᑐᐸᑕᐅᖅᑕᐅᔪᓐᓇᑎᒃ 2014-ᒥ ᓴᓪᑕᐅᐳᑦ ᖃᐸᖅᑕᓪᓗᐊᓂᒃ ᖁᑎᐅᔨᓴᖅᑕᐅᓂᖏᓐᓂᒃ ᐊᓱᓕᑕᐅᖅᑕᐅᓪᓗᑎᓐᓇᑎᒃ ᖁᑎᐅᔨᓴᖅᑕᐅᓂᖏᓐᓂᒃ.

2.2.2 ᓄᓇᖅᑐᓄᔨᓂᖅ ᑎᖏᐊᑦ

ᓄᓇᖅᑐᓄᔨᓂᖅ ᑎᖏᐊᑦ ᓇᓂᑐᐃᓇᕎᑎᐊᖅ ᐊᒥᓱᓕᓂᖅᑐᑦᓯᓕᕐᓂ ᑲᓇᐊᑦᑲᐃᔪᓂ ᐅᓇᑦ (Gratto-Trevor, et al., 2011). ᖃᓄᐃᑎ ᓄᓕᐊᕐᓇᖃᓱᒋᓂᖏᑦ ᐃᑲᑦᑐᐊᑦ ᑎᖏᐊᖃᖅᖃᓇᓂᓄ ᖃᑲᔪᒪᔨᔨᖏᓕᑦ ᑭᓱᐊᓂ ᐊᔪᓚᑕᑭᓴᓇᓱᖏᑦ ᐊᓂᖏᓐᓂ ᓄᓇᒋᔭᐃᔪᓂ ᓴᑦᑐᐊᑦ ᐊᑎᖏᓐᓂᐅᔪᓂ ᖃᑭᑐᖅᑕᐃᓕᐊᓂ, ᑕᐅᑲᑕᐅᖃᑦᑕᖅᓯᒪᔪᓂ ᖃᑲᔪᓴᖅᑎᓄᑦ (Smith, et al., 2012). ᓴᖅᑐᐃᑦ, ᖃᑦᑕᖅᑦᑦ, ᐅᓄᑦᑦᑦ ᐊᓪᒪᓗ ᓄᓇᕗᐊᖅᔪᐊᑦ ᐱᑕᖃᖅᖃᔪᓐᓇᑦ ᑕᐅᖅᓯᓂ (Bray, 1943). ᐊᓯᖏᑦᑕᐅᖅ ᑎᖏᐊᑦ ᓄᓕᐊᕐᖃᑦᑕᕆᓯᐅᑎᐅᓐᓂ ᑎᖏᐊᖃᖅᖃᓇᖏᑦ, ᒪᓂᐊ, ᐃᓓᖅᔪᐊᖅ/ᑐᑦᓭᐅᓴᖅᓱᒃ, ᖃᓂᑦᖃᓭᐅᔪᖅᓱᒃ, ᑐᖏᐃᑎᑦᑎᖅᓱᒃ ᐊᓪᒪᓗ ᐃᓴᖅᖃᖅ ᓄᓇᕗᐊᖅᔪᒃ.

ᓴᑦᓭᖅᑎᐅᑦ ᐅᓂᒃᑲᓭᐅᓭᐅᕆᐅᖅᑐᑦ ᐊᒥᓱᓕᓂᖅᑐᑦᓯᓕᕐᓂᖅᑲᓂᖏᓐᓂᒃ ᓄᓇᖅᑐᓄᔨᓂᖅᑎᓴᓂ ᐊᔨᔪᑦᖏᓐᓂᒃᓂᖏᑦ ᓴᑦᓭᐅᖅ ᖃᑭᑕᓗᓇᖅᑐᓄᑦ, ᐱᓗᐊᖅᑐᑐᒃ ᑕᒪᒃᑯᐊ ᓴᖅᑐᐃᑦ, ᓇᓗᓴᐃᓕᐊᖅᑐᑦ ᐊᒥᓓᓕᐊᓂᐅᓴᓗᓂᒃ ᐃᓕᓴᕐᓂᖅᑐᑦ ᐅᑎᖅᑕᐅᓂᒃ (Carter, et al., 2018). ᓴᑦᓭᖅᑎᐅᑦᑕᐃᑦᑕᐅᖅᑲ ᑕᑯᓚᒃᑭᓕᐅᖅᑐᓄᑦ ᓄᓇᖅᑎᐅᑕᓂᒃ ᐊᒥᔪᓯᓀᓕᐅᖅᑐᓄᓪᓗ ᐅᐱᖓᓇᒃᑯᓂ ᓄᓕᐊᕐᓂᐅᑦ, ᓴᑦᓭᖅᑎᐅᑦ, ᖃᓄᐃᓕᓐᓂᒃᑯᓂᖏᓐᓂᒃ ᑲᑎᕐᓂᖅᑐᑦ ᓄᓇᖅᑐᓄᔨᓂᖅᑎᓴᓂ ᐃᓐᓇᑭᓐᓂᒃᑯᓂᖏᓐᓂᒃ ᑲᑎᕐᓂᖅᑐᑦ ᓴᑦᓭᖅᑎᐅᑦ ᐊᒥᓱᓕᓂᖅᑐᑦᓯᓕᕐᓂ ᑲᑎᕐᓂᖅᑐᑦ ᓴᑦᓭᖅᑎᐅᑦ ᐊᒥᓱᓕᓂᖅᑐᑦᓯᓕᕐᓂ ᐊᓯᖏᑦᑕᐅᖅ (Pectorial-ᓂᒃ) ᐃᓐᓇᐃᖅᑲᒃ ᖃᑲᔪᓴᖅᑎᐅᑦᑕᐅᖅᑐᓂ ᑎᐅᖅᑕᖅᑕᖅᑐᐅᑦᑕᐅᖅᑐᓂᒃ ᐅᐸᐅᖅᑕᖅᑐᐅᑦ ᐅᐸᐅᖅᑕᖅᑐᐅᑦᑕᐅᖅᑐᓂᒃ ᓴᖅᑐᐃᑦᑐᐅᑦᑕᐅᖅᑐᓂᒃ ᐃᓕᖅᑲᓯᐅᑦᑐᐅᑦᑕᐅᖅᑐᓂᒃ ᐊᑕᐃᓴᖅᑐᑦᑕᐅᖅᑐᓂᒃ ᐃᓐᓇᑭᓐᓂᒃᑯᓂᖏ

ᓴᖅᓯᒥᐅᓄᑦ ᐅᖃᐅᓯᖏᖅᑭᓯᒪᔪᑦ ᐃᓱᐊᓕᓵᑎᑦᑐᖓᓂᖅᐳᑦ ᐊᒥᓱᖓᓗᐊᓯᖅᓂᖓᓂᖅᐳᑦ ᐃᓕᖓᑎᑦ ᑕᒪᐅᑯᐊ, ᐱᓗᐊᖏᑐᒥᖅ ᑕᐃᐅᑯᓂᖓᓪᓚᐃ ᐃᒥᖏᑯᑕᐃᓕᓴᓂᐅᖅ ᐊᓪᒪᓗᓐ ᑐᓴᓕᖓᓂᖅᐳᑦ ᐊᓪᒪᓗᓐ ᖃᖏᖏᓴᐅᓂᖅᐳᑦ. (Carter, et al., 2018). ᓄᓇᖓᓂ ᑕᐅᖓᓂ ᒪᒥᔪᓱᐃᑐᖏᖅ ᐅᐱᖓᓱᐅᑯᑦ ᑎᑎᕐᓂᐊᕐᓗᒃᑐᖏᖅᐳᖏᖅ ᑕᐅᔪᓚᖓᑎᐅᓴᑐᒃᑐᓄᑦ ᐊᖏᕐᕋᓂᖅ, ᑲᓇᖓᓇᖏᑦᓴᖓᓚᐅᑎᓕᕐᓕᐊᓂᖓᓂᓕ.

2.3 ᐊᓱᖓᑦ ᐅᒪᔪᐃᑦ

2.3.1 ᓄᓇᒥᐅᑕᐃᑦ ᐅᒪᔪᐃᑦ ᐊᐃᒪᓚᑎᑦᑎᓱᑦ ᐱᒥᓂᖅ

10-ᖓᔪᑦ ᓄᓇᒥᐅᑕᐃᑦ ᐅᒪᔪᑦ ᐱᑕᖏᓱᑦ ᐅᕐᓕᓪᓂᓂᖅ ᐱᑕᖏᓚᑕᖏᓱᒪᔪᑦ ᐃᓚᖏᑐᐊᑦ ᑎᕐᒥᐊᖏᖅᓴᖓᓂᓂ. ᓴᓕᐅᑦ ᖁᑭᖏᒋᓪᓗᐊ ᐃᓕᑕᒪᓴᐅᓱᒪᔪᖏᖅ ᑎᓪᓚᓂᐊᑕᒪᒪᓪᓚᐊᓪᓂᕐᒧᑦ. ᐅᑭᐅᖏᑕᖏᑐᒥ ᑎᓪᓚᓂᐊᑦ ᐊᒥᓱᓂᕐᑕ ᑭᓱᐊᓂᑕᑕᐅᖏᓚ, ᐊᓱᓴᑭᓕᑕᐅᖏᓚᖅᑐᓂ ᐊᖓᔪᖅᑕᐃᓐᓇᓂᖅ ᐃᐱᕐᓚᐅᑦ (ᐸᑯᓂᓗ Parker, 1974).

ᓭᐸᑐᖏᓂᑐᕐᖁᐅᑦ ᑐᐅᑐᐃᑦ ᐱᑕᓕᐊᓪᕿᓭᓕᐅᖏᓱᒪᔪᑦ ᓴᑦᓕᓂ ᑕᐃᓴᓱᒪᓂᐅᑦᓕᓂᓪ ᑭᓱᐊᓂ ᑕᐃᓕᓂ 1955-ᖓᔪᖏᑎᑦᓕᔪᑉ, ᓄᖓᔪᑎᐊᓴᒪᓭᓕᐅᖏᖅᓱᒪᔪᑦ (Sutton, 1932). 1967-ᒥ ᐅᒪᔪᓕᓪᓚᓵᑦ ᐊᓴᖏᖅᑐᐃᑎᐊᓕᓕᐅᖏᖅᑐᑦ ᑕᐅᓭᖓᑦᓴᖅᑐᓂᓪ ᖁᑭᖏᖅᑐᐃᓂᖅ (Manning, 1967; Parker, 1975). ᑕᐃᒪᖓᓂᓪ, ᐊᒥᓱᓂᓂᖏᖅᐸᓚᑦᐅᐅᓕᐅᖏᖅᓴᑦ ᑕᒪᓂᓕᐅᖏᖅᑐᑦ 30,000ᖓᓂᓂᑐᑦ ᑐᐃᑐᐃᑦ, ᐊᖏᕐᔪᓕᐅᖏᖅᑐᒥ 1997-ᒥ. ᓴᓕᓪᓂ ᑐᐃᑐᐃᑦ ᐊᒥᓱᑭᓚᐊᖏᓱᒪᓕᓪᖏᖅᑐᑦ 2013-ᒥ 2015-ᒥᓗ. ᑐᐃᑦᖏᓕᓪᖏᖅᑐᑦ ᑕᐃᓂ 12,300-ᖓᓂᓂᑐᓂᓪ (Campbell & Boulanger, 2015). ᐃᓕᖓᓂ ᐃᖏᑕᐃᐊᖅᑎᓪᓗ ᑎᖓᒥᐊᖏᖅᑎᖓᓗ ᓄᓇᖓᓂ ᐊᐅᓴᖏᓂ ᑐᑐᖏᓱᓂᖓᑦ (Parker, 1975). ᑐᐃᑐᐃᑦ, ᑭᓱᐊᓂ ᐊᒥᓱᓗᐊᑭᓇᖓᑐᑦ; ᑕᐅᔪᓇ ᓄᓇᖓᑎ ᑭᓂᐸᓗᐊᖏᓕ ᓕᓴᐃᓇᓕᐅᓕᐊᖏᖅᓱᓂ, ᑐᐊᐸᐃᓇᐅᓕᓪᓗᐊᖏᖅᓱᓂᓗ. ᑕᐃᓴᓱᒪᓂᓕᑕᐅᖏᓪ, ᐃᖏᑕᐃᐊᑦ ᑎᖓᒥᐊᖏᖅᑎᖓᓚᓂᓂ ᐅᐊᖓᓇᖏᖅᐸᓱᖓᒡᒐ ᐱᖓᑎᓇᖓᓂᓂ ᓄᓇᖓᓂ ᑐᐃᑐᓄᑦ ᓄᖏᓚᐃᐅᕐᓕᐅᓚᖓᑦ (ᓄᓇᖓᓕᒥ ᓄᓇᐃᑦ ᐊᑐᑎᐅᓴᓓᖓᓄᑦ ᐸᖏᓇᐃᓐᓂ ᑲᒥᓱᓇᐅᑦ, 2000). ᐃᓐᓇᓀᖏᖅ ᖁᐅᐱᒪᔪᓴᑎᑦ ᐃᓱᒪᓇᖏᖅᓱᑦ ᐱᒪᓚᐅᐅᓂᖓᓂ ᓄᖏᓚᐃᐅᕐᓕᓂᖅᑐᑦ ᐊᕐᓗᐊᐊᓂᓴᓕᖏᖅᑐᑦ ᑲᓪᓕᐅᑦ ᓄᓇᖓᒡᒐ ᐅᓇᓂᖅᐸᓱᖓᓂᓪ (ᓄᓇᓕᒥ ᓄᓇᐃᑦ ᐊᑐᑎᐅᓴᓓᖓᓄᑦ ᐸᖏᓇᐃᓐᓂ ᐅᒥᓱᓇᐅᑦ, 2012).

ᐊᒥᑭᐃᑦ ᐊᒥᓱᐊᓂᕐᕋᓂᐅᖏᖅᓱᒪᔪᑦ, ᓄᖓᔪᑎᐊᓴᒪᓭᓂᖓᓕᓕᓂ ᑲᓪᓕᐅᑦ ᖁᑭᖏᖅᑐᒋᓂᖓᓂᖅ ᓄᓇᓕᖓᓄᑦ ᐱᓴᐅᖏᑕᖏᓂᓕᐅᖏᓱᖓᓂᖅᑐᓄᑦ ᐊᒥᓱᓂᓕᐅᓕᐊᖏᖅᑐᖏᖅ ᑐᐃᑐᓂᓪ. ᑭᖓᔪᓕᖏᖅᐸᖏᖅ ᐊᒪᑭᐅᖅ ᖁᑯᖏᑕᐅᓕᐅᖏᖅᓱᒪᔪᖏᖅ ᑕᐃᓂ 1937-ᒥ (Manning, 1942). ᐊᒪᑭᐃᓕᓗ ᑎᓪᓚᓂᐊᓕᓗ ᐊᐅᐸᓗᐅᑦ ᑕᑯᓴᐅᓚᔪᐃᑦᑐᓚᓪᓚᐊᓗᓕᑦ. ᓚᖏᑕᐃᓱᓂᓂᖅ ᐱᓂᓱᓂᓚᓂᓂ ᖁᓇᓂᓂ ᒥᑭᒧᖁᑕᐅᓱᒪᔪᖁᓕᐅᖏᖅᑐᖏᖅ ᓴᓕᐅᑦ ᐊᑕᐅᓱᓂᑦ (Roe’s Welcome Soundsd-ᖓᓚᔪᑦ), ᓴᓕᐅᑦ ᖁᑭᖏᖅᑐᒋᓗᐊᓄᑦ.

ᐅᑲᓪᓕ ᐱᑕᖃᕆᔨᖏᓪᓗᓕ ᑎᖅᒥᐊᖅᑎᕕᐅᓂᖏᑦ, ᐱᑕᖃᖅᖢᓱᓪᓕ ᖃᓄᑎᖅᓂᕐᕿᓂᕕᐅᓂᖏᖅᓂ ᓄᓇᒪ. ᑕᒪᑲᓄᐊᑕᐅᕐᖓ ᑎᓇᐊᑦ ᐱᑕᖃᖅᑭᒃᑲᓪᓚᕐᓂᒃ ᑎᖅᒥᐊᖅᑎᕕᖅᖢ. ᐊᑕᐅᓂᕆᑦ ᑕᓄᐅᕐᔪᖅᑲᓪᓕᐅᕐᖓᑦ ᑲᓪᓕᐅᖅᓯᓂᐅᓂᖅᓯᓂ ᑲᑎᒪᔨᖅ ᐱᓕᕆᔨᖅᑕᐅᓂᖅᓯᓂ ᑕᐃᒪᐃᓐᓇ 2015-ᒥ.

2.3.2 ᑕᓚᐅᖅᑎᐅᑕᐃᑦ ᐅᒪᔪᐃᑦ

9-ᖓᐅᑦ ᑕᓚᐅᖅᑎᐅᑕᐃᑦ ᐅᒪᔪᐃᑦ >ᐃᓕᑦ ᐱᑕᖃᓂᑦ ᐃᑲᑦᑐᐊᖅᑎᒥ ᑕᐃᒪᐃᓐᓂᕐᓗ ᐱᑕᖃᓂᖓᓕᓐᓂᑎᒃ ᐃᑲᑦᑐᐊᑦ ᑎᖅᒥᐊᖅᑎᖃᓕᓴᓂ. ᑕᐅᕐᓂ ᓴᓪᓕᐅᑦ ᐅᐊᖏᓇᖅᑲᓂᔪᖅᓴ ᐊᐅᖅᑲᖏᓪᓕᓴᖅᑕᓪᓕ, ᓴᓪᓕᐅᑦ ᑎᒋᓂᒃ Roe’s Welcome Sound-ᒥ ᐅᑭᐃᐅᕐᒪᑐᖅ ᖃᑭᓇᑐᓴᓪᓕ ᖃᑲᓂᔪᓇᓪᓗ. ᖃᑲᓂᔪᐊᑦ ᐅᑦᑲᑐᑦ (ᖃᑲᓂᔪᓕᕐᑐᑦ) ᑕᒋᓂ ᑎᖅᒥᐊᖅᑎᖃᓕᓴᖅᖢ ᑎᓂᖓᓂᖅᖢ. ᐅᓴᐊᐃᓕ ᐱᑕᖃᐃᖖᓇᓂᖓᓐᓂᖅ ᐃᑲᑦᑐᐊᖅᑎᖅᖢ. ᐊᐅᖅᑲᑯᑦ ᐊᒥᓂᕐᑐᑦ ᖃᑭᓇᑐᐃᓕ ᑕᐅᕐᓂ ᐃᑲᑦᑐᐊᖅᑎᖅᖢ, ᐃᑕᖖᓇᑯᓪᓗ ᓂᖃᕈᓴᖃᑦᓯᐅᕐᓂᕐᓄᓪᓚᐊᓂᖅ. (ᓪᑐᐊᖅ ᐱᓪᓚᖃᑎᖅᓂᓪᓗ-Latour et al., 2008; Fᐊᖖᑎᖖ ᐊᓚᒪᓗ ᒪᓗᓚᓇ-Fontaine and Mallory, 2011).

ᓄᓴᖅᖢᓂᓄᖓᓂᖖᓂ ᐊᑎᑎᖅᑐᐅᕐᓂᓂᑦ ᓄᓇᐃᑦ ᐱᓚᒪᓇᐊᔭᕐᑐᐅᑦ ᐅᓄᕐᓄ ᓂᖃᕈᓴᖃᑦᓯᐅᕐᓂᖖᓇᑦᓂᔪᑦ ᐊᐅᖅᑲᓯᓂ (ᕐᖖᑎᑦ-Fontaine, 2011). ᓄᓇᖓᓄᑦᑐᑎᓄᓂᖓᓐᖅ ᑕᐅᕐᓂ ᐃᓗᔪᑎᓄᓂᖓᓐᓂᑎᒃ (ᓄᑎᖅᑎᓄᓂᖓᓐᓂᑎᒃ) ᑎᖅᒥᐊᖅᑎᖃᓕᓴᖅᑎᑦ, ᑕᐅᕐᓂᓗ ᓄᓇᒥ ᖃᑭᒥᐊᓪᓚᐃᑭᒪᑦ ᐊᒪᓚᓗ ᖃᓄᐅᑲᓇᑕᐃᓕᓐᓂᖅ ᐃᓄᒃᑎᐊᖅᑐᑦ. ᐃᓄᐃᑦ ᖃᓄᐅᓕᒃᑐᑐᖅᓯᓂᖅ ᐃᓄᓕᓴᓂᓄᖅ ᐃᓄᓕᓴᓂ ᐊᑎᓄᓴᑕᓪᓚᕆᓂᕐᓂᖓᓂᖅ ᑕᐃᒪᓐᓇᓂᖅ ᐊᖏᔪᖅ 2004-2012-ᒥᖅ, ᐊᒪᓚᓗ ᖃᓄᖅᓯᓴᑎᖅᑕᐅᕐᖓᓂᖓᓂᖅ-ᐊᖓᔪᖅᐅᕐᖓᓂᖅᓯᓗᑎᒃ ᖃᓄᐅᓕᕆᖅᑕᐅᖃᓄᑕᖅᓂᖓᓂᖅ, ᖃᓄᓐᔪᐊᖅᑎᑕᐅᓯᒪᔪᑎᔪᑦ ᖃᓄᐅᓕᒃᔨᔪᑕᐅᕐᓂᖓᓂᖅ ᒪᓕᔪᐊᑎᓕᖅᓂᖅ, ᐊᖏᖏᔪᐃᒪᓐᓇᑦ ᑎᒥᖅᓯᓄᓄᖅ ᖃᓄᐅᓕᖅᑕᐅᖃᓄᑕᖅᓯᒪᔪᓇᑎᒃ 20-ᓂᖅ ᐊᖏᖏᔪᓂᖅ, ᐃᓯᒪᓇᖅᑐᑦ ᑕᐅᕐᓂ ᓴᓂᖏᓕᕐᓂᑦ ᓯᑯᓯᓚᕐᓂᕕᐅᓂᖅᓂᖓᓂᖅ ᐊᓯᓕᑭᓚᐊᖅᓯᒪᓕᖅᓂᑦᑐᖅ. ᐊᒥᓱᓂᕕᐅᓂᖅᓯᓂᖅ ᐃᓄᐃᑦ ᐊᒥᓱᓂᖅᓂᖓᓂᖅ ᐊᓯᓕᑭᓚᐊᖅᓯᒪᔪᓇᑎᒃ ᐊᒥᓱᓂᖅᓂᖓᓂᖅ. ᐊᒥᓱᓂᖅᓂᖓᓂᖅ ᓴᓪᓕᐅᑦᑐᕐᓂᓗ ᖃᓄᐅᓕᖅᑕᐅᖃᓄᑕᖅᓯᒪᔪᓇᑎᒃ ᐊᒥᓱᓂᖅᓂᖓᓂᖅ ᐃᓄᐃᑦ ᑕᐃᒪᐃᓐᓇᑦᑐᖅ.

2.3.3 ᑕᓚᐅᖏᓪᓂᕐᓂᒥ ᑕᓯᑎᐅᑕᐃᑦ ᑕᓚᐅᖅᑎᐅᑕᐃᓕᓐᓂ ᐃᖅᑲᓇᑐᓴᐃᑦ

ᐊᒥᑎᓵᓇᐃᑦ ᑕᓱᓕᓴᐃᑦ, ᑕᓯᐊᖅᓴᐃᓕᓐᓂ, ᑯᓕᓇ, ᑕᓯᐅᑦᓴᓴᐊᓕᓐᓂ ᑎᖅᒥᐊᖅᑎᖃᓕᓴᖅᖢ ᐃᒪᖅᑲᐅᖅᓴᑦ ᐊᒪᓚᓇ ᑕᓚᐅᖅᑎᒥ ᐃᒪᖅᓂ ᐊᒥᓱᓂ ᐊᐅᖅᑎᕐᓂᕐᑐᓂᐅᓂᑲᑦ ᐃᖅᑲᓇᑐᓴᐃᓕᓐᓂᑎᒃ. ᑕᒪᐃᓐᓂᑲᑦ ᖃᓄᐅᓕᕆᔨᖅᑕᐅᖃᓄᑕᖅᓯᒪᔪᓇᑎᒃ ᐊᐅᖅᑲᑦᑐᕐᓂᕐᑐᓂᐅᓂᑲᑦ ᐃᖅᑲᓇᑐᓴᐃᓕᓐᓂᑎᒃ ᐃᓄᐃᑦ ᑎᒥᖅᓯᓄᓄᖅ ᑎᑎᖅᑲᓂᔪᓇᖅᑐᑦ. ᑭᓱᐊᓂᓕᕆᓂᖅ, ᑕᒪᓐᓇᓂᖅᑕᐅᖅᑲᓂᔪᓇᖅᑐᑦ 40-ᖓᐅᓐᓂᑐᑦ ᒪᔪᖅᑭᒃᑐᑦ/ᓱᑐᖓᓂᑐᑦ ᐊᒪᓚᓗ ᑕᓚᐅᖅᑎᐅᑕᐅᖓᓐᓂᖏᓐᓇᖅᑭᒃᑐᑦ ᐃᖅᑲᓇᐃᑦ ᐊᐅᖅᑲᓂᕐᑐᑦ, ᐃᓕᖅᑲᖅᓱᑎᒃ ᐊᑕᐅᓱᖅᑎᑦ ᐊᒥᓱᓂᑐᐊᓕᖅᓂᖅᑕᐅᓱᒪᔪᓕᐅᑦ, ᐱᑕᖃᖅᓄᑦᑎᐊᖅ (ᐊᐃᒃᑐᑐᖅᓱᒪᔪᓕᐅᓂ ᑕᑯᑎᑕᑎᔪᓕᑦ 3; ᑯᑦ ᐊᒪᓚᓗ ᕿᐃᓚᓂ-Coad and Reist, 2004)

2.4 ᐱᑭᑐᖅᓂᖅ ᓄᓇ

ᓄᓇᖅᓄᓂ ᑕᐅᕐᓂ ᐃᑲᓂᑐᐸᑦ ᑎᖅᒥᐊᖅᑎᓇᖅᓄᓂ ᐱᑭᑐᒪᒪᓇᐊᑦᑐᕐᖅ ᑕᒪᑲᑯᓂᖅᓱᒃ ᑭᓚᐢᓇᖅᓂᖅ. ᐱᑕᓕᐊᑦᔨᖅᑐᐊᖅᓯᑎᑲ ᐊᓯᐢᖅᑭᓂᓕᐅᑕᖅ ᐱᑕᖅᓯᐅᒍᓂ ᑕᒪᑲᑯᓂᖅᓄᓂ ᐃᐃᖅᓂᖅ, *ᑕᑭᓱᑯᑦᓵᖅᓂᖅ* ᑕᐃᑲᑯᓂᖅᓄᓗ *ᓄᓇᕆᖅᓯᑦ ᐃᐃᑯᑦᖅᓂᖅ. ᒪᓕᒃᑲᐃᓗ ᓄᓇᕆᐃᑦ, ᐅᕆᖅᐅᖅᓴᐃᓕᓕᓗ,* ᐊᒻᒪᓗ ᑎᖅᓄᐅᖅᓴᐃᑦ <ᓂᕆᓂᖅᓴᐅᓂᕆᖅᑎᖅᑕᕐᖅᑐᑦ. ᕆᑯᒪᔪᑦ ᒪᑭᑕᓱᖅᓂᖅᓴᐃᑦ ᐱᐅᖅᑐᑕᐅᖅ ᓯᕆᓗ ᐱᑕᖅᓯᒧᖅᑎᒋᔭᐅᑦ, ᐃᒃᓯᑕᐃᑦᓗ ᑭᓯᑦᑕᐃᓕᓴᖅᑐᖅ ᐊᐅᓚᓗᑐᖅ, *ᑕᒪᐃᓂᖅᓄᓗ ᑭᓚᐢᓇᖅᓂᐅᓂᖅᓂᕐᓂᓂᐅᓂ ᐊᑎᓯᐊᓕᐅᖅᖁᔭᐅᑦ. ᐱᑕᖅᓄᓘᐅᑦ* ᑲᑎᒪᔨᖅᑐᑦ ᐱᑕᖅᑐᑦ ᐃᐃᖅᓂᖅ ᐃᓕᓴᐃᑦ ᐊᒻᒪᓗ ᐃᓕᓴᐃᓂᖅᓄᓂ ᐊᑎᓯᐊᓕᐅᖅᖁᔭᐅᑦ

ᐊᒥᓱᖅᑭᓗᐊᓕᕆᓂᖅᓄᑦ ᒪᓕᒐᖅᑭᑦ) ᐃᓕᑕᒪᔭᐅᓱᒪᓕᕐᓂ ᐅᕝᕙᓚᓐᓂ ᐃᓕᔭᐅᑐᐃᓐᓇᒪᐊᖅᑲᖅᓴᑦ ᒪᓕᒐᖅᑎᒍᑦ ᑕᐃძᐊ ᑎᖅᒥᐊᖅᑲᐃᖅᑕᒃ ᐱᑕᖅᓱᑦ. (ᐊᐃᔪᖅᑐᓱᒪᓚᔪᓂ ᑕძᑎᕋᑎᔪᓕᑦ 3).

ᐊᐃᔪᖅᑐᓱᒪᓚᔪᓂᒃ ᑕძᑎᕋᑎᔪᓕᑦ 4: ᐅᒪᔪᐃᐃᑦ ᐊᒥᓱᖅᑭᓗᐊᓕᕆᓂᖅᓂᑦ ᑎᖅᒥᐊᖅᑲᐃᖅᓗᒥ

ᑕᐃᔨᐅᓂᖅᑭᑦ ᐃᓄᔪᑎᑐᑦ ᖃᖅᑲᓇᐅᑉᓄᓇᓗ ᐅᖅᑲᔪᓱᖅᑕᑐᓄᓇᓗ ᑎᑎᕋᖅᑭᒪᓕᕐᓂᑦ	ᖃᓄᐃᐃᑕᖅᓱᓂᖅᑭᑦ ᖃᓇᑕᒥ ᐃᓕᑕᒪᔨᐅᒪᓕᐅᖅ		ᐱᑕᖅᑲᓂᖅᑭᑦ ᐃᓄᖅᑐᐊᓱ ᑎᖅᒥᐊᖅᑲᐃᖅᓗᒥ
	ᖃᓄᐃᐃᑕᖅᓱᓂᖅᑭᑦ ᖃᓇᑕᒥ ᐊᒥᓱᖅᑭᓂᕐᓂᕆᑕᐅᓱᒪᓕᐊᓂᑦ ᐅᒪᓕᐊᓂᑦ - COSEWIC-ᓄᑦ ᖃᓄᐅᔨᓴᖅᓱᒪᓕᕆᖅᑭᑦ	ᐅᒪᓕᐊᓂᑦ ᐊᒥᓱᖅᑭᓗᐊᓕᕆᓂᖅᓂᑦ ᒪᓕᒐᖅᓄᓇᓗᒥ	
ᑎᖅᒥᐊᑦ			
Red knot ᓴᖅᑭᖅ ᓱᓕᔪᒪᐊᓕᒃ - ᓗᐊᓗᐊᓪᓚᖅ *Calidris canutus rufa*	ᓄᖓᓕᓕᒫᖅᑐᑦ	ᓄᖓᓕᓕᒫᖅᑐᑦ	ᓇᓗᓇᐃᖅᑕᐅᓱ-ᒪᔪᑦ
Red-necked phalarope ᐊᐅᓪᓗᑐᖅ ᓴᐅᖅᑭᖅ *Phalaropus lobatus*	ᐊᖏᐅᖅᑭᑐᒥᒃ ᐃᓱᒪᓗᑕᐅᔪᑦ	ᐊᖏᐅᖅᑭᑐᒥᒃ ᐃᓱᒪᓗᑕᐅᔪᑦ	ᐱᑕᖅᑐᐃᓐᓇ- ᓕᐊᓪᓚᑦ
Short-eared owl ᒥᑭᓂᖅᓴᖅ ᐅᖅᐱᒃ *Asio flammeus*	ᐊᖏᐅᖅᑭᑐᒥᒃ ᐃᓱᒪᓗᑕᐅᔪᑦ	ᐊᖏᐅᖅᑭᑐᒥᒃ ᐃᓱᒪᓗᑕᐅᔪᑦ	ᐱᑕᖅᑐᐃᓐᓇ- ᓕᐊᓪᓚᑦ
Peregrine falcon ᖁᑭᓐᔪᐊᔪᖅ/ᑲᓴᒃ *Falco peregrinus*	ᐊᒥᓱᔪᓐᓂᖅᐸᓕᐊᖅᑭᑐᒥᑦ	ᐊᖏᐅᖅᑭᑐᒥᒃ ᐃᓱᒪᓗᑕᐅᔪᑦ	ᓇᓗᓇᐃᖅᑕᐅᓱ-ᒪᔪᑦ
ᓂᕐᔪᑎᑦ ᐊᒪᐃᓴᒃᑎᑎᓄᐊᖅᑐᒥᑦ ᐱᐊᑎᕋᒥᓂᒃ			
Caribou, barren-ground population ᓇᓵᑐᖅᖏᑦᑐᕋᒥ ᑐᒃᑐᐃᑦ *Rangifer tarandus*	ᐅᓗᓚᐊᓇᖅᑐᒋ-ᓕᖅᑐᑦ	ᒪᓕᒐᖅᑎᔭᖏᓐᖏᑦᑐᑦ ᓱᓕ	ᓇᓗᓇᐃᖅᑕᐅᓱ-ᒪᔪᑦ
Atlantic walrus (Central/Low Arctic population) ᐊᐃᐃᖅ (ᖁᑎᖅᐸᓪᓕᐊᓂ/ᐊᑎᑎᒃᑐᓂᖏᓐᓂ ᐅᑭᐅᖅᑕᖅ ᐱᑕᖅᓴᓂᖏᓐᓂ *Odobenus rosmarus rosmarus*	ᐊᖏᐅᖅᑭᑐᒥᒃ ᐃᓱᒪᓗᑕᐅᔪᑦ	ᒪᓕᒐᖅᑎᔭᖏᓐᖏᑦᑐᑦ ᓱᓕ	ᓇᓗᓇᐃᖅᑕᐅᓱ-ᒪᔪᑦ
Beluga whale (West Hudson Bay population) ᖁᓇᓗᑲᐃᑦ (ᐱᓐᒪᓇᓐᒪᓂ ᕼᐊᓐᓴᓐᐲᑦ) *Delphinapterus leucas*	ᐊᖏᐅᖅᑭᑐᒥᒃ ᐃᓱᒪᓗᑕᐅᔪᑦ	ᒪᓕᒐᖅᑎᔭᖏᓐᖏᑦᑐᑦ ᓱᓕ	ᓇᓗᓇᐃᖅᑕᐅᓱ-ᒪᔪᑦ

ᑕᐃᔨᐅᓂᖅᑎᑦ ᐃᓄᒃᑎᑐᑦ ᖃᐅᔨᓴᖅᑎᓄᑕ ᐅᖃᐅᓯᐊᒍᑦ ᑎᑎᕋᖅᓯᒪᓂᖅᑎᑦ	ᖃᐅᓄᐃᓕᓴᕐᓂᖅᑎᑦ ᑲᑎᒪᔨᖅ ᐃᓕᓴᕐᓂᖅ		ᐱᑕᖃᖅᓂᖅᑎᑦ ᐃᒃᑐᑐᐊᑦ ᑎᓪᓗᐊᖃᖅᖃᓕᕆᓂ
	ᑲᑎᒪᔨᖅ ᐃᓄᒃᑎᑐᑦ ᐊᐱᕕᒃᓴᓂᖅᑎᑦ ᐃᓕᓴᐃᑦ - COSEWIC-ᑯᑦ ᖃᐅᔨᓴᖃᖅᓯᒪᔪᖅᑎᑦ	ᐅᓪᓗᕈᓯᐅᑦ ᐊᑎᓯᖅᑐᓂᖅᓴᖅᑎᑦ ᓯᕗᓪᓕᖅᑎᑦ	
Bowhead whale (Eastern Canada-West Greenland population) ᐊᖏᓐᓇᑦ (ᑲᓇᖅᓭᓯᒪᓂ ᑲᓇᑕᐅᑦ- ᑕᓪᐅᖅᑐᖅᓂ - ᐊᑯᑭᑐᐃᓪᓗ ᐱᖅᒪᖅᓇᖅᓭᓯᒪᓂ ᑕᓪᐅᖅᑐᖅᓂ ᐊᖏᓐᓇᑦ) *Balaena mysticetus*	ᐊᑦᓴᐅᖅᑎᑐᑎᒃ ᐃᓱᒪᓕᑕᐅᔪᑦ	ᓯᓚᖅᑎᔪᖅᓄᖅᑎᑐᑎᒃ ᓱᓕ	ᓄᓚᓄᐃᑦᑕᐅᓱᒪᔪᑦ
Narwhal ᑐᒃᓕᓕᑦ ᖃᑭᓇᓚᒃᐃᑦ *Monodon monoceros*	ᐊᑦᓴᐅᖅᑎᑐᑎᒃ ᐃᓱᒪᓕᑕᐅᔪᑦ	ᒪᓕᑲᖅᑎᔪᖅᓄᖅᑎᑐᑎᒃ ᓱᓕ	ᐱᑕᖅᑐᐃᓐᓄᓕᐊᓕᓕᑦ
Polar bear ᓄᓇᖅ *Ursus maritimus*	ᐊᑦᓴᐅᖅᑎᑐᑎᒃ ᐃᓱᒪᓕᑕᐅᔪᑦ	ᐊᑦᓴᐅᖅᑎᑐᑎᒃ ᐃᓱᒪᓕᑕᐅᔪᑦ	ᓄᓚᓄᐃᑦᑕᐅᓱ-ᒪᔪᑦ
Ringed seal ᓄᕐᕿᑦ *Pusa hispida*	ᐊᑦᓴᐅᖅᑎᑐᑎᒃ ᐃᓱᒪᓕᑕᐅᔪᑦ	ᒪᓕᑲᖅᑎᔪᖅᓄᖅᑎᑐᑎᒃ ᓱᓕ	ᓄᓚᓄᐃᑦᑕᐅᓱᒪᔪᑦ
ᖃᖅᖁᓘᖅ *Gulo gulo*	ᐊᑦᓴᐅᖅᑎᑐᑎᒃ ᐃᓱᒪᓕᑕᐅᔪᑦ	ᐊᑦᓴᐅᖅᑎᑐᑎᒃ ᐃᓱᒪᓕᑕᐅᔪᑦ	ᓄᓚᓄᐃᑦᑕᐅᓱ-ᒪᔪᑦ
Fishes			
ᓂᐱᒍᖅ ᖃᑯᖅᓯᖅᑐᑦ *Cyclopterus lumpus*	ᐅᓚᕆᐊᓇᖅᑐᒻ-ᓪᓕᖅᑐᑦ	ᒪᓕᑲᖅᑎᔪᖅᓄᖅᑎᑐᑎᒃ ᓱᓕ	ᐱᑕᖅᑐᐃᓐᓄᓕᐊᓕᓕᑦ

2.6 ᑕᒫᓂᕆᐅᑕᑐᖅᐅᓵᖅᑎᑦᑐᑦ ᐅᒪᔨᐃᑦ/ᖃᑯᐸᐱᖃᑭᓚᓵᓄᑦ, ᐱᔭᖅᑐᓪᓚᓵᓄᑦ ᐱᑕᖅᖁᓂᖅᑎᑦ

ᓇᑭᑐᐃᓇᖅ ᐃᓚᓂᐃᑦ ᐱᓕᕆᖅᑕᓂᖅᓯᒪᓂᓚᓂ ᓴᖅᑭᑎᑎᕆᑐᐅᕐᒃᓴᑦ ᐱᑕᖅᑲᓕᔭᓇᖅᓂᖅᓯᒪᓂᓄᑦ ᐱᑕᖅᖃᖅᐸᓪᐅᖅᓂᑦ ᓚᓇᖅᓯᒪᓂᓄᑦ, ᐱᔭᖅᑐᓂᓗᓪᓂᓄᑦ, ᐅᒪᔨᓄᓗᓪᓂᓄᑦ, ᖃᑯᐸᐱᖃᑭᓄᓗᓪᓂᓄᑦ (ᑲᓇᑕᒥ ᓚᐲᓕᖅᑐᓄᑦ ᐅᒪᔨᓄᑦ ᓴᓴᓄᐊᑦᓄᓴᒧᑦ ᑲᑎᒪᔨᖅᓯᒪᓂ, 2011). ᓚᓇᔅᑦ ᑲᕐᒪᒪᐃᑦ ᑎᑦᑯᐊᖅᓱᓕᕆᓂᖅᓴᑦ 14-ᓄᑦ ᓚᓇᕐᓄᑦ ᐱᔭᖅᑐᓂᑦ ᓚᓇᔅᑦ ᐱᑕᖅᖅᑐᑐᖅᑎᑦᑕᐅᔪᒪᓕᒪᓂᖅᑐᑦ ᐱᑕᖅᑲᓕᖅᑐᓂᑦ. ᑕᒪᑯᐊᐃ ᐱᔭᖅᑐᑦ ᓄᓐᔪᑎᓚᕐᓕᐊᓱᒪᓂᖅᑐᑦ ᐊᑦᓴᐅᖅᑎᑐᑎᒃᑐᑎᑎᐅᔪᓄᑦ ᐱᔭᖅᑲᓕᖅᑐᓂᑦᑐᑎᑎᐅᔪᓄᑦ ᓚᓇᔅᑦ. ᖃᑎᖅᑐᒥ ᐱᔭᖅᑐᑦ ᓄᓐᔪᑎᓚᕐᓕᐊᓱᒪᓂᖅᑐᑦ ᐊᑦᓴᐅᖅᑎᑐᑎᒃᑐᑎᖅᑎᑦᑕᐅᔪᓄᑦ ᐱᑕᖅᑲᓕᖅᑐᓂᑦ ᐱᑕᖅᑲᓕᖅᑐᓂᑦ ᐱᑕᖅᖁᓂᖅᑎᑦ ᐃᒃᑐᑐᐊᑦ ᐅᑎᓪᓗᐊᖃᖅᖃᓕᕆᓂ.

3 ᐃᓕᓴᕆᔨᑎᖏᑦ ᐊᑐᓚᓱᐅᕙᖓᓂᕐᓂ

ᑎᓪᓗᐊᓴᖅᖃᓄᖓ ᐱᒋᓴᖅᑎᓯᑎᓂᓴᖅ ᐱᔾᔨᑎᓴᒪᓂᐊᓴᖅᓯᑦ ᐃᓱᒪᑎᕐᐅᑎᓂᕐᓂᓐᖏᑦ ᓴᓗᓂᓐᖏᑦ ᑎᓪᓗᐊᓴᓂᖅ ᓇᔨᑭᓇᖅᑕᖏᓂᓐᖏᓪᓗᓐᓄ, ᑭᓱᐊᓂᓕ ᑕᐃᖅᑯᐊ ᐃᓄᐃᑦ ᐃᑲᔪᓱᐊᖅᑕᕐᑎᑭᓴᖏᓐᓄᑦ ᐊᖏᑎᕐᓯᑦ ᐃᓕᑕᓚᖅᓱᓕᑕᐅᑎᕐᓂᑦ ᐃᓄᐃᑦ ᐃᓕᑎᕐᐅᑦᓭᓚᓂᖏᓂᖅᓂᖅᓂᖅᖓᑦ ᐊᐅᑎᖏᓂᓄᑦ. ᐃᓄᐃᑦ ᖃᐅᔨᓴᒪᔭᖓᖏᓂ ᐊᑐᑎᓕᓚᓚᐊᑐᖓᒪᑕ ᖃᐅᔨᓴᓱᐊᑕᐅᔪᓄᓇᖓᓕᒫᑦ ᓄᓇᒥᐅᑦ, ᐃᒪᓕᖅᓂᖅ, ᓄᓇᒥᐅᑕᕆᓚᐅᔪᓂᓕᓐᓄ ᖃᐅᓂᕐᓴᐅᔪᓂᖅ, ᑐᖓᔪᖓᓚᑎᑎᓐ ᑐᑭᓯᓚᐅᔪᓂᖅᓄᑦᓯᐊᕐᓇᖓᓕᒫᑦ ᐃᓱᒪᓕᐅᕈᑕᐅᔪᓚᐊᖏᖃᓄᖅᑐᓂᖅ. Ċᐅᑯᐊ ᑎᑎᕆᖏᓱᒪᓂᕐᓂ ᐅᖃᐅᔨᓱᖃᖅᓯᑦ ᑕᐃᓵᓱᒪᓂᐅᓕᐅᖅᑐᖅ ᐅᓪᓗᑎᑦᓄᑐᖅ ᓄᓇᕗᑦ ᐊᑐᖅᑕᐅᓚᐅᖏᓪᓗᖅᑐᓂᖅ ᑎᓪᓗᐊᓴᖃᖅᓄᑦ, ᐊᓚᑦᓗᐊᓂᓗ ᓄᓇᕗᑎᓂ, ᐊᒻᒪᓗ ᖃᑭᒥᖅᑭᐊᖅᑕᕐᕕᖃᖅᖢᓂᕐᓂᑎᒍᓐᓇᖅᑐᑦ ᖃᐅᔨᓴᕐᕕᐊᓚᐅᖏᓪᓗᖅᑐᓂᖅ ᐱᓕᕆᔨᖃᖅᑐᓂᖅ, ᑕᑯᔪᖓᓚᐅᖏᓪᓗᖅᑐᓂᖅ ᑕᑯᔪᖓᓚᐅᖏᓪᓗᖅᑐᓂᖅ ᐃᓕᓴᕆᔨᑎᖏᑦ ᐊᑐᖅᑕᐅᓚᐅᖏᓪᓗᖅᑐᓂᖅ, ᑕᐃᓵᓱᒪᓂᐅᓕᐅᖅᑐᖅ ᐃᓄᐃᑦ ᐃᓄᐃᑦ ᐊᑐᖅᑕᐅᓚᐅᖏᓪᓗᖅᑐᓂᖅ ᐃᓕᓴᕆᔨᑎᖏᑦ ᐊᑐᖅᑕᐅᓚᐅᖏᓪᓗᖅᑐᓂᖅ.

ᐊᐅᓪᓚᓇᖅᐳᖅᑕᕐᑐᑐᖅᐃᑦ, ᓄᓇᐃᑦ, ᐃᓕᐃᓪᓗ ᑕᐃᔪᓱᑐᖅᖏᓂ, ᖃᐅᐅᔨᓴᒪᓂᓪᓗ ᓱᓕᒥᖅ, ᐅᓕᔭᐃᓪᓗ ᐱᓕᕆᔨᒪᓂᕐᓂᖅᓂᖅ, ᑕᓕᑲᔪᐃᓐᓇᖅᑐᓂᖅ ᖃᐅᔨᓴᖅᑎᑦᓯᐊᕐᓇᖓᓕᒫᑦ ᐃᓄᐃᑦ ᐃᓄᐃᑦ ᐊᑐᖅᑕᐅᓚᐅᖏᓪᓗᖅᑐᓂᖅ ᐃᓕᓴᕆᔨᑎᖏᑦ ᐃᓄᐃᑦ ᐊᑐᖅᑕᐅᓚᐅᖏᓪᓗᖅᑐᓂᖅ, (ᓱᓕᓪᓗ ᓂᕐᑭᓂᖅ ᐱᑕᓴᐃᓂᕐᓂ, ᐃᓕᖁᓪᓗᓂᖅ ᓴᓇᓂᖅ), ᑕᐃᔨᓱᒪᓂᓂᒋᐃᓐᓇᓪᓗᓐᓄ ᐃᓄᐃᑦ ᐃᓄᐃᑦ ᐊᑐᖅᑕᐅᓚᐅᖏᓪᓗᖅᑐᓂᖅ ᓄᓇᕗᑦ ᑕᒪᖃᖅᑐᖅ ᑕᑯᔪᖓᓕᐅᓪᓗᐊᓂᖅ ᐊᐅᓪᓚᓇᖅᐳᖅᑕᕐᑐᑐᖅᐃᑦ ᐃᓄᐃᑦ ᓄᓇᒥ ᐊᑐᖅᑕᐅᓚᐅᖏᓪᓗᖅᑐᓂᖅᑐᑦ ᐃᓄᐃᑦ ᐃᓄᐃᑦ ᐊᑐᖅᑕᐅᓚᐅᖏᓪᓗᖅᑐᓂᖅᑐᑦ ᐃᓕᓴᕆᔨᑎᖏᑦ ᐃᓄᐃᑦ ᐊᑐᖅᑕᐅᓚᐅᖏᓪᓗᖅᑐᓂᖅᑐᑦ, ᐃᓕᓐᓂᑦ, ᐃᓅᓕᐅᑎᓂᕐᓂᓐᓄᑦ ᐱᓪᓚᑦ, ᐃᓕᓴᕆᔨᑎᖏᑦ ᐃᓄᐃᑦ ᐊᑐᖅᑕᐅᓚᐅᖏᓪᓗᖅᑐᓂᖅ ᐃᓕᓴᕆᔨᑎᖏᑦ ᐃᓄᐃᑦ ᐊᑐᖅᑕᐅᓚᐅᖏᓪᓗᖅᑐᓂᖅ ᐃᓕᓴᕆᔨᑎᖏᑦ ᐃᓄᐃᑦ ᐊᑐᖅᑕᐅᓚᐅᖏᓪᓗᖅᑐᓂᖅ.

3.1 ᐃᓕᓴᕆᔨᑎᖏᑦ ᓄᓇᕗᑦ ᐱᓴᕆᑎᐃᓕᓂᑦ ᐊᑐᖅᑕᐅᐳᓚᐅᔪᓂᖅᓂᑦ ᑐᑭᖃᓴᓂᖅᓂᓪᓗᓐᓄ ᖃᐅᔨᓴᖅᑕᐅᔪᓂᖅᓯᒪᔭᑦ

ᐃᓄᐃᑦ ᓄᓇᕗᑦ ᐊᑐᖅᑕᐅᓚᐅᖏᓪᓗᖅᑐᓂᖅ ᐊᒻᒪᓗ ᓄᓇᕗᒻᒥᐃᓚᖅᑕᐅᓚᐅᖏᑕᐅᑐᖅᑐᖅᓂᖅᑐᖅ ᐱᑕᓚᐊᖓᔪᓚᐅᖏᓪᓗᖅᓯᒪᔭᑦ (ᐃᓄᑕᓚᔨᐅᑎᖅᑲᓇᕐᓂᕐ-INAC, 1976) ᐅᓂᖅᑲᑕᐅᓚᐅᖏᓯᒪᔭᑦ, ᓄᓇᖓᔪᐊᑕᐅᑕᐅᓚᐅᖏᓂᑕᑎᓪᓗ ᑕᐃᔨᓯᒪᓂᐊᓗᖅᓂᖅ ᐱᓚᕐᓚᐊᓂᐃᓚᖅᓂᖅᓂᖅ ᐃᓄᐃᑦ ᓄᓇᖅᑕᓚᓚᐅᔨᓂᖅᓂᕐᓂᓐᓄ ᑕᒪᐃᓂᓂᖅᑐᖅ ᖃᖅᑐᓵᓂᑐᖅ ᓄᓇᕗᑦ ᓄᓇᕗᑦ (ᑕᓚᐅᔨᒪᓗ ᓱᑯᒪᓗ). ᑕᐃᖅᑯᐊ ᓄᓇᕗᒻᒥ ᓄᓇᖖᔪᐊᑕᐊᖖᔪᓯᓚᓂᑦ (Riewe, 1992) ᓴᓇᓯᒪᑎᐊᖅᖃᓂᖅᑐᑐᖅᑐᖅ ᑕᑯᔪᖓᑎᑎᑕᓚᐅᖏᑐᑦ ᐊᑐᖅᑕᐅᓯᒪᓂᖅᓵᖖᔪᓂᕐᓂᓂᖅᑐᖅ ᓄᓇᓂᖅ, ᐃᓄᖕᓄᑦ ᑎᑭᑕᐅᖃᖅᑕᐅᓚᐅᖏᑕᐅᑐᖅᑐᖅᓂᖅᑐᖅ ᐊᖏᔪᒡᒐᒪᖅ, ᐊᑕᐅᓯᔪᒥ ᓄᓇᕗᓕᒥᐅᑕᐅᔪᐅᓚᐅᖏᓂᓂᕐᓂᓂᖅ ᓯᓚ, ᓱᓕᓪᓗ ᐄᓇᑎᑐᑦ, ᑕᐃᖅᑯᓂᖅᓚᓪ ᐊᓯᔾᔨᓂᖅ ᓄᓇᖅᑐᓂᖅ ᓄᓇᓂᖅ ᑎᑭᓯᒪᔭᖅᓴᓕᐅᕐᓇᓂᖅᓂᕐᓂᓂᖅᑐᖅ ᓄᓇᖖᔪᓚᐅᖏᓪᓗᖅᑐᓂᖅᑐᖅ ᐃᓅᔨᔪᑎᑎᖅᑲᓇᕐᓂᕐᓂᓐᓄᑦ ᐱᓕᕆᔨᒪᓂᕐᓂᖅᑐᖅ ᓄᓇᕗᑦ ᐃᓕᓴᕆᔨᑎᖏᑦ ᐊᑐᖅᑕᐅᓚᐅᖏᓪᓗᖅᑐᓂᖅᑐᖅ ᐊᐅᓪᓚᓇᖅᐳᖅᑕᕐᑐᑐᖅᐃᑦ ᐃᓄᐃᑦ ᐃᓄᐃᑦ ᐊᑐᖅᑕᐅᓚᐅᖏᓪᓗᖅᑐᓂᖅᑐᖅ ᐃᓕᓴᕆᔨᑎᖏᑦ ᐃᓄᐃᑦ ᐊᑐᖅᑕᐅᓚᐅᖏᓪᓗᖅᑐᓂᖅᑐᖅ.

ᑐᓴᓚᕆᔨᑦ ᒪᓕᑎᖅᑕᑎᑕᐅᑉᐳᑦ ᐊᒋᒍᑦ *ᑎᑎᕀᖅᓯᒪᓂᖅᑎᓂᒃ 6,* ᐃᓄᐃᑦ ᐃᑲᔪᐊᕆᓴᖅᓂᓄᑦ ᐊᓴᑭᑐᐱᑎᒍᑦ ᑎᑎᕀᖅᓯᒪᔪᕀᓴᐅᕀᖅᓯᒪᑦ ᖃᐅᔨᓪᓚᕆᐊᕐᑐᖅᓴᓂᒃ ᓄᓇᒥᐅᑕᓚᑲᐅᔪᓂᒃ ᐱᓚᓗᐅᔪᓂᒃ ᐃᓄᓐᓄᑦ ᑎᓪᓗᐊᖅᑎᖅᓯᐃᐅᓕᒥᑦ ᐊᐅᒪᓗ ᐅᒪᔪᖅᓴᐃᐅᔪᓂ ᓄᓇᑭᓕᒥ. ᑕᒪᑲᐊ ᐱᑕᖅᑎᑲᑦᑐᓂᓕᑦ ᖃᐅᔨᓪᓚᑲᐅᓂᐊᖅᑕᒥ ᐃᑲᔪᖅᓴᐃᓂᒍᑦ ᐊᐅᑕᑕᐅᓂᕐᓂᓄᑦ ᐊᑐᓂ ᓴᓐᓂᐊᖅᑕᑕᐅᔪᓴᐃᓐᓇᓕᒃ ᓇᓂᑐᐃᓇᖅ ᓄᓇᑭᓕᒥ, ᐃᑲᔪᑕᐅᓗᑎᓗ ᑐᑭᓯᐅᓚᑲᐅᑭᓇᖅᑐᓂᓄᑦ, ᐊᒪᓚᓗ ᑎᑎᖅᑎᓯᑎᑕᐅᓂᐊᖅᑐᓂᒃ ᐱᑕᖅᑎᓂᐊᖅᑎᒥ ᖃᐅᔨᓪᓚᑲᐅᓂᐊᖅᑐᓂᒃ, ᐃᑕᖅᑯᔪᑐᖅᑎᖅᑎᒍᑦ ᐱᓚᒪᓚᐅᓂᖃᖅᑐᓂᒃ ᐃᓄᓐᓄᑦ, ᐃᑲᔪᑕᐅᓗᑎᓗ ᐱᓕᕆᑕᐅᔪᓐᓇᓕᓐᓂᒍᑦ, ᐊᑐᖅᑕᐅᔪᓐᓇᓂᐅᑕᓐᓄᑦ ᐃᑕᑕᒪᔭᐅᓗᑎᓗ ᐃᓄᐃᑦ ᐅᖃᖅᐳᐊᒍᑦ ᑕᐃᒍᖅᑕᐅᓚᐅᖅᑐᓂᓚᖅᐅᓂᖅ ᐊᑐᖅᑕᐅᓗᑎᓗ ᑕᒫᓗ ᓴᓐᓂᐊᖅᑕᑕᐅᔪᓴᐃᓐᓇᓕᒃ. ᐃᓕᓴᐃᓂᖅ ᐊᐅᑕᑕᐅᓂᕐᓂᓄᑦ ᐃᓕᓴᕐᓂᖅᑐᓂᒃ ᐊᑐᖅᑕᐅᓗᑎᓗ ᐃᓄᐃᑦ ᐊᐅᑕᑕᐅᓂᕐᓂᓄᑦ ᐃᓕᓴᕐᓂᖅᑐᓂᒃ ᐃᓄᐃᑦ ᐅᖃᖅᐳᐊᒍᑦ ᑕᑯᔪᓐᓇᖅᑐᑦ ᐃᓄᐃᑦ ᐊᐅᑕᑕᐅᓂᕐᓂᓄᑦ ᐃᓕᓴᕐᓂᖅᑐᓂᒃ ᐃᓄᐃᑦ ᐊᐅᑕᑕᐅᓗᑎᓗ ᐃᓄᐃᑦ ᐊᐅᑕᑕᐅᓂᕐᓂᓄᑦ ᐃᓕᓴᕐᓂᖅᑐᓂᒃ

ᐅᑭᐅᓚᓴᖏᓂᓂ ᑕᒪᓂ 1902-1903 ᐊᐪᓂᐊᖅᓂᐊᓗᖏᓪᓗᑦ ᖃᓄᒪᓂᐊᓗᖏᓪᓗᑦ ᑎᑭᑕᐅᓕᓚᖅᑐᑦ, ᑐᖅᔪᖅᑐᐊᔪᓐᓗᑎᒃ ᑕᒪᖅᑎᓂᐊᖅᓯᑕᐃᓇᓂᐅᖅᑐᑦᓯᒃᑕᒪᐅᖅᑐᑦ. ᐊᖅᐳᓂᐊᖅᑐᑦᓯᐊᖅᑏᑦ ᐊᐪᓂᐊᕈᑎᓂᒃ ᑎᑭᑕᐅᓕᓚᖅᑏᑦ ᐃᓇᓂᓂ ᐃᓕᓴᐅᔪᓂᒃ ᑎᑭᑕᐅᓕᓚᖅᑏᑦ ᐃᓇᓂᓂ ᑎᑭᑎᑎᒋᒪᐅᖅᑐᑦ ᓇᐅᔨᖅᑎᐅᓂᒃ, ᐅᒃᑯᓱᓕᖅᑎᐅᓂᓗ ᑕᒪᓂ 1908-ᒥ ᓴᖅᒪᓄᑦ, ᓴᓇᔨᐅᑎᑕᐅᒋᐊᖅᓯᑎᒃ.


1924-ᖁᑎᖅᑐᓗᒍ, ᐄᐊᑦᓴᖅᐋ ᑲᒡᕕᓂᒃᑯᑦ ᓂᐅᕐᓄᓕᖅᑎᒃᑯᑦ ᓄᓇᕗᑦ ᐅᓂᑎᒋᒪᓕᕐᓂᒃ ᐊᒃᕐᑐᖅᓴᐊᖅᑎᑦᓯᐅᒋᐊᖅᑐᑦ ᓴᖅᒪᓄᑦ ᐃᓄᐃᑦ (ᖃᑭᑭᖅᑕᓪᓕᖅᑎᑦ [Baffin-ᒥᐅᑦ] ᓂᒋᐊᓂᒥᐅᑕᓂᒃ) ᓄᒃᑎᖅᑕᐅᓕᐅᖅᓯᒪᖅᒪᑕ ᓂᐅᕐᓄᖅᑎᒃᑯᑦᓂᓗ. ᑎᓚᖏᓂᐊᖅᑲᑦᓂᖅ ᐱᒻᒪᓚᐊᑐᓕᓕᐅᖅᑐᖅ ᑭᓇᐅᔨᓕᐅᑕᐅᓴᐅᓇᖅᓂᔪᑦ, ᓂᐅᕐᓄᐅᑎᖅᓴᓄᑦ ᐄᐊᑦᓴᖅᐋᒃᑯᑦ ᓂᐅᕐᓄᖅᑎᒃᑯᖏᓂᒃ (ᑕᒥᒃ, ᖃᖅᓴᓂᒃ, ᐊᓯᖏᓂᓂᔪᓐ). 1940ᑦ ᐊᑦᕿᔪᐃᑦ ᐃᓯᖅᐸᓯᖏᓪᓗ, ᐊᒥᐊᓕᒃᒥᐅᑦ ᒥᑦᑕᖁᐅᓕᐅᓕᓕᐅᖅᑐᑦ ᓯᓚᑎᑦᑎᐊᖏᓪᓗ ᓴᖅᑐᐅᑦ. ᑭᖑᓂᐊᔪᑦ ᑲᓇᑕᐅᑦ, ᓴᓕᒪᒃᑯᓂᓇᑦ ᐃᐅᓚᑎᓕᕐᓂᒃ ᐊᐅᓚᑎᓕᕐᓂᒃᑯᑦ ᑕᒪᒃᑯᓂᖅᓯᖏᓪᓗ ᓴᖅᑐᐅᑦ, ᐊᐪᓂᐊᕈᑎᓂᒃ, ᑭᓇᐅᔨᓕᐅᑕᐅᓴᐅᓇᖅᓂᔪᑦ ᐃᓇᓂᓂ ᐊᐪᓂᐊᕈᑎᓂᒃ ᐃᓇᓂᓂ ᐃᓕᓴᐅᔪ

(ᑕძᔪᐢᓴᖅ 6). ᐃᓄᖅᑲᖅᐸᓪᓕᐅᖅᑐᑦ ᓄᓇᐃᑦ ᓄᓇᖓᔪᐊᖅᓂᖅᑎᑕᐅᓕᖅᑐᑦ ᐃᑲᖅᑐᐊᖅᑎᑦ ᐊᒥᓱᓕᓴᐃᑦ ᐅᒪᔨᖅᓂᒃ ᐅᓪᓗᖅᓯᓕᕆᓂᖅ ᐊᓪᓚᖅᑎᖓᓂᖅ ᐅᕐᖓᓂᒃᓂᑦ ᑐᐱᖅᓯᖓᓂᖅ ᑕᐅᖓᓂ ᑎᕐᒥᐊᖅᑲᖅᓯᐅᔨᓄᓇᖕᖓᓂᒃ. (ᑐᑭᓯᒥᐊᖅᓯᐅᓱᒪᔨᖅ: ᐅᐃᓕᓴᒃ ᑭᐊᒡᒃ-William Kemp, ᑐᖓᓕᕆᔨᑦ ᐃᓕᓴᖅᔨᑐᖅᑲᖅᑎᒍ ᐱᖅᔨᑎᓕᓂᖅᑲᖅᓂᖕᖓᓂᒃ ᑐᑭᓯᒥᐊᖅᑎᖓᓪᓚᑦ, ᑕძᔪᐢᓴᖅ 6).

ᐱᒡᒪᓚᐊᓗᖕᒃ ᐃᓕᓴᖅᔨᑐᑕᖅᖓᓂᒃᓂᒃ ᐊᑐᖅᑕᕐᕋᓂᖕᖓᓂ ᓇᓗᓇᖓᓂᑐᑦ ᐊᒥᓱᓕᕿᐅᓂᖕᖓᓂᒃ ᐅᓪᓗᖅᓯᓕᕆᓂᖕᖓᓂᒃᓂᓪᓗ

ᐊᐃᖅᑐᖏᓪᓗᐊᓂᖅ ᑕᑯᑎᑎᑦᓴᓪᓕᐊᓂ 5: ᐃᓄᐃᑦ ᐃᓕᖏᑯᓯᑐᖏᖅᓯᖅᑐᓂᖅ ᑕᐃᔪᓯᑦ ᓄᓇᖅ ᐃᒪᖏᓕᓄᖅᓂ ᑎᖅᒥᐊᖏᖃᖅᒥ (ᐋᖏᑭᒥᐊᖏᑕᐅᑎᐊᖏᖃᓄᖅᑐᖏᑐᑦ ᐃᓄᐃᑦ ᐱᐅᓯᑐᖏᖅᓯᖅᑐᓂᖅᑐᑦ <<ᑎᑎᔨᖏᑯᑦ ᑎᑎᖏᖃᑯᑎᖅᓯᓄᖅᑐᖅ)


ᖏᖅᑕᓄᓇᑎᑐᑦ ᑎᑎᖏᑎᔪᑦ	ᐃᓄᖅᑎᑐᑦ ᑎᑎᖏᑎᔪᑦᑦ	ᓯᓇᐅᓂᖅᓱ/ ᖏᓄᐃᑕᖅᓱ-ᓂᖅᓱᓕᓄᖅᑐᑦ	ᐃᑕᑕᒪᔨᐅᓯᒪᓕᑕᖏᑐᑎ ᑦ ᑕᐃᔨᐅᓂᖅᑭᑦ	ᐅᓂᖅᑲᓚᐊᖅᖃᓂᓕᕆᑎᑦ
Qimiarjulairut	ᖃᑭᒋᐊᖅᓪᓚᐃᕈᑦ	ᖃᑭᒋᐊᑦᓂᖅ		ᐃᓯᐊᓂᑦᑐᖅ ᑐᐊᕐᓴᖅᑭᓇᐅᕐᓂ ᖏᖅᖃᓴᖅᓱᓂᓂᖅ
Ujarajjuat	ᐅᔨᕆᓴᐊᑦ	ᓄᐅᖅ	ᐃᑲᑦᑐᐊᕐ ᓄᐅᐊ	ᐅᔨᕆᐃᑦ, ᐅᖅᑯᓯᖅᓴᔨᐃᑦ ᖏᑲᑯᓂᖅᖅᓴᐃᑦ
Ikkattuaq	ᐃᖅᑲᑦᑐᐊᖅ	ᓯᓗᔨᖅᕕᓯᐅᓂᖅᓱ	ᐃᑎᓴᐊᖏᓴᖅᒥᒃ ᐅᔨᕆᑯᓴᐊᖏᒍᑦ	ᐃᑲᵒᓂᐅᓴᑦ
Ikkattuap kuunga	ᐃᖅᑲᑦᑐᐊᕐ ᑯᖅᓱ	ᑯᖅ	ᐊᑎᑎᖅᓂᖅᓱᓂ (ᓇᖅᖃᖏᖅᓱᓂ) ᐃᑲᑦᑐᐊᕐ ᐃᑕᑕᒪᔨᐅᓕᓂᓂ ᑕᐃᔨᐅᓴᖅ ᑕᓯᐊᓕᖅᒥᒃ	ᑕᐃᔨᐅᓴᖅ ᑕᐃᒧᒃ ᑎᓂᖅᓱᓂᖅᓱᓕᓂ ᑎᑭᐅᒪᓂᖅᓱᓕᓂ ᑯᖅᑕᖏᓂᖅᓱ ᑕᐅᐅᖅᓱ ᑕᓪᐅᐅᕐ ᕿᖅᓱᓕᓂ
Mamitsuittuq	ᒪᒥᑦᓯᐃᑦᑐᖅ	ᑯᖅᔪᓂᖅ		ᒧᓴᓇᐅᓴᖅ ᑕᐃᒪᖅᓱᑦ ᒪᒥᑐᓇᖅᑐᖅᖃᖅᑐᖅ ᓯᑯᓴᐃᑐᑎᒥ. ᐃᖏᓕᑐᒪᓪᓚᓕᓕᒃ ᑕᒧᓂ.
Itijuarjuk	ᐃᑎᓴᐊᖏᓴᒃ	ᖏᓄᖏᓕᖅᖏᓄᖅ		ᑦᖅᑐᓴᖅ ᖏᓄᖏᓕᖅᖏᓄᖅ. ᑕᑯᒃᖏᐅᑎᐊᖅᑐᒥ ᓇᓕᓇᐃᒃᑯᑕᓪᓚᐅᕐᓴᒃᑐᖅ ᐅᖅᓱᓯᑐᑐᒥᒃ.
Nunattaunngi	ᓄᓇᑦᑕᐅᖅᖏᑎ	ᑕᖅᒧᒪᖏᐅᖏᓄᑕᖅᓱ-ᑐᖅ		ᓇᓴᓱᒃᓴᑎᑎᐊᕐᓇᖅᖏᑦᑐᑐᖅ ᑕᖅᒪᖏᓂᐅᓂᖅᓱᒪᑦ.



ⵆⵔⵔⵙⵜⵣⵖⵍⵅⵔⵍⵡⵛ: ⵆᵃᵚⵍⵍⵍᵇⵚⵛ. 2014



ⵆⵔⵔⵙⵜⵣⵖⵍⵅⵔⵍⵡⵛ: ⵐⵙⵍⵛ ⵍⵐᵚⵐⵛ ⵣⵆⵛᵍⵇⵔⵅⵛᵍⵜᵚⵜᵇ ⵑⵔⵖⵣⵖᚴⵔⵅᵍⵆⵛ. ⵍⵐⵔⵖⵑⵆⵛᵍᵇⵚᵚⵐⵛ 1976



ⵆⵔⵔⵙⵜⵣⵖⵍⵅⵔⵍⵡⵛ: 1992 ⵐⵙⵓⵍⵍ ⵐⵙⵚⵣⵖⵔⵅᵍⵇⵛ. ⵖⵅ-Riewe 1992

ⵛⵚⵅᵇⵑⵍᵍᵇ 6: ⵛⵍⵍᚔⵑⵔᵇ ⵍⵐⵍⵛ ⵣⵆⵛᵍⵇⵛⵛⵜⵚⵔⵛ ⵐⵙⵍᵐᵇ ⵑⵛⵔⵅᵎ ᵛⵔⵔᵍᵇⵔᵓⵓⵣⵔᵇ.

3.3 ᐃᓕᓐᓇᐅᑎᖅ ᐃᓄᐃᑦ ᐊᑐᖏᑦᑕᕐᓂᖅ ᓄᓇᒥᒃ

ᐅᑦᓗᒥᐅᓕᖅᑐᖅ, ᓴᑦᓕᕐᒥᐅᑦ ᓯᓕ ᓂᕆᑭᒃᓴᓂᒃᓯᒪᔪᖅ ᐊᒥᕐᓂᓴᓗ, ᕿᑭᙵᓂᓴᓗ ᐅᒪᔨᓂᒃ, ᐃᓄᔨᐱᑎᒃᓴᖅᑭᓐᓄᐃᓂ ᐊᑐᖅᑐᐃᓗᒃ. ᐃᓄᔨᐱᑎᒃᓴᓂᒃ ᐊᖓᔪᓇᓯᖏᑕᕐᓂᓂᖅ ᐅᒪᔨᖏᖅᖃᙱᐅᓂᖅᑐᓂᒃ ᒪᓕᒃᐸᒃᑐᑦ ᐱᓇᓯᐊᖅᐸᙵᓂᖅᑐᓂᒃ ᓴᓂᓕᕐᓯᓂ ᕿᑭᙵᓂᓴᓗ ᐅᒪᔨᓂᒃ ᐃᓕᓴᕐᓯᓂᒃ ᓴᓂᓕᕐᓯᓂᒃ, ᑲᖏᖅᓯᐅᑦᑕᕐᓂᕐᒥᒃ, ᐊᖓᔪᓇᓯᕐᐊᖏᑕᕐᓂᕐᒥᔪᓗ ᓄᓇᒥᐅᑕᓂᒃ ᐊᒻᒪᓗ ᑕᕆᐅᑦᒥᐅᑕᓂᒃ ᐅᒪᔨᓂᒃ, ᓯᑦᓗ ᓇᓄᑦᓂᒃ, ᑐᐃᑐᓂᒃ, ᓇᑎᑎᓂᒃ, ᓴᕿᓇᓚᖅᓂᒃ ᐊᒻᒪᓗ ᐊᐃᐃᖅᓂᒃ. (). ᓴᓂᓕᕐᒥᐅᓂᓕ ᐊᐅᓪᓚᕆᔨᐃᓂᒃᓴᖅᓴᐅᕐᓂ ᑕᐅᕐᓂᒃ ᐃᓚᖓᔾᔪᑎᑦ ᐃᑲᑕᑐᐊᑦ ᑎᓐᒥᐊᖏᖅᓴᐃᓚᓄᑦ, ᑲᖏᖅᓯᐅᑦᑕᕐᓂᕐᒥᑦ ᒪᓂᓕᕆᐊᖅᓂᕐᒪᑦᓯᐅᓂᒃ ᐱᓕᕐᑕᐅᓂᒃ ᐅᒪᔨᓂᒃ, ᓯᑦᓗ ᐊᓂᒃᑲᓐᓂᕐᒪᑦ ᐃᓕᓴᕐᓯᓂᒃ ᐊᑐᖅᑐᐃᓗᒃ ᐱᓕᕐᑕᐅᓂᒃ ᐅᒪᔨᓂᒃ ᐊᐃᐃᖅᓂᒃ ᐊᒻᒪᓗ ᐊᐃᐃᖅᓂᒃ. ᐃᓕᓴᕐᓯᓂᒃ ᐊᑐᖅᑐᐃᓗᒃ ᐊᑐᖅᑐᐃᓗᒃ ᐊᑐᖅᑐᐃᓗ

4 ᑐᖏᓚᐅᑉᐳᓪᓕ ᐊᓚᒪᓗ ᐱᓴᖏᑕᑉᐳᓚᓪᓕ

4.1 ᑕᑉᑐᑎᓚᐅᑉᐳᓪᓕ

ᑕᑉᑐᑎᕐᑲᖏᑐᓪ ᑎᖕᒥᐊᖏᑲᕐᓇᖕᒥ ᐃᓚᐃᖅᓭᓕᖕᒥᓚᓪᓕᐅᓂᒃ:

ᐃᑲᑐᑐᐊᖏᒥ ᑎᖕᒥᐊᖏᑲᕐᓇᒃ ᐊᑉᐳᓕᑕᑉᑉᐅᖏᖅ ᐊᑯᓂ ᓯᑉᐅᓂᑲᓚᒍᑐ ᐱᑕᖏᑲᐃᖅᓭᐅᓕᑦᓴᒥᑕ ᐊᓚᒪᓗ ᓴᓗᐅᖅᓭᑉᐳᓯᓂᐅᓐᓴᒥᑕ ᓄᓇᓕᓕᒫᓂᖅ, ᐃᓚᐃᓪᓗ, ᐅᓚᔪᓕᒻᒧᓗ, ᐃᓄᖕᓄᑦᓗ ᐱᔪᖅᓭᐅᑎᖏᑲᖕᒥᖅᓄᑦ ᐊᓚᒪᓗ ᖃᑭᐃᖅᓇᐅᑉᐳᓯᓂᐅᓐᓴᒥᑕ, ᐊᑐᖏᑕᑦᓗᑎᒃ ᐃᓄᐃᑦ ᖃᒃᐅᓱᒪᕐᑐᖏᖅᓂᒃ ᑕᒪᐃᖅᓂ ᐃᓯᒪᒃᓴᖏᑐᐊᑎᓕᑐᖏᖅᓂᒃ.

ᑕᒪᒃᑯᐊ ᐊᑯᓂ ᓯᑉᐅᓂᑲᓚᒍᑐ ᑕᑉᑐᑎᓚᐅᑉᐳᓪᓕ ᐊᑐᓕᑎᓂᕐᓗᑦ ᐃᒃᑕᑐᐊᖏᑲᕐᓇᖕᒥᒃ ᐱᖓᓱᓯᐃᓕᑦᖃᒃᖓᓚᐅᖅᓭᓕᖕᒥᒃ ᐃᓯᒪᓂᐃᓯᑐᐊᖏᑲᖕᒥᖅᓄᑦᓗᑎᒃ ᐊᑐᖏᑕᑦᓗᑎᒃᑐᓂᒃ ᐊᑐᖏᑕᑦᓗᑎᒃᑐᓂᒃ ᐃᓯᒪᓂᐃᓯᑐᐊᖏᑲᖕᒥᖅᓄᑦᓗᑎᒃ ᐊᑐᖏᑕᑦᓗᑎᒃᑐᓂᒃ.

- ᓴᓐᓂᐊᖏᑲᑕᐳᓗᑎᒃ, ᐅᓚᔪᐃᑦ ᐅᓚᔪᖏᑲᖓᐊᓂᐅᓇᓂᓂᓐᓗ ᓄᓇᐃᑦ, ᓴᓐᓂᐊᖏᑲᑕᐳᓗᑎᓴᓗ ᑎᖕᒥᐊᑦ ᐊᓚᒪᓗ ᐅᓚᔪᐃᑦ ᐊᒥᓱᓂᕐᑐᓗᐊᑦᖃᖕᒥᑦ;
- ᓴᓐᓂᐊᖏᑲᑕᐳᓗᑎᒃ ᐃᓄᖏᑲᖓᐊᐅᒡᒪᐳᖕᒥᑐᑦᓴᖏᑲᐃᑦ ᓄᓇᓕᓕᒫᓂᖅ ᐊᓚᒪᓗ ᐃᑕᖕᒥᑯᓱᑐᖏᑲᖕᒥᔪᑦ ᐊᑐᖕᒥᑕᐳᐅᖅᑐᑦᓴᖏᑲᐃᑦ (ᐊᑐᖏᓗᑎᒃ *ᑎᑎᕐᖕᒥᓱᒪᓂᕐᓂᑐᓂ 2-ᒥ – ᐃᓄᐃᑦ ᐃᑲᔪᓱᐊᖏᓴᕐᓂᓄ ᐊᓂᕐᑭᑋᕐᔅ*);
- ᐃᓄᐃᑦ ᐱᔪᓇᐅᑎᓂᕐᓂᓂᒃ ᐊᑐᑭᓇᓗᑎᒃ ᐊᓚᒪᓗ ᐊᓯᐅᓇᓱᖏᑕᑭᓇᓗᑎᒃ, ᐊᓚᒪᓗ ᐅᔨᕕᖏᑐᓗᑎᒃ ᐃᓄᐃᑦ ᐃᑲᔪᖕᒥᑕᐳᓭᒡᒪᖏᑕᓇᓗᑎᒃᑐᓂᒃ (ᐊᑐᖏᓗᑎᒃ *ᑎᑎᕐᖕᒥᓱᒪᓂᕐᓂᑐᓂ 2-ᒥ – ᐃᓄᐃᑦ ᐃᑲᔪᓱᐊᖏᓴᕐᓂᓄ ᐊᓂᕐᑭᑋᕐᔅ*);
- ᐃᓕᑕᒪᓚᐅᓗᑎᒃ ᐃᓄᐃᑦ ᖃᒃᐅᓱᒪᕐᑐᖏᖅᓂᒃ ᐱᓯᒪᓚᐅᑉᐳᓪᓕᖕᒥᒃᑐᓂᒃ ᐱᕕᖏᑲᐃᖅᓭᐅᓕᑦᓴᒥᑕ, ᐊᑐᖏᑕᑎᓂᕐᓗᑦ, ᖃᒃᐅᓱᒪᕐᑐᖏᖅᓂᒃᑐᓂᒃ ᐃᓄᐃᑦ ᐃᑲᔪᖕᒥᑕᐳᐅᖅᑐᑦᓴᖏᑲᐃᑦ (ᐊᑐᖏᓗᑎᒃ *ᑎᑎᕐᖕᒥᓱᒪᓂᕐᓂᑐᓂ 2-ᒥ – ᐃᓄᐃᑦ ᐃᑲᔪᓱᐊᖏᓴᕐᓂᓄ ᐊᓂᕐᑭᑋᕐᔅ*);
- ᐃᓕᑕᒪᓚᐅᓗᑎᒃ ᐃᓄᐃᑦ ᖃᒃᐅᓱᒪᕐᑐᖏᖅᓂᒃ ᐱᓯᒪᓚᐅᑉᐳᓪᓕᖕᒥᒃ ᐱ

ᐊᓚᒪᓗ (ii) ᖃᓄᐃᓕᓕᐅᕈᐊᓇᓐᓂᖓᑦ ᐊᐅᓕᕆᓂᖅᑎᑦ ᐊᒃᓱᕈᑎᕈᖃᖅᑎᑦᐊᕐᕆᖅᑐᑦᓂᒃ ᐊᓚᒪᓗ ᐅᓗᓚᐊᓇᕈᑎᐊᓇᓇᖅᑐᑦᓂᒃ ᑎᓯᒥᐊᖃᖅᖃᙱᓐᓂᑦ (ᐃᓗᓕᓯᑎᑦ ᐊᙱᖅᑐᑦᖃᓱᒪᓕᓯᑎᓇᓂᓕᓂᓂᒃ 5.0 – ᐊᐅᓕᕆᑎᐅᐅᓕᓯᑎᓇᓂᓇᓂᓂᓂ

	ᖃᐅᔨᓴᖅᓯᕐᒥᐅᑦ, ᓇᐅᑦᑎᖅᑐᕈᑎᓂᐅᑉ ᐊᓕᒪᓗ ᖃᐅᔨᓚᑎᑕᐅᓂᖅᑳᖃᐅᔨᖅᑕᕐᓗᑎᑦ ᐱᓕᕆᐊᐅᔪᓇᖅᓯᒪᓂᖅ.
	ᐱᓴᓱᖅᑕᐅᔪᒃᑳᖅ 2.3: ᓄᓇᓕᖅᓂ ᐃᓄᐃᑦ ᖃᐅᔨᓚᓂᖅᑳᐅᖃᕐᓂᐅᕐᒪᑕ ᐱᓕᕆᐊᐅᔪᓇᖅᓯᒪᓂᖅᑐ ᑎᖏᒥᐊᖃᖅᑲᖏᒥ.
ᑐᕐᖏᓚᓴᖅ 3: ᐃᓄᐃᑦ ᖃᓄᑐᐃᓇᖅ ᐱᓕᕆᔨᖅᓯᒪᓂᓄᑦ ᐊᐅᑐᖅᑕᐅᓗᐊᓱᓄᒃᑳᐃᑦ ᐱᐅᔪᑐᖃᐅᔨᓂ ᐊᓕᒪᓗ ᐃᓕᓴᓄᓯᖃᖅᓴᖏᓐᓇᓂ ᐱᓕᕆᐅᓂᖅᑎᑦ ᓄᓇᓕᖅᓂ	ᐱᓴᓱᖅᑕᐅᔪᒃᑳᖅ 3.1: ᐅᓴᓇᖅᑐᓗᑎᓄ ᑕᒪᐃᓂᓄ ᐃᓄᐃᑦ ᒪᓕᕐᑎᐊᖃᑕᕆᐊᖃᖅᓂᖅᑎᓂ ᐱᔪᓇᐅᑎᖃᖅᓚᐊᖃᖅᓂᕐᒥᑦ ᐊᖏᐅᐳᒪᑎᑕᐅᔪᓂᖅ ᒪᓕᕐᓂᓂᖅ. ᐱᓴᓱᖅᑕᐅᔪᒃᑳᖅ 3.2: ᐊᑐᓕᖅᑎᑎᐅᓕᕐᓕᐊᓂᖅ ᐱᐅᓂᖅᕿᓂᖅ ᐱᐅᓯᕆᓴᐅᔪᓇᖅᑐᓂᓄᑦ ᐊᐅᑐᖅᑕᐅᓗᐊᓱᓄᒃᑳᐃᑦ, ᓯᕐᓂᖅᑎᖅᑕᐅᑭᓄᒃᑕᐅᔪᓂᓄᑦᓴᓐᓂᓕᕐᓂ ᐱᐅᓯᑐᖃᖅᓚᖅᐅᔪᓂᖅᑐᓂ, ᐃᓕᓴᓄᓯᖃᖅᓴᖏᓐᓇᓕᓐᓂᓗ ᐊᓕᒪᓗ ᐃᓕᓴᓄᓯᖃᖅᓴᖏᓐᓇᓕᓐᓂᓗᓂᓗ ᐃᓕᓴᓄᓯᖃᖅᓴᖏᓐᓇᓕᓐᓂᓗᓂᓗ ᐃᓄᐃᑦ ᐱᓕᕆᐅᓂᖅᑎᑦ	ᐱᓴᓱᖅᑕᐅᔪᒃᑳᖅ 3.3: ᐃᓕᐅᑦᖃᑕᕐᓗᑎᓄ ᐊᐃᓴᑐᖅᓯᒪᓂᖅᑎᓂ ᐱᓕᕆᐊᖏᓪᓕᕐᓕᐊᓂᖅᑐᓄ, ᐱᑎᐊᖅᑎᑕᐅᓯᒪᔪᓂᓕᕐᓂ ᐃᑲᔪᖅᓯᑕᐅᓂᕐᒪᑦ ᑎᖏᒥᐊᖃᖅᑲᖏᒥᑦ ᓴᐳᓂᓂᐊᖅᖢᖅ ᐊᓕᒪᓗ ᐊᓐᓇᐊᓂᓗᓂᖅ ᓄᓇᕐᓂᐅᓂᖅᓂᖅ.

4.3 ᖃᐅᔨᑎᐊᖃᖅᑕᕐᓂᖅ

ᑲᓇᑕᒥ ᐅᒪᔪᓕᕆᓚᐅᑲᓯ ᐊᐊᖏᔪᑕᒧᐊᖅ ᓇᐅᔨᑎᖅᑐᓚᐊᖅᑐᖃᖅᑕᖅᓴᓕ ᒪᓕᐢᓯᓗᑎᐅ ᐊᑐᐃᓇᐅᔨᔪᓇᖅᓯᒪᓂᖅᑎᓂᐅᑉ ᑭᓇᐅᔨᓂᖅ ᐊᐅᒪᓚᐅᔨᖅᑕᐅᓂᖅᑐᓂᓗ ᐊᐅᓚᑦᑎᐊᖅᑐᓂᐅᑉ ᖃᐅᔨᓴᓂᐊᖅᑐᓂᓗ ᐊᐅᓚᑦᑎᐊᖅᑐᓂᐅᑉᓗ ᐊᐅᓚᑦᑎᐊᖅᑐᓂᐅᑉᓗᓂᓗ ᐃᓄᐃᑦ ᐱᓕᕆᐅᓂᖅᑎᑦ 10 ᐊᖏᔪᐃᑦ ᓇᓱᖅᓱᓚᑕ. ᖃᐅᔨᑎᐊᖃᖅᑕᐅᖃᖅᑐᖃᖅᑕᕐᓗᑎᓄᑦ ᑎᑎᕋᖅᓯᒪᓂᖅᑐᓂ ᖃᐅᔨᓴᓂᐊᖅᑐᖃᖅᑕᐅᓚᑎᓄ ᐊᐅᓚᑦᑎᐊᖅᑐᓂᖅᑎᑦ ᐱᓕᕆᐊᖏᓪᓕᕐᓕᐊᓂᖅᑐᓄ ᑎᑎᕋᖅᓯᒪᓂᖅᑎᑦ, ᖃᐅᔨᓴᐅᓯᒪᔪᓂᖅᓴᓗ ᓇᐅᔨᑎᖅᑐᓚᐊᖅᑐᖃᖅᑕᕐᓗᑎᓂᐅᓕᕐᓂᖅᑐᓂᓄᑦ ᐊᓕᒪᓗ ᖃᐅᔨᓴᖅᑎᖏᓪᓕᕐᓂᖅᑐᓂᓄᑦ. ᑲᓇᑕᒥ ᐅᒪᔪᓕᕆᓚᐅᑲᓯ ᐊᓕᒪᓗ ᐊᐅᓚᑦᑎᐊᖅᑐᖃᖅᑕᐅᓚᑎᓄᑦ ᑕᐃᑲᖏᓕ ᓴᐅᓂᐊᖅᑐᖃᖅᑕᐅᓚᑎᓄᑦ ᐃᓕᓴᓄᓯᖃᖅᓴᖏᓐᓇᓕᓐᓂᓗᓂᓗ ᐊᐅᔪᓂᐊᖅᑐᖃᖅᑕᐅᓚᑎᓄᑦ ᑕᒪᐃᓇᓐᓂᓕ ᐱᓕᕆᐊᖏᓪᓕᕐᓕᐊᓂᖅᑐᓄ ᐊᐅᓚᑦᑎᐊᖅᑐᓂᖅᑎᑦ.

5 ᐊᐅᓪᓚᕐᓂᕆᖅ ᐃᓗᒪᒃᓴᖅᓗᐅᕙᑎᒃᓴᐃᓪᓕᕐᓂ

ᐃᓐᓇᐅᔾᔨᖅ, ᐱᓕᒪᐅᔾᔨᓂᒃ ᑐᐊᕕᖅᓯᒋᐊᓂᒃᑐᓐᓇᓐᓂ ᐅᓗᓚᐊᓇᖅᑐᑎᖅᑐᕐᓂᖅ ᐃᑲᓂᑐᐊᑉ ᑎᖏᒥᐊᖅᕕᖅᖃᓕᕐᓗᓂ. ᓄᓇᕗᑎᖏᓪᓗᑦ ᑕᐅᖓᓂ ᐊᐅᑎᒋᐅᓂᕗᓂᓗᓪᓚᓕᒪᓕᒫᓕᒐᓗᑦ, ᑎᖏᒥᐊᖅᕕᖅᖃᐃᐅᓂᖅᓱᓗ ᐃᓄᖅᕕᕐᔨᐃᑦᑐᑦᓗᓂᓂ. ᑭᓚᐊᓂ ᓚᓂ ᒪᑯᐊ ᐊᐅᑕᓪᓂᕕᑦ ᐃᓚᒪᒃᓴᖅᓗᐅᕙᑎᒃᓴᐃᓪᓕᕐᓗᐅᕐᖓᑦ, ᐊᐅᑐᖅᓴᕐᓄᓇᕗᑦ ᑎᖏᒥᐊᖅᕕᖅᖃᖏᓐᓂᖅ.

ᑲᖏᐃᑦ ᐊᒥᓚᓂᕐᓂ ᓄᓇᐃᓪᓗᓗ ᓇᔨᖅᐸᐅᑕᓂᕐᓂ

ᑲᓇᑕᒥ ᐅᒪᔨᑦᓯᓗᓴᐅᑯᑦ ᑎᖏᒥᐊᖅᕕᖅᖃᓕᕆᐅᓕᕐᖅᓯᒪᔨᑦ ᑕᒪᓂ 1959-ᒥ, ᐊᒥᓚᓗᐊᖅᐸᓕᐅᕐᓂᕐᑎᓪᓗᓕᓪᓕᓕᓂ ᑕᐃᔨᓚᒪᓂ ᑎᖏᒥᐊᖅᕕᖅᖃᐃᐅᕐᓕᕆᐅᕐᖅᑐᑦ ᓄᓇᕗᓂ, ᑲᓇᑕᐅᐸ ᐅᑭᐅᖅᑕᖅᑐᑉᓂᓕᓂ, ᑕᐃᔨᓚᒪᓂᓗ ᐱᑕᖅᑲᓕᐅᖅᓯᓂᒃ ᑲᖏᐃᑦ ᖃᑲᓂᕐᓂᓐᓂ 28,000 (Cooch & Barry, 1957). ᑲᖏᐃᑦ ᑎᖏᒥᐊᖅᕕᖅᐸᑦ ᓯᓚᑕᓂᓯᓯᒋᓕᖅᑐᑦ, ᑕᐅᖓᓂ ᖃᓄᖅᓱᓂᓐᓂᒥ (Ell Bay-ᒥ (Kerbes, et al., 2014). ᐃᓐᓇᐃᖏᓪᓗᖅᐸᒥ ᖃᑲᐅᔨᓂᖅᑕᐅᑎᒃ ᐊᒥᓚᓂᕐᓂ ᐊᖅᑎᒍᒥ 2014-ᒥ, ᓇᓚᐅᑦᖅᑕᐅᓚᐅᕙᖅᑐᑦ ᐱᑕᖅᑎᓚᑎᐅᓂᖅ 689,700 ᐅᓪᓗᖅᖃᖅᑐᑦ ᑲᖏᐃᑦ ᐃᑲᑐᑦ ᑯᓐᓄᓂ (ᐅ. ᓪᓚᓗᐊᖅ-J. Leafloor, ᑲᓇᑕᒥ ᐅᒪᔨᓚᓇᐅᓪ, ᑐᐃᐅᒪᖅᑕᐅᑎᓚᓐᓇᕐᓂ). ᓄᓇᕗᐊᓚᒋᒪᒥ ᐃᓚᒪᑕᓪᓗᑕᓂ, ᑕᒪᐃᑯᐊ ᑲᖏᐃᓗᓗᐃᑦ ᐊᒥᓚᑉᕕᐊᖅᓚᒪᓕᕐᓂ 50-ᓂᐅ ᐊᓂᔪᖅᑐᓂ ᐊᖅᑎᒍᓂᒃ, ᒪᓐᓇᐅᓚᖅᑐᖅᓗ “ᐊᒥᓴᓚᓗᐊᓚᖅᓯᑎᐅᒃ” ᑲᓇᑕᒥᓗ ᐊᒥᐊᑲᓚᑲᒥᐅᕐᓗ ᓄᓇᕗᑦᓂᓚᓚᕗᑦ. ᓄᓐᓗᓗᐊᑕ ᐊᑯᓐᓂᖅᓱᓂᓐᓄᓚᖅᑐᑦ ᑲᖏᐃᑦ, ᑕᐃᐅᑯᐊᓗ ᑲᖏᐃᑦ ᐅᓗᖅᑲᓚᑦ ᓴᓪᓚᐅᐸᑦ ᖁᑭᖅᑕᓐᓗᓂᒃ, ᐊᒥᓚᓂᖅᑎᖅᑎᓚᖅᑕᐅᑎᒃ 12.6 ᒥᓚᔪᓐᓂᒃ ᐱᕈᐊᓂᓚᓚᓂᒃ ᑲᖏᐃᑦ (ᐊᓚᓴᐅᓴᑦ, ᐱᓚᓇᖅᑎᓐᓂᓪᓗ-Alisauskas, et al., 2018a), ᑕᐃᐅᑯᐊᓗ ᐊᓚᓐᓂᑦ ᓂᓪᓚᖅᓗᐃᑦ (Ross-ᖁᓚᑦ) ᐊᒥᓚᓂᖅᑐᓚᖅᑐᑦ ᑕᒧᓂᒃ 1.6 ᒥᓚᔪᓂᒃ ᐱᕈᐊᓂᓚᓚᓂᒃ ᑲᖏᐃᑦ ᑲᑎᒪᓂᖅ, ᐊᖅᑎᔪᓕᐅᖅᑐᓂᒃ 2006-ᒥᒃ 2015-ᔪᑦ (ᐊᓚᓴᐅᓴᑦ, ᐱᓚᓇᖅᑎᓐᓂᓪᓗ-Alisauskas, et al., 2018b).

ᐊᒥᓚᓗᐊᓚᓂᓂᓐᓂᑦ ᐃᓚᒪᓗᑕᐅᐅᓚᖅᑐᑦ ᐊᐅᑐᖅᑕᐅᔨᓂᓐᓂᑐᓄᓂ ᐅᑭᐅᖅᑕᖅᑐᑎ ᓚᓴᖅᐸᓚᐅᓂᓐᓂᑦ, ᐊᑎᑎᐅᔪᓐᓂᑦᓗ ᓄᓇᐃᓂ ᐅᒪᔨᓴᖅᓴᖅᑭᐅᓂᓐᓂᑦ, ᓂᕆᑭᓴᖅᓯᐅᑦᓴᖅᑭᑎᓐᓂᑐᓄᓂ ᐊᒥᓚᓗᐊᖅᑐᓄᓂ ᑲᖏᐃᑦ ᓂᓪᓚᓐᓇᓚᓂᓗ (Batt, 1997). ᓴᓪᓚᐅᑦ ᖃᑭᖅᑕᒧᐊᓂ ᖃᐅᔨᓴᖅᑕᐅᓚᐅᓚᓂᓐᓂᑐᓄᓂ ᐅᓂᐅᑦᓚᑦ ᐅᖅᑲᐅᓯᖅᓴᓕᑦ ᑲᖏᐃᑦ ᓂᓪᓚᕆᓴᖅᑕᖅᑐᓄᓂ ᐊᑎᑎᐅᔪᑎᐅᓂᓐᓂᑐᓄᓂ ᓄᓇᓂ ᐃᑲᑦᑐᑦ ᖁᑦᓱᓂᓂ (ᑲᑐ, ᐱᓚᓇᖅᑎᓐᓂᓪᓗ-Carter, et al., 2018; Fᐋᓐᑲᓐ ᐊᓗᒪᓗ ᒪᓗᓐᓇ-Fontaine and Mallory, 2011). ᓄᓇᐃᓂ ᐊᓯᔨᖅᓯᒪᓕᖅᓴᓂ ᑕᐃᑲᑦᑎᑐᑕᓗ ᐊᔨᔨᖅᓂᖅᓂᖅᑕᖅᓯᑎᒃ ᕿᐊᑎᓪᓚᐃᑦ ᑕᓯᐅᔾᔨᐊᖏᓪᑐᑦ ᓄᓇᐃᓂ ᓯᓚᖅᐸᓯᓯᓐᓂᑎᓂᓂᒃ ᐊᑎᑎᐅᔪᑎᐅᓂᓐᓂᑐᓄᓂ ᓄᓇᓕᒫᓂᒃ (ᑲᑐ, ᐱᓚᓇᖅᑎᓐᓂᓪᓗ-Jano, et al., 1998; ᑯᑕᓂᐊᓐ ᐊᓗᒪᓗ ᑕᐋᐅᓚᓴᑦ-Kotanen and Jefferies, 1997). ᐊᐅᑎᓚᓇᔨᔨᐊᖅᑯᑦ ᒧᐃᐅᔾᔨᖅᐸᓚᐊᓚᖅᑐᑦ ᑎᖏᒥᐊᖅᕕᖅᖃᐃᐅᕐᓚᖅᑐᑦ ᓯᕆᖅᑎᖅᑕᐅᐅᓕᓚᐊᓂᓐᓂᑐᓄᓂ ᐊᒥᓴᓗᓚᖅᑐᓄᓂ ᑲᖏᐃᑦ ᓂᓪᓚᕆᓴᖅᓯᐅᓂᓐᓂᑐᓄᓂ, ᓂᓚᐃᐅᓗᐊᓚᖅᑕᐅᓂᓐᓂᑐᓄᓂ, ᑐᑭᓯᐅᔾᔨᓚᓂᑎᖅ ᖁᓄᖅᑲᖅᑕᐅᔾᔨᓂᓐᓂᑐᓄᓂ ᐊᖅᑐᒃᑐᓂ ᑲᖏᐃᑦ ᑎᖏᒥᐊᖅᕕᖅᖃᐃᐅᓂᓐᓂᑐᓄᓂ, ᐅᓪᓗᕆᐊᖃᓕᕆᓂᑦ ᐃᓚᒪᓕᕆᓂᑦ ᐊᑎᑎᐅᔪᑎ

ᓂᓪᓕᐅᕐ ᐊᓂᔪᓇᖅᓯᕐᓚᐅᖅᓴᓗ

ᓂᓪᓕᐅᕐ ᐊᓂᔪᓇᖅᓯᕐᓚᐅᖅᓴᓗᓄᑦ, ᐃᖅᑭᔭᐃᖅᓗᐅᑎᓗᐊᖅᓗ, ᐊᓂᖅᑎᓄᓪᓗ ᐱᓵᕆᐅᖅᑭᓵᓄᑦ, ᑕᑯᐅᐅᓪᓕᓚᐊᖅᑎᕆᓕᖅᑐᑦ ᐊᓂᔪᓇᖅᓂᕐᒥᓂᖅᑎᑦ ᓄᓇᐃᐊᓄ ᐅᓪᓚᕐᓇᖅᑐᖅᐳᖅᑐᑦ ᐃᑲᑎᑦᑎᓇᖅᑐᖅ. ᐊᓂᔪᓇᖅᓯᕐᓚᐊᓵᑦ ᑕᓗᐅᖅᑭᑦ ᐊᓪᓚᓗ ᑕᓂᓵᓂᖅᑐᓂ, ᓄᓇᐅᕐᑦ ᐃᓗᐊᔪᑦ ᖃᑯᐊᖓᔭᖅᑐᓂᖅᑐᓂ, ᐊᐱᖅᑲᑕᖅᓯᓂᖅᑐᓂ ᐊᓪᓚᓗ ᓂᓪᓕᓚᓗ ᑕᓂᓵᓂᖅᑐᓂ ᖃᐅᔨᓴᖅᑕᐅᖅᑲᑕᖅᓂᒪᓕᒃᑲᒃ ᓇᓂᑐᐃᓇᖅ ᐅᑭᐅᖅᑕᖅᑐᑎᓪᓗ ᐅᓪᓚᕐᓇᖅᑎᓂᐅᖅᓴᓂᖅᑐᓂ, ᖃᐅᓇᐃᓕᓂᓵᓂᖅᑐᓂᓗ ᐅᓪᓚᕐᓇᐃᓕ ᑕᓪᒥᑯᓇᓂᖅᑎᓂᖅᑐᖅᓂᖅᑐᑦ. (CAFF, 2010). ᓂᐅᓗ ᐅᐸᑐᑎᓗᔪᒍ, ᑐᓴᐃᔪᖅᓴᓕᓂᓂᖅᓴᐊᐅᖅᑕᓕᓂᖅᑎᓕᓗᔪ ᕿᐊᑦᓴᓵᐲᕐ ᑕᓚᐅᕆᓗᐊᓂ, ᓇᓇᐃᓕ ᖃᑭᓕᒥᐅᓂᖅᓴᐊᐅᖅᑎᓂᕈᓕᒪᑕ (Lunn, et al., 2016) ᖃᐅᔨᓴᐅᖅᑎᓂᖅᓯᓕᓓᕐᓂ ᑎᑎᕋᑕᐅᔨᓕᓓᑎᓗᒃ ᐊᑐᖅᑕᐅᓗᑎᒃ, ᐊᒥᓯᓂᖅᓴᐊᓴᓕᓕᓂᓂᖅᑐᓂ ᓇᓇᐃᓕᑦ ᑎᓪᓗᐊᑦ ᐅᑦᓗᖅᑐᓵᓂᒃ ᓂᕆᖃᖅᓴᓂᖅᓴᐊᐅᖅᑕᓕᓂᖅᑐᓂ ᓴᑦᑕᐅᑦ ᖃᐅᖅᑕᕐᓗᐊᓂ ᐊᓪᓚᓗ ᐊᒃᓯᑐᓴᕐᕕᐊᑎᒃ (Smith, et al., 2010).

ᐊᓯᖏᓐᓂᒃᑕᐅᖅᑲ ᖃᐅᔨᓴᐅᓯᓚᖅᑐᓂ ᓄᓇᒥ ᐱᐅᓯᓗᓴᐅᓕᖅᑐᓗᓂᒃ ᑎᓪᓗᐊᖅᑎᓂᖅᓴᓐᓂᒥ, ᓯᓴᓗ ᓴᓂᓗᐊᖅᑎᕆᕐᓂᖏᓐᓂᒃᓂᓂᒃ ᓄᓇᐃᑦ, ᐃᒃᑲᒃᓯᓕᒡᓕᐊᓂᖏᓐᓂᒃ ᐊᓪᓚᓗ ᓄᓇᐃᑦ ᐃᓕᖏᑦ ᓯᕐᓴᖅᑎᖅᓯᓚᓕᓂᖏᓐᓂᒃᓂᒃ. ᑕᓚᒃᑯᐊ ᐅᓚᔨᑦᑎᓂᖅᓴᓕᓕᓂᖅᑐᓂ ᐊᓯᖏᓐᓂᖅᓯᓚᕐᓇᖏᓐᓂᖅᑐᓂ ᐊᑐᖅᑕᐅᓗᑎᓂᒃ ᓄᓇᐃᖅᑲᓇᖅᑐᖅᓂᒃᓂᒃ ᐊᑐᖅᑕᐅᓂᖅ ᐅᑭᐅᖅᑕᖅᑐᑐᒥ (Smol & Douglas, 2007; Kokelj, et al., 2005) ᐊᓪᓚᓗ ᑕᐃᓚᐃᓕᓓᕐᓂᓕᕆᓗᑎᒃ ᐱᓵᕆᐅᐅᓂᒍᑦ ᓯᓚᐅᑦ ᐊᓯᖏᓐᓂᖅᓴᓐᓂᕆᓗᐊᖅᑎᓂᖅᓴᓐᓂᒃ. ᓴᓐᓂᕐᕿᐅᑦᑕᐃᑦᑕᐅᖅᑲ ᑕᑯᓴᖅᑕᑦᑕᖅᑲᓂᖅᓴᐊᐅᖅᑕᓕᓂᖅᑐᓂ ᓄᓇᐃᖅᑲᓇᖅᑐᖅᓂᒃᓂᒃ ᐊᐅᓚᑦᑎᓂᖅᑐᓂ ᐊᑐᖅᑕᐅᓗᑎᓂᒃ ᐃᓕᓴᐃᑦ, ᓇᒍᖏᐅᖅᓴᖅᑎᑦᓂᖅᑐᓂᓗᓂᒃ ᐃᓕᓴᐃᑦ. (Carter, et al., 2018).

ᐃᓄᐃᑦ ᓄᓇᓕᑦ ᐃᓕᐅᖅᑲᑦᓂᖅᑎᑦ ᖃᐅᔨᓴᒪᑎᑕᐅᖅᓯᖏᓐᓄᓪᓗ

ᓴᓐᓂᕐᕿᐅᑦ ᐃᓱᒪᓕᐅᑎᑎᕐᒥᓂᒃ ᐅᖅᓴᐅᓱᖅᓴᖅᓱᒪᓴᑦ ᐱᔨᔪᑎᖅᓴᖅᑐᓂᒃ ᑎᖏᑦᓯᑦ ᐊᑐᖅᑕᐅᖅᓴᑕᖅᑐᓂᖅᑐᓂᒃ ᖃᐅᔨᓴᐅᓴᖅᑐᒃ ᐊᓪᓚᓗ ᐃᓄᐃᑦ ᖃᐅᔨᓴᐅᓴᒪᓕᐊᑐᖅᓴᖅᑐᓂᒃ ᐊᑐᖅᑕᐅᓗᑎᓂᒃ, ᑕᒪᐃᓂᒃ ᖃᐅᔨᓴᐅᓴᖅᑎᑕᖅᑐᓂᒃ ᐊᑐᖅᑕᐅᓗᑎᓂᒃ, ᑕᒪᐃᓄᑦ ᖃᐅᔨᓴᐅᓴᖅᑎᑕᖅᑐᓂᒃ ᐊᑐᖅᑕᐅᓗᑎᓂᒃ ᓇᓗᓇᐃᖅᑐᓂᒃ ᐃᓕᓴᐃᑦ, ᓇᒍᖏᐅᖅᓴᖅᑎᑦᓂᖅᑐᓂᓗᓂᒃ ᐃᓕᓴᐃᑦ. (Carter, et al., 2018).

ᐃᓄᐃᑦ ᑭᓇᐅᔪᑐᐃᓇᐃᑦ ᐱᓕᕆᓂᖅᑐᓂ

ᓄᓇᒥ ᐅᓴᖃᖅᑎᐅᑕᐅᓂᒃ ᐅᓴᖃᖅᓂᐊᖅᑕᑭᓇᖅᓯᓂᖅᑐᖅ ᐃᓕᓴᕆᓴᐅᓕᖅᑎᓕᕆᓚᐅᓕᖅᑐᖅ ᓇᓂᑐᐃᓇᖅ ᑲᑎᕐᓗᒍ ᐊᑐᖅᑕᐅᓗᑎᓂᒃ ᐅᓴᖃᖅᓂᐊᖅᑕᑭᓇᖅᓯᓂᖅᑐᓂᒃ ᓄᓇᒥ ᐊᑐᖅᑕᐅᓗᑎᓂᒃ ᐃᓄᐃᑦ ᐊᑐᖅᑕᐅᓗᑎᓂᒃ ᐊᑐᖅᑕᐅᓂᖅᓴᐅᓂᖅᑐᓂᒃ ᑕᒪᐃᓄᑦ ᐊᑐᖅᑕᐅᓗᑎᓂᒃ ᐊᑐᖅᑕᐅᓂᖅᓴᐅᓂᖅᑐᓂᒃ

ᖃᓂᖅᐸᐅᓪᓕᒻᒥᐊᑐᓴᖅᑎᑦᓯᐊᖅᑐᑦᓴᓗᒃᑐᖅ ᑐᑭᓯᓴᐅᑐᐊᕈᑎᖅ ᖃᓄᖅᑎᑐᖅ ᑭᓇᐅᔪᓕᐅᕈᑕᐅᔪᓂᖅᓯᓂ ᓄᓇᒥᐅᑕᐃᑦ ᑕᒪᖅᑯᐊ. ᐅᔪᕿᐃᑦ ᖃᖅᑯᓂᖅᓴᐃᑦ ᓴᓇᐯᐊᐃᖅᓴᔪᐃᑦ ᐱᒻᒪᓕᐊᓄᖅᑎᖅ ᐊᑐᑎᖃᖅᖢᖅᑐᑦ ᔪᓄᖅᖃᓂᐅᔪᓂᖅ ᐊᐃᑦᑐᐃᓄᑕᐅᖅᑕᑭᓇᓂᖅᓯᓕᓂ ᐅᔪᕿᖏᓂᑐᓂᖅ, ᐅᔪᕿᖏᓂᐊᖏᓂᖅ ᓯᖅᑕᓂᑎᓚᖏᓂᖅ, ᑕᒪᖅᑯᐊ ᐱᑕᖅᑭᑕᐅᓕᕆᓂᖅᓯᓂᖅᖃᓂᖅ ᑭᓪᓕᖅᑯᓂᖅᓴᐃᑦ ᓄᓇᖃᖅᓯᓂᖅ.

ᖃᖅᓂᒃᖃᓕᖅᑕᓂᖅᑐᑦ 43 ᐅᒥᐊᖏᐊᑦ ᐃᐅᖅᖃᑕᖅᑐᑦ ᓂᒋᐊᑕ ᑕᓚᐅᕆᓪᓗᐊᔪᑦ ᓴᑦᓂᑕᒃᑲᓐᓂᖅ, ᑭᓪᓕᖅᑯᓂᖅᓴᐃᑦ ᐅᓯᓪᖅᑐᑦ, ᐃᓕᓴᐅᑎᖅᑐᑦ ᐅᖅᑭᐅᖅᑐᑦ ᐅᓯᓪᖅᑐᑦ ᐃᓄᐃᑦᑐᓂᖅ, ᑕᒪᖅᑯᐊ ᐅᒥᐊᖏᐊᑦ ᐅᓯᓪᖅᑐᑦ ᐃᓄᐃᑦᑐᓂᖅ, ᐅᔪᕿᖏᓂᐊᖏᓂᖅᓴᓐᓂᑦ. ᐅᓗᓚᐊᓇᖅᑐᖅᑎᒃᑕ ᑯᐃᓯᐃᐅᔪᖅᑎᖅᑭᓇᓂᖅ ᐅᖅᓯᐊᓗᖅᑐᖅ ᖃᖅᓂᒃᖃᓕᖅᑕᓂᖅᑐᑦ ᐃᐅᖅᑐᑦ ᐃᓕᐃᑦ ᐃᐅᖅᑐᑦᑎᐊᖅ ᐱᔪᓯᐅᓯᓕᐅᐅᖅᓴᖅ ᐃᓕᓴᐅᑎᖅᑐᑦ ᐃᐅᖅᑐᑦᑎᐊᖅ (ᒪᑦᓕᖅᑕᓂᓂᖅᑎᓕᓂ) ᐊᐅᓪᓚᕆᓂᖅᑎᓕᓂ.

ᓴᑦᓕᖅᑎᐅᑕᐃᑕᐅᖅᓴᖅ ᐅᖅᖃᐅᓯᖅᑯᖅᓯᒪᔪᑦ ᓴᓇᓴᐅᖅᑯᑦᔨᓂᖅᓴᖅ ᕿᐊᓵᑕᖅᑯᑦ ᐊᖅᑯᑎᐅᖅᓴᒥᑐᖅ, ᑕᐅᐅᖅᓴᒃ ᑎᖏᒥᐊᖅᑲᖅᓴᓐᓇᒪᑦ. ᐊᖅᑯᑎᐅᖅᓴᒥᑐᖅᓴᖅ ᐊᑦᑭᐅᐅᒪᓴᓇᖅᑐᖅᓴᖅ ᐃᐅᖅᓴᐃᑦᐊᐸ ᑯᖅᓴᒃ, ᑎᑭᐅᐅᒪᓗᓂ ᐃᓕᓯᐊᓂᑦ ᐊᐃᐃᓚᐅᑦ ᐊᖅᑯᑎᐅᖅᓴᒥᑐᖅᓴᖅ ᐃᐅᖅᓴᐃᑦᐊᐸ. ᐊᖅᑯᑎᐅᖅᑕᑎᐊᖅᓴᖅᑲᐅᓕᕆᑎᑦ ᑎᖏᒥᐊᖅᑲᖅᓴᓐᓇᒪᑦ, ᐊᐅᓱᓂᖅᑲᐅᓕᕆᓂᖅᓴᖅ ᐃᓕᓴᐅᓕᕆᓂ ᐊᐅᓕᒪᓗ ᐳᓕᖅᑕᐅᓕᕆᒪᖅ ᐱᓕᓚᔪᓂᖅᓴᖅ, ᓂᐅᐃᓂᖅᑎᓕᓂ ᐃᐅᖅᓴᐃᑦᐊᐸ ᐊᐅᓚᐃᓕᓴᐅᑎᖅᑐᑦ ᐊᑦᑭᐅᐅᒪᓴᓇᖅᑐᖅᓴᖅ ᐃᐅᖅᓴᐃᑦᐊᐸ. ᐊᖅᑯᑎᐅᖅᑕᑎᐊᖅᓴᖅᑲᐅᓕᕆᑎᑦ ᑎᖏᒥᐊᖅᑲᖅᓴᓐᓇᒪᑦ, ᐊᐅᓵᐸᓂᖅᑲᐅᓕᕆᓂᖅᓴᖅ ᐃᓕᓴᐅᓕᕆᓂ ᐊᐅᓕᒪᓗ ᐳᓕᖅᑕᐅᓕᕆᒪᖅ ᐱᓕᓚᔪᓂᖅᓴᖅ ᐃᓕᓴᐅᓕᕆᓂᑦ ᓂᐅᐃᓂᖅᑎᓕᓂ ᐊᐅᓚᐃᓕᓴᐅᑎᖅᑐᑦ ᐱᓕᓚᔪᓂᖅᓴᖅ ᐃᓕᓴᐅᓕᕆᓂ ᐃᓕᓴᐅᑎᖅᑐᑦ ᐊᐅᓚᐃᓕᓴᐅᑎᖅᑐᑦ ᐱᓕᓚᔪᓂᖅᓴᖅ ᐃᓕᓴᐅᓕᕆᓂᑦ ᑎᖏᒥᐊᖅᑲᖅᓴᓐᓇᒪᑦ, ᐊᐅᓵᐸᓂᖅᑲᐅᓕᕆᓂᖅᓴᖅ ᐃᓕᓴᐅᓕᕆᓂ ᐊᐅᓕᒪᓗ ᐳᓕᖅᑕᐅᓕᕆᒪᖅ ᐱᓕᓚᔪᓂᖅᓴᖅ ᐃᓕᓴᐅᓕᕆᓂᑦ ᐊᐅᓚᐃᓕᓴᐅᑎᖅᑐᑦ ᐱᓕᓚᔪᓂᖅᓴᖅ ᐃᓕᓴᐅᓕᕆᓂ ᐃᓕᓴᐅᑎᖅᑐ

ᐊᕐᓇᒃᑐᖅᓯᒪᔪᓂᒃ ᑕᑯᑎᑎᑦᓴᓪᓕᐊᓂ 7: ᓇᐃᔅᓴᖅᑕᐅᓯᒪᔪᓂ ᐊᐅᓪᓚᑎᓂᖅᐳᓐᓂᒃ ᐃᓱᒪᒃᑲᖅᐅᓯᕈᑎᒃᑲᐃᑦᓗ ᐊᓚᒐ ᖃᓄᐃᓕᓴᐅᕈᑎᒃᑲᖅᓯᒪᓕᐅᑦ.

ᐊᐅᓪᓚᑎᓂᕆᓂᒃ ᐃᓱᒪᒃᑲᖅᓯᓯᐅᕈᑎᒃᑲᓕᑦ	ᒪᓕᒃᑐᑦ ᑐᕆᖃᕐᓯᒃᑐᐅᔪᓂᖅᐳᓐᓂᒃᓗ ᐊᐅᖃᕐᓯᒃᑐᑦ ᑕᑯᑎᑎᑦᓴᓪᓕᐊᓂᖅᐳᓐᓂᒃ 6-ᒥᑦᑐᓂᒃ	ᐊᐅᓪᓚᑎᓂᕐᒧᑦ ᖃᓄᐃᓕᓴᐅᕈᑎᒃᑲᓕᑦ
ᐱᓕᓐᓇᑎᒃ ᐊᓯᖏᓐᓇᒪᓂᖅ ᑎᖅᒥᐊᖅᑎᖃᐅᔾᔨᒪᓂᒃ ᑲᓇᑕᓕᒪᒥ ᐅᒪᓕᐊᖅᑎᖃᐅᓂᒧᑦ ᐅᖏᓐᑐᑕᐅᓐᓇᑎᒃ ᐱᐅᓂᑦᔪᓐᓇᖅᑐᒃ ᐊᐅᓕᑕᑎᖃᐅᔾᔨᓐᓇᖅᐳᑦ ᐊᓚᒐ ᐊᖅᓯᓐᓇᖅᑐᒃ ᐅᒪᖅᑐᒃ ᐅᒪᔨᓕᒪᓕᖅ, ᐱᓐᐊᖅᑐᖅᓴᓐᓇᓕᖅᑐᑦ.	1.1, 1.2, 1.3, 1.4	6.1 ᐱᐅᓯᓚᖅᒥᒧᑦ ᓄᓇᒥᐅᑕᓂᒃ ᐊᐅᓕᑕᑎᓂᖅ 6.2 ᐅᒪᔨᓕᓚᓂᖅᑐᒃ ᐊᓚᒐ ᐅᒪᔨᕐᑲᖅᖃᐅᔾᔨᓐᓇᖅᐳᓂᒃ ᐊᐅᓕᑕᑎᓂᖅ 6.3 ᓇᐅᑕᑎᖅᑐᐃᓚᖅᑐᕐᖃᖅᐅᖅᑲᕐᓂᕆᓂᖅᐳᓂ ᖃᓄᐅᔨᕐᖃᖅᐳᓂᒃᓂᓐᓂᒃ 6.4 ᐃᓄᖅᑯᑎᑭᓚᐅᔨᓂᒃ ᖃᓄᐅᔪᒪᑎᕈᑎᒃᑲᓂᖅᐳᓐᓂᒃᓴᓕᖅᑐᖅ ᐊᓚᒐ ᖃᓄᐅᓕᓚᐅᖅᓯᓂᒃ ᐊᐅᓕᑕᑎᐊᖅᓯᓂᖅ
ᑲᖏᖅᑐᐃᓐᓇᖅᑐᒃ ᐃᓕᓴᕐᓂᒃ ᐊᓯᖏᓐᓇᒪᓂᖅ ᓄᓇᓕᒫᓂᖅ	1.2	6.2 ᐅᒪᔨᓕᓚᓂᖅᑐᒃ ᐊᓚᒐ ᐅᒪᔨᕐᑲᖅᖃᐅᔾᔨᓐᓇᖅᐳᓂᒃ ᐊᐅᓕᑕᑎᓂᖅ 6.3 ᓇᐅᑕᑎᖅᑐᐃᓚᖅᑐᕐᖃᖅᐅᖅᑲᕐᓂᕆᓂᖅᐳᓂ ᖃᓄᐅᔨᕐᖃᖅᐳᓂᒃᓂᓐᓂᒃ
ᓄᓇᕗᑦ ᐃᓐᓇᐅᑎᓂᖅᐳᓂ ᓄᓇᕗᑦ ᐃᓐᓇᖅᑐᒃᑲᓂᕐᒥᔪᓐᓇᖅᑐᖅᓯᒪᓂᖅᐳᓂᒃ ᐊᓚᒐ ᐃᓐᓇᖅᑐᒃᑲᓂᕐᒥᓂᖅᐳᓂᒃ ᐅᑭᐅᖅᑕᐅᓂᖅᐳᓂᒃ	2.1	6.4 ᐃᓄᖅᑯᑎᑭᓚᐅᔨᓂᒃ ᖃᓄᐅᔪᒪᑎᕈᑎᒃᑲᓂᖅᐳᓐᓂᒃᓴᓕᖅᑐᖅ ᐊᓚᒐ ᖃᓄᐅᓕᓚᐅᖅᓯᓂᒃ ᐊᐅᓕᑕᑎᐊᖅᓯᓂᖅ
ᓄᓇᒥᐅᑕᓂᒃ ᐅᓴᕐᓂᓕᐊᑲᓇᖅᐳᑦᔪᓐᓇᖅᑐᕈᑎᒃᑲᓂᖅᐳᓂ ᐊᓚᒐ ᐱᓐᓇᐅᑎᐊᖅᑕᐅᔾᔨᓐᓇᖅᐳᓂ ᖃᐅᔨᓂᖅᑐᑦ ᐅᓴᕐᓇᐃᑦ ᑲᓇᖏᐊᑲᓴᑦ ᓄᓇᒥᐅᑦᑐᓂᖅᐳᓂ, ᐊᑭᐊᓂᑦᑐᑦ ᑎᖅᒥᐊᖅᑎᖃᐅᔨᓴᓕ	1.2, 1.3, 3.2, 3.3	6.3 ᓇᐅᑕᑎᖅᑐᐃᓚᖅᑐᕐᖃᖅᐅᖅᑲᕐᓂᕆᓂᖅᐳᓂ ᖃᓄᐅᔨᕐᖃᖅᐳᓂᒃᓂᓐᓂᒃ 6.4 ᐃᓄᖅᑯᑎᑭᓚᐅᔨᓂᒃ ᖃᓄᐅᔪᒪᑎᕈᑎᒃᑲᓂᖅᐳᓐᓂᒃᓴᓕᖅᑐᖅ ᐊᓚᒐ ᖃᓄᐅᓕᓚᐅᖅᓯᓂᒃ ᐊᐅᓕᑕᑎᐊᖅᓯᓂᖅ

ᐊᐅᓪᓚᕐᓂᕆᔨᖅ ᐃᓱᒪᓕᕆᔨᐅᑎᑦᓯᓕᕐᓂ	ᒪᓪᑐᑐᑦ ᑐᕆᓚᓴᐅᔪᓂᖅ/ᐱᓕᕆᖅᑕᐅᔪᒪᓂᖅᑭᖅᓂᓗ ᐊᓚᖅᑐᕆᖅᓯᒪᔪᓂ ᑕᔾᑎᑎᑦᑎᐊᑎᖅᓴᖅᑭᖅᑐᓂᖅ 6-ᒥᑦᑐᓂᖅ	ᐊᐅᓪᓚᕐᓂᕆᓕᒍᑦ ᕆᓇᐃᓪᓚᐅᑎᑦᓯᓕᕐᓂ
ᓴᓪᕎᒥᐅᑦ ᐃᓱᒪᓗᑎᕆᔨᓂᖅᑭᑦ ᐊᑐᖅᑕᐅᓂᖅᓴᐅᕆᑕᓚᐊᕆᔨᓂᖅᑭᓂᖅᓂᖅ ᐃᓇᐃᑦ ᕆᓇᐅᓴᓴᑐᕆᖅᑭᑦ ᑕᒪᖅᔾᔨᑦ ᕆᓇᐅᓴᕆᖅᑕᐅᔪᖅᑐᑐᖅᓂᖅ ᐊᓕᒪᓗ ᐊᐅᓪᓚᕐᓄᐅᕆᖅᑕᐅᔪᓂ ᐃᓱᒪᓪᓕᐅᕈᑕᐅᔪᓚᐅᓪᓂᓂᓂ	2.1	6.4 ᐃᓇᕆᔨᑎᑎᑭᐅᔪᓂᖅ ᕆᓇᐅᓴᒪᑎᑦᑎᓂᖅᓴᐅᑭᓚᐊᕆᔨᓂᖅᓂᖅ ᐊᓕᒪᓗ ᕆᓇᐅᒪᖅᐅᑭᓚᐊᐅᓪᓂᖅ ᐊᐅᓪᓚᕐᓄᐅᓂᖅ 6.3 ᓇᐅᑦᑎᖅᑐᐃᓴᖅᑐᕆᓚᐊᕆᔨᓂᖅᑎᑦᓯᓕᕐᓂᖅ ᕆᓇᐅᓴᓴᕆᓚᐊᓄᐅᓂᖅᓂᑦᓗ
ᓴᑦᓪᕎᒥᐅᑦ ᐃᓱᒪᓗᑎᕆᔨᖅᑐᑐᑦ ᐃᑲᔪᓱᐊᖅᓴᓂᖅᑕᐅᔪᐊᕆᔨᖅᑎᑦᓯᓕᕐᓂᓄᐅᓂᖅ ᐃᓪᓚᐅᓂᖅᓴᐅᑎᖅᑎᑦᑐᑕᖅᑐᓂᖅ ᑭᓇᐅᔾᔪᓪᓕᐅᔨᓇᐅᓴᖅᑎᑦᑕᖅᑐᓂᖅ ᐊᓱᓄᐅᓂᓗᑐ ᐱᓕᑦᓪᑕᐅᑦᔨᓇᐅᔨᐅᓴᖅᑐᑐᖅᓂᖅ ᐊᐅᓪᓚᕐᓂᕆᓂᓄᐅᓂᖅ ᑎᓕᒥᐊᕆᖅᓇᓕᒃ	2.2	6.4 ᐃᓇᕆᔨᑎᑎᑭᐅᔪᓂᖅ ᕆᓇᐅᓴᒪᑎᑦᑎᓂᖅᓴᐅᑭᓚᐊᕆᔨᓂᖅᓂᖅ ᐊᓕᒪᓗ ᕆᓇᐅᒪᖅᐅᑭᓚᐊᐅᓪᓂᖅ ᐊᐅᓪᓚᕐᓄᐅᓂᖅ
ᐊᖅᑐᖅᑕᐅᓴᒪᓂᕐᓂᑦ ᓴᓪᒪᐅᑦ ᐊᓴᓵᖅᑦᓪᓕᐊᓂᕐᓚᐅᓗᓄᑦ ᑎᓕᒥᐊᕆᔨᓇᓕᒥ ᐊᐅᓪᓚᕐᓂᕆᓕᒍᑦ	1.2	6.3 ᓇᐅᑦᑎᖅᑐᐃᓴᖅᑐᕆᓚᐊᕆᔨᓂᖅᑎᑦᓯᓕᕐᓂᖅᓂᑦ ᕆᓇᐅᓴᓴᕆᓚᐊᓄᐅᓂᖅᓂᑦᓗ
ᐅᒥᐊᕐᔪᐊᕆᔨᖅᓗᐊᓴᒍᓪᓕᕐᓂᖅ ᑭᓕᑦᓪᕎᒥᐅᓄᑦ ᓄᓇᕗᑦᓂᓇᓕᐅᓕᖅᑐᓂᖅ ᐊᓪᓚᐃᓗ ᐅᓗᑎᐊᓇᑦᓂᖅᑎᑦ ᐅᖅᓴᐊᓗᖅᒥᒃ ᑯᓚᓴᓚᐅᔪᓕᐅᓂᖅᑎᑦ ᐃᒪᐃᑦ	1.2, 3.2, 3.3	6.3 ᓇᐅᑦᑎᖅᑐᐃᓴᖅᑐᕆᓚᐊᕆᔨᓂᖅᑎᑦᓯᓕᕐᓂᖅᓂᑦ ᕆᓇᐅᓴᓴᕆᓚᐊᓄᐅᓂᖅᓂᑦᓗ 6.4 ᐃᓇᕆᔨᑎᑎᑭᐅᔪᓂᖅ ᕆᓇᐅᓴᒪᑎᑦᑎᓂᖅᓴᐅᑭᓚᐊᕆᔨᓂᖅᓂᖅ ᐊᓕᒪᓗ ᕆᓇᐅᒪᖅᐅᑭᓚᐊᐅᓪᓂᖅ ᐊᐅᓪᓚᕐᓄᐅᓂᖅ
ᐊᖅᑯᓂᕆᔨᓚᐅᖅᑎᑦᓯᓕᕐᓂᖅ ᕿᑎᖅᑦᓪᕎᒥᐅᑦ ᑎᓕᒥᐊᕆᔨᓇᓕᒥᓂᖅᓚᐅᓂᖅ	2.3, 3.1, 3.2, 3.3	6.4 ᐃᓇᕆᔨᑎᑎᑭᐅᔪᓂᖅ ᕆᓇᐅᓴᒪᑎᑦᑎᓂᖅᓴᐅᑭᓚᐊᕆᔨᓂᖅᓂᖅ ᐊᓕᒪᓗ ᕆᓇᐅᒪᖅᐅᑭᓚᐊᐅᓪᓂᖅ ᐊᐅᓪᓚᕐᓄᐅᓂᖅ

ᐊᐅᓪᓚᕐᓂᕆᔨᖅ ᐃᓗᒪᖏᖅᓯᐅᑕᑎᖏᓐᓄᑦ	ᒪᓪᒃᑐᑦ ᑐᕋᒪᓱᐅᔪᓂᒃ/ᐱᓴᓰᖅᑕᐅᔪᒪᓂᖅᑭᖅᓂᓗ ᐊᕐᓇᒃᑐᖏᓯᒪᔪᓂ ᑕᔾᑎᕋᑎᒃᔪᓚᑎᖏᓐᓄᑦᑭᖅᑐᓂᒃ 6-ᓯᑦᑐᓂᒃ	ᐊᐅᓪᓚᕐᓂᕆᔨᑦ ᖃᓄᐃᓪᓕᐅᑕᑎᖏᓐᓄᑦ
ᐊᒥᓱᓂᖏᓐᓄᐅᓪᓕᖅᓂᕐᓂᑦ ᐊᖓᔪᓱᕐᐊᑭᓱᓕᖅᑐᑦ ᐊᑭᓚᖅᑐᓂᒃ/ᐊᐅᓚᑭᓱᓕᖅᑐᑦ ᐊᖏᔪᑎᖏᕈᑎᐊᖏᓂᖏᓐᓄᐅᓪᓕᖏᓐᓇᖅᓯᑦ	2.3, 3.1, 3.2	6.4 ᐃᓄᖅᔪᑎᑎᓴᐅᔪᓂᒃ ᖃᐅᐅᓕᒫᑦᑎᓂᖏᓐᓄᐅᓴᓕᐊᖏᖅᖏᓂᖅ ᐊᓪᓚᓗ ᖃᐅᐅᒪᓴᐅᓴᓕᐊᓕᓐᓂᒃ ᐊᐅᓚᕐᓂᑎᖅᓂᖅ

6 ᐊᐅᓪᓗᕆᐅᔪᓂᖅ ᖃᓄᐃᓕᓚᐅᕈᑎᒃᓴᐃᓪ

ᑲᓇᑕᒥ ᐅᓪᓚᔨᓕᓴᖅᑎᓂ, ᐃᓕᑎᓪᔨᑐᓂ ᐃᖏᓂᐅᖏᓐᓄᑦ ᐊᐅᓪᓗᕆᐅᔪᐅᖏᓐᓄᖏᓪᓗᒍ ᑲᑎᓚᔨᖏᓪᓕᕆᓂ, ᐊᐅᓪᓗᕆᐅᔪᐅᓂᐊᖅᑐᑦ ᐃᑲᖏᑐᐊᖏᓐ ᑎᐅᒋᐊᖏᖃᖏᓐᓇᖅ ᐱᓇᔪᒍᑎᖃᖅᑐᑦᓂᖅ ᐊᐅᓪᓗᕆᑎᓂᖏᓪᓗᒍ ᖃᓄᐃᓕᓚᐅᕈᑎᒃᓴᐃᓪᓂᑲᓪᓚᔨᖏᑦᑐᓂᖅ ᒪᓕᒃᑐᑐᓂᖅ ᒪᓂᒃᑐᓂᖅ: ᐱᐅᔪᓇᓴᒥᒍᓪᓗ ᓄᓇᒥᐅᑕᓂᖅ ᐊᐅᓪᓗᕆᔨᐅᓂᖅᑐᓄᓪᓗ (ᑎᑎᕋᖅᔪᒪᔪᓂᖅ 6.1-ᒥ), ᐅᒪᔪᑦᓚᓂᖏᑦ ᐊᓪᓚᒍ ᐅᒪᔪᖅᖃᕐᓂᐊᕐᓂᖅᓂᖅ ᐊᐅᓪᓗᕆᑎᓂᕐᓂᖏᑦ (ᑎᑎᕋᖅᔪᒪᔪᓂᖅ 6.2-ᒥ), ᓇᐅᕆᑎᖅᑕᐃᓴᖅᑐᖃᖅᖃᕐᓂᐊᕐᓂᖅᓂᕐᓂ ᖃᐅᔨᓴᖃᕐᓂᐊᕐᓂᖅᓂᕐᓗ (ᑎᑎᕋᖅᔪᒪᔪᓂᖅ 6.3-ᒥ), ᐊᓪᓚᒍ ᐃᓄᖅᑎᑎᕐᓄᑦᓂᖅᓴᐅᔨᓂᖅ ᖃᐅᔨᓚᑎᕆᑎᓂᖅᓂᖅ ᐊᓪᓚᒍ ᖃᐅᔨᓴᓴᐅᔨᓇᓕᒫᓂᖅ ᐊᐅᓪᓗᕆᑎᓂᕐᓂᖏᑦ (ᑎᑎᕋᖅᔪᒪᔪᓂᖅ 6.4-ᒥ). ᖃᓄᐃᓕᓚᐅᕈᑎᒃᓴᐃᓪ ᑕᓚᖅᑯᐊ ᐃᑲᔪᑕᐅᓂᐊᕐᓂᖅ ᑕᐅᑐᓗᓇᖅᑐᓂᖅ, ᑐᕋᓚᓇᖅᐅᔪᓂᖅ ᐊᓪᓚᒍ ᐱᔨᓇᖅᑕᐅᔪᒪᓂᐊᕐᓂᖅᑐᓄᖅ ᑕᖅᑯᐊᓂᖅ ᐊᐅᓪᓗᕆᑎᓂᖏᓪᓗᒍ, ᐱᓕᒪᒃᖁᖅᓯᐅᑎᑐᓂᖅᓂᖅᑐᓄᖅ ᐃᓯᒪᓂᖅᓯᐅᑎᑐᓂᖅᓂᖅ ᐊᐅᓪᓗᕆᑎᓂᖏᓪᓗᒍ ᐃᓕᓴᕐᓂᖅᑐᓄ

ᑕᐃᔨᓯᑐᖏᓐᓄᑦ ᓄᓇᓄᑦ ᐊᑐᖏᑕᐅᓕᖅᓯᒋᐊᓕᒻᒥᓪᓗ ᐃᓄᐃᑦ ᐅᖏᐅᓯᑐᖏᓐᓄᑦ (*ᐃᓄᐃᑦ ᐃᑲᔪᓯᐊᕐᓯᖏᓐᓄᑦ ᐊᖏᕈᑎᑦ 2.1.6*). ᑲᓇᑕᒥ ᐅᒪᔨᓕᓚᕆᓄᑦ ᐃᖏᓂᐅᖏᓇᖅᑐᓪᓗ ᐊᐅᓕᕆᑎᕆᐅᖏᖅᑕᐅᓂᖓᓄᑦ ᑲᑎᒪᓕᕆᓂᖅ ᑐᓴᓯᓂᖏᖅᑐᑦ ᐃᓄᖏᖅᑐᐅᓕᐅᖏᖅᑐᑐᖏᖅᑕᐅᓕᖅᓯᓄᑦ, ᐃᓄᓐᓄᑦᓗ ᐱᒻᒪᓚᐅᔨᓄᑦ ᐃᓄᖏᖅᑐᐊᕐᓯᓚᐅᔨᖏᓐᓄᑦ ᒪᑕᓯᑎᑐᑦ *6.4*-ᒥᑦ *6.7*-ᖓᓂ *ᐃᓄᐃᑦ ᐃᑲᔪᓯᐊᕐᓯᖏᓐᓄᑦ ᐊᖏᕈᑎᑎᓄᑦ*, ᐱᓕᕆᐊᖏᑦᑐᖓᒥᑦ, ᐊᐅᓕᖅᓯᒋᐊᓕᒻᒥᓪᓗ ᐊᐅᓕᕆᓂᖅ ᐊᐅᓕᕆᓂᖅ

ᑲᑎᒪᓕᕆᓂᖅ ᐅᒪᔨᓕᓚᕆᓄᑦ ᐃᓄᖏᖅᑐᐅᓕᖅᓯᒋᐊᓕᒻᒥᓪᓗ ᐊᐅᓕᕆᓂᐅᓕᖅᓯᒋᐊᓕᒻᒥᓪᓗ ᐊᐅᓕᕆᓂᖅ ᐊᐅᓕᕆᓂᖅ ᐊᐅᓕᕆᓂᖅ

6.1.3 ᐃᓄᐃᑦ ᑕᐅᖅᑐᑦ ᑕᐃᔨᓯᑐᖃᖅᖢᑎᑦ ᓄᓇᒧᑦ ᐃᒪᖁᓕᓪᓗ

ᐱᖁᖏᖁᖅᑎᑕᐅᕐᓂᖅ ᐊᑕᐃᓗᑦ *ᐃᓄᐃᑦ ᐃᑲᔪᐊᕐᖏᓐᓄᑦ ᐊᖏᑎᑲᑎᑎᑐᒧᑦ (ᑎᑎᖁᖏᖅᓂᒃ 2.1.6-ᒥ)* ᑎᑎᕋᖁᑕᓚᐊᖁᖅᓂᖏᖅᓂᒃ ᐃᓄᖅᓄᑦ ᑕᐅᖅᑐᑦ ᑕᐃᔨᓯᑐᖃᖁᓚᐅᕐᔪᓂᒃ ᓄᓇᒧᑦ ᐃᒪᖁᓄᓐᓂᑕᐅᕐᔪᓂᒃ. ᓄᓇᒧᑦ ᑕᐃᔨᓯᑐᖃᖅᑎᑦ ᑎᑎᕋᖁᑕᐅᓕᐅᖅᔪᒪᔪᑦ ᒪᕐᑐᐃᖅᑕᖁᔪᓂᒃ ᐊᐅᓚᓂᒃ (2006-ᒥᓗ 2012-ᒥᓗ). ᑲᓇᑕᒥ ᐅᒪᔪᓕᓚᖅᓯᑦ ᐃᖁᓂᐅᖁᓕᐅᖅᑐᑦ ᐊᐅᓚᖅᑎᓂᑎᐊᖅᑲᐃᓐᓇᐊᖁᖁᑦᑐᑦ ᐊᔪᖅᑭᓂᒃ ᖃᐅᔨᒪᓕᐅᓂᖅᑭᓂᒃ ᑲᑎᕋᖁᑕᐅᓂᖅᑎᑦᑎᐊᖅᑲᓂᒃ ᖃᐅᔨᒪᓕᐅᖁᔨᐅᓚᑦᑕᑦ ᑕᐃᒃᑯᓗᒃ.

6.2 ᐅᒪᔪᓕᓚᓂᒥᒃ ᐊᖓᒪᓗ ᐅᒪᔪᖁᖅᖁᐅᓴᖏᓐᓂᑦᓂᒃ

6.2.1 ᐊᔪᕐᓇᖅᑕᐅᓂᖏᓐᓄ ᐃᓕᓴᕆᐅᔪᒪᔪᖅᖢ ᖃᐅᔨᐃᑦᑐᓂᖏᓐᓄᓂ ᓴᓇᓂᒃᑯᑎᐊᖅᑐᐊᔭᓚᐅᐊᖅᔪᓂ ᑎᖏᒥᐊᖅᖁᖁᓕᐅᓂᖅᓗᑦ, ᐊᔪᖏᑕᐅᖅᖢ ᐅᒪᔪᐃᑦ ᐊᒥᓱᖏᓪᓗᐊᓕᖅᑐᑦ ᐊᖓᒪᓗ ᐅᒪᔪᐃᑦ ᐃᒃᓴᖅᑕᐅᓴᖅᑲᓕᖅᑐᓂᒃ ᐊᖅᑭᔪᓕᒫᒃ ᓴᓇᓂᒃᑯᑎᐊᖅᑐᐊᔭᓚᐅᐊᖅᔪᓂᒃᑯᑎᐊᖅᑲᓂᒃ. ᑲᓇᑕᓕᒫᒥ ᐅᒪᔪᖁᖅᖁᐅᓴᖏᓐ ᐱᐅᓂᖅᓴᓂ ᓴᓇᓂᒃᑯᑎᐊᖅᑐᐊᔭᓚᐅᐊᖅᔪᓂᒃᑯᑎᐊᖅᑲᓂᒃ ᑕᒪᖁᖅᖢ ᐅᒪᔪᐃᑦ ᐊᖓᒪᓗ ᐊᖏᔪᓕᐅᓇᖁᓂᖅᓴᓂᒃᑯᑎᐊᖅᑲᓂᒃ ᐊᑐᖁᔭᓂᒃ ᐃᓄᐃᑦ *ᐃᓄᐃᑦ ᐃᑲᔪᐊᕐᖏᓐᓄᑦ ᐊᖏᑎᑲᑎᑎᑐᒧᑦ*. ᒪᓕᑐᖁᓕᖅᑕᒪᓕᖁᓂᖅᓴᖏᓐᓄᑦ ᓄᓇᓕᐃᑦ ᑲᓇᑕᓕᒫᒥ ᐅᒪᔪᖁᖅᖁᐅᓴᖏᓐᓄᑦ, ᑕᐃᓚᐃᓕᓕᖅᑐᑦ ᐅᒪᔪᐃᑦ ᐃᓄᐃᑦ ᐊᖓᒪᓗ ᐃᓄᐃᑦ ᐃᒃᓴᖅᑕᐅᓴᖅᑲᓕᖅᑐᓂᒃ ᐊᐅᓚᓂᒃ ᐃᓄᐃᑦ ᐃᓄᐃᑦ ᐊᖏᔪᓕᐅᓇᖁᓂᖅᓴᓂᒃᑯᑎᐊᖅᑲᓂᒃ ᐊᑐᖁᔭ

6.2.2 ᐊᒥᓱᓗᐊᓕᕐᓂᖅᓯᒪᓕᒃ ᑲᖏᐅᔪᓂᖅ

ᐊᒥᓱᓗᐊᖅᑐᓖᓕᓚᐊᔪᓕᕐᓂᖅᓯᒪᓕᒃ, ᓇᒍᑎᑭᓗ, ᐊᕐᑎᑎᑦᓗ ᐊᐅᑐᖅᓯᕐᓕᐊᓂᖅᓯᒪᓕᒃ ᑲᖏᐅᔪᓂᖅ ᑕᕐᕿᓴᖅᓯᐅᔪᐊᖅᑕᕐᓂᖅᓯᒪᓕᒃ ᐃᔪᓗᐊᓂ ᐃᑲᒋᐊᖅᑐᐃᖅᓯᒪᓕᒃ ᓯᓚ ᑐᕿᓯᐅᓴᓕᖅᑎᐊᖅᑭᑦ, ᕿᓯᐊᓂ ᐅᒪᔨᓚᓴᖅᑯᑦ ᑕᒪᐃᓐᓇᖅᑐᑦ ᐃᓯᒪᓴᖅᓯᐅᑎᖅᑎᐊᖅᑐᑦ ᐊᐅᓚᑕᐅᔨᓴᖅᑐᓂᖅᓯᒪᓕᒃ ᐊᐅᓚᑕᐅᕿᖅᑐᐃᓐᓇᖅᑐᑦ ᖃᓄᐃᓕᓚᑦᑕᖅᑐᓂᖅᓯᒪᓕᒃ, ᐊᒻᒪᓗ ᐊᑐᖅᑕᐅᔨᓴᖅᑐᑦ ᖃᓄᐃᓕᓚᐅᕿᖅᑐᐃᓐᓇᖅᑐᑦ. ᑲᖏᐃᑦ ᐊᒥᓱᓗᐊᓕᕐᕿᒪᑕ ᖃᐅᔨᓴᖅᑕᐅᓯᒪᔨᓂᓗ ᑎᑎᕋᖅᓯᒪᓕᒃᑐᑦ ᐊᐅᑐᖅᑕᐅᓕᕐᓂᖅᓯᒪᓕᒃ ᓇᓇᖅᑐᑦ ᑎᓯᒥᐊᖅᑲᓴᖅᓗ, ᖃᐅᓂᓴᖅᑐᓂᓂᓴᖅᓗ ᑲᖏᖅᓯᐅᕋᓴᖅᑐᓂᓂᓴᖅᓗ (ᑲᑐ, ᐱᓕᓚᖃᑎᖅᓯᖅᓵᓗ-Carter, et al., 2018; Fᐊᓐᑎᓐ ᐊᒻᒪᓗ ᐃᓗᓐᓚ-Fontaine and Mallory, 2011). ᐃᓯᒪᖅᓯᑕᐅᓗᑎᒃ ᖃᓄᐃᖅᓯᐅᔪᓴᐅᔨᓐᓇᖅᓯᓯᒪᔪᑦ ᑲᓇᑕᒥᓗ ᐊᒥᐊᓕᑲᒥᐅᓂᓗ ᓇᓀᓴᓕᑕ ᖁᑎᖅᑭᓯᐊᓂ ᑲᖏᐃᑦ 1999. ᑲᓯᖅᑕᐅᖅᑲᑕᕐᓂᖅᓯᒪᓕᒃ ᐊᒥᓯᑕᖅᓯᒪᒪᔪᖅᓯᑎᒃ, ᑲᓯᖅᑐᐃᑦ ᐊᒥᓯᑕᖅᑭᑦᓕᐊᑎᐊᔪᓕᕐᓂᖅᓯᒪᓕᒃ ᐊᒥᓯᑕᖅᑭᑦᓕᐊᑎᐊᖅᑕᐅᓇᓗᔪᑦ, ᐊᒥᓱᓐᓂᕐᑎᐊᔪᓕᕐᓂᖅᓯᒪᓕᒃ ᐊᒻᒪᓗ ᑭᓇᐅᔪᓕᐅᑕᐅᔨᓴᖅᑐᓂᓂᓴᖅᓯᒪᓕᒃ (Carter, et al., 2018). ᑭᓯᐊᓂᓕᑕᐅᖅ, ᐃᓕᓴᐅᔪᖅᑲᓴᖅᑐᑦ ᐃᓕᓕᖅᓯᓯᒪᒋᔪᓕᒃ ᖃᓄᐃᑦ ᐱᒻᒪᓚᐅᓂᖅᑲᓕᕐᓂᖅᓯᒪᓕᒃ ᐊᕐᑎᐅᔪᑦᓗ, ᐊᕐᑎᑎᐅᑕᖅᑭᓚᓇᓕᓗ ᐅᓚᓴᖅᑐᑦ ᐃᓕᓴᖅᑲᓕᕐᓂᖅᓯᒪᓕᒃ ᐊᐅᑐᖅᑕᐅᓕᕐᓂᖅᓯᒪᓕᒃ ᐃᓕᓴᖅᑲ

(ᓇᓗᓇᐃᖅᑕᐅᓯᒪᓕᖅᑐᓂᒃ ᐅᒪᔨᐃᑦ ‘ᓄᖓᓕᑎᓕᕆᓂᖅᓗᒃ’ ᐅᕐᓀᓗᓐᓂᑦ ‘ᐅᓗᓚᐊᓇᖅᑐᑦᓯᓂᖅᓗᒃ’) ᐅᕐᓀᓗᓐᓂᑦ ᐊᐅᓪᓚᑎᓂᒧᑦ ᐸᖁᐅᑎᓂ ᐅᖁᑲᐅᓯᐅᒪᔪᑎᑲ (ᐅᒪᔨᓂᒃ ᑕᐃᔭᐅᓪᓚᑐᓂᒃ ‘ᐊᖏᐅᕐᓂᑕᒥᒃ ᐃᓱᒪᓕᑕᐅᓂᖅᓗᒃ’) ᑐᕆᖓᓂᐊᖅᑐᓂᒃ ᐊᒥᓱᑦᑐᓂᒃᓴᓂᒧᑦ ᖃᓄᐃᓪᓕᐅᑎᑲᖅᑕᖓᖅᓂᒃ ᑕᒪᒃᑯᐊ. ᑕᒪᒃᑯᐊ ᑎᑎᕆᖅᓱᒪᓂᐊᖅᑐᑦ ᐅᓂᒃᑲᕆᓂᐊᖅᑐᑦ ᓇᓂᖅᑕᕆᓂᖓᓂᒃ ᐅᒪᔨᐃᑦ ᐱᑕᖅᐃᓇᑭᓇᓂᐊᕐᓗᒍ.

ᖃᐅᔨᒪᔭᐅᓂᖓᓂ ᓇᔨᓴᓇᕐᓂᖓᓂ ᓇᒋᕐᕕᐊᓂᕐᓂᖓᓂ ᐅᒪᔨᐃᑦ ᐊᒥᓱᓕᕐᓂᑦᑐᑦ ᐃᑲᓂᑐᐊᖅᑎ ᑎᓕᒥᐊᖅᑲᕕᖅ ᐃᑲᔪᓂᖅᑲᖅᑐᑦ ᓴᓇᔭᐅᔪᓇᕐᓂᖓᓂᓐᓂ ᐊᑐᓕᖅᑎᑕᐅᔨᓇᓂᖓᓂᓐᓂᒃ ᐊᒥᓱᓕᖅᑐᓂᒃᑯᑦᑐᓂᒃᑯᐊᑕᑎᓂᖅᓴᐅᓂᒡᒎ ᐊᒥᓱᓕᖅᑐᓂᒃᑯᑦᑐᓂᒃᑲᐅᔭᓂᒃ. ᑎᓕᒥᐊᖅᑕᕕᖅᑎᓂᖓᓂ ᓴᓗᓂᒥᐊᖅᑕᐅᔨᓇᓂᖓᓂᒃ ᐅᒪᔨᐃᑦ ᐱᑕᖅᐃᓇᑭᓇᓂᐊᕐᓂᒃ ᐊᒥᓱᓕᖅᑐᓂᒃᑯᑦᑐᓂᒃᑲᐅᔭᓂᒃ ᓄᓇᓕᒫᒃ ᐊᒥᓱᓕᖅᑐᓂᒃᑯᑦᑐᓂᒃᑲᐅᔭᓂᒃ ᑕᐃᔨᓄᑦᓯᒪᓕᖅᑐᓂᒃᑯᑦᑐᓂᒃᑲᐅᔭᓂᒃ ᐅᒪᔨᐃᑦ ᐱᑕᖅᐃᓇᑭᓇᓂᐊᕐᓂᒃ ᐊᒥᓱ

ᐃᓄᐃᑦ ᑎᖏᑕᐊᖅᓯᐅᑦᑐᑕᑭᓐᓇᖅᓂᖓᑦ, ᐊᓯᖏᑭᓐᓂᒃᓗ ᑎᖏᑕᐊᖅᓂᒃ ᓂᖅᑭᒃᓴᓗᐅᑦᒧᐊᓂᒃ, ᒪᑕᖏᓂᐊᖅᑎᓪᓗᓂ ᓄᓇᕗᒻᒥ ᓄᓇᕗᑦᑎᓄᑦ, ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓯᒪᓂᖏᓐᓄᓄᑦ ᐃᑲᔪᓯᐊᖅᑕᑭᑎᒃᓴᖏᓐᓄᑦ ᑎᑎᕆᖅᓯᒪᔪᓂᒃ, ᑲᑎᒪᐅᑕᐅᕐᓂᖏᓐᓄᑦ ᑎᖏᑕᐊᓕᓚᓂᖅᓴᓂ ᒪᓕᑐᐃ *MBCA*-ᓂᒃ ᑕᒪᐃᔨᖏᓐ ᐃᓕᑎᕆᐅᓂᖏᓐᓂᒃ ᒪᓕᑌᓪᓗᑐᐃᐅᔪᓂᒃ.

6.2.6 ᑎᖏᓯᑦ ᐃᖏᕆᕐᕉᓂᖏᑦ ᑕᐅᕐᓂ

ᐊᕆᕎᒍᑦ ᐃᓕᐃᓐᓇᖓᓂ, (ᓯᑦᕆ ᑎᖏᑕᐊᖅᑲᓂᕆᒥ), ᑎᖏᓯᑦᑲᑭᓐᓇᖅᓂᖓᑦ ᕿᑯᓕᐅᑦᑎᐊᖅᑐᑦ ᐃᑲᑦᑐᐊᑦ ᑎᖏᑕᐊᖅᑲᕐᓇᖅᓂᒃ ᐊᐅᓕᑕᐅᓂᐊᖅᑐᑦ ᖃᑲᓂᓕᕆᑕᐃᓕᓕᒫᓚᐊᖅᑲᕐᓂᖅ ᐊᓚᒪᓗ ᑎᖏᓯᑦᑲᖅᑕᐃᓕᓕᒫᓚᐊᖅᑲᕐᓂᖅ ᐊᓚᑐᖅᑕᐅᓴᕆᕕᐃᓂᖏᓐᓂᒃᓂ ᓄᓇᕗᒻᒥ ᐊᓚᒪᓗ ᐊᓚᑐᖅᑕᐅᓴᕆᕕᐃᓂᑐᓂᖏᓐᓂᒃᓂ ᑎᖏᑕᐊᑦ (ᓯᑦᓗ ᑎᐱᕐᓕᓕᐊᓕᓴᕆᑦᓂᖏᓐᓂᒃᓂ, ᐅᓪᓗᖅᑕᓂᖅᑎᓪᓗᓂ, ᐊᓚᒪᓗ ᑎᖏᓐᓄᒃᓴᐃᓕᓕᒫᓚᐊᖅᑲᕐᓂᖏᓐᓂᒃᓂ ᐃᓚᓴᖅᓯᒪᓂᖏᓐᓂᒃᓂ), ᐊᓚᒪᓗ ᑎᖏᓐᓄᒃᓴᐃᓕᓕᒫᓚᐊᖅᑲᕐᓂᖏᓐᓂᒃᓂ ᐃᓚᓴᖅᓯᒪᓂᖏᓐᓂᒃᓂ ᐊᓚᑐᖅᑕᐅᓴᕆᕕᐃᓂᑐᓂᖏᓐᓂᒃᓂ ᐃᓚᓴᖅᓯᒪᓂᖏᓐᓂᒃᓂ ᑎᖏᑕᐊᑦ ᐃᓚᓴᖅᓯᒪᓂᖏᓐᓂᒃᓂ ᐃᓚᓴᖅᓯᒪᓂᖏᓐᓂᒃᓂ ᑎᖏᑕᐊᑦ

(1) ᒪᓕᑭᐊᖏᖅᑐᑦ ᑕᐃᵇᑯᓂᖅᓱ ᐅᑭᐅᖏᑕᖅᑐᒥ ᐅᒥᐊᖅᓯᐊᑦ ᓯᑕᖅᑎᓇᑦᑕᐃᓕᓕᒫᓇᐊᖏᖅᓂᖅᓯᒪᓂᑦ ᒪᓕᑎᓪᓗᑐᑦ ᐊᑎᓴᓄ ᐅᑭᐅᖏᑕᖅᑐᑉᐣ ᐃᒪᖅᑯᑎᖅᓯᓂᖅ ᓯᑕᖅᑎᓇᑦᑕᐃᓕᓕᒫᓇᐊᖏᖅᓂᖅᓱᒪᓂ ᒪᓕᑎᐃᑦ *(1985)*;
(2) ᐅᒥᐊᖅᓯᐊᐸ ᐃᖏᒃᐸᓪᓕᐊᖏᑕᖅᓂᖅᓯᒪᓂᖅᓂᖅ ᐃᒪᖅᓗᑯᖅᓯᒪᓂᖅᓂᖅ ᑯᕐᓴᑦᑕᐃᓕᓕᒫᓇᐊᖏᖅᓂᖅᓯᒪᓂᖅᓂᖅᓂᖅ, ᐊᓯᖏᓐᓄᑕᐅᑦᑕᐃᓕᓕᒪᓪᓗᑎᒃᓗ ᐃᒪᐃᑦ ᐅᒥᐊᖅᓯᐊᐸ ᐊᑕᐅᒍᑦᑐ ᐅᓪᓚᑕᖅᑐᑦ ᑯᕐᓴᑦᑕᐃᓕᓕᒫᓇᐊᖏᖅᓂᖅᓯᒪᓂᖅᓂᖅᓂᖅ ᐃᑎᕐᕕᖅᑕᐃᓕᓕᒫᓇᐊᖏᖅᓂᖅᓯᒪᓂᖅᓂᖅᓂᖅ ᐃᑎᑐᑦ ᐃᓕᓴᐅᑎᖅᑐᑦ ᐊᑐᐅᑦ ᑎᓐᓕᖅᑎᖅᓯᐅᑎᓂᖅᓂᖅᓂᖅ ᑭᓱᓕᕐᓂᖅᓂᖅᓂᖅ ᐃᑎᑐᑦᓂᖅᓂᖅᓂᖅ;
(3) ᑐᓴᖅᓴᐅᔪᓂᖅᑎᑎᑕᐃᓕᒪᑎᐊᖅᓯᓗᑎᖅ (ᓯᑲᓗ ᐊᐅᓕᐅᑦᑕᑦ ᐃᖏᑎᕆᑲᒃᓕᓕᐊᑭᓐᓇᖅᓯᒪᓂᖅᓂᖅ ᓯᖅᑲᓕᓂᖅᓴᓴᐅᒃ, ᖃᐅᔨᐊᑭᑎᖏᑦᓕᓗ ᐊᑐᖅᑕᐅᑕᐃᓕᓕᓗᑎᖅ) ᐅᒥᐊᖅᓯᐊᐸᓂᖅ ᐅᖓᑕᓯᖏᓕᓐᓂᒃᑎᑦᑕᑎᐊᖅᑲᖅᑐᓂ ᐊᑕᐅᓯᖏᓐᓄᒃ 1 km ᑭᒋᒥᑕᓂᖅ ᑎᓐᓕᐊᖅᑲᖅᓯᒃ (ᐅᓗᓚᐊᓇᖅᑐᒃᑎᓕᑭᓇᓂᖅᓂᑦ ᐃᓱᒪᒋᔭᐅᓗᑎᖅ); ᐊᖏᒪᓗ
(4) ᑐᑭᓯᓂᐊᖅᓇᐅᖅᑕᑦᓗᑎᖅ ᑲᓇᑕᒥ ᐅᒪᓱᓕᓚᕕᖓᑦ ᐃᖅᑲᓗᓂᓂ, ᐱᓕᓚᐊᓚᓂᐊᖅᑕᖏᓐᓄᖅᓂᖅ ᑲᔪᓯᑎᑎᑭᓚᐅᔭᖓᓂᖅᓂᓂ, ᐅᖅᑲᐅᓯᐅᓂᐊᖅᓚᒪᑕ ᐅᒥᐊᖅᓯᐊᐸ ᓇᐅᑲᑦᑐᖅᓂᖅᓂᖅᓂᖅ, ᐊᖏᒪᓗ ᖅᓇᓂᖅ ᐅᓰᒪᔪᓂᖅ ᑕᒪᐃᓐᓄᖅᑑᓗᓂᖅᓂᖅᓂᖅ ᑐᓴᖅᓴᐅᒃᑎᑦᑕᑎᐊᖅᑲᖅᓯᒪᓂᖅᓂᖅᓂᖅ ᑐᖅᓴᖅᓴᐅᒃᑎᑦᑕᑎᐊᖅᑲᖅᓯᒪᓂᖅᓂᖅᓂᖅ.

ᐊᖅᑎᒍᐸ ᐃᓕᐃᓐᓇᖅᓱᓂ (ᓯᑲᓗ ᑎᓐᓕᐊᖅᑲᓂᖅᓱᖅᓱᓂ) ᑕᒪᖅᑎᒃ ᐅᒥᐊᖅᓯᐊᑦ ᐃᒪᓐᓇᐅᓕᖅᓱᖅᓱᓕᓐᓂᖅᓯᒪᓕᒫᓂᐊᖅᑕᖏᓐᓄᑦ ᖃᓄᖅᓕᓗᑎᐊᖅᑕᐃᓕᓕᒪᓪᓗᑎᖅᑲᖅᓯᒪᓂᖅᓂᖅᓂᖅ:

(1) ᑎᑭᓯᒪᔪᓇᐊᖅᖃᖏᑦᑐᑦ ᐱᓕᓇᓂᖅᓯᒪᓂᖅᓂ ᐅᑦᓗᑲᖅᑲᐃᐅᕐᓂᖅᑐᓂ ᑎᓐᓕᐊᖅᓇᑦ (ᑕᖅᑭᒥ ᐊᒻᒪᓕᐸᓂᖅ ᑎᑭᑦᓗᑐ-ᐊᑲᓯᒪᑦ ᑎᑭᑦᓗᑐ);
(2) ᐅᖓᓱᓯᓂᖅᑲᓇᐊᖅᑲᖅᑐᑦ 500 m ᑭᑕᓂᖅ ᖃᓄᖅᓕᓕᑕᓕᓕᑎᑕᐅᓗᑎᖅ ᑎᓐᓕᖅᑲᖅᓯᐅᓂᖅᓂᖅ ᐊᖏᒪᓗ/ᐅᕐᓕᓗᑦᓂᖅᓂᑦ ᓇᓇᒥᐅᑕᖅᑲᖅᑐᓂᖅ ᑎᓐᓕᐊᖅᓂᖅ (ᐊᕐᑎᑎᓕᓚᕆᔨᐊᐊᖅᑯᑦ 2016).

6.3 ᓇᐅᑦᑎᖅᑐᐃᓱᖅᑐᖏᐊᕐᖓᑦᑕᓂᖅᑐᑦ ᖃᐅᔨᔪᖏᐊᕐᓂᖅᓂᑦᓗ

ᐋᖅᑭᒪᒪᑦᓕᐊᖅᑐᓂᖅ ᐊᑐᑎᖅᑲᑦᑎᐊᖅᑐᓂᖅᓗ ᓇᐅᑦᑎᖅᑐᐃᓱᖅᑐᖏᐊᕐᖓᑦᑕᑦᑲᖅᑕᖅᓂᖅᑐᑦ ᐱᓕᓚᐅᖅᓴᐃᑦ ᑲᑐᔪᐅᔪᓚᐊᖅᑎᑲᖅᓯᐅᓂᖅᓂᖅ ᖃᓄᖅᓇᓕᐅᑦᑕᐅᓕᖅᑐᓂᖅᓂᖅ, ᐱᓕᓚᐊᖓᔪᓕᓐᓂᒃᑎᖅ ᐃᓕᓐᓂᑦᑐᑦ ᖃᓄᖅᓇᔪᔪᖅᑎᑦ ᐊᓯᖏᓐᓂᒃᓗ ᐃᓱᐅᔪᓂᖅᑲᖅᓯᐅᓂᖅᓂᖅᓂᖅ ᑐᓂᓯᔪᓐᓇᖅᑐᖅ ᐊᓯᖏᓐᓂᒃᓗ/ᐅᕐᓕᓗᑦᓂᖅᓂᑦ ᐃᓱᒪᓕᐅᑦᑕᐅᓕᖅᑐᓂᖅᓂᖅᓂᖅ ᑐᓂᓯᔪᓐᓇᖅᑐᖅ ᐊᑐᐅ

(5) ᓂᓪᓕᐅᕐ ᐊᓂᔪᓇᖅᑐᑦᓯᐊᓂᖓᓄᓇᓕᒃ ᐊᐅᑐᖅᑕᐅᓂᒪᓕᒃᑐᑦ ᓄᓇᕗᑦ, ᓄᓇᒥᐅᑕᐃᓐᓇ, ᐊᐢᑎᑎᐅᑕᖏᑦᑐᓂᓇ, ᐅᒪᔨᑦᑕᐊᓇ ᐊᓂᔪᓇᖅᓂᒪᓕᒃᑐᖅᓂᒃ ᐊᑐᓂᓇ, ᐊᒥᓂᑉᑐᓐᓂᐊᕐᓇᖅᑐᖅᓂᒃ ᐊᓯᒪᓇ ᐅᒪᐃᖅᓇᑉᓇᖅᑐᖅᓂᒃ.

ᐃᓕᓴᕐᕿᓘᑦᑕᑦ ᐊᐅᓪᓚᑎᑕᐅᕐᕿᑐᑦᑕᐅᔪᑦ ᑲᑎᒪᔨᕆᓪᓕᐊᑦ ᐃᓂᓂᒪᖅᓯᐅᑦᑕᐅᓇᖅᑐᑦ ᐊᐱᖓᓪᓚᐅᔪᖅᓯᒪᓕᖅᑐᑦ ᐊᓯᒪᓇ ᐅᒪᐃᖅᓇᑉᓇᖅᑐᖅᓂᒃ ᐃᓕᓴᖅᓇᑉᓇᖅᑐᖅᓂᒃ.

ᐃᓕᓴᕐᕿᓘᑦᑕᑦ ᐊᐅᓪᓚᑎᑕᐅᕐᕿᑐᑦᑕᐅᔪᑦ ᑲᑎᒪᔨᕆᓪᓕᐊᑦ ᐃᓂᓂᒪᖅᓯᐅᑦᑕᐅᓇᖅᑐᑦ ᐊᐱᖓᓪᓚᐅᔪᖅᓯᒪᓕᖅᑐᑦ ᐊᐱᖓᓪᓚᐅᔪᖅᓯᒪᓕᖅᑐᑦ ᐊᐅᓪᓚᑎᑕᐅᕐᕿᑐᑦᑕᐅᔪ

ᐃᓄᐃᑦ ᑎᒥᖅᑎᖕᓂᓄᑦ, ᓴᖅᒪᓂᖅᓄᑦ, ᐊᖏᕐᕕᑎᓂᑲ ᓴᓇᖃᖅᑳᑐᖅ. ᖃᐅᔨᓴᓪᓕᐅᑎᐊᕐᓂᖅᓴᐅᕙᓐᓄᑦᓚᖅ ᑎᖕᒥᐊᖃᖅᓴᖅ ᓴᖅᒥᒃᑎᑕᒥᓄᑦ ᐃᓄᓇᑎᖅᓂᖕᓄᑦᑲ ᐊᓗᖕᓂᓄᑦ ᐱᓕᕆᐊᖓᔪᖅᑐᓂᐊᖅᑐᖅ ᐊᕐᓇᒍᑦᒪᖅ ᑲᔪᓇᒥᑦᑲ ᐊᐅᓚᑎᑦᑎᐊᓂᖕᓂᓄᑦ ᐱᓕᕆᐊᖓᔪᖅᑐᓂᐊᖅᑐᖅᑐᖅ ᐊᐅᓚᑎᑦᑎᐊᓂᖕᓂᓄᑦ ᐊᐅᓚᑎᑦᑎᐊᓂᖕᓂᓄᑦ ᐃᓕᓴᕐᓂᐊᕐᓂᖅ ᑲᑎᒪᓂᕐᓂᖅ.

6.4.1 ᓄᓇᓕᒃ ᐅᓴᐅᓇᖕᓂᖅᓴᐅᕙᓐᓄᑦᓚᖅᑎᑦᓂᖕᓂᒃ ᑐᐱᓇᐅᒪᔭᕆᐊᖃᖅᑎᑦᓂᖕᓂᒃᓚ

ᑲᔪᓇᖕᓇᒥᑎᐊᖃᓂᐊᕐᑐᖅᓄᑦ ᐊᐅᓚᑎᑦᒥᑎᐊᕐᓇᖕᓂᒃ ᑎᖕᒥᐊᖃᖅᑎᖕᒥᑦᑐᖅ, ᐃᖃᓗᐅᖃᖅᑎᖕᓂᑦ ᐊᐅᓚᑎᑦᑎᓴᐅᖃᖅᑎᑦᓂᖕᓂᒃ ᑲᑦᓂᖕᓂᖅᓄᑦ ᓄᑖᒪᓕᕆᓂᖕᓂᒃ ᓇᓕᓪᓗᑎᒃᓄᑦᒥᑎᐊᖃᖅᑎᑦᓂᖕᓂᒃ ᓇᓕᓪᓗᑎᒃᓄᑦᒥᑎᐊᖃᖅᑎᑦᓂᖕᓂᒃ ᑲᑦᓂᖕᓂᖅᓄᑦ ᐊᐅᓚᑎᑦᑎᐊᖃᖅᑎᑦᓂᖕᓂᒃ ᐊᐅᓚᑎᑦᑎᐊᖃᖅᑎᑦᓂᖕᓂᒃ ᑎᖕᒥᐊᖃᖅᑎᖕᒥᑦᑐᖅ, ᒪᓕᒐᓂᑎᒃ ᓴᖅᑭᑎᓂᖕᓂᒃ ᐊᐅᓚᑎᑦᑎᐊᖃᖅᑎᑦᓂᖕᓂᒃ ᐊᐅᓚᑎᑦᑎᐊᖃᖅᑎᑦᓂᖕᓂᒃ ᑎᖕᒥᐊᖃᖅᑎᖕᒥᑦᑐᖅ, ᒪᓕᒐᓂᑎᒃ ᓴᖅᑭᑎ

ᐃᓗᐊᓂᒃ ᓄᓇᕗᑦ ᑎᒍᑎᓕᕆᔨᒃᑲᓂᕐᒥᐅᑦ (*ᐃᓄᐃᑦ ᐃᑲᔪᖏᖅᓯᒪᔭᖅ ᐊᖏᕈᓐᓇᖅᑐᓂ ᑎᑎᕆᔨᒃᑲᓂᕐᒥᐅᑦ 6.8.1*).
ᑐᑭᓯᐅᑎᕿᓕᕐᓂᖅᓴᐃᑦ ᐱᔪᓐᓇᕆᐊᓗᖅᑐᑦ ᐊᑭᑎᕿᓕᕐᒪᔪᓂᒃ ᐊᖏᕈᓐᓇᖅᑐᓂᒃ ᐱᑕᖅᐃᓐᓇᕆᔨᓂᒃ
ᐳᓪᓕᖅᑐᓪᓗᒍᑦ, ᐅᓗᓇᐃᔨᐃᒪᐅᔨᓐᓇᕐᓂᖏᓐᓄᓪᓗ ᐃᓗᐊᓂᒃ ᐊᖏᔪᓂᒃ ᑎᒍᑎᓕᕆᔨᒃᑲᓂᕐᒥᐅᑦ ᖃᐅᔨᓴᖅᑎᑦᑎᓂᒃ.

ᑕᒪᖅᑭᒃ ᓴᓇᔨᐅᓕᒫᑦ ᐊᑐᐃᓴᖅᓯᐃᑦ (ᑎᑎᕆᖅᓯᓕᒫᑦ, ᑐᓴᖅᓴᖅᓯᐃᑦ, ᑕᕆᓴᔨᑎᑲᓪᓗ) ᐃᓕᓴᐅᔪᐊᑭᑎᑦ,
ᑐᑭᓯᐅᑕᑎᓇᓕᒫᑎᑲᓴᓐᓂᐅᓚᑦ ᐃᐅᔨᓂᒃ ᑎᒍᑎᓕᕆᔨᒃᑲᓇᖏᑦᑑᒃ ᐊᑐᐃᓇᔨᔨᔨᕐᓂᐅᔪᖅᑐᑦ ᐃᓄᐃᑦ ᐅᕆᐅᔨᓕᕆᔨᒃᑲᓂᕐᒥᐅᑦ (*ᐃᓄᐃᑦ
ᐃᑲᔪᖏᖅᓴᖅᓯᒪᔭᖅ ᐊᖏᕈᓐᓇᖅᑐᑎᑦ ᐃᓗᓕᖏᓐᓂ 6.2.1*). ᑲᓇᑕᒥ ᐅᒪᔨᓕᓚᕐᓴᐅᔪᓂ ᐊᐅᓕᕆᑎᓴᐅᕆᔨᑲᑕᐅᔪᓂᕐᒧᑦᓗ ᑲᑎᒪᓴᓭᖏᓐᓂ
ᐃᓄᐃᑦ ᐅᕆᑲᐅᔨᑐᕆᖏᓐᓂᒃ ᐃᑲᔪᕆᔨᐃᔪᓂᕐᒥᐅᑦ ᐊᑐᕆᑎᓴᐅᔪᓂᖏᓐᓂᒃ ᑎᒍᑎᓕᕆᔨᒃᑲᓇᖏᑦᑑᒃ. ᔨᕐᓗ
ᐅᖅᑐᑎᑎᓗᔪ, ᑕᒪᖅᔪᓂᖏᓐ ᓄᓇᖓᓕᓚᕆᓚᐅᔨᒪᔨᔪᓂᒃ, ᑕᔪᐊᖅᓴᓂᒃ ᑎᑎᕆᖅᑲᖅᑲᑐᓂᒃ ᓇᓗᓇᐃᖅᔪᑕᓂᒃ,
ᑐᑭᓯᐅᑭᐅᑦᑐᖅᓴᓂᓪᓚᓐᓂ ᓴᓇᔨᒪᔨᓂᒃ ᑎᑎᕆᖅᑲᖅᓯᒪᔨᖏᓐᓄᐊᖅᑐᑦ ᐃᓄᐃᑦ ᓄᓇᓂᒃ ᑕᐃᔪᔨᓐᓇᕐᓂᖏᓐᓂᒃ ᐊᑐᕐᓗᑎᒃ.
ᑕᒪᖅᑭᒃ ᓴᓇᔨᐅᓕᒫᖏᓐᓂᐊᖅᑐᑦ ᑕᒪᖅᔪᐊ ᐊᑐᕆᓂᐊᖅᑎᑦᑕ ᐃᓄᐃᑦ ᖃᐅᔨᓴᒪᔨᑐᕆᖏᓐᓂᒃ, ᐱᓗᐊᖅᑐᑦ ᐱᔨᓴᖅᓯᐃᑦ
ᐃᓄᐃᑦ ᓄᓇᓂᒃ ᐃᓐᓇᐃᓐᓇᕐᓂᒃ ᐊᑐᕆᑎᓴᐅᔪᓂᖏᓐᓂᒃ ᐊᒻᒪᓗ ᐃᓕᓴᐅᔪᓂᖏᓐᓂᒃ (*ᐃᓄᐃᑦ ᐃᑲᔪᖏᖅᓴᖅᓯᒪᔭᖅ
ᐊᖏᕈᓐᓇᖅᑐᑎᑦ ᐃᓗᓕᖏᓐᓂ 6.8.4*) ᖃᐅᑎᖃᐊᖅᑐᕆᔨᖃᑎᒌᓕᐊᖅᑐᓂᒃᓗ ᑲᓇᑕᒥ ᐅᒪᔨᓕᓚᕐᓴᖏᓐᓄᐊᖅ, ᐅᓴᖅᑐᑎᑎᓕᐊᖅᓯᓕᐊᖅᓯᒪᓕ
ᑕᒪᓕᖅᓴᓕᔨᖅᓚᐊᖅᑲᖏᓐᓂᖏᓐᓂ ᐅᒪᔨᓕᓚᕐᓴᐅᓂᒃ ᓄᓇᕗᑦᑐᓂᒃᓗ ᓄᓇᒥᐅᑕᓚᕆᐅᔨᓂᒃᓗ ᖃᐅᔨᓴᒪᑎᑦᑎᓂᒃ,
ᐱᑕᖅᖃᐅᑦᑕᖅᓂᖏᓐᓂ ᑐᑭᓯᐅᑭᐅᑦᑕᐅᔨᓂᒃ ᓴᓇᔨᐅᓕᒫᓂᖏᓐᓄᐊᖅᑐᑦ (*ᐃᓄᐃᑦ ᐃᑲᔪᖏᖅᓴᖅᓯᒪᔭᖅ ᐊᖏᕈᓐᓇᖅᑐᑎᑦ ᐃᓗᓕᖏᓐᓂ 6.8.5*).

ᑲᓇᑕᒥ ᐅᒪᔨᓕᓚᕐᓴᐅᔪᓂ ᑕᔪᐊᖅᑲᓂᓕᖅᓴᓕᒪᑎᑐᒃ ᖃᐅᔨᓴᒪᔨᔭᕐᑎᑎᑕᕆᐅᔨᑐᓂ ᐃᑲᑕᑐᑦ ᑎᒍᑎᓕᕆᔨᒃᑲᓇᖏᓐᓄᐊᖅᑐᑦ, ᐊᑐᖅᑲᑕᓂᓂᐊᖅᑐᓂᒃ
ᓂᐅᑦᕿᐊᓄᑦ ᑐᑭᓯᐅᑭᐅᑦᑕᐅᔨᓂᖏᓐᓄᐊᖅ ᐱᒪᓚᐃᖏᓐᓄᐊᖅ ᑕᒪᖅᔪᓂᖏᓐᓂᒃ ᐃᓕᓴᐅᔪᓂᖏᓐᓂᒃ ᐃᓕᓴᐅᔪᓂᖏᓐᓂᒃ ᐃᓄᐃᑦ ᐃᑲᔪᖏᖅᓴᖅᓯᒪᔭᖅ
ᐊᖏᕈᓐᓇᖅᑐᑎᑦ ᐃᓗᓕᖏᓐᓂ 6.8.1-ᓂᒃ ᐃᓕᓴᐅᔪᓂᖏᓐᓂᒃ. ᖃᐅᔨᓴᒪᑎᑦᑎᓂᒃᑐᑭᓯᐅᑭᐅᑦᑕᐅᔨᓂᒃ ᐊᑐᐃᓇᔨᔨᔨᕐᓂᐅᔪᖅᑐᑦ
ᖃᐅᔨᓴᒪᓕᕆᔨᒃᑲᓂᕐᒥᐅᑦ ᐃᓄᐃᑦ ᐃᑲᔪᖏᖅᓴᖅᓯᒪᔭᖅ ᐊᑐᐃᓇᔨᔨᔨᕐᓂᐅᔪᖅᑐᑦ ᐃᓄᐃᑦ ᐃᓕᓴᐅᔪᓂᖏᓐᓂᒃ ᐊᑐᐃᓇᔨᔨᔨᕐᓂᐅᔪᖅᑐᑦ

ᖃᐅᒪᑎᑎᑭᐊᖏᑎᓪᓗ ᑐᖅᖁᓪᓚᕐᐳ ᐅᓂᒃᑲᖅᑕᕐᓂᖅ ᐱᑕᖃᐃᓐᓇᔨᓂᕐᒥᒃ ᓄᓇᒥᒃ ᐱᓵᐋᑎᖃᖅᑐᑐᓂᒃ ᐱᓕᕆᓇᒪᓕᕆᓂᒃ, ᐊᑯᓂᐅᔪᓐᓇᖅᑐᓂᒃ ᓴᓱᒪᓗ ᓇᖅᒥᓂᖅ ᐱᓕᕆᐊᖓᑕᐃᓇᓂᖏᓐᓂᒃ ᓄᓇᕗᒻᒥᐅᓄᓂ ᒪᒃᑯᑐᓄᓂ (*Δ.ᓄᓂᐃᑦ Δᑲᓱᓯᐊᓵᖅᓯᒪᓂᓄᓂ ᐊᖏᑕᑏᑦ Δ.ᑐᓕᖏᑦ 9.2.1*). ᐊᐋᑎᓕᕆᔨᓴᓱᐊᑲᑯᑦ Δᓕᒥᔭᐅᔪᓂ ᑲᓇᑕᐅᓕᖅᑐᓂᒃ ᐱᓕᕆᔨᐅᔪᓂᒃ ᐱᐋᓪᓕᐊᔾᔨᐊᑎᓵᖅᓴᐃᓂ ᑲᒪᒥᓕᕆᓂᖓᓂᒃᑐᓂᒃ, ᑐᖁᖓᓚᒪᓕᒡ ᑕᒪᑲᑯᑦ ᐃᓄᐃᓂᒃ ᓄᓇᕗᒻᒥ ᐊᔪᕈᓯᖏᓐᓇᖅᑐᓂᒃᑐᓂᓄᓂ ᐱᓕᕆᐊᖏᓵᖅᑕᕐᓂᖓᓂᒃᑐᓂᒃ ᑲᑎᒪᔨᓴᖅᑎᑦᑎᓂᒃ ᑲᑎᒪᔨᓴᖅᑎᑦᑎᓂᒃ ᑲᑎᒪᔨᓴᖅᑎᑦᑎᓂᒃ ᐱᓕᕆᔨᐅᔪ

ᑎᖃᐊᖅᑎᕕᐅᑦ ᓄᓇᖅᓯᓂ. >ᓪᓕᕐᑐᓪᓗᒥᒃ ᐱᓪᓗᐊᖓᔪᓄᓇᕐᓂᖅᓯᒪᓕᖅᑎᕕᖃᖅ ᐊᔪᖅᑭᓂᐅᓪᓚᑦᑎᓂᐊᕐᓂᖅ ᓚᐃᔾᔨᖅᑕᐅᔾᔨᓕᖅᑎᓂᐅᓪᓗᐊᖅᑐᑦ ᐱᓕᕐᓂᖅᑕᐅᔪᓕᖅᑎᓂᒃᑯᓕᖅᑎᓂᐊᕐᓂᖅ ᐊᑦᑕᒍᑦ ᓄᓇᕗᒻᒥ ᐊᐅᓪᓕᖅᑕᐅᔾᔨᓕᖅᑎᓂᒃᑯᓕᖅᑎᓂᐊᕐᓂᖅ ᓄᓇᕗᒻᒥ ᐅᒪᔨᓪᓚᑦᑎᓂᒃ ᓪᓕᕐᑐᓪᓗᒥᒃ ᓄᓇᕗᒻᒥ ᐃᓄᖅᑲᐃᐅᕐᓇᐅᖅᑕᐃᓇᐅᑦᑐᒃ ᐊᒻᒪᓗ ᓄᓇᕗᒻᒥ ᐃᓄᖅᑲᐃᐅᕐᓇᐅᖅᑕᐃᓇᐅᑦᑐᒃ ᐊᒻᒪᓗ ᓄᓇᕗᒻᒥ <ᖃᓄᐃᓂᑦᑎᓂᒃ ᐱᓪᓗᐊᖓᔪᓄᓇᕐᓂᖅᓯᒪᓕᖅᑐᓂᒃ ᖃᐅᔨᓴᖅᑎᓪᓚᑦᑎᓂᐊᕐᓂᖅ ᓄᓇᕗᒻᒥ ᐊᑐᖅᑕᕐᓗᑎᓪᓚᑦᑎᓂᐊᕐᓂᖅ ᓄᓇᕗᒻᒥ. ᐃᓄᐃᑦ ᑕᒪᐃᓐᓇᐃᓄᒃ ᐃᓄᒃᑯᑦᑕᐅᔾᔨᓕᖅᑎᓂᐊᕐᓂᖅᓯᒪᓕᖅᑎᓂᒃᓴᖅ

ᓄᓇᕗᑦᑕᖃᖏᓐᓇᖅᑐᕈᐊᓂᒃ ᐊᓯᖏᑐᓂᓗ ᐱᖁᑎᖃᖏᓐᓇᖅᑐᓇᖃᖅᑐᒃ ᐃᓕᓂᒃᑯᑦ ᖃᐅᔨᓴᕈᑎᐊᓚᐊᖃᖅᓂᖏᑐᓄᑦ ᐱᐅᔨᓂᒃ ᐋᖅᑭᐅᓚᑦᑎᐊᓂᖓᓂᒃ ᐊᓕᓚᓗ ᐱᐅᖏᑦᑐᓂᒃ ᐊᒃᑐᐃᓂᖃᖅᓂᐊᖏᑦᑐᑐᓂᒃ ᐊᐅᑎᒥᓂᒃ ᐊᐅᓗᐊᓂ.

6.4.6 ᐃᓄᖓᓄᑦ ᐱᖁᑎᑎᑭᐅᔨᓴᑦ ᓄᓇᐃᑦ

ᐃᓄᖓᓄᑦ ᐱᖁᑎᑎᑭᐅᔨᓴᑦ ᓄᓇᖓᑎᑦ ᑎᖏᑕᐊᖅᑲᖅᓇᖏᒥ ᐊᐅᓪᓚᑕᐅᒃᑐᑐᑦ ᒪᓕᒃᑕᐅᑦᓗᑎᒃ *ᑎᑎᕋᖅᓯᒪᓂᖏᑦ 4-ᒥ* ᐃᓄᐃᑦ ᐃᑲᔪᐊᓴᖅᓂᖏᓄᑦ ᐊᖏᑦᑭᑎᖏᑦ ᐅᖃᖅᖢᓯᒪᖏᒪᑕ ᑕᐃᒃᑯᐊ ᓄᓇᒥ ᐱᓪᒪᓗᐅᓂᖅᑲᕋᖏᑦᑐᖏᑦ ᐊᔪᐳᐅᑎᑦᑕᐅᖃᖅᑲᕋᖅᑲᖅᑎᑦᑎᓂᖏᓐᓂᒃ. ᐃᓗᓕᖏᑦ *4-*ᖏᓐᓄᒋᔪᑦ ᖃᓄᐃᓕᓕᐅᖅᑎᐅᓂᖏᓐᓂᒃ ᑲᒪᕐᕉᒃᑐᑦ ᐃᓄᐃᑦ ᐃᓕᓴᕐᕕᖏᓐᓂᒃ ᐊᐅᓚᑦᑎᐊᓂᖓᓂᒃ ᐃᓄᖓᓄᑦ ᐱᖁᑎᑎᑭᐅᔨᓴᑦ ᓄᓇᖁᔭᑎᖏᑦ ᑎᖏᑕᐊᖅᑲᖅᓇᖏᐅᓂᕐᒥᓂᖏᑦ, ᓇᓗᓇᐃᔨᖅᑕᐅᓂᒋᓪᓗᓐᓄᑐᓗ ᐊᖏᑦᑭᑎᑎᑭᐅᓇᐃᓕᓂᕐᒪᔪᓂᒃ ᐊᔪᖄᖅᑎᑎᑭᐊᕆᔭᖏᓐᓂᖏᓄᑦ ᐊᑐᖅᑕᐅᔪᓇᖅᑐᑦ ᐊᓕᓚᓗ ᐊᓂᖏᑦᑐᓂᒃ ᐊᑐᐸᓇᕐᓂᖏᑦᑐᓂᒃ ᐃᓂᓕᓴᕐᕕᖏᓐᓂᒃᓄᑦ (ᐅᒪᔪᓕᓚᐅᓂᒃ ᐳᖅᑐᖅᑐᐃᐅᖏᑦᑐᓄᑦ, ᐱᓕᓚᐅᐅᔪᓄᑦ ᐊᓕᓚᓗ ᐊᖏᑦᑭᑎᓂᒃ ᐊᐅᓚᑦᑎᐊᓂᖓᓂᒃ ᐃᓂᓕᓴᕐᕕᖏᓐᓂᒃᓄᑦ ᐅᑎᓂᖅᑐᐃᐅᔪᓄᑦ ᐊᓕᓚᓗ ᐊᓂᖏᑦᑐᑎᑎᒍᑦ ᐱᓕᓚᓕᖅᓄᑦ ᐊᐅᓚᑦᑎᐊᓂᖓᓂᒃ ᐃᓅᓯᖅᑐᓂᒃ ᑎᖏᑕᐊᖅᑲᖅᓇᐃᓕᒃ ᑎᑎᕋᖅᓯᒪᖏᒪᑕ ᐱᓕᓚᓕᖅᓄᑦ ᐅᑎᓂᖅᑐᐃᐅᔪᓄᑦ ᐃᓅᓯᖅᑐᓂᒃ ᑎᖏᑕᐊᖅᑲᖅᓇᐃᓕᒃ ᑎᑎᕋᖅᓯᒪᖏᒪᑕ ᐃᓄᖓᓄᑦ ᐃᓅᓯᖅᑐᓂᒃ ᐃᓄᐃᑦ ᐊᓕᓚᓗ

- ᐅᓴᖅᑐᑎᐊᖅᓂᖅ ᖃᐅᔨᒪᑎᑎᒋᔪᓕᒃᑯᑦ ᐊᑐᐃᓐᓇᐅᓂᖏᓐᓂᒃ ᐃᓄᖃᖅᖃᐃᐅᒪᔨᒃᑐᓂ ᓴᓪᓕᓂ;
- ᐅᓴᖅᑐᑎᐊᖅᓂᖅ ᐅᒥᐊᔪᐊᒃᑯᔨᖅᑕᑐᑦ ᖃᐅᔨᒪᓂᖏᓐᓂᒃ ᐱᔪᓐᓇᐅᑎᖅᑲᐊᖅᑕᓇᖅᓂᖏᓐᓂᒃ ᐃᐃᒍᑕᓐᓇᖅᖏᓐᓄᑦ ᐃᑲᑐᑐᐊᖅᑐᑦ;
- ᐅᓴᖅᑐᑎᐊᖅᓂᖅ ᑎᓂᓗᔨᖅᑕᑦ ᖃᐅᔨᒪᓂᖏᓐᓂᒃ ᐱᔪᓐᓇᐅᑎᖅᑲᐊᖅᑕᓇᖅᓂᖏᓐᓂᒃ ᐃᓗᐊᓂᓐᓇᓂᕐᐊᑎᒃ ᐃᑲᑐᑐᐊᖅᑐᑦ ᑎᓂᒥᐊᖅᑲᕐᓱᓚᓂᒃ;
- ᐅᓴᖅᑐᑎᐊᖅᓂᖅ ᓴᓐᓂᒥᑎᐅᑦ ᐃᓄᓕᒪᓂ, ᓇᓂᒥᓂᓗ ᐱᓕᓱᓐᓀᓂ, ᖃᐅᔨᒪᓴᓱᐊᖅᑲᖅᓂᖏᓕ ᐱᔪᓐᓇᐅᑎᖅᑲᐊᖅᑕᓇᖅᓂᖏᓐᓂᒃ ᐃᓕᓂᖏᓐᓂᒃ ᐱᓕᓱᔪᓐᓇᖅᓂᔭᑦ ᑎᓂᒥᐊᖅᑲᖅᓂᖅ;
- ᐱᔨᒥᓴᐃᓂᖅ ᓴᓐᓂᒥᑎᐅᑕᓂᒃ ᐃᓄᓕᓵᓂᓗ ᐅᖃᐅᔨᓕᖅᑕᕐᓯᔪᐅᓂᖏᓐᓂᒃ ᒪᓕᑐᖅᓂᒃ ᓯᕐᕃᐃᔨᓂᒃ ᐱᓕᓱᐊᖅᑐᓂ ᑎᓂᒥᐊᖅᑲᖅᓇᐅᓂᕐᒥ ᐅᖓᓚᓄᖅᑐᓂ ᐱᔨᔪᑎᖃᖅᑐᓂ ᑎᓂᒥᐊᖅᓂᖅ ᑎᑭᓯᓂᖅᑐᓂ ᐅᖃᐅᓕᕐᓴᐅᔪᓐᓇᖅᑐᓄᑦ (ᑕᓯᓗᑎᑦ ᐊᐃᖅᑐᖅᓯᒪᓂᖏᓐᓂᑦᑐᓂ 10-ᒥ ᐅᖃᖅᖃᐃᖅᓴᐃᓄᑦ ᑭᓇᖅᓯᓂᖏᓐᓂᑦ); ᐊᓚᓗ
- ᐱᔨᒥᓴᐃᖅᓯᓂᖅ ᑕᐃᒪᓐᓕᑦ ᖃᐅᔨᓕᒋᐊᖅᑕᐅᖅᑕᑦᓂᖏᓐᓂᒃ ᑎᑭᓯᒪᐃᐅᖅᑕᑦᓂᖏᓐᓂᑦ ᐅᖅᖁᓗᓂᑦ ᐱᓕᓱᐃᖏᓐᓂᒃ ᖃᐅᔨᓕᒋᐊᖅᐃᐅᑕᑭᓂᖏᓐᓂᒃ (ᓯᓚᖅᓗ ᖃᑎᖅᑕᐊᖅᑕᐅᖅᑕᑦᓂᓗᑎᒃ ᑎᑭᓯᒪᔨᓂ ᑎᑎᕆᖅᑐᓂᖏᓐᓂᑦ ᑎᑎᕆᐃᐃᐅᖅᑕᑕᓇᓯᐃᔨᓂ ᐅᑎᖅᑐᓂᒃ ᑎᑭᓯᒪᔨᓂᒃ) ᑎᓂᒥᐊᖅᑲᐃᓐᒥ.

6.4.8 ᐊᐃᔨᑐᖅᓯᒪᓂᖏᓐᓂᒃ ᖃᔭᐃᑕᐅᑕᐅᖅᑕᓂᑦ ᐱᑎᐊᖅᑎᑕᐅᓯᒪᓕᕐᖅᑐᑦᓗ

ᓯᓚᑎᒥᐅᑕᓇᓴᐅᔨᓂᒃ ᐃᔨᐱᑎᓴᐅᖅᑲᑕᑦᓂᔨᓂᖅ ᐃᓯᒪᑎᓴᐅᓗᐊᖏᓐᓂᖅᓴᓓᑐᐃᓐᓇᖅᓯᓓᑐᖅᑲᐊᖅᓯᓂᒃ ᑕᔨᔪᐃᓐᖓᓕ ᐊᐅᑕᕐᑎᓂᒥᐅᑦ ᐃᓕᓴᐅᓴᐅᓂᖅᑐᓂᒃ ᐃᓕᓴᐅᓴᐅᓂᖅᑐᓂᒃ (ᓯᓚᖅᓗ ᖃᑎᖅᑕᐊᖅᑕᐅᖅᑕᑦᓂᓗᑎᒃ ᑎᑎᕆᐃᐃᐅᖅᑕᑕᓇᓯᐃᔨᓂ ᑎᑎᕆᐃᐃᐅᖅᑕᑕᓇᓯᐃᔨᓂᒃ ᐅᑎᖅᑐᓂᒃ) ᑎᓂᒥᐊᖅᑲᐃᓐᒥ.

ᓯᓚᑎᒥᐅᑕᓇᖅᓯᓂᒃ ᐃᖅᐱᒋᖅᑲᖅᑕᑦᓂᔨᓂᖅ ᐃᓯᒪᒋᖅᐅᓗᐊᖏᓐᓂᖅᓴᓓᑐᐃᓐᓇᖅᓯᓂᒃ ᐃᓯᒪᒋᖅᐅᓗᐊᖏᓐᓂᖅᓴᓓᑐᐃᓐᓇᖅᓯᓂᒃ ᑕᔨᔪᐃᓐᖓᓕ ᐊᐅᑕᕐᑎᓂᒥᐅᑦ ᐃᓕᓴᐅᓴᐅᓂᖅᑐᓂᒃ ᐃᓕᓴᐅᓴᐅᓂᖅᑐᓂᒃ ᐱᑎᐊᖅᑎᑕᐅᓯᒪᓕᕐᖅᑐᑦᓗ ᐱᓕᓱᐃᖏ

- ᓄᓇᕗᒻᒥ ᓄᓇᕐᓂ ᐊᑐᑎᖅᑳᖅᓯᓄᑦ ᐸᖁᓇᐅᑎᓕᕆᓂᖏᑦ ᐊᓚᒪᓗ ᓯᕗᓂᖅᑳᒥ ᐋᖁᑭᐊᖅᑕᐅᔪᓇᖁᓂᖅᓯᓄᑦ ᐱᔪᐊᑎᖁᖅᑐᓂᖅ.

ᐊᓯᖏᓐᓄᓂᑕᐅᖅ ᐱᑭᐊᖅᑎᑕᐅᓯᒪᔪᓇᖅᑐᓂᖅ ᐊᓚᒪᓗ ᐱᕐᖅᑳᓴᐅᑕᓇᖅᑐᓂᖅ ᐃᓐᓇᓂᒃᑯᓂ ᓴᖅᑭᑦᑐᖅᑐᓂᖅ ᓯᕗᓪᓕᕆᐊᑕᑕᐅᑎᐊᖁᓂᖏᑦ ᐊᕐᕕᖅᑯᓯᒪᓂᖅᓯᓄᑦ ᓴᐳᓄᓂᐊᖅᑕᐅᓂᖅᓯᓄᑦ ᑎᖅᑭᐊᑦ ᑎᑭᐸᐸᖅᑐᓂ ᐊᓯᖏᓐᓂᓗ ᐅᒪᔪᐃᓪᓚᒃ, ᒪᓕᓂᖅᓱᒪᓂᖅᓯᓄᑦ ᐃᓚᖅᑯᓯᒪᓂᖅᓯᓄᑦ ᐃᓕᓴᐅᔪᓇᖅᑐᓂᖅ ᐊᑐᖅᑐᓂᖅ ᐊᓯᖏᓐᓂᓗ ᐅᒪᔪᐃᓪᓚᒃ (ᐃᓄᓇᐃᓪ *ᐃᑲᔪᐊᕐᓯᖏᓐᓄᓂ ᐊᖏᕈᑎᓪ ᐃᓗᓕᖏᓪ 3.4.3*).

ᐊᐅᓚᕎᑎᐅᖅᑲᑕᐅᓂᖓᓪᓕᐊᑦ ᑲᑎᒪᔨᖏᓪᓗᓀᑦ ᐅᖅᑲᐅᔪᖅᑲᖅᑕᖅᓯᓂᐊᖅᑐᑦ ᐊᓚᒪᓗ ᑐᐱᓕᐅᖅᑲᖅᑕᖅᑐᑦ ᖃᓄᐃᓕᓴᐅᑎᓗᒃ ᖃᓄᐃᓇᑕᐃᓕᓴᐅᓂᖅ ᐃᓕᓴᐅᑎᓂᖅᓯᓄᑦ ᑕᐃᒪᐃᓐᓇᖅᑐᖅ ᐊᓯᖏᓐᓂᓗ ᐱᔪᐊᑎᖁᖅᑐᓂᖅᓯᓄᑦ ᐊᐅᓚᕎᑎᐅᖅᑲᑕᐅᓂᖓᓪᓕᐊᑦ ᑲᑎᒪᔨᖏᓪᓗᓀᑦ ᐃᓕᓴᐅᑎ

7 ᐊᖏᑭᑕᐅᓯᒪᔨᕆᓂᑦ ᐱᓕᕆᐊᒃᓴᐅᑦ, ᐊᔪᖅᓯᖅᑎᑕᐅᔪᑦ ᐱᓕᕆᐊᕕᐅᔪᓐᓇᖅᓂᖓᔪᑦ ᐊᓘᓚᓗ ᐊᔪᖅᑎᑕᐅᔪᑦ ᑎᐳᓯᒪᕕᐅᔪᓐᓇᖅᖏᓐᓄᑦ

ᑲᑎᒪᐅᑕᐅᕙᒃᑐᓂ ᑎᖏᒥᐊᓕᕆᓂᖓᔪᑦ ᒪᓕᑦᓕᓕᐊᕆᔨᐅᕙᒃᑐᓂ ᒪᓕᑦᓕᐅᑕᐅᕙᓚᒪᑦ ᑐᖓᔫᓕᐅᑎᓪᓗᒍ ᐱᓕᕆᐊᖅᑐᑦ ᐊᖏᔪᒪᓂᒃᓴᖅ, ᐅᓪᓗᓂᑲᓇᖏᑦᓯᓗ, ᒪᓐᓂᖏᑦᓗ, ᓯᑲᑎᖅᑕᐅᕙᑲᑕᐅᓇᖅᑐᖏᓐᓂᖅᑐᖓᓂ, ᐸᖏᖅᑎᖅᑕᐅᕙᑲᑕᐅᓇᖅᑐᖏᓐᓂᖅᑐᖓᓂ, ᐊᓘᒪᓗ ᐃᓂᖅᑕᐅᕙᑲᑕᐅᓇᖅᑐᖏᓐᓂᖅᑐᖓᓂ. ᑕᐃᒃᑯᐊᓗ *ᑲᑎᒪᐅᑕᐅᕙᒃᑐᓂᓂ ᑎᖏᒥᐊᓕᕆᓂᖓᔪᑦ ᒪᓕᑦᓕᓴᐃᓕᕆᔨᐅᓂᖏᓐᓄᑦᑐᖓᓂ ᒪᓕᑦᓕᓐᓄᐃᓕᕆᔨᐅᕙᒃᑐᓂ* ᑐᑭᖏᓐᓄᓐᓂᖅᑕᐅᕙᒃᑐᖓᓂ ᐊᐃᑲᑕᐅᕙᓂᖅᑐᖓ ᓴᓇᔨᐅᔨᓕᖅᑐᖓ ᒪᓕᒃᓴᐅᕙᒃᑐᖓᓂ ᐊᐃᑲᑕᐅᕙᒃᑐᖓᓂ ᑲᑎᒪᐅᑕᐅᕙᒃᑐᖓᓂ *ᑲᑎᒪᐅᑕᐅᕙᒃᑐᓂ ᒪᓕᑦᓕᓕᐊᕆᔨᐅᕙᒃᑐᓂ, ᓴᖅᑭᑎᑎᓂᓗᑕᐅᕙᒃᑐᓂ* ᑲᑎᒪᐅᑕᐅᕙᒃᑐᓂ ᐊᐃᑲᑕᐅᕙᒃᑐᖓᓂ ᐊᔪᖅᑎ

- ᑭᓇᑐᐃᓇᖅ ᑎᖅᑎᕐᕋᓱᐊᕐᓂᖅ.
- ᑭᓇᑐᐃᓇᖅ ᓯᕆᔨᑎᑦᓴᕐᕕᖅ ᑎᒍᓱᖃᑦᑕᓗᐊᕐᕿᓂᓕᕆᓂ ᑎᖅᑎᐊᑦ ᐅᑦᓗᖅᑭᓂᖅ.
- ᑭᓇᑐᐃᓇᖅ <<ᕆᑎᓕᖅᓂᐊᖅᕈᓕᖅ ᐅᒪᓴᒥᒃ ᓯᓕ ᑎᖅᑎᐊᕆᒃ, ᑎᑎᕐᓚᓂᖅᓄᑦᓗᓂᓂ, ᐊᒥᖅᓚᓂᑦᓗᓂᓂ, ᐅᑦᓗᖅᓚᓂᑦᓂᓂ, ᒪᓂᖅᑭᓂᖅᓄᑦᓗᓂᓂ ᑎᖅᑎᐊᑦ.
- ᑭᓇᑐᐃᓇᖅ ᖃᓄᑭᐅᓱᔪᖅᑕᖅᓂᐊᖅᕈᓕᖅ ᐊᖓᔪᓇᖅᓂᕐᒪᔭᓂᓂ ᐊᐅᑎᓕᖅᒥᒃ ᐱᕆᔭᒃᓴᕐᕕᖅ ᐱᓯᒪᔪᖅᑐᓂ
- ᑭᓇᑐᐃᓇᖅ ᑎᒍᐊᓇᓯᒪᔪᒥᓂᒃ ᑲᑎᒪᔨᖅᑎᑦᓴᓂᓂ ᐃᐱᖅᓯᖅᑎᑎᕐᓂᖅᕈᓕᖅᓂᕐᕕᖅᓂᑦᖅ ᓇᒧᑐᐃᓇᑭᖅᑐᑦᖅ
- ᑭᓇᑐᐃᓇᖅ ᐱᓕᕆᔨᖅᕈᓕᖅᓂᐊᖅᕈᓕᖅ ᓯᖏᓪᓕᓇᖅᓂᕐᒪᔭᓂᓂ, ᒪᓂᖅᑭᓂᖅᓄᑦᓗᓂᓂ, ᐅᑦᓗᖅᑭᓂᖅᓄᑦᓗᓂᓂ, ᓇᔨᓴᕆᔨᖅᑕᖅᑭᓂᖅᓄᑦᓗᓂᓂ ᑎᖅᑎᐊᑦ ᓄᓇᐃᑦ.

7.1 ᐊᐅᓪᓚᕆᓚᓄᑦ ᐊᖏᕐᕋᑕᐅᓴᔪᑎᑦ

7.1.1 ᐊᖏᕐᕋᑕᐅᕐᕋᑕᕐᓂᖅᓂᑦ ᐱᔨᓂᓇᐅᑎᑎᑦ

ᐊᑐᕐᕋᑕᐅᓂᖏᓐᓂ *ᑎᖅᑎᐊᕐᕿᑲᓇᓚᓄᑦ ᒪᓕᑦᓴᑐᑐᐃᐃᑦ* ᐊᓪᓚᓗ ᐅᐸᑐᕐᕋᑕᕐᕿᑲᖏᓐᓇ, ᒥᓂᔅᑕᐅᔨᕐᓄ ᐊᐅᑎᓕᓪᓗᓂᔪᓕ ᐊᖏᕐᕈᓐᓇᖅᓱᕐᓄ ᐊᔨᕐᓇᖅᑎᑕᐅᔨᓂᖅ ᐱᓕᓪᓚᐃᔪᓇᓇᖅᑐᓂᖅ ᐊᑐᕐᕋᑕᐅᓄᑦᓗᓂᓂ *ᑎᖅᑎᐊᕐᕿᑲᓇᓚᓄᑦ ᒪᓕᑦᓴᑐᑐᐃᐃᑦ* ᑐᓂᓯᔪᓐᓇᖅᑐᓂ ᐊᖏᕐᕈᓐᓇᖅᑎᑕᐅᔨᓂᖅ ᐱᓕᓪᓚᐃᔪᓇᓇᖅᑐᓂᖅ ᐊᐅᑎᓕᓪᓗᓂᔪᓕ

ᒪᓕᑦᑲᕆᑎᑕᐅᓂᖅᓂ ᖃᓄᐃᓕᓐᓄᓴᖅᓱᕐᕆᑦᑲᕆᑎᑕᐅᓂᖅᓂᒃᓗ ᐊᐅᓚᑦᑎᓂᕐᒥᓂ ᐱᓕᕆᓱᖅᓯᐊᕐᑕᖅᓂᖅᓂᒃ, ᒥᓂᔨᑕᐅᑦ ᐃᓕᓴᕐᕕᓂᕐᑐᑎᑦᑲᕆᑎᑕᐅᓂᖅᓂᒃᑐᖅ ᐊᐅᑐᑦᑲᕆᑎᑕᐅᓕᓴᕐᕆᑦᑲᕆᑎᑦᓂᖅᓂᒃ ᐊᐅᑐᑦᑲᕆᑎᑕᐅᑐᓂᖅᓂᒃ ᐱᓕᕆᓱᖅᓯᐊᕐᑕᖅᓂᒃᑐᖅᓂᒃ ᐱᐅᓂᖅᓯᐊᕐᑕᖅᓂᒃᑐᖅᓂᒃ ᐊᐅᑐᑦᑲᕆᑎᑦᓂᖅᓂᒃᑐᖅᓂᒃ ᐱᐅᓂᖅᓯᐊᕐᑕᖅᓂᒃᑐᖅᓂᒃ ᐊᐅᑐᑦᑲᕆᑎᑦᓂᖅᓂᒃᑐᖅᓂᒃ ᐱᐅᓂᖅᓯᐊᕐᑕ

7.1.3 ᓄᓇᕗᒻᒥᐅᑦ ᐃᓄᐃᑦ ᐱᓕᕆᔨᒃᑲᓐᓂᑦ ᑭᓱᑕᒥ ᐱᔪᒪᓇᐅᑎᖅᑕᑎᒃ

ᑭᓇᐅᔨᓕᕆᔨᑕᑎᑦᓴᓂᒃ ᐱᓕᕆᔨᖏᓚᐅᔪᑦ ᐱᓕᕆᓴᓪᓕᐅᕕᒃᓴᓪᓕᐅᔪᑦ ᐱᓕᕆᔨᒃᑲᓐᓂᑦ ᑎᒃᓕᒋᐊᖅᓴᖅᓗᒃ ᓄᓇᕗᒻᒥ ᐃᓄᒃᓄᑦ ᓇᒃᓕᒋᓂᖅᓴᐃᔨᔪᓂᕐᓄᓪᓗᓂ. ᑕᐃᒪᑦᑕᓪᓕᐊᖅᑲᖅᑎᑕᐅᓚᐅᖅᓯᓂᑦ ᑭᓱᑕᒥ ᐊᐅᓪᓚᕐᓴᐅᑦᑕᖅᑕᑦ ᐊᖏᓇᓂᕐᒋᐊᕐᓴᓪᓗᑎᒃ ᐊᖏᓇᓂᑎᓂᒃ ᐃᖅᑲᓪᓕᐊᖅᓂᕐᓴᓪᓗᓂᓪᓚᓂ. ᓄᓇᕗᒻᒥ ᐃᓄᐃᑦ ᐊᐅᓪᓚᕐᓴᐅᖅᑕᑦ ᐱᔪᒪᓇᐅᑎᖅᑕᓪᓕᐊᖅᑲᖅᓯᑎᒃ ᑭᓱᑕᒥ ᐃᓄᓪᓕᓇᐅᔨᓂᑦ ᐊᖏᓇᓂᖏᑎᒋᐊᖅᓯᓂᑦ, ᐃᖅᑲᓪᓕᐊᖅᑎᑕᐅᔨᓪᓚᓂᓪᓂ ᐊᐅᓪᓚᕐᓴᐅᑦᑕᖅᑕᑦ ᐊᖏᓇᓂᑎᓂᒃ ᐊᖏᓇᓂᖏᑎᒋᐊᖅᓯᓂᑦ, ᐃᖅᑲᓪᓕᐊᖅᑎᑕᐅᔨᓂᑦ ᐊᐅᓪᓚᕐᓴᐅᑦᑕᖅᑕᑦ ᐱᔪᒪᓇᐅᑎᖅᑕᓪᓕᐊᖅᑲᖅᓯᑎᒃ ᐃᓄᓪᓕᓇᐅᔨᓂᑦ ᐊᖏᓇᓂᖏᑎᒋᐊᖅᓯᓂᑦ, ᐃᖅᑲᓪᓕᐊᖅᑎᑕᐅᔨᓪᓚᓂᓪᓂ ᐊᐅᓪᓚᕐᓴᐅᑦᑕᖅᑕᑦ ᐊᖏᓇᓂᖏᑎᒋᐊᖅᓯᓂᑦ (ᓄᓇᓗ ᐊᐅᑎᒥᒃ ᑕᑯᓂᐊᖅᑎᓂᓂᒥᒃ, ᖃᐅᔨᒪᐊᓕᕐᓂᔨᓂᒥᒃ, ᐊᓄᖏᓐᓂᓪᓗ) ᓄᓇᕗᒻᒥ ᐃᓄᓪᓂᒃ ᐱᔪᒪᓇᐅᑎᖅᑕᒪᐊᖅᑲᐅᓇᖅᓯᓂ. ᑐᑭᓄᓂᐊᖏᓚᐅᑎᑕᑦ ᐃᑐᓂᖏᑦ ᑎᑎᕋᖅᑕᖏᓐᓄᑦ 7.1.7-ᒥ ᑐᑭᓄᔨᕋᑎᐊᐳᖅᓂᐅᑲᓇᕐᓂᕐᒧᑦ ᐱᔪᒪᓇᐅᑎᓕᕆᔨᓂᒃ ᐱᓕᕆᓂᖅᑲᖅᑐᑎᓂᒃ.

7.1.4 ᐃᓄᓕᕆᐅᖏᓕᑐᓄᑦ ᐱᓕᕆᔨᖅᑭᑲᓇᕐᓂᖏᓕᑦ ᐱᔪᒪᓇᐅᑎᖅᖏᓕᑐᓄᑎᒃ

ᑕᒪᐃᓐᓂᕕᐊᕋᖅ ᐱᓕᕆᔨᖓᔪᓐᓇᖅᑐᑦ, ᐃᓕᓴᖅᓴᓄᓕᓗ ᑕᐃᐳᑯᓂᖓᓕ ᓄᓇᒧᑦ ᑕᐅᑭᑐᖓᓕ ᑎᑎᕗᓕᔪᓐᓇᖅᑐᓂᖅᑐᑦ ᐊᒻᒪᓗ ᓄᓇᕗᒻᒥᓂᐅᓪᓚᕆᐊᖃᖅᑎᑦ ᑲᔪᓯᑎᑕᐅᓇᓱᑦᑎᒃ ᐱᔪᒪᓇᖅᑎᑕᐅᔪᓂᒃ, ᐱᔪᒪᓇᐅᑎᓴᖅᓯᐊᕐᓴᐅᑦᑕᖅᑐᑦᓴᓕᐅᕐᓂᑦ ᓴᓕ ᐃᓄᓕᓕᖅᑐᖓᓕᑦ, ᑎᓪᓗᐊᑦ ᓂᕆᐳᓴᓕᖅᓯᐅᒋᔭᖅᑐᑦ ᐃᓄᐃᑦ ᓄᓇᕗᒻᒥᓂᕐᓂᖅᓱᓕᒥᒃ (ᑕᒪᓕᓂ ᐃᓕᓴᐅᒃ ᐊᐅᔪᐃᓕᖅᓱᑦ ᑎᑎᕗᓕᔪᓐᓇᖅᑐᓂᖅᑐᑦ, ᐊᓄᖏᓐᓂᓪᓗᓐᓂ) ᓄᓇᕗᒻᒥ ᐃᓄᓪᓂᒃ ᐱᔪᒪᓇᐅᑎᖅᑕᒪᐊᖅᑲᐅᓇᖅᓯᓂ. ᑐᑭᓄᓂᐊᖏᓚᐅᑎᑕᑦ ᐃᑐᓂᖏᑦ ᑎᑎᕋᖅᑕᖏᓐᓄᑦ 7.1.7-ᒥ ᑐᑭᓄᔨᕋᑎᐊᐳᖅᓂᐅᑲᓇᕐᓂᕐᒧᑦ ᐱᔪᒪᓇᐅᑎᓕᕆᔨᓂᒃ ᐱᓕᕆᓂᖅᑲᖅᑐᑎᓂᒃ.


7.1.5 ᐃᓄᓕᕆᐅᖏᓕᑐᓄᑦ ᐱᔪᒪᓇᐅᑎᓴᓪᓕᐅᖏᓕᑦ ᐊᖏᓂᖅᑕᐅᔨᓪᓚᕆᐊᖃᖅᑐᑦ ᑭᓱᑕᒥ ᐱᔪᒪᓇᐅᑎᖅᓴᓕᐅᕐᓂᑎᒃ

ᐃᓄᓕᕆᐅᖏᓕᑐᓄᑦ ᐱᔪᒪᓇᐅᑎᖅᑕᒪᐊᖅᑲᖅᓯᓂᑦ ᓴᐃᑭᐅᔨᓱᑲᓇᕐᓂᕐᒧᑦ ᑎᓪᓗᐊᑦᑕᖅᑲᓇᕐᓂᑦ. ᐃᓄᓕᕆᐅᖏᓕᑐᓄᑦᑕᐅᓕᑦ ᐱᔪᒪᓇᐅᑎᓕᕆᔨᖅᑕᒪᐊᖅᑲᐅᓇᖅᓯᓂᑦ ᓴᐃᑭᑕᒪᐊᖅᑲᖅᓴᓕᐅᕐᓂᑦ ᐊᒻᒪᓗ <<ᓕᕐᑎᐱᓪᓇᖅᓂᖅᑐᑦ ᑐᖅᐃᔨᑕᒥᓂᒃ ᑎᓪᓗᐊᑦᓂᒃ. ᑕᒪᓐᓇ ᐱᓴᖅᓱᐅᓴᓂᑕᕐᓄᓪᓚᖅᑐᑦ ᐃᓄᓕᕆᐅᖏᓕᑐᓄᑦ-ᐊᖏᓇᓂᕐᒋᐊᖅᑲᖅᓱᐅᓱᓂᓪᓚᒃ ᐅᓪᓗᓕᒫᖅᑕᐅᓂᓪᓚᒃ (ᐊᖏᓇᓂᕐᒋᐊᖅᑲᖅᓯᓂᖅ ᐃᓄᓕᓕᐅᑲᓪᓚᖅᑐᑦ ᓄᓇᕗᒻᒥᑕᐅᔨᓪᓚᓂ, ᐱᔪᒪᓇᐅᑎᖅᓴᖅᑎᑎᓪᓗᓐᓂᓪᓂ ᐃᓄᓪᓂᒃ, ᐱᔪᒪᓇᐅᑎᖅᓴᖅᑎᑎᓪᓗᓐᓂᓪᓂ ᐃᓄᓪᓂᒃ, ᐊᖏᓇᓂᖏᑎᒋᐊᖅᓯᓂᑦ ᐃᓄᓪᓂᒃ *5.3.1-ᒥ*; ᐊᐃᔨᑐᖅᓴᓕᓂᖅᑐᓂᒃ 7.1.2). ᑐᑭᓄᓂᐊᖏᓚᐅᑎᑕᑦ ᑎᑎᕋᖅᑕᖏᓐᓄᑦ 7.1.7-ᖅᓴᓂ ᑕᖅᓂᒃ ᑐᑭᓄᔨᕋᑎᐊᐳᖅᓂᐅᑲᓇᕐᓂᕐᒧᑦ ᐱᔪᒪᓇᐅᑎᓕᕆᔨᓂᒃ.

7.1.6 ᐱᓕᕆᔨᓐᓇᖅᑐᖅ ᐊᖏᖅᑕᐅᓂᒪᓕᒃᑯᓂᖓᓪᓗᓂ ᐃᓄᖓᓐᓄᑦ ᐱᖁᑎᑎᕿᐅᔪᓂ ᓄᓇᓄᑦ

ᑕᐃᒃᑯᐊ ᑲᑎᒪᔨᑕᐅᔭᖅᑎᓂᓄᑦ ᑎᖏᐊᓕᕆᓂᖓᓪᓗᓂ ᒪᓕᕆᐃᑦ, ᐊᒪᓚᓗ ᒪᓕᕆᑦᓗᑐᓇᕗᖅᑎᑦ ᐊᑐᖅᑕᐅᑉᓂ ᐃᓄᖓᓄᑦ ᐱᖁᑎᑎᕿᐅᔪᓂᖓᓄᑦ ᓄᓇᓄᑦ. ᒥᓂᑕᐅᔪᖅ ᐊᐅᑎᓕᕆᓇᕐᓂᐊᒃᑯᓐᓄᑦ ᑐᓂᓯᒪᔪᓐᓇᖅᑐᑦ ᐱᔪᓐᓇᑑᑎᒃ ᐃᓄᖓᓕᕆᔨᓐᓇᖅᑐᑦ ᐃᓄᖓᓄᑦ, ᑐᐱᓯᓂᐊᕐᑕᐅᓕᕆᔨᑐᓇᑎᒃ ᑭᐅᓂᒃᑯᐊᑎ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑎᑦᑐᒥᒃ ᐊᑐᖅᑕᐅᑉᓯᐃᓐᓇᖅᑐᑦ ᐃᓄᖓᓄᑦ, ᑐᐱᓯᓂᐊᕐᑕᐅᓕᕆᔨᑐᓇᑎᒃ ᑐᓴᕐᓯᒪᓕᕆᓂᖓᓂᒃ ᐱᐅᓯᕆᔨᖅᑐᑦ ᐃᓄᖓᓄᑦ, ᑭᓇᐅᔨᒪᔨᓕᕆᔨᖅᑐᑦ ᐃᓄᖓᓄᑦ ᐱᖁᑎᑎᕿᐅᔪᓂᖓᓄᑦ, ᑭᓇᐅᔨᒪᔨᓕᕆᔨᖅᑐᑦ ᐃᓄᖓᓄᑦ ᐱᖁᑎᑎᕿᐅᔪᓂᖓᓄᑦ, ᑭᓇᐅᔨᒪᔨᓕᕆᔨᖅᑐᑦ ᐃᓄᖓᓄᑦ ᐱᖁᑎᑎᕿᐅᔪᓂᖓᓄᑦ, ᐱᖁᑎᑎᕿᐅᔪᓂᖓᓄ

ᑲᓇᑕᒥ ᐅᒪᓯᓕᓗᓴᖅᑯᑦ ᑐᓂᓯᔪᓐᓇᐃᖓᑎᓕᑦ ᐱᔪᓐᓇᐅᑎᓂᒃ ᐱᔪᓐᓇᐅᑎᑕᕋᓯᐊᖅᑐᒍᓄᑦ ᐅᒃᑐᖅᓯᒪᔨᒍᓄᑦ, ᑭᓯᐊᓂ ᐱᐅᔨᓂᒃ ᒪᓕᕐᑎᐊᖅᓂᖅᑕᐅᓯᒪᔨᓕᕐᖅᑐᓂᒃ ᑐᑭᓕᐅᖅᑕᐅᓯᒪᓕᕐᖅᑐᓂᒃ ᑐᓂᓴᐅᓯᒪᔨᔪᓕᕐᓗᓐᓂᑦ ᖃᐅᔨᓴᖅᑕᐅᔾᔨᓇᐊᖅᓱᖓᓂᖅᖃᖅᑕᐅᔪᑎᒃ ᑭᓯᐊᓂ, ᐸᖅᑭᖅᓯᒪᓗᑎᒃ ᐱᓕᕐᓇᐊᖅᓱᖓᓂᖅᖃᖅᑕᐅᓗᑎᒃ. ᐊᓯᒪᑕᐅᖅ ᓯᓕ, ᐅᖅᑕᐅᔾᔨᒋᐊᖅᑎᐅᖅᑕᖅᑕᒪᒡᒪᑦ ᒥᓂᑕᒧᑦ ᑕᒪᐃᓐᖅᑐᓂ ᐱᔾᔪᑎᖅᑐᖅᑐᓂ <ᖁᓇᐅᑎᓕᐊᓇᔾᐅᖅᓱᖅᑐᓂ ᐊᓯᒪᓗ ᐊᐅᓕᑕᐅᖅᓱᖓᓖᓐᓂᓐᓄᓂ ᑎᓐᓕᐊᖅᑯᖅᓴᐃᓐ. ᐃᖅᓂᐅᖅᓴᓴᐅᔪᑕᐅᖅᑲ ᐊᐅᓕᓪᑎᓴᐅᖅᑲᑕᐅᓂᒧᑦ ᑲᑎᒪᓐᓇᖅᑕᐅᓂᒃ ᑕᒪᐃᓐᖅᑐᓂ ᖃᐅᔨᓴᖅᑕᐅᔾᔨᓇᐊᖅᓱᖓᓂᖅᖃᖅᑕᐅᓗᑎᒃ ᐊᓯᒪᓗ ᑐᓂᓴᐅᓯᒪᔨᔪᓕᕐᓗᓐᓂᑦ ᑲᑎᒪᓐᓇᖅᑕᐅᓗᑎᒃ ᐱᓕᕐᓇᐊᖅᓱᖓᓂᖅᖃᖅᑕᐅᔪᑎᒃ ᐊᓯᒪᓗ ᑐᓂᓴᐅᓯᒪᔨᔪᓕᕐᓗᓐᓂᑦ ᑲᑎᒪᓐᓇᖅᑕᐅᓗ


ᐱᓕᕆᓚᐅᑎᒋᔭᐃᓂᖓᓂᒃ
ᐅᖃᖅᖢᖅ
ᓄᓇᕗᒻ ᓄᓇᕗᑦ ᐊᑐᑎᒃᓴᖅᑭᓐᓄᑦ ᑲᒥᔪᓇᕗᑦ
a) ᖃᐅᔨᓴᖅᓯᖅᑕ ᓄᓇᕗᑦ ᐊᑐᑎᒃᓴᖅᑭᓐᓄᑦ
ᐸᖅᓇᐅᔪᐊᑦ ᒪᓕᒐᓴᐅᔪᓂᖅ ᒪᓕᖅᓴᓗᒍ
b) ᐱᓕᕆᓚᐅᔪᓇᖅᑐᑎᒃ ᐊᑐᖅᑯᓕᔪᓇᖅᓂᖅᓴᒪᒥᑦ
ᖃᐅᔨᓴᖅᓯᖅᑕᓄᓇᕗᒻ ᓄᓇᕗᑦ
ᐊᑐᖅᖢᑕᐅᔪᓇᖅᓯᖅᑭᓐᓄᑦ ᑲᑎᒪᔨᑦ ᐊᕐᕌᒍᓂᖅᓯᒪᔭᑦ
ᖃᐅᔨᓴᖅᓯᖅᑕ ᐱᓕᒻᒪᕐᓴᒪᓂᖅᑕ
ᐅᑭᐅᖃᖅᑐᑦ ᑲᑎᒪᔨᑦ
ᐊᑐᖅᖢᑕᐅᔪᓇᖅᓯᖅᑭᓐᓄᑦ ᑲᑎᒪᔨᑦ
ᖃᐅᔨᓴᖅᓯᖅᑕ
ᐱᓕᕆᓚᐅᑎᒋᔭᐃᓂᖓᓂᒃ
ᐊᐅᓚᑦᑐᖅᑕ
ᐱᓕᕆᓚᐅᑎᒋᔭᐃᓂᖓᓂᒃ
ᑲᑎᒪᔨᖅᑕᐅᓂᖅ
ᐅᖃᖅᖢᑕ ᐅᒪᔨᑦᑐᕐᓴᐅᑦ
(ᐅᒪᔨᑦᑐᕐᓴᐅᑦ)
ᑐᓂᓗᐅᕐᓂᑦ
ᐱᓕᕆᓚᐅᑎᒋᔭᐃᓂᖓᓂᒃᐊᑐᖅᑯᓕᔪᓇᖅᓯᖅᑭᓐᓄᑦ
ᐅᖃᖅᑐᑎᒃ
ᐊᐅᓚᑦᑐᖅᑕ
ᐊᐅᓚᑦᑐᖅᑕᐅᓂᖅᓴᐅᑦ
ᐊᐅᓚᑦᑐᖅᑕᐅᓂᖅᓴᐅᑦ
ᐱᓕᕆᓚᐅᑎᒋᔭᐃᓂᖓᓂᒃ

8 ᓄᓇᖅᑐᑦ ᑎᖅᑎᐊᖏᖅᓯᐅᔨ ᖃᓄᐃᓇᑕᐅᖏᓐᓄᖅᓯᐅᓂᖅᑐᖅ ᑕᐃᓴᐅᓂᖅᑐᑦ ᐃᓕᑕᒪᓴᐅᓕᓂᖅᑐᓪᓗ

ᐃᑲᑦᑐᐊᖅ ᐊᓐᓇᐅᔨᖅ ᐃᓕᑕᒪᓴᐅᓕᒪᓕᖅ ᑎᖅᑎᐊᖏᖅᓯᐅᓂᖅᑎᑲ; ᑭᓕᐊᓂᓕ, ᐊᑦᕿᔪᓕᒥᓴᖅ ᓴᓐᓂᐊᑦᖅᑕᐅᕐᓇᖅᓯᐅᓂ ᐅᓴᔨᓂᔪᑎᖅᓂᓂᓄᑦ ᓄᓇᒥᐅᑕᒪᓴᐅᔨᓂᓕᓪᓗ ᐱᑐᖅᐅᔨᓂᑦ ᑲᓇᑕᓕᒥᒥ ᐅᓴᔨᖅᑲᓇᕐᓂᐅᑐᓂᓂ ᐱᐅᓂᖅᓴᐃᐅᓇᖅᓴᖅᓂᖅ.

ᓯᕗᓕᓂᖅᑎ ᑕᐃᓴᓕᓴᓂ ᑎᑦᓂᐊᖅᑕᐅᖓᓴᑦ ᑎᖅᑎᐊᖏᖅᓯᐅᓂᖅᑐᑦ ᓱᓴᓕᓚᐅᖅᓴᑦ ᐅᓴᔨᐃᑦ ᓴᓐᓂᐊᑦᖅᑕᐅᑎᖅᓂᐅᖅᓄᑦ, ᑲᑎᒪᔨᓄᑦ ᐃᓚᖓᑕᐅᖅᓯᓂᖅᓄᑦ ᐱᓕᕆᔨᑦ ᐃᓕᓴᐃᑦ ᐃᓕᓴᐅᓂᖅᑐᑦ ᐊᑐᓂᖅᑐᑦ, ᐊᓂᖅᓄᑦ ᐃᓕᓴᖅᑎᖅᓯᐅᔨᓕᖅᑐᑦ ᑕᐅᖅᓯᓂᖅᑐᑦ ᐊᑐᓂᖅᑐᑦ ᐊᑎᓪᓗ ᐃᓕᓴᐃᑦ ᐃᓕᓴᐅᓂᖅᑐᑦ ᐊᑎᓪᓗ ᐃᓚᖓᑕᐅᖅᓯᓂᖅᑐᑦ ᐊᑐᓂᖅᑐ

ᐊᐃᔭᑐᖏᓪᓗᓯᓂᔭ ᑕᔨᑎᕐᑎᓴᓪᓕ 8: ᒪᓕᑌᓴᐅᐅᓪᓕ ᖃᔨᓂᐃᓕᓕᖏᖏᓂᕐᒧᓂ ᑎᑦᔨᐊᖏᑕᐅᓯᒪᓂᕐᒧᓂᐊᕐᓂ ᐃᑦᑐᑐᖏᖅ ᐃᓐᑕᓕᖏᖅ ᑲᓇᑕᓕᒥᒃ ᑎᕐᒥᐊᖏᖃᐃᐅᓂᓪᓗᑦ.

ᒪᓕᑌᐃᑦ ᖃᔨᐃᓕᓕᖏᖏᓂᕐᒧᓂᓪᓗᑦ ᑲᓇᑕᓕᒥᒃ ᐅᒪᔪᖏᖏᓂᕐᒧᓂᓪᓗᑦ (ᐊᕐᕈᑎᑕᑌᓵᓂᔪᐊᖅᑯᑦ 2017b)	ᐃᐃᑦᑐᐊᑦ ᖃᐅᐅᓵᖏᖏᑕᐅᓯᒪᓂᕐᓗ ᖃᔨᐃᓕᓕᖏᓂᕐᓗᓂᕐᓗ ᒪᓕᑌᓕᑌᓴᐅᔪᓂᖅ ᒪᓕᖏᓕᕐᓂᖅ
1. ᑕᒪᐅᓐ ᓄᓇᖏᓗ ᐱᑕᓕᖅ ᒥᑭᓂᖅᑰᖓᓂᖏᑎᑕᐅᔪᒥᖅ 1% >ᓵᖑᑎᖏᑉᓂᑐᓂᖅ ᑲᓇᑕᒪᔨ ᐅᒪᔪᑯᑎᖏᓐᓂᖅ ᐊᐳᐅᓵᕐᖏᖏᓂᓗᓐᓂᑦ ᐅᒪᔪᓂᖅ ᑎᑭᐸᖅᑐᓂᖅ ᑎᖏᒥᐊᖏᓂᖅ ᐅᐊᖓᓐᓂᑦ ᑕᐃᖅᑯᓂᖏᓗ ᐅᒪᔪᓂᖅ ᐊᒥᔨᖏᑎᓂᓐᓂᕐᔪᒪᔪᓂ ᐱᑕᖏᓂᖅ[1].	✓ ᐅᖏᓚᑉᓂᑐᑦ 5% >ᓵᖑᑎᓂᖅ ᑲᓇᑕᒥᐅᓂ ᑲᖏᔫᑦ ᑕᐅᐊᓂ ᐅᓪᓗᖅᑯᔨᑦ ᐃᓗᐊᑦ ᑎᖏᒥᐊᖏᓂᕐᓗᐊᕐᓂ ✓ ᐅᖏᓚᑉᓂᑐᑦ 1% >ᓵᖑᑎᓂᖅ ᓂᓪᓕᓂᓇᐃᑦ (Atlantic brant) ᐅᓪᓗᖅᑭᖅᓇᓪᓂᖏᑦ ᑎᖏᒥᐊᖏᓇᖏᒥ, ᒪᓕᖅᑕᐅᓪᓗᑎᖅ ᑕᐃᒪᖏᓂᖅ ᖃᐅᐅᓵᕐᖏᖏᓂᖅᑕᖅᑯᓂᖏᑦ
2. ᑕᒪᐅᓐ ᓄᓇᖏᓗ ᐊᒥᔨᐊᓗᖏᓂᖅ[2] ᐱᑕᖏᖏᓂᕐᒪᔨᖅ ᑎᖏᒥᐊᖏᓂᖅ ᑎᖏᒥᐊᓪᓗ ᐊᐳᐅᒪᓕᒋᐅᖏᑦᑐᖏᓂᖅ ᐅᐸᐅᓗᓐᓂᑦ ᐅᒪᔪᓂᖅ ᐊᒥᔨᖏᓂᓗᓐᓂᕐᖏᓗᓂᖅ ᐱᑕᖏᓂᖅ ᐊᑐᔨᓂᐅᐅᖏᖅᑕᕐᓗᐊᖏᓗᒃ ᐊᒥᔨᓂᖓᓴᐅᐸᐅᓗᓐᓂᓗᓐᓂᑦ ᑕᒪᔪᓂᖏᓗ ᐅᒪᔪᓂᖅ, ᐊᒥᔨᓂᓪᓕᓐᓂᑦ ᖃᐅᔨᓂᖅᑕᐅᓕᖅᑐᓂᑦ, ᐱᑕᖏᓂᖅᑯᓂᖏᓗ ᐅᒪᔪᓂᖅ ᐊᒥᔨᓂᖓᓴᐅᐸᐅᓗᓐᓂᓗᓐᓂᑦ ᑕᒪᔪᓂᖏᓗ ᐅᒪᔪᓂᖅ.	✓ ᐱᖏᓚᔨᖏᖅᑐᑦ 6-ᖃᓗᔪᑦ ᐅᒪᔪᐃᑦ ᐊᒥᔨᓂᖓᓴᐅᓗᓐᓂᕐᓂᓪᓗᑦ ᑎᑎᕐᖏᖅᑕᐅᔨᒪᓕᒋᐅᖏᑦᑐᖏᓂᖅ ᐊᑕᔪᑦ ᐅᒪᔪᐃᑦ ᐊᒥᔨᓂᖓᓴᐅᓗᓐᓂᕐᓂᓪᓗᖓᓂᖅᑯᑦ ᒪᓕᑌᓴᕐᓂᖅᓯᔪᑦ, ᐊᓚᒪᓗ ᔨᑕᒪᐅᖏᑎᒋᔪᖅ COSEWIC-ᓄᑦ ᑲᓇᑕᒥ ᑲᑎᑕᐅᓪᓗᖅᑕᐅᔨᒪᓕᒋᐅᖏᑦᑐᖏᓂᖅ ᐊᒥᔨᓂᖓᓴᐅᓗᓐᓂᕐᓂᓪᓗᖓᓂᖅᑯᑦ ᐊᒥᔨᔪᖏᓂᖅᑕᖅᑭᓕᕐᓂᐊᕐᓂᓪᓗᖅᑕᕐᑕᐅᔨᒪᓕᒋᐅᖏᑦᑐᖏᓂᖅ ᐅᒪᔪᐃᑦ ᐊᓚᒪᓗ ᔨᑕᒪ ᑕᐃᖅᑯᓄ ᔨᑕᒪᐅᖏᑎᒋᔪᖅ 7 COSEWIC-ᓄᑦ ᐊᒥᔨᓂᖓᓴᐅᓗᓐᓂᕐᖏᖏᓂᖅ ᑲᑎᑕᐅᓪᓗᖅᑕᐅᔨᒪᓕᒋᐅᖏᑦᑐ

ᒪᓕᑦᑐᐃᓕ ᖃᓄᐃᓕᓕᒫᓂᖃᖅᖢᓂᕐᓗᒍ ᑲᓇᑕᓕᒥ ᐅᒪᔨᖃᖅᖢᓂᕐᒪᑦᑐᕐᓂᕐᓗᒍ (ᐊᐅᑎᓕᓚᔨᔨᐊᕆᔭᖅ 2017b)	ᐃᑲᑐᑐᐊᑦ ᖃᐅᔨᔪᖅᑕᐅᓂᒪᓂᖅ ᖃᓄᐃᓕᓕᖅᓂᖅᓂᑦ ᒪᓕᖅᓕᓚᕆᐅᔪᓂᒃ ᒪᓕᖅᓯᒪᓂᖅ
4. ᑕᒪᐅᓇ ᓄᓇᖅᑐ ᐊᐱᔨᖅᖢᕈᓕᒃ ᐅᒪᔨᖃᖅᖢᖏᐅᔨᓇᓂᕐᓗᒍ ᐃᓕᔨᓄᑦ ᐱᒋᐊᓂᓄᑦ ᐅᒪᔨᓄᑦ ᐱᓕᕆᓂᖅᑐᑦ ᐊᓂᔪᓂᒃᑯᑦ ᐊᐱᔨᖅᖢᕈᓕᒃ ᐱᒋᐊᓂᖅᑐᑦ ᐊᐱᔨᖅᖢᕈᓕᒃ ᐊᐱᔨᖅᖢᕈᓕᒃ ᐊᐱᔨᖅᖢᕈᓕᒃ

ᐃᓕᖅᑕᕐᓂᑦ ᑕᒪᐅᐸᐊ ᖃᐅᓪᓗ ᒪᓕᑕᒫᐅᔾᔪᑎᑦ ᐅᒪᔨᖅᖃᓕᓕᐅᕈᓐᓇᕐᓂᖅᑕᖅᑐᓂᓕᒫᑦ, ᑕᒪᓐᓇ ᐃᑲᑦᑐᐊᑦ ᑎᓪᓗᐊᖅᖃᓕᓚᖅ ᐊᐅᓂᐊᓗᑐᑦ ᑲᒪᕕᑎᖅᑕᖅᓯᒪᕐᓂᖅ ᖃᐅᔨᔪᖅᑕᐅᖅᑕᖅᑕᖅᑐᓂᓕᒫᑦ, ᐱᓗᐊᖅᑐᒥᑦ ᐱᔨᔪᑎᖅᖃᑐᓂᖅ ᑲᖏᖅᓯᒪᔪᓂᖅ ᐊᖏᔪᓂᖅᑕᐅᖅᑕᖅᑕᖅᑐᓂᓕᒫᑦ ᐊᑐᖅᑕᐅᓯᒪᓕᑦ ᑎᓪᓗᐊᖅᖃᓕᓚᖅ, ᐊᑐᐃᓐᓇᐅᓕᖅᑐᑦ ᐱᔨᔪᑎᖅᖃᑐᓂᖅ ᐊᐅᓂᐊᓗᑐᑦ ᐊᑐᖅᑕᐅᓯᒪᓕᑦ ᑎᓪᓗᐊᖅᖃᓕᓚᖅ ᑎᑎᕋᖅᑕᐅᓯᒪᔪᓂᖅ ᐊᐅᓂᐊᓗᑐᑦ ᖃᐅᔨᓴᐊᖅᑕᐅᖅᑕᖅᑕᖅᑐᓂᓕᒫᑦ ᖃᐅᔨᓴᕐᔪᐊᑦᑐᖅ ᐊᑎᒪᓗ ᑎᑎᕋᖅᑕᖅᑕᐅᓯᒪᔪᓂᖅ ᖃᐅᔨᓴᕐᔪᐊᑦᑐᖅ ᖃᐅᔨᓴᖅᑕᖅᑕᖅᑐᓂᓕᒫᑦ ᖃᐅᔨᓴᕐᔪᐊᑦᑐᖅ ᖃᐅᔨᓴᖅᑕᖅᑕᖅᑐᓂᓕᒫᑦ.

ᐊᓚᖅᑐᖅᓯᒪᔪᓂᖅ ᑕᒧᑎᑕᑎᕐᔪᑦ 9: ᓇᐃᓐᓴᖅᑕᐅᓯᒪᔪᑦ ᖃᐅᔨᔪᖅᑎᑎᐅᒪᐅᖅᑐᑦ ᓇᐅᑦᑎᖅᑐᐃᔾᔪᖅᑐᑦᑕᖅᑕᖅᑕᖅᓯᒪᔪᖏᓪᓗ ᑎᓪᓗᐊᖅᖃᓚᖅᓴᖅ

ᐊᖏᔪᒍᒥ	ᖃᐅᔨᔪᖅᑎᑦ ᐊᑎᑭᓯᖏᑦ (ᑎᒥᖏᓕᑎᖏᑦᓴᓂᓐᓂᑦ)	ᐱᔨᔪᑎᖅᖃᖅᑐᓂᖅ
1934	Ꮮᓂᒃ - Manning (bᓇᑕᒥ ᓚᓇᑕᓃᔨᑦ bᑐᔨᔪᖅᑎᑎᖏᑦ ᐊᓕᒪᓗ >ᓃᑕᐢᒥᐅᑦ ᑕᓀᔭᑦᖃᖅᖢᓕᑕᓃᔨᖏᑦ)	ᑎᓪᓗᐊᑕᓃᓂᖅ/ᓚᓇᒧᐃᓚᐅᑦᓴᑎᖅ
1936	>ᓰ, Ꮮᓂᒃ - Bray, Manning (bᓇᑕᒥ ᓚᓇᑕᓃᔨᑦ bᑐᔨᔪᖅᑎᑎᖏᑦ ᐊᓕᒪᓗ >ᓃᑕᐢᒥᐅᑦ ᑕᓀᔭᑦᖃᖅᖢᓕᑕᓃᔨᖏᑦ)	ᑎᓪᓗᐊᑕᓃᓂᖅ/ᓚᓇᒧᐃᓚᐅᓴᑎᖅ
1952-1953	ᑯᑦᔪ - Cooch (bᓇᑕᒥᐅᑕᓂ ᐅᒪᔨᑕᓃᔨᑦ, ᐊᐸᑎᑕᓃᔨᖏᔨᐊᖅᑯᑦ)	ᖃᐅᔨᔪᖅᓂᖅ – bᖓᖅᓂᖅ
1953, 1956-1957	<ᐅᓃ - Barry (bᓇᑕᒥᐅᑕᓂ ᐅᒪᔨᑕᓃᔨᑦ, ᐊᐸᑎᑕᓃᔨᖏᔨᐊᖅᑯᑦ)	ᖃᐅᔨᔪᖅᓂᖅ – ᓂᖅᑕᖅᓇᖅᓂᖅ (Brant)
1961	Lᑭᓂᔪ - MacInnes (ᔨᑐᓂᐸᓕᑎᖓ ᐱᖓᑎᓇᖓᓂᓂ ᐊᑲᑎᐅᓃᔨᑦ)	ᖃᐅᔨᔪᖅᓂᖅ – ᓂᖅᑕᖅᓇᖅᓂᖅ (Cackling)
1966	ᑯᐱᔪ - Kerbes (bᓇᑕᒥᐅᑕᓂ ᐅᒪᔨᑕᓃᔨᑦ, ᐊᐸᑎᑕᓃᔨᖏᔨᐊᖅᑯᑦ)	ᖃᐅᔨᔪᖅᓂᖅ ᐃᒪᖅᒥᓕᐊᑐᑕ ᑎᓪᓗᐊᖅᑐᖅ
1972-1973	ᑯᐱᔪ - Kerbes (bᓇᑕᒥᐅᑕᓂ ᐅᒪᔨᑕᓃᔨᑦ, ᐊᐸᑎᑕᓃᔨᖏᔨᐊᖅᑯᑦ)	ᐊᓪᓚᑕᐅᖅᑕᐅᓂᖓᓐᓄᑦ bᖓᖅᑲᖅᓴᐃᑦ
1979	ᓰᑦ - Reed, ↄ>ᐊ - Dupuis, (bᓇᑕᒥᐅᑕᓂ ᐅᒪᔨᑕᓃᔨᑦ, ᐊᐸᑎᑕᓃᔨᖏᔨᐊᖅᑯᑦ)	ᐊᓪᓚᑕᐅᖅᑕᐅᓂᖓᓐᓄᑦ bᖓᖅᑲᖅᓴᐃᑦ
1984	Lᖅᓂᐅᑦ - MacNeil (ᒥᖅᖠᐃᖅᓯᖓᓚᕐᓃᖅᑯᑦ bᓇᑕᒥ)	ᖃᐅᔨᔪᖅᓂᖅ ᓚᓇᒥᖅ
1987	ᖃᐅᔨᔪᓕᐅᖅᓂᑕᑦ	ᖃᐅᔨᔪᖅᓂᖅ ᓚᓇᒥᐅᑕᓂᖅ
1991-1995	ᑲᔪᖓᐅᑦ - Caswell (bᓇᑕᒥᐅᑕᓂ ᐅᒪᔨᑕᓃᔨᑦ, ᐊᐸᑎᑕᓃᔨᖏᔨᐊᖅᑯᑦ)	ᖃᐅᔨᔪᖅᓂᖅ bᖓᖅᓂᖅ

ᐊᖏᕕᒋ	ᖃᐅᔨᓴᖅᑎᑦ ᐊᑎᖅᓯᖏᑦ (ᑎᒥᖏᑎᖏᓪᓗᓂᖅᓂᑦ)	ᐱᓱᐊᑎᖃᖅᑕᐅᓂᒃ
1997	ᑯᐱᔪ - Kerbes (ᑲᓇᑕᒥᐅᑕᓂ ᐅᒪᓱᓕᓐᓄᑦ, ᐊᐋᑎᓕᓐᓄᖏᓱᐊᖅᑯᑦ)	ᐊᐳᔪᓕᐅᖅᑕᐅᓂᖏᓐᓂᓪᓗᓂ ᑲᖓᕐᖃᖅᖁᓘᑦ
1999-2001	ᕿᓗᓐ - Bazin (ᑲᓇᑕᒥᐅᑕᓂ ᐅᒪᓱᓕᓐᓄᑦ, ᐊᐋᑎᓕᓐᓄᖏᓱᐊᖅᑯᑦ)	ᖃᐅᔨᓴᕐᓂᖅ ᑲᖓᕐᓂᒃ
2003-2004	ᓴᒥᑦ - Smith (ᖃᐅᔨᓴᕐᓂᕐᒧᑦ ᓴᓇᕐᑭᑎᑕᖏᓐᓄᐊᖓᔪᓂᒃᑐᓂᓪ, ᐊᐋᑎᓕᓐᓄᖏᓱᐊᖅᑯᑦ)	ᖃᐅᔨᓴᕐᓂᖅ ᓗᖏᓴᐊᐋᖏᓕᓚᖓᓂᒃ
2005-2008	ᑲᔪᖁᐃᑦ Caswell (ᑲᓇᑕᒥᐅᑕᓂ ᐅᒪᓱᓕᓐᓄᑦ, ᐊᐋᑎᓕᓐᓄᖏᓱᐊᖅᑯᑦ)	ᖃᐅᔨᓴᕐᓂᖅ ᑲᖓᕐᓂᒃ
2009-2020	ᓚᖕᓂᐊᕐ - Leafloor (ᑲᓇᑕᒥᐅᑕᓂ ᐅᒪᓱᓕᓐᓄᑦ, ᐊᐋᑎᓕᓐᓄᖏᓱᐊᖅᑯᑦ)	ᖃᐅᔨᓴᕐᓂᖅ ᑲᖓᕐᓂᒃ
2011	ᓚᖕᓂᐊᕐ - Leafloor (ᑲᓇᑕᒥᐅᑕᓂ ᐅᒪᓱᓕᓐᓄᑦ, ᐊᐋᑎᓕᓐᓄᖏᓱᐊᖅᑯᑦ)	ᖃᐅᔨᓴᕐᓂᖅ ᐃᒪᕐᓗᒍᓱᐊᓂᕐ ᑎᓖᒥᐊᕐᓂᒃ
2014-2020	ᕿᐅᔪ - Rausch (ᑲᓇᑕᒥᐅᑕᓂ ᐅᒪᓱᓕᓐᓄᑦ, ᐊᐋᑎᓕᓐᓄᖏᓱᐊᖅᑯᑦ)	ᖃᐅᔨᓴᕐᓂᖅ ᓗᖏᓴᐊᐋᖏᓕᓚᖓᓂᒃ

ᑲᓇᑕᓯᒪᓕ ᐅᒪᓱᖃᖏᓴᐅᖓᖓᓱᖃᖅᑐᑦ ᐃᑲᓱᓂᖃᖏᓂᐊᖃᖅᑐᖅ ᔪᐳᓐᓂᐊᖏᓂᒧᑦ ᐃᑕᖃᑯᓗᑕᓴᐅᐅᓱᓂᒃ ᑕᐃᒪᓐᓕᒃ ᐱᓪᒪᓇᐅᑎᑕᐅᓕᐅᖅᑐᓂᒃ ᐃᓇᓐᓄᑦ, ᑕᒪᑲᑯᐊ ᓗᑦᓗ ᐃᓗᐃᖃᖏᖃᖏᓐᓂᒃᑐᓂ, ᓄᓇᕆᔨᐅᕈᒪᐅᓂᖏᓐᓂᒃᑐᓂ ᐊᓪᒪᓗ ᐱᐅᓗᑐᖃᖅᑎᑎᒍᓕᑦ ᐃᓇᓐᓄᑦ ᑕᖓᒪᖃᓕᐅᖅᑕᑐᖃᖅᑎᑐᓂᒃ, ᐃᑦᓂᐅᑦᖃᐃᖅᑯᑦ ᐊᐅᓪᓚᕆᔨᐅᖅᑎᑕᐅᓂᒃᑐᓂ ᑲᑎᒪᔨᕿᓗᓂᓄᑦ ᐱᐅᓂᖅᓴᓂᒃ ᔪᐳᓐᓂᐊᖏᑕᐅᓂᕆᐊᖅᑲᓂᖓᓂᓂᒃᑐᓂ. ᐱᑕᖃᓄᑕᐅᓂᖓᓂᒃ ᐱᓱᕈᐅᓗᒪᓂᖓᓂᓪᓗ ᑕᐃᖅᑯᐊ ᐃᓄᐃᑦ ᐃᑲᔪᓗᐊᖅᓴᖓᓂᓄᑦ ᐊᖓᑎᓂᔨᑦ ᐱᓱᕈᓴᖅᓴᓐ ᑐᖓᓚᐃᐅᓴᖅᓴᓂᒃᑐᖅ ᐊᐅᓚᑦᓴᖅᓴᓐ ᐱᑕᖃᓄᑕᐅᓂᖓᓂᒃᑐᖅ ᐊᐅᓚᑎᓪᓗᓴᖅᓴᓐ ᐱᑕᖃᐃᓐᓇᓪᓗᖅᓴᓐ ᓄᓇᕗᖅ ᑕᒪᐃᓐᓇᖅ ᐃᓄᐃᑦ ᐃᑲᔪᓗᐊᖅᓴᖓᓂᓄᑦ ᓄᓇᕗᖅ ᑕᒪᐃᓐᓇᖅ ᑐᖓᓚᐃᐅᓴᖅᓴᓐ ᐱᑕᖃᓄᑕᐅᓂᖓᓂᒃᑐᖅ ᐱᓱᕈᓴᖅᓴᓐ ᐱᑕᖃᐃᓐᓇᓪᓗᖅᓴᓐ ᓄᓇᕗᖅ ᑕᒪᐃᓐᓇᖅ ᐃᓄᐃᑦ ᐃᑲᔪᓗᐊᖅᓴᖓᓂᓄᑦ

ᖃᐅᔨᓂᑕᐅᖃᕐᓂᕆᔨᖃᕐᓂᖅᓯᒪᔪᓂᖅ ᑐᑭᓯᑎᕆᔨᖃᕐᓂᖅᓯᒪᔪᓂᖅᓱᓕᒃ ᑐᖓᔪᖑᓚᐅᒃᑯᓂ, ᑭᖑᓂᓕᒫᖅ ᓂᕆᔨᖃᕐᓂᖅᓯᒪᔪᓂᖅᓱᓕᒃ ᐊᐅᓚᑎᑦᓯᐅᖃᕐᓂᖅᓱᑦᔪᓂ ᐅᑎᒪᔨᑦᓯᐅᖃᕐᓂᖅᓯᒪᔪᓂᖅ ᓂᕆᔨᖃᕐᓂᖅᓯᒪᔪᓂᖅᓱᑦᔪᓂ ᐃᓕᓴᐅᒪᔨᓂᖅᓱᒪᔪᓂᖅ ᐊᐅᓚᑎᑦᓯᐅᖃᕐᓂᖅᓯᒪᔪᓂᖅ ᑎᒥᖅᓯᐅᑎᖃᕐᓂᖅᓯᒪᔪᓂᖅ, ᐃᓕᓴᐅᒪᔨᑦᓯᐅᖃᕐᓂᖅᓯᒪᔪᓂᖅ ᖃᓄᐃᓕᒫᓂᖅᓱᓕᒫᓂᖅ ᐊᐅᓚᑎᑦᓯᐅᖃᕐᓂᖅᓯᒪᔪᓂᖅ ᑎᒍᑎᓪᓗᒍ.

ᑐᑭᓯᓂᐊᖅᑕᐅᓂᖅᓱᒪᔪᓂᖅ ᑭᓇᐅᖃᕐᓂᖅᓱᒪᔪᓂᖅ ᐊᓯᔨᐅᖃᕐᓂᖅᓱᒪᔪᓂᖅᓱᑕᐅᖃᕐᓂᖅ ᑲᒪᒋᔭᒪᐊᖃᕐᓂᖅᓯᐊᒪᓕᕆᓂ ᐊᓯᔨᐅᖃᕐᓂᖅᓯᐊᖏᓪᓗᒍᓂ. ᐊᑐᓂᔪᓯᐅᓂᖃᕐᓂᖅᓯᑕᐅᖃᕐᓂᖅᓯᓯᒪᔪᓂ ᐃᓯᒪᔪᓇᔨᓂᖅᓯᐊᓯᔨᐅᑕᐅᔨᔪᓯᐅᓂᖅᓯᒪᔪᓂ/ᓯᑎᖃᕐᓂᖅᓯᐅᕐᓂᖅᓯᒪᔪᓂ, ᐊᓚᒪᓗ ᑐᔭᑐᖃᕐᓂᖅᓯᐅᕐᓂᖅᓯᒪᔪᓂ ᓄᓇᕗᑦ ᐃᓯᒪᔪᓇᔨᓂᖅᓯᐊᖏᓪᓗᒍᓂ. ᐊᒥᓯᑦᓴᖃᓂᖅ ᖃᐅᔨᓂᐊᖅᑕᐅᓂᖃᕐᓂᖅᓯᑕᐅᖃᕐᓂᖅᓯᓂᐊᖅᑐᓂ ᖃᐅᔨᓚᓕᐅᓂᐊᖅᓯᓂᐊᖅᑕᒪ ᐃᓯᒪᔪᓇᔨᓂᖅᓯᒪᔪᓂ, ᐱᓚᒪᓚᐅᓂᖅᓯᐅᓇᖅᑐᓂᖅ ᓴᐳᓂᐊᖅᑕᐅᑎᐊᖅᓯᓂᐊᕐᑕᒪᖅ ᐊᓚᒪᓗ ᐱᑕᖃᐅᔪᓇᒪᐊᖅᓯᓂᐊᕐᑕᒪᖅ ᐃᓯᒪᔪᓇᔨᓂᖅᓯᐊᖏᓪᓗᒍᓂ ᐱᓕᕆᓂᐅᔪᓇᖅᑐᓂᖅ ᐊᓕᕆᓂᐅᔪᓇᖅᑐᓂᖅ ᐃᓕᓴᐅᒪᔨᑦᓯᐅᖃᕐᓂᖅᓯᐊᖏᓪᓗᒍᓂ ᐱᓕᕆᓂᐅᔪᓇᖅᑐᓂᖅ ᐊᓕᕆᓂᐅᔪᓇᖅᑐᓂᖅ ᐃᓕᓴᐅᒪᔨᑦᓯᐅᖃᕐᓂᖅᓯᐊᖏᓪᓗᒍᓂ ᐱᓕᕆᓂᐅᔪᓇᖅᑐᓂᖅ ᐊᓕᕆᓂᐅᔪᓇᖅᑐᓂᖅ ᐃᓯᒪᔪᓇᔨ

9 ᐊᐢᓂᖅᑕᐅᖅᑕᐃᓕᓕᒫᑎᓂᖅ, ᐊᐢᓂᐊᖅᑐᓕᓂᖅ ᐊᒻᒪᓗ ᐅᓗᓚᐊᖅᑐᓕᓂᖅ

ᐊᐳᑎᓕᓚᓂᕆᒃ ᑐᐊᕕᕐᓇᖅᑐᖏᖅᓯᑦ, ᐊᑎᑐᖅ ᐅᖏᖏᕕᐅᓂᐊᖅᓴᑦ ᐅᑯᐊ
ᑲᓇᒋᒥ ᐊᐳᑎᓕᓚᓂᕆᒍᑦ ᑐᑐᐃᕕᕐᓇᖅᑐᖏᓕᓚᓗᓂᓂᖅ ᖃᐅᔨᓂᕆᑎᕕᖅ
ᓄᓇᕗᑦᑎᒍ ᓄᓇᕐᒥᒍ:

1-867-920-8130.

ᑲᓇᒋᒥ ᐅᒪᔨᓕᓚᓗᖏᓕ ᓴᓴᐅᓂᐊᖅᑎᐅᓂᐊᖅᑐᑦ ᐊᐢᓂᐊᓕᓚᓚᑯᑕᐅᔭᓚᐊᖅᖃᖏᓂᖅᓂᐅᓂᖅ ᐊᒻᒪᓗ ᐅᑐᓚᐊᓇᖅᑐᖏᖅᖃᖏᓂᖅᓂᐅᓂᖅ ᐃᓄᓕᒻᒥᓄᑦ, ᖃᐅᔨᓗᒪᓂᑎᖅᑕᖏᑦᑐᑎᖅ ᓄᓇᒧᖓᐅᑭᐅᑐᖅ ᓂᐅᕗᑦᕈᓯᓚᕐᓂᓗ ᐅᑐᓚᐊᓇᖅᑐᓂᖃᖅᑐᖅ, ᖃᐅᔨᓴᓕᓚᐊᖅᑐᖅᑕᖏᑦᑐᖅᓯᒪᓂᖅ ᓂᐅᐅᑎᔨᐅᐊᖅᑐᖅᑕᖏᑦᑐᖅᓯᒪᓂᖅ ᐅᑐᓚᐊᓇᖅᑐᓂᖃᖅᑐᖅ, ᐊᐢᓂᐊᓕᓚᐅᑎᓕᓚᐊᖅᑐᖅᑕᖏᑦᑐᖅᓯᒪᓂᖅ ᐊᑐᖅᑕᐅᔭᓇᖅᑐᖅ. ᐊᒻᒪᒪᐅᔭᖅ, ᑲᓇᒋᒥ ᐅᒪᔨᓕᓚᓗᖏᓕ ᐱᓕᓚᖏᓕᓂᓗ ᑕᒪᐃᓇᓕᒋᐊᖅ ᐅᔨᓴᖅᑐᑎᐊᖅᐸᑦᑎᖅᑕᕐᓂᐊᖅᑐᑦ ᓴᓴᐅᓂᐊᖅᑎᐅᓂᐊᖅᑐᖅ ᐃᓕᖓᓂᖅ ᐊᐢᓂᐊᓕᓚᖅᑕᐃᓕᒪᓚᐊᖅᑐᖅᑕᖏᑦᑐᖅᓯᒪᓂᖅ, ᐅᓗᓚᐊᓇᖅᑐᖅᑎᖅᓯᓇᖅᑐᖅᓯᒪᓂᖅ ᐃᓕᖓᓂᖅ, ᐱᓕᓚᖅᑎᑭᖅᑕᖏᓂᖅᓂᓗ. ᑭᓯᐊᓂᕕᑕᐅᖅ, ᑭᓇᐃᐃᓇᖅ ᐃᓂᒃ ᐃᓂᐃᓕᓗᓂᓂ ᐱᓇᓱᑎᐊᖅᑕᕐᓚᐊᖅᑎᓪᓗᓐᓂ ᐃᓕᖓᓂᒃ ᖃᐅᔨᓴᖅᑐᑎᐊᖅᐸᑦᑎᖅᑕᕐᓂᐊᖅᑐᖅ ᐅᓗᓚᐊᓇᖅᑐᖅᑎᖅᓯᓇᖅᑐᖅᓯᒪᓂᖅ ᐃᓕᖓᓂᖅ, ᐱᓕᓚᖅᑎᑎᑭᖅᑕᕐᓂᐊᖅᑐᖅᑕᖏᑦ

9.1 ᓇᓄᐃᑦ ᑐᖅᑯᑕᐅᓴᖅᓂᕐᑎᑦ

ᑭᓇᐯᑐᐃᓇᖅ ᑐᓴᐁᑎᓯᖅ ᓇᓄᖅᑎᒃ ᑎᖏᑕᐊᖅᑲᐃᖏᒥ, ᐱᓕᕆᑎᖅᖅᓯᓂ ᑐᐊᐃᓇᖅᑐᑐᒃ ᐱᓗᖅᑯᑕᐅᖏᑕᑐᑕᒪᑐᓄᓐᓂᑦ, ᐅᖅᐅᓯᖅᑕᐊᖅᖅᓴᖅ ᐊᓕᖅᓇᑭᓐᓂᑐᐊᖅᐸᑦ. ᐱᑐᐊᖅᑐᑎᒃ ᐅᑯᐊ ᐅᖅᐅᑎᕆᐅᓕᖅᓴᕆᐅᑦ ᓄᓇᕗᑦ ᑲᖏᒪᐃᑦ ᐅᒪᓕᒪᓂᓴᖏᑦ ᓴᓪᓕᓂᓂ (867-925-8823) ᐊᓚᒪᓂ ᑕᐃᖅᑯᐊ ᐊᖏᑎᓕᕆᓂᓴᕐᔪᐊᖅᑯᑦ ᓯᓕᐅᓪᓂ ᐊᓯᓴᖅᐸᓪᓕᐊᓕᕆᓂᓴᖏᑭᓐᓄ ᐅᒪᓕᕆᓂᓴᖅᑯᑦ ᐱᓕᖅᓇᐅᑎᓕᕆᓂᓴᖅᓱᓚᐅᓂᑦ (ec.nupermisscf-cwspermitnu.ec@Canada.ca). ᓴᖅᑯᑕᐅᓂᖏᑦ ᑐᓂᐅᐅᓂᑎᒃ ᐅᒪᔪᐃᑦ ᑎᒥᖅᑐᖅᑎᖅᑎᓪᓂᑦ ᑐᐊᐃᓇᑐᖅᑎᓪᓂᑦ, ᒪᓕᑎᓂᓂᖅ ᓯᓇᐃᔪᖅᑐᖅᑎᓪᓂᓗᒍ, ᐱᓕᕆᑎᖅᑐᑦᑐᒥᓪᓚᖅᑯᑦ ᑐᓴᐁᑎᓯᖅᑐᖅᑎᓪᓂᑦ ᒪᓕᑎᐊᖅᑐᖅᑐᖅᓴᖅ ᐃᓗᓕᔪᖅᑐᖅᑐᑦ ᑕᐃᖅᑯᓂᖅᓱ *5.6.55*-ᖅᑎᓂᓕᑐᓂᖅ, ᓄᓇᕗᒻᒥ ᓄᓇᕐᓂᓴᒍᓂ ᐊᖏᑎᑐᑎᓂ. ᓇᓪᓕᐅᖅᑯᒧᐊᓂᖅᑐᐅᐊᖅᑐᑦ ᐊᐃᐃᓴᑦ ᐊᖓᔪᓇᓯᖅᑎᑦ ᑲᑐᔾᔨᖅᑎᑎᖅᑎᓪᓂᑦ ᓇᓪᓂᖅᑐᑦᓯᖅᓯᓪᓂ ᑎᖏᑕᐊᖅᑲᐃᖏᒥ, ᒪᓕᓚᑐᑎᒃ ᐃᑐᓪᓕᖅᑎᓪᓂᖅᑐᖅᑐᖅ *12.3-ᒥᖅᑐᑐᒥᖅ*, ᐃᓄᐃᑦ ᐃᖅᓯᓯᖅᓴᖏᑎᓄᑦ ᐊᖏᑎᑐᑎᒥᖅ. ᐃᓄᐃᑦ ᑎᖏᑕᐊᖅᑲᐃᔪᖏᓚᐅᓕᐊᑦ ᐊᓕᑐᓵᑦᖅᓴᖅᑕᐅᓯᒪᔭᕆᐊᖅᑲᐃᖅᑐᑦ ᓇᓄᖅᑐᑐᓇᓕᓂᖏᓐᓂᖅ ᐅᑭᓴᖅᑐᑐᓇᓕᓂᒥᖅᑎᑦ ᐊᓚᒪᓂ ᖃᐅᔨᓴᕆᐊᑐᓇᓕᓂᖅᑎᑦ ᐅᖃᖅᑎᖅᑐᑦ ᐅᖅᖸᓐᓂᑦ ᓇᓄᖅᑎᓂᖅ ᓇᐅᑎᖅᑐᑐᒃᓴᖅᓴᖅᑐᒃ ᐃᓕᓐᓂᐅᒥᖅᑎᖅ.

10 ᐊᐃᓕᖅᑎᑕᐅᓯᒪᓕᐊᓂᖅ

ᐊᐅᓪᓚᑕᐅᓂᖏᓐᓂᒃ ᐊᐃᖅᑕᑕᐅᔭᕆᐊᓕᓪᓚ ᒪᓕᒃᑕᐅᔪᓂᑎᒃ *ᑲᑎᒪᔨᐅᑕᐅᓯᖓᓂᖏᓐᓂᓄᓪ ᑎᖏᒋᐊᓕᕆᓂᖅᓴᓪ ᒪᓕᑎᓕᒋᐊᖅ ᐊᓚᒪᓪ ᑎᖏᒋᐊᖅᑐᙵᓄᓪ ᒪᓕᑎᓪᓕᔪᓐᓇᖅᑐᑦ, ᐊᐅᑎᓕᕆᔨᖅᓴᐊᐅᑦ ᐅᒪᔨᓕᕆᔨᓐᓂᖅ ᓴᖏᑎᓂᖅᑐᑎᑕᐅᓯᓕᖅ >ᑭᖅᑕᓕᕆᐅᑦ. ᑎᒃᑦᐊᖅᑕᐅᓂᒋᔨᓪᓗᒍ ᓄᓇᓕᖅᑎ ᐅᒪᔨᓕᕆᔨᓐᓂ ᐊᓚᒪᓪ >ᑭᖅᑕᓕᕆᐅᑦ ᐊᐅᓪᓚᑕᑎᑦᐅᑎᑕᐅᓯᓕᖅ ᒪᓕᑎᓂᓂᒃ ᒪᓕᒃᑕᐅᑎᑎᖅᑕᑭᓐᓇᖅᑐᖅ ᑕᐃᒃᑯᓂᓴᖅ ᒪᓕᑎᓂᓂᒃ *ᑲᑎᒪᔨᐅᑕᐅᓯᖓᓂᖏᓐᓂᓄᓪ ᑎᖏᒋᐊᓕᕆᓂᖅᓴᓪ ᒪᓕᑎᓂᓂᒃ ᐊᓚᒪᓪ *ᑎᖏᒋᐊᖅᑎᙱᓄᓪ ᒪᓪᓗᓕᕆᓂᓪᓗᖓᓂ.

ᐊᐅᑎᓕᕆᔨᖅᓴᐊᐅᑯᓂ ᐅᒪᔨᓕᕆᔨᓐᓂ ᓇᐅᑎᖅᑐᐃᐃᐅᐅᑐᑦ ᒪᓕᒃᑕᐅᓂᖏᓐᓂᒃ ᐊᖏᑭᑕᐅᕆᓂᒃᓯᒪᓂᖅᓴᓪ ᒪᓕᑎᓴᕆᐅᑎᑕᐅᔪᓪ ᐊᓚᒪᓪ ᐱᔨᓇᐅᑎᖅᑲᑕᐅᔨᓂ ᑐᓂᔅᐅᔪᓕᔨᓂ ᒪᓕᒃᑕᐅᓕᓪᓗᓂᒃ *ᑲᑎᒪᔨᐅᑕᐅᓯᖓᓂᖏᓐᓂᓄᓪ ᑎᖏᒋᐊᓕᕆᓂᖅᓴᓪ ᒪᓕᓴᖓᓂ ᐊᓚᒪᓪ *ᑎᖏᒋᐊᖅᑎᙱᓄᓪ ᒪᓪᓗᓕᕆᓂᓪᓗᖓᓂ ᑕᐃᒪᖓᓂᒃ ᑲᔪᓯᓂᐊᖅᑐᓂᒃ ᐊᓚᒪᓪ ᖃᐅᔨᓴᖅᑕᐅᔨᔪᖅᑎᖅᑲᐊᖅᑕᖅᓯᑦ ᖃᐅᔨᓴᓇᓇᖅᓯᒪᒃ. ᐊᐅᑎᓕᕆᔨᖅᓴᐊᐅᑯᓂ ᐅᒪᔨᓕᕆᔨᓐᓂ ᒪᓕᓴᖅᑎᓂᑎᓯᒧᓇᓂ ᓇᓂᑐᐃᓇᖅ ᑕᒪᐃᓐᓂᒃ ᑕᐃᒃᑯᓂᓴᖅ *ᑎᖏᒋᐊᖅᑎᙱᓄᓪ ᒪᓪᓗᓕᕆᓂᓪᓗᖓᓂ ᑕᐃᒪᖓᓂᒃ, ᐱᓕᕆᓂᐊᖅᑐᓂᒃᓴᖅᑎᑐᓂᒃ, ᐊᓚᒪᓪ ᐱᖁᓇᖅᑐᓂᒃᓴᖅᑎᑐᓂᒃ ᖃᐅᔨᓴᖅᑕᐅᔨᔪᖅᑎᖅᑲᐊᖅᑕᖅᓯᑦ ᓇᓗᓇᐃᔨᓂᒃ ᐱᖁᓇᖅᑐᓂᒃ ᐃᓕᓴᕐᓂᖅ ᐃᓚᖓᑕᐅᓂᒃ ᐊᐅᑎᓕᕆᔨᖅᓴᐊᐅᑯᓂ ᐱᖁᓇᖅᑐᓂᒃ ᐃᓚᖓᑕᐅᓂᒃ ᐊᐅᓪᓚᑕᑎᑦᐅᑎᑕᐅᓯᓕᖅ ᒪᓕᑎᓴᕆᐅᑎᑕᐅᔪᓪ ᐊᓚᒪᓪ ᐊᐅᓪᓚᑕᑎᑦᐅᑎᑕᐅᓯᓕᖅ ᒪᓕᑎᓂᓂᒃ

ᐅᕐᓂᓗᓐᓂᑦ ᓇᖏᑦᓂᖅ ᐅᖅᑲᐅᑎᕆᐅᓗᑎᑲ ᓇᑦᓚᐊᖏᓄᑐᐃᓇᖅ ᐃᖅᓂᐅᖅᕿᑲᓐᓂᑦ ᐊᐅᓚᑦᑎᕆᐅᖅᑲᑕᐅᓂᖅᓗᑦ ᑲᑎᒪᔨᖅᑕᖏᓐᓄᑦ ᓴᖅᓂᓚᓂ.

11 <ᕿᓇᐅᔨᓴᓂ ᐊᑐᓕᖅᑎᑕᐅᓂᖅᓯᓂᑦ

ᑲᓇᑕᒥ ᐅᒪᔪᓕᓗᓐᓂ ᐊᓚᒪᓗ ᐃᖏᓂᐅᖏᕐᕕᖅᑲᑕᐅᓂᖏᓪᓗᑦ ᑲᑎᒪᓐᕿᓭᓕᖅᓯᖓᓂ ᐊᑐᓕᖅᑎᑎᑎᓂᐊᕐᕕᖅᑐᑦ ᐊᐅᓕᕿᑎᓂᖏᓪᓗᑦ <ᕿᓇᐅᑎᓂᒃ ᑕᒪᓐᓇ ᐊᑯᓂᐅᓂᖅᑲᓂᐊᕐᕕᖅᑐᒪᑦ ᖃᑯᓗᓄᑦ ᐊᖏᕿᔪᓄᑦ. ᐊᑐᓕᖅᑎᑕᐅᓂᖓᓂᑦ ᒪᓕᖅᓂᐊᕐᕕᖅᑐᑦ ᐊᖏᕿᔪᑕᒪᓴᖅ ᐊᓕᓚᐅᖅᓴᖓᓂᖅᓂᒃ ᐊᓚᒪᓗ ᐱᓇᐅᔨᖅᓴᖅᑕᐅᓯᒪᓕᖅᓂᖅᓂᒃ. ᑲᓇᑕᒥ ᐅᒪᔪᓕᓗᓐᓂ ᐊᓚᒪᓗ . ᐊᐅᓕᕿᑎᓂᐅᖏᕐᕕᖅᑕᐅᓂᖏᓪᓗᑦ ᑲᑎᒪᓐᕿᓭᓕᖅᒥᓂ ᐊᑐᖏᑦᕿᓂᖅᓴᐅᓂᐊᕐᕕᖅᑐᑦ ᐋᖏᕿᑎᑕᕿᓂᖅᓴᖓᓂᖅᓂᒃ ᐊᐅᓕᕿᑎᓂᖏᓐᓄᑦ.

ᐊᐅᓕᕿᑎᓂᐅᖏᕐᕕᖅᑕᐅᓂᖏᓪᓗᑦ ᑲᑎᒪᓐᕿᓭᓕᖅᒥᓂ ᐊᑐᓕᖅᑎᑎᑎᓂᐊᕐᕕᖅᑐᑦ ᒪᓕᖅᓄᑎᓂ ᑕᒪᐃᓐᓄᓂ ᐱᓕᕿᑲᓇᕆᐊᓕᓚᖅᓯᒪᓕᖅᓂᖅᓂᒃ ᐃᓕᓴᕐᓂᒥᓄᑦ (*ᐃᓄᓂᑦ ᐃᑲᓴᓱᐊᕐᓂᖅᓂᓄᑦ ᐊᖏᕐᕉᑎᑦ ᐃᓗᓕᖅᓂᖅᑦ 2.1.9*). ᑲᓇᑕᒥ ᐅᒪᔪᓕᓗᓐᓂᑕᐅᕆᓂᒃ ᖃᑲᐅᕐᐊᖏᑕᓂᕐᓂᐊᕐᕕᖅᑐᑦ ᐊᑐᓕᖅᑎᑕᐅᓂᖏᓪᓗᑦ <ᕿᓇᐅᑎᑦ ᑕᓕᒪᐃᑦ 5 ᐊᖏᕿᔪᐃᑦ ᐊᖏᕐᕉᑕᐅᓕᐅᖏᓂᖏᓪᓗᑦ ᐊᓚᒪᓗ ᕈᐃᓂᐊᔪᑦ ᖃᑯᓗᓄᑦ 10 ᐊᖏᕿᔪᐃᑦ ᓴᓲᖏᓐᓂᒃ, ᒪᓕᖅᓄᑎᓂ ᐊᐅᓕᕿᑎᓂᖓᓂᑦ ᐱᓕᕿᑲᓇᕆᐊᓕᓚᖅᓯᒪᓕᖅᓂᖅᓂᒃ ᐃᓕᑕᒪᓴᐅᓯᒪᔪᓂᖅᓂᒃ ᑕᐃᑲᓂ ᐊᐃᓴᑐᓂᖏᓪᓗᑦ ᑕᑯᑎᑕᑎᓴᐊᑎᓴᖅᓯᓂᑐᓂᖅᓂᒃ 10-ᒥ ᐃᓄᓕᖅᑎᓴᑐᓂᖅᓂᒃ. ᒡᖅᑯᐊ ᑎᑎᕆᖏᓐᓂᖅᑎᑦ ᐊᐅᓕᕿᑎᓂᖓᓂᑦ <ᕿᓇᐅᑎᓂᖅ ᐊᓯᓴᓄᓕᐅᕐᓴᖅᑎᑦᑕᓂᑦ ᐊᖏᕿᔪᐃᑦ ᐱᓕᕿᑲᓇᕆᐊᓕᓚᖅᑕᖅᑎᑦᑕᓂᖅᓂᒃ <ᕿᓇᑕᐅᐊᖅᑐᓂᖅᓂᒃ, ᐃᑲᔪᓂᓪᓗᑎᒃᑲᓂᖅᓂᒃ ᓇᐅᕆᓂᖅᑐᓂᖅᓂᒃ ᐊᖏᕿᔪᑎᓂᖅᓂᒃ ᐱᓕᕿᑲᓇᕆᐊᓕᓚᖅᓯᒪᓕᖅᓂᖅᓂᒃ ᐱᓕᕿᑲᓇᕆᐊᓕᓚᖅᓯᒪᓕᖅᓂᖅᓂᒃ ᐃᓕᑕᒪᓴᐅᓯᒪᔪᓂᖅᓂᒃ ᐃᓕᑕᒪᓴᐅᓯᒪᔪᓂᖅᓂᒃ ᐊᐅᓕᕿᑎᓂᖓᓂᑦ ᐊᖏᕿᔪᐃᑦ

ᐊᕐᓇᐅᑎᓯᒪᔨᓂᖅ ᑕᐃᑎᑎᕐᓂᔨᓪᓚᓂ 10: ᑕᓪᓕᒪᓄᑦ ᐊᖏᔪᒍᓄᑦ ᐊᐅᓚᑦᑎᓂᖏᓪᓗᑦ ᐸᖃᐅᐅᑎᕐᖓᖅ ᐊᑐᓵᖅᑎᑎᕐᒋᓪᓕᐊᓂᖅ.

ᐱᓕᕆᐊᓂᖅ	ᐊᖏᖅᓯᒪᓂ				
	1	2	3	4	5
ᐅᖃᐅᓯᖃᕈᓂᖅ ᒥᓂᕈᓴᓂᖅ ᐊᒋᑎᓕᕆᓂᖏᓪᓗᑦ, ᐃᓚᒪᓕᓴᖏᓐᓄᑦ, ᑕᒪᐃᓂᖅ ᐱᔭᕇᑎᖃᕈᓂᖏᓪᓗᑦ ᐸᐅᓕᓕᕆᓂᖏᓪᓗᑦ; ᐅᓯᓇᖅᑐᑎᐊᖏᓪᓗᓂᒃ ᐃᓕᒪᒃᖁᓇᓕᐅᑕᐅᑎᐊᖃᖅᑕᕈᓐᓇᓂᒃ ᖃᐅᔨᐃᑐᑐᐃᓐᓇᐃᓂ ᐃᓄᐃᑦ ᖃᐅᐅᔨᓕᕆᖃᖅᑎᑦ ᐅᖃᐅᓯᓕᕆᓂᖏᓪᓗᑦ ᐅᖃᐅᓯᓕᕆᓂᖏᓪᓗᑦ ᐅᑎᒪᔨᐅᖏᓪᓗᓂ (*ᐃᓄᐃᑦ ᐃᑲᔪᐊᕐᒪᖏᓐᓄᓂ ᐊᖏᕐᕈᑎᑦ ᐃᓗᓕᖏᓪᓗᑦ 3.3*).	X	X	X	X	X
ᐱᓴᖅᑕᐅᖃᖅᑕᕈᓐᓇᓂᒃ ᖃᐅᓄᐃᓕᓕᐅᕈᑕᐅᓂᖏᓪᓗᑦ ᐊᐅᓚᑦᑎᓂᐅᔨᖃᖅᑕᐅᓂᖏᓪᓗᑦ ᐅᑎᒪᔨᓴᓪᓕᓐᓄᑦ ᐅᖃᐅᓯᓕᕆᓂᖏᓪᓗᑦ ᐃᓄᐃᑦ ᐃᑲᔪᐊᕐᒪᖏᓐᓄᑦᑕᕐᓂᐊᕐᓂᖏᓪᓗᑦ ᐃᓄᐃᑦ, ᐃᓕᓴᖃᓐᓂᒃ ᐅᖃᐅᓯᓕᕆᓂᖏᓪᓗᑦ ᐊᖏᕐᕈᑎᑦ ᐃᓗᓕᖏᓪᓗᑦ ᐃᓄᐃᑦ ᐃᑲᔪᐊᕐᒪᖏᓐᓄᑦ ᐊᖏᕐᕈᑎᑦ ᐃᓗᓕᖏᓪᓗᑦ, ᐃᓕᓴᖃᓐᓂᒃ ᐅᖃᐅᓯᓕᕆᓂᖏᓪᓗᑦ ᐊᐅᓚᑦᑎᓂᐅᔨᖃᖅᑕᐅᓂᖏᓪᓗᑦ ᐃᓄᐃᑦ ᐃᑲᔪᐊᕐᒪᖏᓐᓄᑦᑕᕐᓂᐊᕐᓂᖏᓪᓗᑦ ᐃᓄᐃᑦ, ᐃᓕᓴᖃᓐᓂᒃ ᐅᖃᐅᓯᓕᕆᓂᖏᓪᓗᑦ ᐊᖏᕐᕈᑎᑦ ᐃᓗᓕ

ᐱᓕᕆᐊᓂᖅ	ᐊᖏᕿᒍᑎ				
	1	2	3	4	5
ᑎᑎᕋᖅᑕᐅᖏᑦᑐᓗᒃ ᐅᓂᒃᑲᖅᑕᐅᖏᑦᑐᓗᑎᑐᓗ ᒪᓕᑦᔨᓂᒃ ᓯᕋᐃᔨᖏᖅᑐᓂᒃ ᐱᓕᕆᔨᖏᖅᓴᖅ.	X	X	X	X	X
ᐃᓕᐅᖏᑦᑐᓗᑎᒃ ᐊᕐᕉᑐᖏᓯᒪᓂᖅᓴᓕᓕᐊᖏᑐᓂᒃ ᐱᑎᐊᖏᑎᑕᐅᓕᓕᐊᖏᑐᓂᓗᓐᓂᒃ ᐃᑲᔨᖏᓯᐃᓂᕐᒧᑦ ᓴᓗᓐᓂᐊᑕᐅᓂᖅᓱᓂᒃ ᑎᖓᒥᐊᖏᑦᑕᐅᓂᖅᓱᒪᓂᒃ ᑎᖓᒥᐊᖏᑦᕼᐃᐅᔾᔨ, ᓄᓇᖓᓂᓂᓗ ᐊᖏᓗᐊᓂᑐᑐᓂᒃ.	X	X	X	X	X
ᑎᑎᖏᑦᔪᓂᑎᑐᓗᒃ ᐃᓐᓇᐅᔾᔨᓗᒃ ᐱᑕᖏᑦᑐᓂᖅᓴᓂᒃᑐᓂᒃ ᐃᓕᓴᕐᓂᐊᕐᓂᖅ ᐊᑐᐊᓂᖅ, ᐱᓕᕆᖏᑎᑎᓂᖅᓴᓂᒃᑐᓂᒃ ᐅᑭᐅᖅᑕᕐᓂᖅ ᐊᑐᐊᓂᖅ ᐃᓚᖓᑕᐅᓂᖅᓴᓂᒃ ᐊᑐᐊᓂᖅ ᐱᖓᓱ

11.1 ᐊᐅᓪᓚᕐᒧᑦ ᓴᖅᓇᐅᔨᑦ ᐃᖅᑭᒋᐊᖅᑕᐅᔪᓇᖅᓯᒪᓂᑦ

ᑲᓇᑕᒥ ᐅᓪᓚᔨᓕᓚᐅᔨᒻᒪᑦ ᐃᖅᑭᒋᐊᖅᓯᔨᓇᖅᑐᖅᓴᖅ ᐊᐅᓪᓚᕐᓂᒧᑦ ᓴᖅᓇᐅᑎᒧᑦ ᓴᖅᑲᒪᑐᐃᓐᓇᖅ. ᑭᓇᐅᒻᒋᑐᐃᓐᓇᐃᓐᓂ ᓴᙱᓚᑦᑐᑦ ᐃᓇᐃᓐᑲᓐᑦᑦᑐᑦ ᐊᐅᑐᖅᑕᐅᓯᒪᔨᒧᑦ ᑕᐅᒻᒥᐃᓐᓂᑦ ᐱᒋᐊᖅᑐᑦᑕᐅᑎᒃᓴᖅᑐᑦ ᐅᑎᖅᓴᐅᐃᑐᑐᖅᓴᖅ ᐃᖅᑭᒋᐊᖅᑕᐅᑎᒃᓴᐅᒧᖅᓴᖅ ᐅᑎᖅᓴᐅᑎᖅᑕᐅᑎᒃᓴᐅᒧᖅᓴᖅ ᐅᑎᖅᓴᐅᑎᖅᑕᐅᓴᐅᒧᑦ ᑲᑎᖅᓴᐅᑎᖅᑕᐅᓴᐅᒧᑦ ᐱᑎᐊᖅᑎᑕᐅᑎᖅᓴᐅᒧᑦ ᑲᑎᖅᓴᐅᑎᖅᑕᐅᓴᐅᒧᑦ ᐊᐅᓪᓚᕐᓂᒧᑦ ᐸᖅᓴᓕᐅᑐᖅᓴᖅ ᑲᑎᖅᓴᐅᑎᖅᑕᐅᓴᐅᒧᑦ ᐃᓇᐃᓐᑲᓐᑦᑦᑐᑦ ᐊᐅᓪᓚᕐᓂᒧᑦ ᐸᖅᓴᓕᐅᑐᖅᓴᖅ ᑲᑎᖅᓴᐅᑎᖅᑕᐅᓴᐅᒧᑦ ᐃᖅᑭᒋᐊᖅᑕᐅᓴᐅᒧᑦ ᐃᓇᐃᓐᑲᓐᑦᑦᑐᑦ ᐊᐅᓪᓚᕐᓂᒧᑦ ᐸᖅᓴᓕᐅᑐᖅᓴᖅ ᑲᑎᖅᓴᐅᑎ

12 ᐱᓕᕆᖅᑎᑦᑎᐊᑭᓐᓇᖅᓂᖅ

ᑲᔪᓯᓐᓇᕐᑎᐊᖅᓂᐊᑭᑎᑎᒃ ᑕᒃᑯᐊ ᐊᐅᓪᓚᕐᑎᓂᖅᓗᑦ <ᖅᓇᐅᑦᓴ, ᑐᖏᕆᕃᖅᓚᕆᓚᖓᓂᐊᖅᓴᑦ ᐱᓕᕆᖅᑎᑦᑎᐊᑭᓐᓇᖅᓂᖓᓂᓂᒃ ᑐᑭᓯᓂᐊᑭᑕᐅᓐᖓᓂᖓᓂᓂᒃ ᑲᑭᓯᑎᐊᑭᑕᐅᖅᓇᖅᑐᑦ ᐅᒪᔪᓕᐊᓚᒃᑯᑦ, ᓴᖅᓪᓕᐊᓂᖅᓯᓚᖅᑐᑦ ᐱᓕᕆᖅᑎᑦᑎᐊᑭᓐᓇᖅᓂᖓᓂᓂᒃ ᑲᑭᓯᑎᐊᑭᑕᐅᖅᓇ

ᒪᓕᑕᕆᐅᔪᓂᖅ ᒪᓕᒋᐊᖏᖅᓯᒪᓕ ᐊᓘᓚᓗ ᑕᒪᐃᓐᓂᖅ ᑲᓇᑕᓕᒫᓕ ᒪᓕᑕᕆᐅᔪᓂᖅ, ᓄᓇᕗᔪᐊᖏᑎᒃᓄᑦᓗ ᐊᖏᑭᑐᐅᔪᓂᖅ ᐊᓘᓚᓗ ᑲᑎᒪᔨᐊᑭᑕᐅᔭᖅᑐᑦᓂᖅ ᐊᓘᓚᓗ ᐃᓕᓴᕆᓂᖅᑐᑦ ᑕᒪᐃᓐᓇᖅᑐᒃ ᑕᒪᓐᓇᖅᑐᒥ ᐊᐅᓚᑦᑎᑕᕐᓂᖏᑦ ᓄᓇᕗᒻᒥ ᐅᒪᔨᓂᖅ, ᑕᐃᔨᓂᖏᓚᓗ ᒪᓕᒋᐊᖏᖅᓯᒪᓂᖅᑐᒃ ᓄᓇᕗᒻᒥ ᓄᓇᑦᓂᔭᒃᑯᑦ ᐊᖏᑭᑐᐅᑎᓂᖅ. ᐃᓕᓴᕐᓂᖅ ᐱᓗᐊᖏᑐᒥᑦ ᑐᓴᕆᖏᖅᓴᑦ ᐱᓕᕆᔨᑦᑎᓐᓂᖅ ᓇᓕᔪᐊᓂᖅ ᐃᓕᓴᕐᓂᖅᑐᒃᓴᖅᑐᑦ ᐊᐅᓚᑦᑎᐊᕐᓂᖏᓐᓂᖅ, ᒪᓕᒋᐊᖏᖅᓯᒪᓂᖅ ᒪᓕᑲᓱᐊᕆᔭᖅ, ᐊᓘᓚᓗ ᐊᑐᖏᑕᕐᓇᖏᖅᓯᒪᓂᖅᑐᒃ ᖃᓄᖅᓯᐅᒃᑐᓂ ᐱᓕᕆᔨᑦᑎᓐᓂᖅ ᓄᓇᕗᒻᒥ ᓄᓇᑦᓂᔭᒃᑯᑦ, ᐃᓅᐱᑎᕆᐃᓐᓇᓗ ᐱᓘᓪᓚᐅᑎᓪᓗᐊᓂᖅ ᐃᓄᐃᑦᑐᓂᖅ ᐊᐅᓚᑦᑎᐊᕐᓂᖏᓐᓂᖅ ᐃᓄᐃᑦᑐᓂᖅ ᓄᓇᕗᒻᒥ.

ᓄᓇᕗᓂ ᑲᖏᒪᐃᑦ ᐱᓕᕆᔨᖏᓕᒃ ᐃᓕᓴᕐᕕᓴᕐᓂᑎᑦᓗ ᐱᓘᓪᓚᐅᑎᓪᓗᐊᓂᖅᓴᓗ ᐃᓕᓴᕆᐅᓪᓚᕆᔨᕐᓂᓵᓂ ᐱᓕᕆᔨᐊᖏᖅᓴᓂᖅ ᐊᑐᓕᖅᑎᓪᓗᐊᓂᖅᓯᖅᓴᓗ ᐊᑐᐊᖃᐃᐃᑦ, ᐊᑐᐃᓴᕐᓂᖅ ᐱᓕᕆᔨᖏᓕᒃ ᐊᓘᓚᓗ ᐱᔨᓂᑎᐅᑦᑎᓂᖅ ᐃᓅᖅᓯᒪᔪᖅ. ᑕᒪᕐᕕᐊ ᐱᔨᓂᑎᐅᖅᑕᕐᓂᓂ ᑐᖃᖅᑐᓪᓚᕆᐊᓂᖅᓯᓂᖅ ᐃᓕᓴᕐᕕᓴᕐᑐᐃᓐᓇᖅᑐᓂᖅ, ᐅᖅᑲᐅᓴᑐᖅᑕᓚᐅᔪᓂᖅ, ᐅᖅᑲᐅᓴᑐᖅᑕᓚᐅᔪᓂᖅ, ᐱᓘᓪᓚᐅᓂᖅᑲᖅᑎᓪᓗᐊᓂᖅ ᐊᐅᓚᑦᑎᐊᕐᓂᖅ ᖃᐅᔨ ᐱᓕᕆᖏᖅᓴᓂᖅ ᓄᓇᕗᒻᒥᐅᓂ. ᓄᓇᕗᓂ ᑲᖏᒪᐃᑦ ᐱᓕᕆᔨᖏᓕᒃ ᐃᓕᓴᕐᕕᓴᕐᓂᑎᑦᓗ ᐱᓘᓪᓚᐅᑎᓪᓗᐊᓂᖅᓴᓗ ᐃᓕᓴᕆᐅᓪᓚᕆᔨᕐᓂᓵᓂ ᐱᓕᕆᖃᑎᖅᑭᓯᓚᐅᖅᓂᖅ ᖃᐅᔨᓴᖅᑐᓂᖅ ᐃᓅᖅᑲᖏᐅᑦᑐᖅᓯᒪᔪᖅᓴᓗ ᐱᔨᓂᑎᐅᖅᑕᕐᓂᖅᓯᖅᓴᓗ ᓄᓇᕗᒻᒥᐅᓂ, ᐃᓕᑎᒥᓂᖅ ᓄᓇᕗᒻᒥ ᐊᐅᓚᑦᑎᐊᕐᓂᖅᓯᖅᓴᓗ ᓄᓇᕗᒻᒥ ᑲᓇᑕᒥ ᑲᖏᒪᐃᑦ ᑎᑎᕋᖅᑕᐅᓂᖅ ᐊᐅᓚᑦᑎᐊᕐᓂᖅᓯᖅᓴᓗ ᐃᓕᓴᕐᕕᓴᕐᓂᑎᑦᓗ ᐱᓘᓪᓚᐅᑎᓪᓗᐊᓂᖅᓴᓗ ᐃᓕᓴᕆᐅᓪᓚᕆᔨᕐᓂᓵᓂ ᐱᓕᕆᔨᐊᖏᖅᓴᓂᖅ ᐃᓅᖅᑲᖏᐅᑦᑐᖅᓯᒪᔪᖅᓴᓗ ᐱᔨᓂᑎᐅᖅᑕᕐᓂᖅᓯᖅᓴᓗ ᐃᓅᖅᑲᖏᐅᑦᑐᖅᓯᒪᔪᖅᓴᓗ ᐱᓘᓪᓚᐅᑎᓪᓗᐊᓂᖅᓴᓗ ᐃᓕᓴᕆᐅᓪᓚᕆᔨᕐᓂᓵᓂ ᐱᓕᕆᔨᐊᖏᖅᓴᓂᖅ ᓄᓇᕗᒻᒥ, ᐃᓕᑎᓐᓄᓂᖅ ᓄᓇᕗᒻᒥ ᓄᓇᓕᖅ, ᐃᓕᑎᓐᓄᓂᖅ ᓄᓇᕗᒻᒥ ᑲᓇᑕᒥ ᑲᖏᒪᐃᑦ ᐃᓅᖅᑲᖏᐅᑦᑐᖅᓯᒪᔪᖅᓴᓗ ᐃᓅᖅᓯᐅᒃᑐᓂ ᐱᔨᓂᑎᐅᖅᑕᕐᓂᖅᓯᖅᓴᓗ ᐃᓅᖅᓯᐅᒃᑐᓂ ᐱᔨᓂᑎᐅᖅᑕᕐᓂᖅᓯᖅᓴᓗ ᐃᓕᓴᕐᕕᓴᕐᓂᑎᑦᓗ ᐱᓘᓪᓚᐅᑎᓪᓗᐊᓂᖅᓴᓗ ᐃᓕᓴᕆᐅᓪᓚᕆᔨᕐᓂᓵᓂ ᐱᓕᕆᔨᐊᖏᖅᓴᓂᖅ ᓄᓇᕗᒻᒥ

13 ᐅᖃᓕᒫᖅ ᑐᑭᓯᐊᖃᕐᓂᐅᓯᒪᓂᖅᐸᑦ

ᐊᓕᓴᐅᔨᒃᑲ-Alisauskas, R. T., Dufour, K. W. & Leafloor, J. O., 2018a. Midcontinent lesser snow goose Chen caerulescens caerulescens. In: A. D. Fox & J. O. Leafloor, eds. *A global audit of the status and trends of Arctic and Northern Hemisphere goose populations.* Akureyri: CAFF International Secretariat.

ᐊᓕᓴᐅᔨᒃᑲ-Alisauskas, R. T., Dufour, K. W. & Leafloor, J. O., 2018b. Ross's goose Chen rossii. In: A. D. Fox & J. O. Leafloor, eds. *A global audit of the status and trends of Arctic and Northern Hemisphere goose populations.* Akureyri: CAFF Internation Secretariat.

ᐋᓕᓴᵒ-Allison, L., 1977. *Migratory bird sanctuaries in the Northwest Territories - a background paper,* s.l.: Unpubl. rep., Can. Wildt. Serv., Edmonton. 3 vols. 370 pp.

ᐊᖏᓀᐹᒃᑎ ᑎᖃᒥᐊᓕᓚᓂᖅᓴᑦ ᑲᑎᒪᔨᑦ-Atlantic Flyway Council, 2011. *Atlantic Brant Management Plan,* s.l.: s.n.

Bᐋᒋᑦ-Batt, B. D., 1997. *Arctic ecosystems in peril: report of the Arctic Goose Habitat Working Group.,* s.l.: Arctic Goose Joint Venture Special Publication.

Bᐅᒋᑦ-Bird, J., 1953. *Southampton Island,* Ottawa: Edmond Cloutier, Queen's Printer and Controller of Stationery.

Bᐅᓀ-Bray, R., 1943. *Notes on the birds of Southampton Island, Baffin Island and Melville Peninsula.* Ottawa, ON: The Auk.

CACNMP, 1990. *A review of the boundaries of Bird Sanctuaries in the Northwest Territories,* Ottawa, ON: s.n.

CAFF, 2010. *Arctic Biodiversity Trends - Selected indicators of change,* Akureyri, Iceland: CAFF International Secretariat.

ᑲᓇᑕᒥ ᓄᖅᓯᓕᖅᑐᓄᑦ ᐅᒪᔨᓕᓀᐸᑦ ᑲᑎᒪᔨᖏᑦ-Canadian Endangered Species Conservation Council. 2011. *Wild Species 2010: The General Status of Species in Canada, s.l.: National General Status Working Group.*

ᑲᑦᐳᑦ ᒥ. ᐊᓱᒪᓂ ᐳᓴᕐᐅᐃ, ᔨ-Campbell, M. & Boulanger, J., 2015. *Preliminary Report on The Long-Term Abundance Fluctuations of the Southampton Island Caribou Herd - 1978 -2015,* Arviat, NU: Nunavut Department of Environment, Wildlife Research Division.

ᑲᑐ, ᐱᓕᓚᖅᑎᖏᑦᓂᓂ-Carter, N. A. et al., 2018. *Inuit knowledge about light geese in the Kivalliq region, Nunavut.,* s.l.: s.n.

ᑯᑦ Bᐃ ᐊᓱᒪᓂ ᖃᐃᔨᑦ-Coad, B. W. & Reist, J. D., 2004. *Annotated list of the Arctic Marine Fishes of Canada,* s.l.: Canadian Manuscript Report of Fisheries and Aquatic Sciences 2674.

ᑲᓕᓐ-Collins, H., 1956. *Archaeological investigations on Southampton and Coats islands, N.W.T.* , s.l.: National Museum of Canada Bulletin No. 42.

ᑯᑦ-Cooch, F. & Barry, T. W., 1957. *Proposed Migratory Bird Sanctuaries Southampton Island, Keewatin,* s.l.: CWSC 3498.

ᐊᕐᑎᓕᓚᐅᖅᓯᐊᕆᔨᑦ ᓂᓪᑕᐅᑦᓗ ᐊᓂᔪᓄᖅᓴᑦᓕᐊᓯᓕᓚᐅᖏᓪᓗᑦ-Environment and Climate Change Canada, 2016. *Environment and Climate Change Canada's input to the Nunavut Planning Commission regarding Key Habitat Sites for migratory birds in the Nunavut Settlement Area, May 2016 revision,* Yellowknife, NT.: Canadian Wildlife Service.

ᐊᕐᑎᓕᓚᐅᖅᓯᐊᕆᔨᑦ ᑲᓇᑕᒥ-Environment Canada, 2013. *Bird Conservation Strategy for Bird Conservation Region 3 Prairie and Northern Region: Arctic Plains and Mountains,* Yellowknife, Northwest Territories: Canadian Wildlife Service, Environment Canada.

Fᐊᓪᒥᖏ ᓂ-Flemming, S. A., Calvert, A., Nol, E. & Smith , P. A., 2016. *Do hyperabundant Arctic-nesting geese pose a problem for sympatric species?,* s.l.: Environ. Rev. 24.

Fᐋᐟᑎᐟ ᐊᒻᒪᓗ Iᓗᓚ-Fontaine, A. and Mallory, M., 2011. *Detection and Classification of Land Cover Classes of Southampton Island, Nunavut, Using Landsat ETM+Data,* Prairie and Northern Region: Occasional Paper No. 119, Canadian Wildlife Service.

Gᓴᑕᐟ-Gaston, A. J., Decker, R., Cooch, F. G. & Reed, A., 1986. *The distribution of larger species of birds breeding on the coasts of Foxe Basin and northern Hudson Bay, Canada.,* s.l.: Arctic Vol. 39 No. 4.

ᓴᖅᑐ-ᑐᓚᐊᐳ-Gratto-Trevor, C. et al., 2011. *Trends in Canadian shorebirds.,* Ottawa, ON.: Canadian Biodiversity: Ecosystem Status and Trends 2010, Technical Thematic Report No. 13. Canadian Councils of Resource Ministers.

ᐃᓄᓕᓚᐅᑐᖃᖅᑎᑦ-INAC, 1976. *Inuit Land Use and Occupancy Project, 1974 (Milton Freeman Research Ltd.),* s.l.: Department of Indian and Northern Affairs Canada.

ᐊᓄ ᐊᒻᒪ ᐱᐅᐊᑭᓚᓴ-Jano, A. P., Jeffries, R. L. & Rockwell, R. F., 1998. *The detection of vegetational change by multitemporal analysis of LANDSAT data: the effects of goose foraging.,* s.l.: J. Ecol.

Kerbes, R. H., 1975. *The nesting population of Lesser Snow Geese in the eastern Canadian Arctic: a photographic inventory of June 1973,* s.l.: Can. Wildl. Serv. Rep. Ser. No. 35. 47 pp.

ᑯᐱᓴ-Kerbes, R. H., Meeres, K. M. & Alisauskas, R. T., 2014. *Surveys of Nesting Lesser Snow Geese and Ross's Geese in Arctic Canada, 2002 – 2009.,* U.S. Fish and Wildlife Service, Washington, D.C. and Canadian Wildlife Service, Ottawa, Ontario: Arctic Goose Joint Venture Special Publication.

ძPᑕᐣ, ᐱᓕᓗᖏᑎᖏᑦᓂᓗ Kokelj, S. V. et al., 2005. *The Influence of Thermokarst Disturbance on the Water Quality of Small Upland Lakes, Mackenzie Delta Region, Northwest Territories, Canada,* s.l.: Permafrost and Periglac. Process. 16.

ძᑐᓂᐊᵒ ᐊᒡᒪᓗ ᔨᐊᐅᓇᖅ-Kotanen, P. & Jefferies, R. L., 1997. *Long-term destruction of sub-arctic wetland vegetation by lesser snow geese,* s.l.: Écoscience 4.

ᓕᑐᐊᖅ, ᐱ. Latour, P. e. a., 2008. *Key Migratory Bird Terrestrial Habitat Sites in the Northwest Territories and Nunavut (3rd Ed.),* s.l.: Occasional Paper No. 114. Canadian Wildlife Service.

ᓕᵒ, ᓂ-Lunn, N. J. et al., 2016. *Demography of an apex predator at the edge of its range: impacts of changing sea ice on polar bears in Hudson Bay,* s.l.: Ecological Applications 26(5).

ᐃᓂᵒᖅ, ᑎ-Manning, T., 1942. *Remarks on physiography, Eskimos, and mammals of Southampton Island,* s.l.: Can. Geog. Jour. 24 (1):.

ᐃᓂᵒᖅ, ᑎ-Manning, T., 1967. *A report on the transfer of barren-ground caribou from Coats Island to Southampton Island, NWT,* Ottawa: CWSC Rept. 1143, Canadian Wildlife Service.

ᓂᕆᑕ, ᓯ-Nissley, C., 2016. *Assessing the impact of Lesser Snow Goose and Cackling Goose competition on breeding Atlantic Brant,* s.l.: University of Delaware.

ᓄᓇᐅᒻᒥ ᓄᓇᕐᓂ ᐊᑐᑎᐹᖏᑦᓄᓂ ᐸᖏᐅᕕᓂ ᑲᒥᓯᓇᕐᓂ-Nunavut Planning Commission, 2000. *Keewatin Regional Land Use Plan,* s.l.: s.n.

ᐸᑯ, ᔨ-Parker, G., 1974. *A population peak and crash of lemmings and snowy owls on Southampton Island, NWT,* s.l.: Can. Field-Nat. 88.

ᐸᑯ, ᔨ-Parker, G. R., 1975. *An investigation of caribou range on Southampton Island, Northwest Territories,* Ottawa, ON.: Canadian Wildlife Service Report Series No. 33.

ᐸᑯ, ᐃ-Parker, G. R. & Ross, R. K., 1973. *Notes on birds of Southampton Island, Northwest Territories.* s.l.

ᓀᐅ, ᓀ-Riewe, R., 1992. *Nunavut Atlas,* s.l.: Canadian Circumpolar Institute and Tungavik Federation of Nunavut.

ᖁᑦ-Ross, M. V., Alisauskas, R. T., Douglas, D. C. & Kellett, D. K., 2017. Decadal declines in avian herbivore reproduction: density-dependent nutrition and phenological mismatch in the Arctic. *Ecology,* 98(7).

ᓴᐸ, ᓯ-Sharp, C. M. & Abraham, K. F., 2010. *Status of Atlantic Brant breeding at East Bay, Southampton Island: 30 years later,* Peterborough, ON.: Poster to the North American Arctic Goose Conference.

ᓴᒥᑦ, ᐸ-Smith, P. A., Elliott, K. H., Gaston, A. J. & Gilchrist, H. G., 2010. *Has early ice clearance increased predation on breeding birds by polar bears?,* s.l.: Polar Biol.

ᓴᒥ, ᐸ-Smith, P. A., Johnston, V. & Rausch, J., 2012. Southampton and Coats Islands. In: *Arctic Shorebirds in North America: a decade of monitoring.* Berkeley(California): Studies in Avian Biology (no. 44), University of California Press.

ᓴᒍ, ᔨ-Smol, J. P. & Douglas, M. S., 2007. *Crossing the final ecological threshold in high Arctic ponds,* s.l.: PNAS.

ᐅᓴᐃᔨᑦ ᑲᓇᑕ-Statistics Canada, 2016. *Coral Harbour, HAM [Census subdivision], Nunavut and Nunavut [Territory] (table). Census Profile. 2016 Census Census of Population,* Ottawa: Statistics Canada Catalogue no. 98-316-X2016001.

ᓴᑎᕐᓴᐅ, ᓂ ᐊᓚᓗ Lᑯᐊᒥᑲ, Ṗ-Stephenson, S. M. & McCormick, K., 1986. *Draft Harry Gibbons Bird Sanctuary Management Plan,* s.l.: Canadian Wildlife Service.

ᓴᑎᐟ, ᔨ-Sutton, G., 1931. *The blue goose and lesser snow goose - Southampton Island, Hudson Bay,* s.l.: Auk, Vol. 48.

ᓴᑎᐟ, ᔨ-Sutton, G., 1932. *The Exploration of Southampton Island, Hudson Bay,* s.l.: Memories of the Carnegie Museum, Vol. 12.

ᕽᐢ ᓴᑐᐟ-Van Stone, J. W., 1960. *Notes on the Economy and Population Shifts of the Eskimos of Southampton Island,* Toronto, ON: Department of Anthropology, University of Toronto.

ᑎᑎᕋᕆᐃᑦ ᐃᓕᓴᕆᓂᖅᓯᒪᔪᑦ

ᑎᑎᕆᕼᐃᕐ ᐃᓕᓴᕼᓐᓂᕐᒥ A: ᒪᓕᑦᑐᕼᑎᒍᑦ ᐅᓂᒃᑲᖅᑕᐅᑎᐊᖅᓂᕐᒥ ᓄᓇᐃᕐ ᓇᓂᓐᓂᕐᒥ

ᒪᓕᑦᑐᕼᑎᒍᑦ ᐅᖅᑲᐅᓗᐅᓗᒪᕐᒉᑦ ᐃᔨᔪᐊᖅᑕᐅᓗᒪᓪᓕᑐᑎᒃ ᓇᖏᒥᓂᖅ ᑎᑎᑕᖅᓗᒪᓂᕐᒥ ᑕᐃᒃᑯᓇᖏᑦ ᑎᖏᒥᐊᖅᑲᕆᓐᓄᑦ ᒪᓕᑦᓪᓗᑕᑦᑲᐅᔨᓂᒃ (*C.R.C., c. 1036*; 1958) ᐃᒪᐢᓇᐃᓕᓴᕐᓗᔨᓂᒃ:

“ᕼᐊᐅᓕ ᑭᔪᓵ ᑎᖏᒥᐊᖅᑲᕆᕆ

ᑕᒪᖏᒃ ᐃᓕᑭᕆᐅᔨᑦ ᓄᓇᖏᐢᓄᑦ ᓴᖅᓕᐅᔨᔪ ᖃᑭᖅᑕᓪᓕᐊᓄᑦ ᐃᒪᖏᑭᓄᓪ ᕼᐊᑦᓴᓵᔨᔪᐊᖅᑎᓪᓕᐊᓂᑦ, ᐊᕆᒃᑐᖅᓯᒪᓕᖏᓐᓄ ᑭᐢᓕᓚᐅᔨᔪ, ᐃᓄᓇᓯᐊᖅᑎᒥ, ᐃᓕᑭᕆᐅᓕᖏᑭᓂ ᐱᓪᓕᐊᖅᑐᒥᒃ ᐅᓕᒃᑲᖅᑕᐅᑦᓂ ᐃᒪᐢᓇᐃᓕᓴᕐᓗᓂᕆᓕᖅᑐᐃᓕᖅᑐᒃ:

*ᐱᑭᖅᐊᕆᖅᑲᓕᓂ ᑕᐃᓐᖏᓐᓂᓕᓂᒃ ᐃᒪᖅᑲᕆᐊᐅᖏᓐᓂᓕᓂᒃ ᐃᑲᑭᑐᐊᖅᑎᒥ, ᑐᑭᑦᑐᐊᖅᑎᒥ ᐃᑲᑭᑐᐊᖅᑎᒥ ᑐᑭᑦᑐᐊᖅᑎᒥ ᐃᑲᑭᑐᐊᖅᑎᒥ ᐃᒪᐢᓇᐃᓕᓴᕐᓗᓂᕆᓕᖅᑐᐃᓕᖅᑐᒃ

ᖃᖅᖢᐅᖅᑐᑎᒥ (East Bay-ᒥ) ᐊᓚᓗ ᐃᒃᑐᐊᖏᒥ (ᕼᐃᐅᓇ ᒧᐸᓴᒥ) ᑎᓴᒥᐊᖃᖏᕐᓱᒃ -

ᑭᓴᓪᓕᖅᐸᒥᒃ ᑎᑎᕋᖅᓯᒪᓕᖅᑐᑦ ᐊᐅᓪᓚᑎᓂᕐᒧᑦ ᐸᖏᓐᓄᑦᓯᐊᕐᓂᖅᓴᐃᑦ ᖃᖅᖢᐅᖅᑐᑎᒥᒃ ᐊᓚᓗ ᐃᒃᑐᐊᖏᒥᒃ ᑎᓴᒥᐊᖃᖏᕐᓱᓐᓂ ᐅᒪᔨᓕᓚᔪᖏᐊᑲᑯᓗᒃ ᐊᓐᓕᖅᑕᐅᔨᕐᓴᐅᑕᐅᓪᓗᓂᒃ

QAQSAUQTUUQ (EAST BAY) AND IKKATTUAQ (HARRY GIBBONS) MIGRATORY BIRD SANCTUARIES -

Final Draft Management Plans for Nunavut Wildlife Management Board's Approval

ᓄᓇᕗᑦ ᐅᒪᔨᓕᓚᔪᖏᐊᑲᑯᓗᒃ ᑲᑎᒪᓂᓵᓐᓚᓱᓐᓂᒃ ᑲᑎᒪᓂᓐᓇᓐ, ᓇᔭᐅᑎᓕᒃ RM 001-2020, ᐃᖃᓗᐃᑦ, 11 ᐄᓕᔪ, 2020

NWMB RM 001-2020, Iqaluit, 11 March 2020

ᐃᖏᓂᐅᖏᖅ ᐊᐅᓚᓕᔪᖏᐊᑲᑯᓗᒃ ᐃᓕᓴᕐᓂᖅ

IRNIURVIIT AREA CO-MANAGEMENT COMMITTEE

ᐅᓂᖅᑲᓕᐊᓂᖅ ᑕᒪᐃᓐᓂᖅ ᑐᑭᓯᑎᕆᑎᑭᐊᕐᓂᖅ
Presentation Overview

- ᖃᐅᔨᒪᑎᕆᑎᓴᐸᑎᖅᓴᐃᓕᕆ ᓴᓐᓂᐊᖅᑕᐅᓯᒪᓂᖏᓐᓂᖅ ᓄᓇᐃᕆ ᓄᓇᕗᓴᒥ
- ᑕᑯᖅᓴᐅᑎᕆᑎᔪᕆ ᐃᓗᓕᖏᓐᓂᖅ ᑎᑎᕆᖅᑕᐅᓯᒪᓂᖏᓐᓂᖅ ᓯᓕ ᐊᐅᓪᓕᕆᓂᕐᑎᓂᕐᒧᑦ ᐸᓇᐅᑎᖅᓴᓂᖅ



ᒥᑎᕕᖅ Common Eider

- Information on protected areas in Nunavut
- Show you the content of the Draft Management Plans



ᑐᖕᕕᑎᑎᓵᖅ Ruddy Turnstone

ᖃᐅᔨᒪᑎᑦᑎᓕᐊᕈᑎᑦ ᑲᓇᑕᒥ
ᐅᒪᔨᖃᖅᖁᐅᑎᑕᐅᔨᓂᒃ ᐊᓚᒪᓗ
ᑎᖏᒥᐊᖃᖅᖁᐅᑎᑕᐅᔨᓂᒃ ᓄᓇᕗᓚᒥ

Information on National Wildlife Areas and Migratory Bird Sanctuaries in Nunavut

ᐊᓂᑎᓪᓗᖅᐸᐃᑦ ᑲᓇᑕᒥ ᓄᓪᓕᐅᕐᓂᓗ
ᐊᓯᔾᔨᖅᐸᓪᓚᐊᓂᖏᓐᓂᑦ ᑲᓇᑕᒥ
Environment and Climate Change Canada

ᐱᔾᔨᑕᐅᓂᖅᑎᑦ ᐱᓚᑦᖃᑎᒋᔭᑕᐅᓂᖅᓂᕐᒧᑦ:


- ᓴᐳᓄᐊᖅᑕᐅᓯᒪᔪᑦᐊᖃᖅᓂᖅᑎᓄᑦ
- ᖃᐅᔨᓴᖅᑕᐅᖃᑕᑦᐊᖅᓂᖅᑎᓄᑦ
- ᑐᑭᓯᓇᖅᓯᑎᑕᐅᔪᑦᐊᖃᖅᓂᖅᑎᓄᑦ

ᓄᓇᑕᒥ ᐅᒪᔨᖃᖅᓴᑦ ᓴᐳᔾᔪᐅᓯᒪᓕᕐᓇᖅᑎᑦ
ᑎᖅᑎᐊᖃᖅᓴᑦ ᓴᐳᔾᔪᐅᓯᒪᓕᕐᓂᖅᑎᑦ

Purpose of Network

- Conservation
- Research
- Interpretation

National Wildlife Areas (NWA)
Migratory Bird Sanctuaries (MBS)



ᐊᖑᓕᓂᖅᑎᑦ 12 ᒥᓕᒃᖑᓪ
ᐄᐃᐊᖅᑕᓂᖅ ᑕᒪᖏᖅ ᓄᑎᓪᓗᓕᑦ
ᐊᖑᓕᖑᓕᑦ
12 million hectares in total


ᐊᓂᑎᓕᓵᕐᓯᐊᖏᑦ ᑲᓇᑕᒥ ᓯᓚᐅᔪᓐᓇ ᐊᓱᓴᖅᐸᔪᓕᐊᓂᖓᓂᑦ ᑲᓇᑕᒥ
Environment and Climate Change Canada

	ᑎᖅᒥᐊᖅᑲᓇᐃᑦ ᓴᓗᐅᔨᒪᕐᓇᖅᑐᑦ	ᑲᓇᑕᒥ ᐅᒪᔨᖅᑲᓇᐃᑦ ᓴᓗᐅᔨᒪᕐᓇᖅᑐᑦ
ᓴᓗᓂᐊᕐᑕᐅᓂᖅᓗᑦ ᑐᖏᓚᖅᖃᑐᑦ	ᑎᖅᒥᐊᑦ	ᑕᒪᖅᑎᑲ ᐅᒪᔨᐃᑦ
ᓴᓗᓂᐊᖅᑕᐅᔨᓂᖅᑭᓇᓄᑦ ᐊᑐᖏᓂᖅᑎᑦ	ᐅᓪᓗᖅᑲᖏᓐᓇᖅᑐᑦ /ᐃᓕᓴᖏᓐᓇᖅᑐᑦ/ ᓄᓕᐊᖏᓐᓇᖅᑐᑦ	ᐊᖏᕿᔭᓕᒫᖅ
ᒪᓕᒐᖅᑎᔪᑦ	*ᑎᖅᒥᐊᖏᓪᓕᓚᓂᓂᕆᑲᖅ ᑲᑎᒪᖏᖏᖅᑲᑕᐅᔨᒪᔨᓂᔨᑦ ᒪᓕᒐᐃᑦ*	*ᑲᓇᑕᒥ ᐅᒪᔨᓕᓚᓂᓂᖏᓪᓗᑦ ᒪᓕᒐᐃᑦ*

	MBS	NWA
Protection Focus	Migratory birds	All wildlife
Protection Period	Nesting/breeding season	Year-round
Legal Basis	*Migratory Birds Convention Act*	*Canada Wildlife Act*

ᐃᓄᐃᑦ ᐊᖏᑐᕐᑕᐅᓯᒪᓂᖅᐸᓂᖅ ᐱᓕᕆᓂᕐᑕᐅᔨᓐᓇᖅᑐᓂᒃᑲᓂᕐᓗ ᐊᖏᕈᓐᓇ (2008; 2016)

Inuit Impact and Benefit Agreement (2008; 2016)





ᐃᓄᐃᑦ ᐊᖏᑐᖅᑕᐅᓯᒪᓂᖅᐸᓄᓄᑦ ᐱᓱᒋᑦᑕᐅᔪᓐᓇᖅᑐᓂᓗ ᐊᖓᔪᖅᑎᑎ (2008; 2016)

Inuit Impact and Benefit Agreement (2008; 2016)

- ᐃᓕᒪᓕᐅᑭᑕᐅᓱᖏᑐᑦ ᐱᔭᕆᑎᖏᖅᑐᑦ ᑎᓪᓗᐊᖏᖅᓴᖅᓂᒃ ᐊᒻᒪᓗ ᐅᒪᔨᖏᖅᓴᖅᓂᒃ ᓴᖅᑭᔨᒥᒃ ᐃᓄᐃᑦ ᖃᐅᔨᒪᔭᖏᖅᖢᑦ ᑕᐃᒃᑯᓂᖅᓴ **ᐃᓄᐃᑦ ᖃᐅᔨᒪᔭᑐᖏᖅᓂᒃ**

- ᐃᓕᑕᒫᔭᐅᓕᕆᑦ ᐊᐅᓪᓚᑎᐅᖏᖅᑕᐅᔭᒫᐊᖏᖅᓂᖅᑭᑦ ᐃᓄᐃᑦ ᑎᓪᓗᐊᖏᖅᓴᖅᓂᒃ ᐊᒻᒪᓗ ᐅᒪᔨᖏᖅᓴᐅᔨᓂᒃ ᐱᒻᒪᒫᐅᓂᖅᑭᑦ ᐊᖓᔪᖅᑎᑎᐅᓕᖅᑐᓂ: **ᐊᐅᓪᓚᑎᐅᖏᖅᑕᐅᓂᖏᓐᓄᑦ ᑲᑎᒪᔨᖁᓪᓗᑦ**


- Decisions for Migratory Bird Sanctuaries and National Wildlife Areas strongly influenced by **Inuit Qaujimajatuqangit**

- Guarantees co-management of Migratory Bird Sanctuary and National Wildlife Areas
  Important feature: **Area Co-Management Committees**

ᐊᐅᓪᓗᕆᐅᖅᑕᐅᓂᖏᓐ ᑲᑎᒪᔨᕋᓪᓕᐅᑦ
Area Co-Management Committees (ACMCs)

- 9-ᖃᓄᑦ ᐊᐅᓪᓗᕆᐅᖅᑕᐅᓂᖏᓐ ᑲᑎᒪᔨᕋᓱᒃᑐᖅᑎᑦ ᓄᓇᕗᒻᒥ, ᐊᐅᓪᓗᕆᐅᖅᑕᐅᓯᒪᔪᑦ 13-ᖃᓄᓂᒃ ᓴᓇᔨᓂᐊᖅᑕᐅᓕᒫᖅᓂᐊᓂᒃ ᓄᓇᓂᒃ
- ᑲᑎᒪᔨᖅᓯᖃᓪᓚᑎᒃ ᑕᒪᓕᒪᓂᒃ 5 ᐃᓄᖃᓂᒃ ᖁᓂᑭᖏᑐᓂ ᓄᓇᓕᓂᖃᖅᓯᒪᔪᓂᒃ ᐊᓪᒪᓗ ᐃᓕᖁᖃᓗᑎᒃ ᐊᑕᐅᓯᕆᑦ 1 ᑲᓇᑕᒥ ᐅᒪᔨᓕᕆᔨᒋᑦ ᐱᓕᕆᓚᖏᑐᓂ (ᐊᓯᑎᓕᕆᔨᒋᐊᓂᒃ ᓯᓚᐅᒪᓗ ᐊᓯᔨᕐᓴᒪᓕᐊᓂᖏᓐᓂᒃ)

- 9 ACMCs were created in Nunavut to co-manage the 13 protected areas
- Made up of 5 people from the associated community and 1 from Canadian Wildlife Service (Environment and Climate Change Canada)



ᐊᐅᓪᓕᑎᕆᐅᖏᑕᐅᓂᖅᓯᒪᓂ ᑲᑎᒪᔨᖏᓪᓕᓂ
Area Co-Management Committees (ACMCs)

- ᐃᖅᓂᐅᕆᖃᑦᓂᓪ ᐊᐅᓪᓕᑎᕆᐅᖏᑕᐅᓂᖅᓯᒪᓂ ᑲᑎᒪᔨᖏᓐᓇᖅᑐᓂ ᑲᑎᒪᔨᖅᓯᓪᓕᐅᖅᓴᓂ 2012-ᒥ
- ᑕᓪᓕᒪᓂᒃ 5 ᑲᑎᒪᔨᖏᖅᓴᓂ ᓴᓪᓕᖅᑎᐅᑕᓂᒃ
  - ᓄᐊ ᑲᓪᓚᒃ (ᐃᒃᓯᐊᐅᑕᖅ)
  - ᐊᓂ ᓂᖏᐅᓐᓚᓐ
  - ᔭᐃᓴ ᑲᓪᓚᒃ
  - ᐅᐃᓕ ᐊᑕᒪᓴ
  - ᐊᑕᐅᓯᖅ ᐃᓄᖏᖏᑦᑐᖅ
- ᐊᑕᐅᓯᖅᑎᒃ 1 ᑲᓇᑕᒥ ᐅᒪᔨᓕᓚᔨᒃᑯᑦ ᐱᓕᓚᐃᖏᓐᓂ ᑲᑎᒪᔨᖏᓚᐅᙱᑦ: ᐸᐅᓵ-ᐃᐊᐟ ᑐᐅᐊ (ᐃᒃᓯᐊᐅᑕᐅᐸ ᑐᖏᓕᓕᖏᓂᒃ)
- ᐃᑲᔨᖅᑕᐅᔨᖅ ᐅᒧᖏᒐ ᑲᓇ ᓂᖏᐅᓐᓚᓐ (ᑭᐊᓕᓕᖅᒥ ᐃᓄᐃᑦ ᑲᑐᐸᔨᖅᑎᑦᑎᑦ)

- Irniurviit ACMC was formed in 2012
- 5 members from Coral Harbour
  - Noah Kadlak (Chair)
  - Annie Ningeongan
  - Louisa Kudluk
  - Willie Adams
  - Vacant
- 1 member from Canadian Wildlife Service: JF Dufour (Vice-Chair)
- Supported by Ron Ningeongan (KIA)



2012



2018

ᐃᖏᓂᐅᕐᕕᐊᑦ
ᐊᐅᓪᓗᓯᐅᖅᑕᐅᓂᖓᓪᓚᑦ
ᑲᑎᒪᔨᖏᓪᓕᖅ


IRNIURVIIT AREA
CO-MANAGEMENT
COMMITTEE

ᒪᖏᕈᒃ ᑎᖅᑭᐊᖅᑲᓇᐃᒃ
ᓴᐳᓂᐊᖅᑕᐅᓯᒪᖕᓇᐅᔭᓕᑎᒃ:

- ᖃᓄᖅᓴᐅᖅᑐᖅᑎ
- ᐃᓄᓕᑐᐊᖅᑎ

Two Migratory Bird Sanctuaries:

- Qaqsauqtuuq
- Ikkattuaq



ᐊᐅᓪᓗᓂᐅᖅᑕᐅᓂᖏᑦ ᑲᑎᒪᔨᖏᓪᓕᓪᓂ
Area Co-Management Committees (ACMCs)

ᐊᐅᓪᓗᓂᐅᖅᑕᐅᔾᔨᓂ ᑲᑎᒪᔨᖏᓪᓕᓯᓪᓂ
ᐃᒪᓐᓇᐃᓕᐅᖅᑕᕐᓂᓕᕆᖅᑐᑦ:

- ᐅᖅᐅᓴᔨᕆᕕᖅᑎᐅᖅᑕᕐᓂᔭᓐᓄᑦ ᒥᓂᔾᑕᒧᑦ ᑕᒪᐃᓐᓇᓂᕆᕕᖅ ᐱᔾᔪᑎᖅᖅᑐᓂᑦ ᑎᓪᒥᐊᖅᕿᓐᓂᓂᑦ/ᐅᒪᔪᖅᕿᓐᓂᓂᑦ ᐊᐅᓪᓗᓂᑎᓂᕐᒥᑦ
- ᖃᐅᒥᖅᑕᐊᖅᑕᕐᓂᔭᓐᓄᑦ ᐱᔪᓐᓇᐅᑎᑦᓴᑕᕐᓂᔪᓐᓇᕐᓂᖏᑦ ᐅᑦᑐᖅᑕᐊᖅᑎᓂᑦ ᑐᑦᔪᖅᑐᐅᑎᓂᑦ
- **ᓴᓇᖅᑕᕐᓂᔭᓐᓄᑦ ᐊᐅᓪᓗᓂᑎᑦᔪᑎᑦᓴᓂᑦ ᐸᖅᓇᐅᑎᓂᑦ**



ACMC will:

- advise the Minister on all aspects of MBS/NWA management
- review permit applications
- **develop Management Plans**



ᐊᐅᓪᓗᒍᓐᓇᖅ ᐸᖅᓇᐅᑎᖏᑦ
Management plans


- ᐱᔨᓇᐅᑕᐅᕙᒃᑐᑦ ᐊᐅᓪᓗᒍᑦᑏᐅᖅᑕᐅᓂᖅᓗᒍ ᑲᑎᒪᔨᕆᓪᓕᑦ ᐅᖃᐅᔨᖅᑭᓇᖅᓂᖏᓐᓂᒃ ᑕᑯᑎᑦᑎᔨᓇᖅᓂᖅᓗ ᑕᐅᑐᑎᓴᖏᓐᓂᒃ ᖃᓄᐃᓕᓐᖏᓴᕆᐊᖃᖅᓂᖏᓐᓂᒃ ᓴᐳᓂᐊᖅᑕᐅᔨᒪᔨᑦ
- ᓇᓗᓇᐃᔭᐃᔨᐅᕙᓪᓗᑎᒃ ᒪᓕᒃᑕᐅᓂᐊᖅᑐᓂᒃ ᐃᔨᒪᓕᐅᖅᑎᐅᕙᓐᓂᖅ ᐱᔪᒥᑎᖃᖅᑐᓂᒃ ᓴᐳᓂᐊᖅᑕᐅᔨᒪᔨᓂᒃ ᓄᓇᓂᒃ
- ᐅᓂᒃᑲᖅᐸᒃᑐᑦ ᐱᓕᒪᓗᐅᓂᖏᓐᓂᒃ ᐃᓕᓐᖅᔨᑐᑎᖅᑐᑦ ᐊᕐᕌᒍᖅᑐᑦᓯᓗ ᖃᖅᑎᖏᓐᓂᒃ ᖃᓄᖅ ᐱᓕᓐᖅᔨᓇᓐᖏᓐᓂᒃ ᓄᓇᖏᑦ

- Allows the Area Co-Management Committees to share their vision of the protected area
- Guides decision making for the protected area
- Describes important cultural and environmental aspects
- States which activities are permitted and not permitted





ᐊᐅᓪᓗᐊᓂᖅᓴᐃᓪᓕᐅᑎᓪ: ᐱᓕᕆᐊᖓᓂᒃᓕᐊᖅᑕᐃᓇᓱᓂᖅ Management Plans: *Steps*



ᐊᐅᓪᓗᒍᓂᐊᕐᓗᒃ
ᐸᖁᐅᓯᐅᖅᐸᑦᓂᐊᓪᓚᐅᖁᓂᖏᑦ
ᐊᒻᒪᓗ ᐊᖏᑎᑐᑎᒃᑲᒧᐃᓱᖏᑦ

Management Plan Preparation and Approval

ᐃᓄᐃᑦ ᐊᑲᑐᖅᑕᐅᔪᓕᓂᖅᑭᓂᓄᑦ ᐱᓱᕆᓪᓕᖅᑐᐃᓐᓇᑦᑐᓂᓪᓗ ᐊᖏᑭᑭᑎ
Inuit Impact and Benefit Agreement

ᐊᐅᓪᓚᑦᑎᐹᐊᖅᓴᓄᑦ ᕿᖏᐅᑎᑦ ᐱᓕᕆᔨᐅᓂᖅ

- 3.5.4 ᑎᑎᕋᖅᑕᐅᓗᑎᑉ ᐊᖏᕐᓂᖅᑕᐅᓗᑎᑉ ᐊᐅᓪᓚᑦᑎᐹᐊᖅᓴᓄᑦ ᕿᖏᐅᑎᑦ, ᐅᓄᐊ ᐊᐅᓪᓚᑦᑎᕆᐊᖃᕐᓗᑦ ᑲᑎᒪᔨᕋᓱᑦ ᐃᓕᓴᕐᑎᑕᐅᔪᑦ ᐃᓄᐃᑦ ᖃᐅᔨᓴᑐᖃᖅᑕᖏᑦ ᑐᓴᕐᓂᓕᕆᓕᕐᑕᐅᔪᑦ ᐃᓚᒋᔭᐅᔪᓐᓇᑦ.
- 3.5.5 ᐅᓄᐊ ᐊᐅᓪᓚᑦᑎᕆᐊᖃᕐᓗᑦ ᑲᑎᒪᔨᕋᓱᑦ ᖁᒥᖅᑭᐊᖃᖅᑎᖅᑎᖅᓯᓗᑎᑉ ᐅᖃᖅᖃᐊᖃᖅᑎᖅᓯᓗᑎᑉ ᐊᖑᑎᓗᒍ ᐃᓄᐃᑦ ᑲᑎᒪᔨᕋᓱᑦ ᐃᓄᐃᑦ ᑲᑎᒪᔨᕋᓱᑦ ᐃᓕᓴᕐᑎᑕᐅᔪᑦ, ᐊᒻᒪᓗ ᓄᓇᕗᑦ ᑐᓴᖏᓐᓄᑦ ᑎᒥᖓ, ᐱᖁᔪᓕᐅᓴᖏᓂᖓᓂ ᐅᖁᑎᑎᑐᑎᒍ ᐊᐅᓪᓚᑦᑎᐹᐊᖅᓴᓄᑦ ᕿᖏᐅᑎᑦ.
- 3.5.6 ᒪᓕᕆᐊᓕᖅ ᐱᓇᖁᐅᓂᐊᖃᖅᑐᓂ ᕿᖏᐅᑎᓂ ᐊᒻᒪᓗ ᑭᓇᐅᖁᖃᖅᑐᑎᓂᖅ, ᐅᓄᐊ ᐊᐅᓪᓚᑦᑎᕆᐊᖃᕐᓗᑦ ᑲᑎᒪᔨᕋᓱᑦ ᑎᑎᕋᖅᑕᐅᓗᑎᑉ ᐊᖏᕐᓂᖅᑕᐅᓗᑎᑉ ᕿᖏᐅᑎᑦ ᒪᓕᕆᓪᓗᒍ ᖃᓄᖅᑭᐅᓂᖅᑐᑦ ᐃᓄᐃᑦ ᐊᑲᑐᖅᑕᐅᓗᑎᑉ ᐊᖏᕐᓂᖅᑕᐅᓗᑎᑉ ᐊᒻᒪᓗ ᖁᒥᖅᑭᐊᖃᖅᑎᖅᓯᓗᑎᑉ ᐅᖃᖅᖃᐊᖃᖅᑎᖅᓯᓗᑎᑉ ᐅᓄᐊᓕᖅᑐᑦ ᐃᓄᐃᑦ ᐊᑲᑐᖅᑕᐅᓗᑎᑉ ᐊᖏᕐᓂᖅᑕᐅᓗᑎᑉ ᐅᖃᖅᖃᐊᖃᖅᑎᖅᓯᓗᑎᑉ ᐅᓄᐊᓕᖅᑐᑦ.

MANAGEMENT PLAN PREPARATION

- 3.5.4 In preparing Management Plans, the ACMCs shall carefully consider any *Inuit Qaujimajatuqangit* brought forward by a member.
- 3.5.5 The ACMC shall consult the relevant RIA, and NTI, before completing the draft Management Plan.
- 3.5.6 Subject to its work plan and budget, the ACMC may prepare the Management Plan according to whatever process it deems appropriate and it may consult as it deems appropriate.

ᐃᓄᐃᑦ ᐊᖏᑕᖅᑕᐅᓯᒪᓂᖅᐸᑦᓄᓇᓕᒃ ᐱᓕᕆᔨᑦᑕᐅᔭᕆᓇᖅᑕᑦᓂᓪᓗ ᐊᖓᔪᖅᑎᑦ
Inuit Impact and Benefit Agreement

3.5.7 ᐅᓄᐊ ᐊᐅᓪᓚᑦᑎᐊᑎᒃᓴᓄᑦ ᐸᖁᓇᐅᑎᑦ ᐃᓪᓚᖅᓂᐊᖅᑐᑦ ᓇᓗᓇᐃᔭᑦᑎᐊᖅᓯᒪᔨᓂᒃ ᐅᓄᓂᖓ:

(a) ᐱᓵᔨᑎᖓᑦ ᑲᓇᑕᒥ ᐅᒪᔨᖅᑲᓐᓇᑦ ᐅᕝᕙᓕᓂᑦ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᖓᒥᐊᓄᑦ ᓴᓵᐳᓕᒫᒃ;

(b) ᐊᐅᓪᓚᑦᑎᐊᑕᐅᔨ ᑐᖏᖓᓐᖓᑦ ᐊᒻᒪᓗ ᑐᑭᓯᐅᑉᐸᑎᑦᑎᓕᕆᓂᖅᐸᑦᑐᑦ;

(c) ᓄᓇᖓᓐ ᐊᒻᒪᓗ ᐱᐅᔨᑐᖅᖓᓐᓂ ᐃᒻᒪᑲᓪᓚᒃ ᖃᓄᐃᓕᖓᓂᖓᓐ ᐊᒻᒪᓗ ᐃᓗᓕᖓᑦ ᐃᓗᐊᓂ ᑦᑲᒃᑯᐊ ᑲᑎᒪᔨᖏᑦ ᐅᕝᕙᓕᓂᑦ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᖓᒥᐊᓄᑦ ᓴᓵᐳᓕᒫᒃ ᐊᐅᓚᓂᖓᑦ;

(d) ᐊᑐᐊᖏᐃᑦ ᐅᓄᐊᓗ ᑐᑭᒧᐊᔪᑕᐅᓂᐊᖅᐸᑦᑐᑦ ᐊᐅᓚᑦᑎᐊᑕᐅᔨᓄᑦ ᑲᑦᓄᓂᖓᓐ ᖃᓄᐃᓕᖓᑦ ᓂᕿᑎᓄᑦ ᖃᓄᐃᓕᖓᑦ ᐅᓪᓗᕆᔨᑕᐅᓂᐊᖅᐸᑦᑐᑦ ᐊᐅᓚᑦᑎᐊᑕᐅᔨᓄᑦ ᓴᓵᐳᓕᒫᒃ ᓴᓵᐳᓕᒫᒃ;

(e) ᐅᓪᓗᖓᑦ ᐊᑐᓚᖅᑎᑕᐅᕐᒪᑦᓴᖓᑦ ᐊᐅᓚᑦᑎᐊᑕᐅᔨᑦ ᐸᖁᓇᐅᑎᖓᑦ ᖃᓄᐃᓕᓚᐅᖅᑕᐸᑦᑐᑦ ᑭᓯᓂᖓᑦ.

3.5.7 The Management Plan shall include a description of:

(a) the purposes of the NWA or MBS;

(b) management goals and objectives;

(c) the natural and cultural history and the context within which the NWA or MBS operates;

(d) policies that will guide the management of the NWA or MBS;

(e) a schedule to implement Management Plan action items.

ᐃᓄᐃᑦ ᐊᖏᑐᖅᑕᐅᓯᒪᓂᖅᐹᓪᓕᐊᓂᑦ
ᐱᓵᓴᓕᑕᐅᔨᓴᓕᖅᑐᓄᓪᓗ ᐊᖅᑎᕈᑎ
Inuit Impact and Benefit Agreement

3.5.8 ᑕᐃᒃᓂᓂ ᓄᓇᑕᒥ ᐅᒪᔨᖅᑎᒋᐊᑦ ᐅᓵ�??ᓕᓂᓂ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᖅᒥᐊᓄᑦ ᓴᓗᐃᑦᐅᓯᒪᔪᓂ ᐃᓕᖅᑐᖅᑐᑦ ᐃᓄᐃᑦ ᓄᓇᒋᐊᑎᖅᑎᑦ, ᐊᐅᓕᑦᑎᐹᑎᖅᓴᓄᑦ ᐸᒋᓇᐅᑎᑦ ᒪᓕᑦᑎᐊᑦᑎᐊᒋᓂᖅᑐᑦ ᐊᒻᒪᓗ ᐅᖃᖅᓯᒪᓗᑎᒃ ᓇᓪᓕᐊᓂᖅᑐᓂᒃ ᐃᓄᐃᑦ ᓄᓇᒋᐊᑎᖅᑎᓯᓂ.

3.5.8 Where an NWA or MBS includes IOL, the Management Plan shall reflect and address any special issues arising from the presence of the IOL.

ᐃᓄᐃᑦ ᐊᖏᑐᖅᑕᐅᓯᒪᓂᖅᐹᓂᑦ ᐱᓕᕆᓂᖅᑕᐅᔪᓐᓇᖅᑐᓄᑦᓗ ᐊᖅᑭᑦᑎᐊᕐᓂᖅ

Inuit Impact and Benefit Agreement

ᐊᐅᓪᓚᑦᑎᐊᑎᒃᓴᓄᑦ ᕿᖏᐅᑎᑦ ᐊᖅᑭᑦᑕᕆᔨᒪᓂᖅᓄᖅ

3.6.1 ᐊᐅᓪᓚᑦᑎᖃᑎᒋᔭᖅᑐᑦ ᑲᑎᒪᔨᕆᓪᓚᑦ ᐊᑐᓪᖏᑦᑕᐅᔾᔨᓗᑎᒃ ᐱᓕᕐᓂᑎᐊᖅᓯᒪᔪᑦ ᐊᐅᓪᓚᑦᑎᐊᑎᒃᓴᓄᑦ ᕿᖏᐅᑎᑦ ᑕᒃᓯᓄᖅᓱᒃ ᓄᓇᕗᒻᒥ ᐅᒪᔪᓕᕐᓇᔨᕐᔪᐊᑲᓯᓄᖅ ᐊᖅᑭᑦᑕᐅᓗᑎᒃ ᒪᓕᒐᓗᔫ ᐊᖅᑭᑦᑎᐊᓂᖅ 5.2.34 (c) ᐊᒻᒪᓗ 5.3.16 *ᓄᓇᕗᒻᒥᒃ ᓄᓇᕈᑎᑎᓄᑦ ᐊᖅᑭᑦᑎᐊᓂᖅ.*

3.6.2 ᐅᓯᐊ ᐊᐅᓪᓚᑦᑎᖃᑎᒋᔭᖅᑐᑦ ᑲᑎᒪᔨᕆᓪᓚᑦ ᑐᓂᔪᓂᕐᒧᑦ ᐊᑐᑎᔪᓄᑦ ᐊᕐᕌᒃᑐᖅᓯᒪᔪᑦ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖃᑎᒋᔭᖅᑐᑦ ᐊᒻᒪᓗ ᓄᓇᕗᑦ ᑐᓴᖅᓱᒃ ᑎᒥᖅᑏ ᑕᕐᕋᖅᓱᓂ ᐱᓕᕆᖃᑎᒌᓕᕐᒪᑦᑐᑦ ᐊᐅᓪᓚᑦᑎᐊᑎᒃᓴᓄᑦ ᕿᖏᐅᑎᑦ ᐅᓯᐊᓗ ᐊᐅᓪᓚᑦᑎᓕᕐᒪᑐᓂ ᕿᖏᐅᑎᑦ ᓄᓇᕗᒻᒥ ᐅᒪᔪᓕᕐᓇᔨᕐᔪᐊᑲᓯᓄᑦ.

MANAGEMENT PLAN APPROVAL

3.6.1 The ACMCs shall recommend completed Management Plans to the NWMB for approval in accordance with sections 5.2.34(c) and 5.3.16 of the *NLCA.*

3.6.2 The ACMC shall provide the relevant RIA and NTI with a copy of the completed Management Plan when it sends the Plan to the NWMB.

ᐃᓄᐃᑦ ᐊᑲᑐᖅᑕᐅᓯᒪᓂᖅᐹᓂᓄᑦ ᐱᓵᓴᓕᐅᔪᓐᓇᖅᑐᓄᓪᓗ ᐊᖏᑎᑕᐅᑎ
Inuit Impact and Benefit Agreement

3.6.3 ᑕᐃᒪᓗ, ᒪᓕᒃᓗᒍ ᐃᓱᒪᓕᐅᓇᓵᓕᐊᔨᑦ ᐅᖃᖅᓱᒪᔪᑦ ᐊᖏᑎᑕᐅᑎᑦ 5.2.34(c) ᐊᒻᒪᓗ 5.3.17 ᑎᑭᒃᓗᒍ 5.3.23 ᓄᓇᕗᒻᒥᒃ ᓄᓇᑕᑭᑎᓄᑦ ᐊᖏᑎᑕᐅᑎᑦ, ᐅᓂᐊ ᓄᓇᕗᒻᒥ ᐅᒪᔪᓕᓇᓵᖅᓯᐊᕐᓂᖅᓂᑦ ᐅᓴᐅᓂᑦ ᐊᑲᒃᖢᑕᐅᑦ, ᑕᒪᖏᒃ ᐅᓴᓵᓐᓂᑦ ᐃᓕᓐᓄ, ᐱᔨᑦᑎᐊᖏᒪᔪᖅ ᐊᐅᓕᑎᑎᐊᕐᕕᓄᑦ ᐸᖃᐅᑎᑦ ᐊᒻᒪᓗ ᐸᖃᐅᑎ ᐅᑎᖏᓗᓂ ᑕᒃᓂᓄᖑᓐᓂᖏᖏᑦ ᖃᑎᒪᔨᕆᓕᑦ ᐃᓂᒪᑭᐅᑎᐊᒃᓂᖏᑦᑐᑦ, ᐊᒃᑐᑎᔪᑦ ᐊᐅᓕᑎᑎᐊᕐᕕᓄᑦ ᖃᑎᒪᔨᕆᓕᑦ ᐃᓂᒪᑎᑎᐊᖏᑕᐅᑦ ᐸᖃᐅᑎᑦ ᐊᒻᒪᓗ ᑐᓂᖃᓂᓂᑕᑦ ᑕᒃᓂᓄᖑᓐᓂᖏᖏᑦ ᐅᓴᐅᓂᑦ

3.6.3 If, in accordance with the decision-making process set forth in subsection 5.2.34(c) and sections 5.3.17 through 5.3.23 of the *NLCA,* the NWMB or the Minister rejects, in whole or part, a completed Management Plan and the Plan is returned to an ACMC for reconsideration, the relevant ACMC shall re-consider the Plan and re-submit it to the NWMB.

ᐃᓄᐃᑦ ᐊᖏᑐᖅᑕᐅᔪᓕᒐᓱᐊᓂᓄᑦ ᐱᓵᓴᓪᓗᑕᐅᔨᓴᓇᖅᑐᓄᓐᓂᓗ ᐊᓱᐊᑭᑎ
Inuit Impact and Benefit Agreement

3.6.4 ᒪᓕᒃᓵᔪᒍ 8.4.13 ᓄᓇᕗᔪᓕᖅ ᓄᓇᑕᐹᑎᓄᑦ ᐊᓱᐊᐹᑎᑦ, ᐃᓚᒪᑎᕆᐅᔨᑦ ᐊᐅᓪᓕᐊᑎᐅᕆᓴᓄᑦ ᕿᖏᐅᑎᑦ ᒪᓕᒃᓂᓂᐊᖏᖅᑐᖅ ᐊᑐᑕᖏᖅᑐᖅ ᐊᐅᓪᓕᐊᑎᐅᔨᓂᖅ ᐊᐅᓪᓕᐊᑎᖏᑐᒥᑦ ᐅᑎᒪᔨᕐᓂᑦ, ᐊᑐᖏᓂᑦ ᐊᑐᑕᖏᖅᑐᔨᑦ ᐱᔨᒪᔨᑦ ᐊᔪᒥᓱᓄᑦ ᐅᕝᕙᓚᓂᑦ ᑎᒥᐅᔨᓄᑦ.

3.6.4 In accordance with section 8.4.13 of the *NLCA,* approved Management Plans shall be based on the recommendations of the relevant ACMCs, taking into account the recommendations of other interested persons or bodies.

3.6.5 ᑕᐃᒪᓗ ᒥᓂᑦ ᐊᖏᑐᖅᕿᒍ ᐊᐅᓪᓕᐊᑎᐅᕆᓴᓄᑦ ᕿᖏᐅᑎᑦ, ᒥᓂᑦ ᐱᒋᐊᖏᑎᓂᓪᓂᐊᖏᑕᖓ ᑲᔪᓯᖏᓐᓇᐊᓕᓕᑦ ᑕᑯᔪᐊ ᐊᑐᑎᖅᖁᑦ ᐊᑐᓕᖅᑕᐅᔪᐊᓕᑦ ᕿᖏᐅᑎᑦ.

3.6.5 Once the Minister has accepted a Management Plan, the Minister shall proceed forthwith to do all things necessary to implement the Plan.

ᐃᓗᓕᓐᖏᑦ ᑎᑎᕐᖅᑕᐅᓂᓕᕆᔨᑦ ᓄᓕ ᐊᐅᓚᑦᑎᐊᕐᓂᖅᓴᓂᑦ ᓴᖅᑭᐅᖅᑎᑦ ᖃᖅᑳᐅᖅᑐᕐᒪᑦ ᐊᒻᒪᓗ ᐃᑲᑦᑐᐊᕐᒪᑦ ᑎᓪᓗᐊᖅᑲᕐᔪᓂᑦ

Content of the Draft Management Plans for the Qaqsauqtuuq and Ikkattuaq Migratory Bird Sanctuaries

ᐃᓗᓕᓐᖏᑦ ᐊᐅᓪᓚᑎᑦᑎᐸᒋᑎᖅᑲᐃᑦ ᓯᖏᐅᑎᒃ
Content of the Management Plans

- ᐅᓂᖅᑲᑦᑎᐊᖅᑐᑦ ᑎᖏᒥᐊᖅᖏᓐᓇᑦ ᖃᓄᐃᓕᓐᖑᓂᖏᓪᓂᖅ
- ᐊᓭᑎᖑᓂ ᓄᓇᒥᐅᑕᐃᓕᓴᐅ ᐅᒪᔪᐃᓕᓴᐅ ᐱᑕᖅᖏᓂᖏᓪᓂᖅ ᑕᒪᓂ
- ᐱᐅᔪᑐᖅᖅᑎᒍᑦ ᐊᓭᑎᒥᐅᑕᖏᓐᖏᑦ ᔪᓇᑕᖅᖏᓂᖏᓪᓕᓴᐅ
- ᑕᐅᑐᒪᓱᐅᔪᑦ, ᑐᖃᓱᓱᐅᔪᑦ ᐊᓚᒪᓗ ᐱᓕᓱᐊᓂᔪᒪᓂᐊᖅᑐᑦ
- ᐊᐅᓪᓚᑎᓂᕐᒧᑦ ᐃᓕᒪᖏᖅᔪᐅᑭᑎᖏᑦᑲᐃᑦ - ᔪᓇᑐᐃᓵᖏᑲᐃᑦ ᐃᓕᒪᖏᖅᔪᐅᑭᑕᐅᓂᖏᓪᑎᑦ ᐊᐅᓪᓚᑎᓂᕐᒧᑦ ᑎᖏᒥᐊᖅᖏᓐᓄᓂᓂ

- Description of the Migratory Bird Sanctuaries
- Ecological resources
- Cultural resources
- Vision, goals and objectives
- Management Considerations – things to consider in managing the Migratory Bird Sanctuaries





ᐱᓪᓕᐅᓴᖅ **Black Guillemot**



ᐃᓗᓕᓐᖏᑦ ᐊᐅᓪᓚᑎᐱᑎᖅᓴᐃᑦ ᐸᖕᓇᐅᑎᓂᑦ
Content of the Management Plans

- ᐊᐅᓪᓚᑎᓂᖕᒪᑦ ᖃᓄᐃᓕᓐᖓᓂᐊᖕᓂᖏᑦ - ᖃᓄᐃᓕᓐᐅᖕᓂᐊᖕᒪᖓᑕ ᑎᑭᐅᑎᕐᒪᓂᖕᒪᑦ ᑐᖓᒪᓗᐅᔪᓄᑦ ᐊᓱᒪᓗ ᐱᓕᕆᐊᖅᐊᓂᔪᒪᓂᐊᖅᑐᑐᓄᑦ
- ᐊᖏᕈᑕᐅᓯᒪᔪᑦ ᐱᓕᕆᐊᖅᔨᓇᖕᓂᖅ ᑎᑭᓯᒪᖕᓇᐅᔪᖏᓐᓄᑦ ᐱᔨᓇᐅᑎᖃᕆᐊᖅᑐᑦ ᐊᓱᒪᓗ ᑭᓇᐅᔭᖓᑦ ᐱᔨᓇᐅᑎᖃᕆᐊᖅᖢᓂᖅᑐᑦ
- ᓄᓇᐃᑦ ᖃᓄᐃᓕᓐᖓᔪᒥᒃ ᑎᒃᓯᐊᖅᑕᐅᓯᒪᓂᖏᑦ: ᐃᔨᓇᐅᔪᖅ ᑎᓪᒥᐊᖅᖃᕐᖢᐅᑎᑕᐅᔪᖅ
- ᐊᐅᓪᓚᑎᐱᑎᖅᓴᐃᑦ ᐸᖕᓇᐅᑎᓂᑦ ᐊᑐᓪᖅᑎᑦᑕᐅᓂᖏᑦ ᐊᓱᒪᓗ ᐱᓕᕆᐊᖅᑐᑐᓂᖏᑦ

- Management Approaches – what we will do to achieve the goals and objectives
- Authorized Activities and Access – what activities require a permit, who requires a permit
- Site designation: currently a Migratory Bird Sanctuary
- Management Plan implementation and collaboration







ᖃᓄᖅ ᐱᓂᓪᓕᐊᒋᐊᓕᓚᐅᖅᓯᒪᓐᒪᓐᓂᑦ
Historical Background





ᖃᓄᐃᓕᐅᖅ 1959-ᓴᑎᓪᓗᒍ ᓴᓇᓂᐊᖅᑕᐅᓇᓯᑎᒃ ᖃᓴᐃᐃᑦ, ᓂᕆᓪᓇᐃᓪᓗ

“ᖃᓄᐃᓕᐅᖅ” - ᖃᓄᐃᓕᐅᖅᑐᓪᓗᒍᒃ ᑕᓯᕋᐊᖅ ᑕᒪᓂ

ᑎᑎᕋᖅᑕᐅᓭᓪᓗᐊᓂᖅᑎᑦ ᓯᓚ ᐊᐅᓪᓚᑦᑎᓗᒃᓴᐃᑦ ᐸᕋᐅᓕᖅᑎᑦ

- Established in 1959 to protect Snow Geese and Brant from disturbance
- “Kouksauktow” – Bay of red-throated loons
- Draft management plan in 1986



ᖃᓄᐃᓕᐅᖅ Red-throated loon

ᖃᓄᖅ ᖃᐅᔨ
Snow goose

ᐃᒃᑲᑦᑐᐊᖅᑎ ᑎᓪᓗᐊᖃᕐᓇᒃ
Ikkattuaq

- ᐱᓪᓕᖅᑎᑕᐅᓚᐅᖅᓯᒪᔪᖅ 1959-ᓴᑎᓪᓗᒍ
  ᓴᐅᓂᐊᖅᑕᐅᓇᓯᓚᑎᒃ ᒃᓴᐃᐃᑦ, ᖃᐅᔨᑦ, ᓂᕆᓪᓇᐃᓪᓗ
- ᐄᐃᐅᓕ ᑭᐱᓪᓴ ᐅᓇᐃᓪᓄᒃ
- ᑎᑎᕋᖅᑕᐅᓭᓪᓗᐊᓂᖅᑎᑦ ᓯᓚ ᐊᐅᓚᑦᑎᓗᒃᓴᐃᑦ ᐸᕋᐅᓕᖅᑎᑦ

- Established in 1959 to protect “Blue” Goose, Snow Goose and Brant from disturbance
- Harry Gibbons Unainnuk (c. 1900-1954)
- Draft management plan in 1986

ᓄᓇᖅᑐᑦ ᑕᐅᓂᓂᒃ ᖃᓄᐃᓕᓐᖅᓯᓂᖅᑐᑦ
Landscape





ᐃᑲᑦᑐᐊᖅᑎ ᑎᖅᑎᐊᖃᖅᖁᓕᑲ
Ikkattuaq

- ᐊᖏᕐᓂᓕᑲ 1,433-ᓂᑲ ᓴᓂᒧᑦ ᑐᑭᒧᑉᓗ ᑭᐸᑦᓴᓐᖅᑐᑎᑲ ᑕᒪᐃᓐᓂ ᑭᓯᒥᑕᓕᖅᓂᑲ, ᐊᓚᒪᓗ 190 ᓴᓂᒧᑦ, ᑐᑭᒧᑉᓗ ᑕᒪᐃᓐᓂᑲ ᑭᓯᒥᑕᓂᑲ ᑕᓗᐅᖏᒥ ᐃᒪᐃᑦ
- ᐳᐊᓴᐅᐸ-Boas ᑯᖏᓚ ᓄᓇᐅᐸ ᐃᓗᑐᓂᖏᓐᓄᑦᑐᖅ ᐊᓚᒪᓗ ᐊᑭᐊᓄᓴᓭᑎᑲ ᑎᓂᖏᓂᐊᓗᓄᖅᑐᑦ ᓭᓴᒥ ᑕᐅᓂᓐᓂᐸ ᐃᑲᑦᑐᐊᐸ ᑕᓗᐅᖏᓐ
- ᓄᓇᖏᓚ ᖃᖃᓴᖃᖏᓕᑦᑐᖅ (ᑕᒫᓂ <60m ᑭᑕᒥᑲ ᐊᖏᕐᓂᖏᒥ) ᐃᒪᐃᓴᖅᖁᖃᑦᑎᐊᖏᕐᑐᐊᓗᓇᓗᓂᓗ

- 1,433 km^2, includes 190 km^2 of marine waters
- Boas River delta and the adjacent tidal flats in the Bay of God’s Mercy
- Land is flat (<60m) & poorly drained

ᖃᖅᖁᓴᐅᖅᑐᖅᑎ ᑎᖅᑎᐊᖃᖅᖁᓕᐅᓂᖏᓐᓄᑦ
Qaqsauqtuuq

- ᐊᖏᕐᓂᓕᑲ 1,124-ᓂᑲ ᓴᓂᒧᑦ ᑐᑭᒧᑉᓗ ᑭᐸᑦᓴᓐᖅᑐᑎᑲ ᑕᒪᐃᓐᓂ ᑭᓯᒥᑕᓕᖅᑎᑲ, ᐃᓕᖃᖅᓯᓂᓗ 286-ᓂᑲ ᑭᐸᑦᓴᓐᓂ ᑕᒪᐃᓐᓂᑲ ᑕᓗᐅᖏᒥ ᐃᒪᖁᓂᑲ
- ᖃᖅᖁᓴᐅᖅᑐᖅᑎ, ᑕᑭᓂᖃᖅᑎᑎᕐᒥᑲ 50-km ᑭᓯᒥᑕᓕᖅᑎᑲ ᑲᖏᕐᓄᖃᖅᖁᖃᖅᓄᖅ
- ᓄᓇᖏᓚ ᖃᖃᓴᖃᖏᓕᑦᑐᖅ (ᑕᒫᓂ <120m ᑭᑕᒥᑲ ᐊᖏᕐᓂᓕᑲ) ᐃᒪᐃᓴᖅᖁᖃᑦᑎᐊᖏᕐᓄᐊᓗᓇᓗᓂᓗ

- 1,124 km^2, includes 286 km^2 of marine waters
- East Bay, a 50-km-long inlet
- Land is flat (<120m) & poorly drained




ᓄᓇᖅᑐᑦ ᑕᐅᔭᓂᒃᑯᓂᖅ
Landscape






ᐃᖃᓪᓗᐊᖅᑎ ᑎᖅᑎᐊᖅᑐᖃᖅ
Ikkattuaq

- ᑕᒪᕆᒃᓴᒃ ᑕᒪᓐᓇ ᓄᓇ ᑲᓇᑕᐅᔨᑦ ᓴᓇᒪᓐᓂᒃᑯᐊᓂᖅ ᓄᓇᕐᔪᑎᑎᕆᔨᓐᓂ
- ᐊᑕᐅᓯᖅ ᑭᓪᓚᐸᓐᓂᕐᒥᖅ ᐃᓄᖅᓄᑦ ᓄᓇᕐᔪᑎᑎᕆᐅᓯᖅ
    - ᓄᓇᐅᔨᑦ ᖃᖅᓴᓂᑕᓂᕐᓄᑦ ᐱᓯᓇᐅᑎᖃᖅᑐᐊᓯᖅ
- Mostly federal crown land
- 1 parcel of Inuit Owned Land
    - Surface rights





ᓄᓇᖅᑐᑦ ᑕᐅᔭᓂ ᖃᓄᐃᓕᖅᑎᑦ

Landscape






ᖃᖅᓴᐅᔪᒃ ᑎᖅᑕᐊᖃᖅᓯᐅᓂᖅᓯᓂ

Qaqsauqtuuq

- ᑕᒪᖁᑲᓴᐃᑦ ᓄᓇᖅᑦ ᑕᒫᓂ ᐃᓄᖅᓄᑦ ᓄᓇᕐᒪᑎᖅᑲᐅᔪᑦ
  - ᓄᓇᐅᔨ ᖃᖅᓱᓂᑐᑐᓂᒃ ᐱᓵᓇᐅᑎᖃᕐᓯᐅᔪᖅ
- ᑕᕆᐅᖁᒥ ᐃᒪᐃᑦ ᐊᓚᓗ ᖁᑭᖅᑕᑯᓗᒃ - ᑲᓇᑕᐅᔨ ᒪᓪᒪᖅᓵᓄᑦ ᓄᓇᒥ
- Mostly Inuit Owned Land
  - Surface rights
- Marine waters and Qikiqtakuluk - federal



ᐊᐢᑎᖏᓂ ᓄᓇᒥᐅᑕᐃᓪᓗ ᐅᒪᔨᐃᓪᓗ ᐱᑕᖅᖃᑐᑦ ᑕᒪᓂ
Ecological Resources

ᐃᒃᑲᑦᑐᐊᖅᒥ ᑎᖏᒥᐊᖅᓴᒃ
Ikkattuaq

- ᓄᓕᐊᖏᐅᓴᖏᓂᖓ ᑕᒪᓂ ᑲᓇᑕᓕᒫᒥ ᐱᓪᒪᓚᐅᓂᖅᑐᖅ ᐱᑕᖅᐃᓕᓇᕐᓇᖏᓐᓂᓕᓂᖅᐳᑦ (5% ᐳᕆᓕᑎᒋᑐᑦ) ᓂᖏᓕᓇᐃᓪᓗ (1% ᐳᕆᓕᑎᒋᑐᑦ)
- ᐱᓪᒪᓚᐊᓗᔭᖅ ᑎᑭᓯᒪᑲᐃᓇᖏᖅᑕᑦᑕᖏᓂᓕᖏᑦ ᐊᓯᖏᓕᓄᑦᓂᐊᓂᖏᓕ ᓯᖁᖅᐸᓯᐅᓂᖏᑦᑎᓯᔪᓯᑦ ᑎᖏᒥᐊᑦ
- ᐅᖓᑕᓂᒋᒥᔪᑦ >50 ᐊᐠᔨᑦᖏᑦᑐᑎᑦ ᑎᖏᒥᐊᑖᕆᐊჲᑦ ᑕᒪᓂᓯᖓᖏᒥᔪᑦ
- ᖃᓄᐅᔪᑦ 13 ᐅᒪᔨᐃᑦ ᐊᒥᓯᖏᓗᔭᐊᓕᖏᓂᖏᑦ

- Breeding habitat for nationally significant populations of Snow Geese (5%) and Brant (1%)
- Important stopover site in fall for shorebirds
- >50 species of birds
- 13 species at risk



ᓂᖏᓕᓇᐃᑦ **Brant**



ᖃᖏᒍ **Snow Goose**

ᐊᐢᑎᖏᓂ ᓄᓇᒥᐅᑕᐃᓪᓗ ᐅᒪᔨᐃᓪᓗ ᐱᑕᖅᖃᖅᑐᑦ ᑕᒪᓂ
Ecological Resources

ᐃᒃᑲᑐᐊᖅᒥ ᑎᖏᒥᐊᖅᖃᕐᓂᖅ
Ikkattuaq

- 19-ᖏᔪᑦ ᐊᓴᒋᖏᑐᑐᑎᑦ ᐊᒪᒪᒃᑎᑦᑎᐸᒃᑐᑦ ᐱᐊᕆᖏᑦᓂᒃ ᓄᓇᒥᐅᑕᐃᑦ ᐅᒪᔨᐃᑦ ᑕᒪᓐᓇᑦ
- ᓇᓄᖅᓯᖅ ᐊᐅᓴᒃᑯᑦ ᐊᓚᒪᓗ ᑕᒪᓂ ᐃᒡᓗᖅᓴᐅᓯᖏᓪᓗᑎᒃ ᓇᓄᕐᓄᑦ
- ᓇᓪᓴᖅᑐᖅᖏᑦᑐᕈᐅᑕᓄᑦ ᑐᒃᑐᖅᕆᖅᓴᐅᐸᒃᑐᖅ, ᐃᓕᖏᑦᓗ ᓄᕐᕆᕆᐅᖅᑕᑦᑕᖅᓯᒪᔪᓗᑎᒃ
- ᓇᑦᓯᖅᑕᖅᖅᑐᖅ ᐊᓚᒪᓗ ᐊᕐᕆᒃᑕᖅᖅᑐᖅ

- 19 species of mammals
- Polar bear summer and denning habitat
- Barren-ground caribou summer range, portions used as calving grounds in the past
- Seals and Bowhead whale



ᓇᓄᖅ **Polar Bear**



ᑐᕐᑐ **Caribou**

ᐊᐢᑎᖏᓂ ᓄᓇᒥᐅᑕᐃᓪᓗ ᐅᒪᔨᐃᓪᓗ ᐱᑕᖅᑐᖅᑐᑦ ᑕᒪᓂ
Ecological Resources

ᖃᖅᓴᐅᖅᑐᑎᒥ ᑎᖏᒥᐊᖅᖁᕕᐅᓂᖏᓂ
Qaqsauqtuuq

- ᓄᓕᐊᖁᕕᐅᔪᖅᑐᖅ ᐱᓕᓚᐅᕕᔾᒥᒃ ᑲᓇᑕᒥ ᐊᒥᓱᓄᓕ ᑲᖏᖅᓄᓕ (2% ᐳᓴᓂᑎᖏᓂᑕᓄᓕ), ᒥᑎᖁᓄᓕ (5% ᐳᓴᓂᑎᓄᓕ) ᓂᖁᓕᖁᖁᓄᓪᓗ (1% ᐳᓴᓂᑎᑦᑕᓐᓄᓕ)
- ᑲᓇᑕᐅᐸ ᐅᑭᐅᖅᑕᖅᑐᑐᓂᓗᒃᑮᖏᓂ ᐊᖏᕐᓭᖏᖅᖁᖅ ᒥᑎᖅᑕᖃᖁᕕᐅᔪᓐᓂᖅ
- ᐅᖏᓪᓕᓂᒃᓕᕆᔨᓕᑦ >90 ᐊᐢᔨᑲᖏᑦᑐᑎᑲᑦ ᑎᖏᐊᓚᐱᐅᑦ ᑕᒪᓂᓕᒃᓴᖏᖅᓕᕆᔨᓕᑦ
- Breeding habitat for nationally significant populations of Snow Geese (2%), Common Eider (5%) and Brant (1%)
- Arctic Canada’s largest single colony of Common Eiders
- >90 species of birds



ᒥᑎᐹᖅ Common Eider



ᓴᐅᖁᖅ Red Phalarope

ᐊᐢᑎᖏᓂ ᓄᓇᒥᐅᑕᐃᓪᓗ ᐅᒪᔨᐃᓪᓗ ᐱᑕᖅᖃᖅᑐᑦ ᑕᒪᓂ
Ecological Resources

ᖃᖅᖁᐅᖅᑐᖅᒥ ᑎᖏᒥᐊᖅᖁᐅᓂᖏᓂ
Qaqsauqtuuq

- ᖁᓕᐅᔨᑦ 22 ᐅᒪᔨᐃᑦ ᐊᒥᓱᖏᓪᓗᐊᓕᖁᓂᖏᑦ
- ᓄᓇᖁᓇᑦ ᐃᒡᓗᖅᖁᐅᓴᐅᑐᑦ (ᔪᑎᖅᖁᐅᓴᐅᑐᑦ) ᐊᐅᔪᓄᑦᓗ ᓂᖏᔪᖅᔪᐅᖁᐅᓴᐅᑐᖅ ᓄᓇᖁᓇᑦ
- ᓄᓪᓚᐃᐅᓴᐅᑐᑎᒥᐅᑕᓄᑦ ᑐᑲᑐᖅᖁᐅᓴᐅᑐᖅ, ᓄᖁᓗᐃᐅᓴᓄᓗ, ᓄᔨᖅᑕᐅᓴᐅᑐᖁᓗ ᐊᐅᔪᐅᓂᖏᓂ
- ᖁᓇᓗᒃᑖᓄᑦ ᐅᖃᓕᖅᑲᖅᔪᒪᔪᑦ (ᐃᕆᓪᓂᖁᓂ), ᐊᒥᔪᖅᑕᖅᔪᑎᒃᓗ, ᓂᖏᔪᖅᔪᐅᖁᐅᓴᐅᔪᓂᓗ ᑕᒪᑲᔪᓇᖏᒃ
- ᐊᐃᐃᖅᑕᓕᑲ ᐊᐅᔪᑲᔪᑦ ᑕᒪᐢᓇ

- 22 species at risk
- Polar bear denning and summer foraging habitat
- Barren-ground caribou calving and summer range
- Beluga calving, molting and foraging habitat
- Walrus summer foraging habitat



ᐊᐃᐃᖅ Walrus



ᖃᓪᓚᖅ Beluga

ᐊᐢᕈᑎᖏᓂ ᓄᓇᒥᐅᑕᐃᓪᓗ ᐅᒪᔨᐃᓪᓗ ᐱᑕᖃᖅᑐᑦ ᑕᒫᓂ
Cultural Resources



ᐃᓄᐃᑦ ᓴᓪᓕᐅᕐᓂ ᖃᐅᖅᑕᒪᓗᐊᖏᖅᑕᑦᑕᕐᓂᑯᓕᖅᓴᓂ ᑕᐅᓴᓴᒪᓕᕆᐅᓕᖅᑐᓂ ᐊᕐᕿᒍᓂ


- ᐊᒥᓱᒪᒪᓕᕆᐅᕐᓂᒃ ᖃᐅᔨᒪᓴᐅᔨᕐᑎᑦ ᑕᐃᔾᓱᒪᓂᓂᑕᖃᕐᓂᐊᕐᓂᖅ, ᐃᓄᖃᕐᓂᐊᕐᓚᐅᕐᓂᖏᓐᓂᕐᓂᖅ, ᐃᓕᓴᕐᓯᑐᖃᕐᓂᖅᓴ ᓇᓗᓇᐃᑯᑕᐃᑦ ᐱᑕᖃᖅᑐᑦ ᐊᑐᓗ ᐱᓴᓯᑎᐃᓂᖃᕐᓂᐊᕐᓂᖅᓱᓂ ᑕᐅᕐᓂᓐᓇ ᓄᓇᖓ

- ᑕᐃᔾᓱᒪᓂᓂᑕᐃᑦ ᖃᖓᓂᓂᑕᐃᑦ ᒥᑭᓂᑕᐃᑦ ᑎᑭᑎᐊᖃᕐᓂᐊᕐᓚᐅᕐᖅᑐᑦ ᑕᐅᕐᓂᓄᓐ



Inuit have lived on Southampton Island for thousands of years

- Numerous known archaeological sites, cultural features & artifacts in the area
- Historic trap lines



ᐊᐢᓂᑎᓂᓄ ᓄᓇᒥᐅᑕᐃᓪᓗ ᐅᒪᔨᐃᓪᓗ ᐱᑕᖅᑲᖅᑐᑦ ᑕᒪᓄ Cultural Resources



- ᐊᒥᔨᐊᓗᐃᑦ ᔨᓕ ᐃᓄᖅᕿᐅᓂᒪᐅᖅᑐᖅᑐᒃᓗᐃᑦ ᔨᓕ ᑎᑎᕿᑕᐅᔨᒪᖏᑦᑐᑦ
- ᐊᑕᐅᔨᕿᒃ 1 ᑎᑎᕿᑕᐅᔨᒪᓕᖅᑐᖅ ᒪᓕᒐᖅᑎᒍᑦ ᓄᓇᖏ ᑕᐅᓂᓄ ᖃᐹᖅᑕᑯᓗᒃ
- ᒪᕆᐣ 2 ᑎᑎᕿᑕᐅᔨᒪᓕᖅᑐᖅ ᒪᓕᒐᖅᑎᒍᑦ ᓄᓇᕗᑦ ᖁᓂᒃᑯᖏᓐᓂᑦ ᐃᓕᔨᐊᕆᔨᒃᑯᑦ


- Many archaeological sites remain unregistered
- 1 registered site on Qikiqtakuluk
- 2 registered sites near Itijuarjuk





ᑕᐃᔨᓯᒪᓂᑐᓄᖅᑎᒃ ᐃᓐᓇᒧᑦᓗ ᐃᓄᖅᓄᑦ ᓄᓇᖅᑎᑦ ᐊᑐᖅᑕᐅᖅᑕᕐᑐᖅᑐᑦ ᐃᓅᔨᐸᑎᖅᓴᓄᐅᓕᖅᓂᕐᒧᑦ

Past and current subsistence land uses



- Goose hunting and egg collection (early spring)
- Seal and walrus hunting
- Polar bear hunting
- Char fishing from lakes and rivers
- Fox trapping

- Camping
- Char fishing continued (using nets)
- Summer fishing camps were set-up at mouth of the rivers
- Seal, caribou and geese were traditionally hunted from canoes along the shoreline
- Sea mammal hunting from boats
- Collecting berries

- Fox trapping
- Caribou until the 1950s
- Fishing nets set under ice to catch char
- Hunting at the flow edge (seal, walrus)

- Securing meat for winter by hunting walrus, caribou, polar bear, and beluga whales
- Fishing with nets under new ice
- Meat caching
- Fox trapping

ᑕᐃᓴᓯᓂᑐᖅᓯᒪᖅ ᐃᓐᓇᒧᑦᓗ ᐃᓄᖅᓄᑦ ᓄᓇᖅᑎᑦ ᐊᑐᖅᑕᐅᖅᑕᕐᑐᖅᑐᑦ ᐃᓅᔨᐸᑎᖅᓴᓂᐅᕙᖅᓂᕆᒧᑦ

Past and current subsistence land uses

ᐅᓪᓗᒥᐅᓕᖅᑐᖅ, ᐊᑐᖅᑕᐅᕙᒃᑐᖅ ᓯᓚ ᐊᐅᓪᓚᐸᖅᓯᒪᓂᖅᓴᓂᐅᕙᖅᓂᕆᒧᑦ ᐊᖅᑐᓯᒪᔪᓴᖅᐱᐅᕙᖅᓂᕆᒧᑦᓗ ᐃᓚᓂᑯᓂᖅ ᖃᐅᑎᑯᓂᖅ

- ᐊᖅᓯᓇᓯᖅᑏᑦ ᖃᑐᐸᖅᑎᑎᒋᖅᓂᑦ ᐃᒡᓗᓕᕆᓚᖅᑐᑦ ᑕᐅᕙᓂ ᐃᖅᑦᑐᐊᒥᒃ ᐊᑐᖅᑕᐅᖅᑕᕐᑐᖅᑐᑦ ᐊᖅᓯᓇᓯᖅᑐᓄᓕᓂ, ᐱᓗᐊᖅᑐᒥᖅ ᓇᓂᐊᖅᓯᒪᔪᓄᓕᓂ



Today, used for camping & harvesting activities periodically


- HTO cabin located in Ikkattuaq used to support harvesting activities, in particular polar bear hunting




ᐊᐅᓪᓚᓂᖅ ᐃᓱᒪᖏᖅᓯᐅᑭᑎᖏᓚᐃᓪ
Management Considerations

ᐊᐅᓪᓚᓂᖅ ᐃᓱᒪᖏᖅᓯᐅᑭᑎᖏᓚᐃᓪ
ᐱᔨᔾᑎᖅᖅᓴᓪ ᑕᐃᒃᑯᓂᖓ
ᖃᐅᔨᒪᔨᐅᔨᓗᐊᖅᖅᑐᓂᒃ ᐃᓗᐊᓂ
ᐊᓚᓗ ᐊᕐᓗᓐᓯᓂᑐᓂᒃ,
ᓴᓴᓂᐊᖅᑕᐅᓱᒪᓂᓯᓂᓂ ᓄᓇᐃᓪ
ᐋᖏᐅᒪᓪᑎᐊᖅᑐᒥᒃ
ᐊᐅᓪᑕᐅᔨᓂᓕᖅᓯᐊᖅᒪᑕ

Management considerations are things we need to be aware of within, and surrounding, the protected area in order to effectively manage it





ᐊᐅᓪᓗᓂᔭᕐᓗᑎ ᐃᓕᓴᖅᓯᐅᑕᑎᓴᐃᑦ Management Considerations

ᐱᑕᖅᖃᑐᖅᓯᓪᓗᒍ ᐱᒻᒪᓚᐅᔨᓂᖅ ᐄᓐᓇᕆᐅᑎᑭᓕᓐᓂᒃ ᐅᓗᓚᐊᓇᑕᐅᔨᓂᖅ, ᑭᓯᐊᓂᓕᑕᐅᖅ:

1. ᑎᓪᓗᐊᖅᖃᕐᐅᓂᓪᓗᑎᑦ ᐃᓕᑕᑦᓴᐅᓂᓪᓚᑦ ᑎᒃᓯᐊᖅᑕᐅᓯᒪᓂᖅ
    - ᓴᐳᓐᓂᐊᖅᑕᐅᓯᒪᑦᑎᐊᒃᑲᓂᓚᐊᖅᖃᖅᑐᖅ
    - ᑎᓪᓗᐊᕐᓂᓐᓇᐅᔨᓐᓂᖅᑐᖅ
2. ᑲᔪᐃᑦ ᐊᒥᓯᓂᑦ ᐊᒻᒪᓗ ᓄᓇᒋᖅᑕᑦᓂᓂᓪᓚᑦ
    - ᐊᒥᓯᓗᐊᓕᖅᑐᑦ
    - ᐅᒪᔨᕐᓂᒃ ᐊᓯᖏᓐᓂᒃ ᐊᒃᑐᖅᓯᓀᓕᓕᐊᖏᒪᑕ ᐱᔭᕇᑕᐅᓗᑎᒃ ᓇᒻᒪᖏᑦᑐᒥᒃ

No significant or immediate threats, but:

1. MBS designation
    - need better protection
    - not just birds
2. Light geese population and habitat
    - too many
    - impact on other species



ᑐᑦᑐ Caribou





ᖃᓕᓗᒃ Beluga





ᑲᖏᒍ ᖃᐅᔨᖅ Snow goose

ᐊᐅᓪᓕᓂᐅᓐᓴᑦ ᐃᓱᒪᖏᖅᓯᐅᑕᑎᖏᓚᐃᓪᑦ
Management Considerations

3. ᓯᓚᐅᐟ ᐊᓯᔨᖅᐸᑦᓪᓕᐊᓂᖅ
   - ᐊᖏᑐᖅᑕᐅᓴᑦᓪᓕᐊᓴᑦ ᐅᐠᔨᕐᓇᖅᓯᖅᓯᒪᓕᑕ
   - ᐊᐅᓪᓕᓂᐅᓂᖅᖃᕐᑎᑦᑕᓄᑦ
4. ᓄᓇᓗᑦ ᐃᓄᐃᑦ ᐃᓪᓕᐅᕆᓪᓚᐊᖅᖅᑐᑦ ᐊᓖᒪᓗ ᖃᐅᔨᒪᕆᓪᓚᐊᖅᖅᓯᑎᖏ
   - ᖃᐅᔨᔨᖅᑕᐅᔪᓂᖅ
   - ᐃᓪᓕᑎᕐᐅᕆᓪᓚᐊᖅᖃᕐᓂᕐᓂᖏ ᐊᑐᖅᑕᐅᓗᑎᖏ ᐃᓄᐃᑦ ᖃᐅᔨᒪᔨᑐᖅᖃᕐᑐᓂ ᓄᓇᓕᕐᓂ ᐃᑲᔪᕐᓂᖅᓂᕐᓂᐊᖏᓪᓕᑕ

3. Climate Change
   - effects are noticed
   - out of our control
4. Community participation and awareness
   - research
   - inclusion of Inuit Qaujimajatuqangit
   - local benefits







ᐊᐅᓪᓗᓂᔭᕐᒪᑦ ᐃᓕᒪᖏᖅᓯᐅᑕᑎᖏᖅᓱᐃᓪᓗᑦ
Management Considerations

5. ᑭᓇᐯᓐᓄᑐᐃᓐᓇᖅ ᐱᓕᕆᐊᖓᔪᓐᓕᐊᔪᑦ
   - ᐅᒥᐊᕐᔪᐊᖏᖅᖢᓕᓕᐊᓕᕐᓂᖅ
   - ᐊᒥᓯᑭᖅᖢᓕᓕᐊᓂᖓᑦ ᑎᑭᑦᖢᓕᓕᖅᑐᑦ
   - ᐊᑭᓴᑎᑦᑎᔪᓂᖅ ᐊᖓᔪᓇᓯᖓᐊᑭᐅᔨᔪᓐᓂᕐᒪᑦ
6. ᖃᐅᔨᒪᔭᐅᔭᕆᐊᓕᑦ ᐱᑕᖃᖓᓐᓂᖓᑦ
   - ᐱᐅᓯᑐᖃᕐᒥᔪᑦ ᐊᔪᑎᖓᓄᓂ ᓄᓇᒥᐅᑕᐃᓕᓐᓗ ᐅᒪᔨᐃᓕᓐᓗ ᐱᑕᖃᕐᓂᖓᑦ, ᐱᓗᐊᖅᑐᒥᐅ ᐃᓄᖃᕐᓇᐅᔪᓕᒪᐅᕐᓂᖓᓂᓂᑲ ᖃᐅᒪᔨᑐᖃᐃᓐ
   - ᑕᕆᐅᒥᐅᑕᓂᑲ ᐅᒪᔨᖓᑦ, ᐃᖃᓗᓕᕆᐃᓕᓐᓗ ᐊᕐᔪᑎᖓᑦᑐᑦ

5. Human Activities
   - shipping
   - increased access
   - sport hunting
6. Information missing
   - Cultural resources, especially archaeology
   - marine habitat, fish







ᐊᐅᓪᓗᓂᔭᕐᓗᑦ ᐃᓱᒪᖏᖅᓯᐅᑕᑎᖏᓚᐃᓪᓗᑦ (ᖃᖅᖏᖏᖅᑐᕐᕆᓂᖅ) Management Considerations (*Qaqsauqtuuq only*)



ᐅᓪᓚᐊᓇᑕᐅᓂᖅᐳᑦ ᒥᑎᕕᖅᖏᓚᐊᓄᑦ ᖃᐅᖅᑕᓯᔪᓐᓇᖅ

- ᓄᓂᔪᖏᕐᓂᕐᒥᒃ ᑎᖅᒥᐊᕐᓄᑦ ᐊᓂᓂᐊᖅᑕᓚᖏᐅᓂᒃᑐᑦ
- ᓇᓄᐃᑦ


Threats to eider colony at Qikiqtakuluk

- Avian cholera
- Polar bear




ᓇᓄᖅ **Polar Bear**

ᑕᐅᑐᒪᓗᓴᐅᔪᑦ, ᑐᕐᒪᓗᓴᐅᔪᑦ, ᐃᓂᖏᑕᐅᔪᒪᔪᓪᓚ
Vision, Goals, and Objectives



ᑕᐅᑐᓗᐅᔪᖅ ᑎᓐᒥᐊᖅᑲᐃᓐᓇᓂ ᓴᓂᓂᐊᖅᑕᐅᔪᓚᐃᓐᓇᓂ
Vision for the MBSs



“ᑎᓐᒥᐊᖅᑲᓕᑦ ᓴᓂᓂᐊᖅᑕᐅᔪᓚᐃᓐᓇᑎᑦ ᐊᐅᓪᑕᐅᓂᐊᖅᑐᑦ ᐅᓱᐹᖅᑐᑦᑎᐊᕈᓚᓂᕐᒧᑦ ᐊᑯᓂᐅᓂᐊᖅᑐᖅ ᓴᓂᓂᐊᖅᑕᐅᓂᖏᓐᓄᓂ ᓄᓇᖏᑦ ᐊᓴᒪᓗ ᐃᒪᐃᑦ ᑕᒪᐃᓐᓄᓂ ᐱᑕᖅᑲᓗᑎᒃ ᐅᒪᔪᕐᓄᓂ, ᐃᓄᐃᑦ ᐱᔪᓐᓇᐅᑎᖅᑲᓕᖏᓐᓄᓂ, ᐱᑕᖅᐃᓐᓇᕐᓂᖏᓐᓄᑐᓗ ᑕᐃᒪᓐᓂ ᐃᓅᓂᕆᓂᖅ, ᐃᓕᒪᓴᕐᔪᐊᑦᑎᓐᓇᕐᓂᑦ ᐊᑐᖅᑕᓂᕐᓗᑎᒃ ᐃᓄᐃᑦ ᖃᐅᔨᒪᔪᑐᖅᑕᖅᓂᑦ ᑕᒪᐃᓐᓂᕐᒪᑦᑎᐊᖅ ᐃᓕᓴᐅᔪᑦᑎᐊᕐᓂᖅᑕᖅᓂᒃ.”


“The MBSs will be managed to ensure the long-term conservation and protection of the land and water for all wildlife, Inuit rights and their heritage while considering Inuit Qaujimajatuqangit in all decisions.”


© Jason Caswell

ᑎᓴᓐᓂᐊᖅ Arctic Fox




ᐊᐅᓪᓂᐹᓂᒃ ᑐᖏᓐᓂᒃ
Management Goals



*“ᑎᓕᒋᐊᖅᑲᓇᑦ ᓴᓇᓂᐊᖅᑕᐅᓂᒡᓕᐊᖅᓯᒪᔪᓂᑦ ᐊᐅᓚᑕᐅᓂᐊᖅᑐᑦ ᐅᖃᖅᑐᑎᐊᕈᒪᓂᖅᓯᑦ ᐊdᓂᐅᓂᐊᖅᑐᖅ ᓴᓇᓂᐊᖅᑕᐅᓂᖅᓂᓇᓂ ᓄᓇᖅᑦ ᐊᓚᒐ ᐃᒪᐃᑦ ᑕᒪᐃᓇᓂᑦ ᐱᑕᖅᑐᑎᓂᑦ ᐅᓪᓗᓂᑦ, ᐃᓄᐃᑦ ᐱᔨᓇᐅᑎᖅᑲᖅᖢᓂᓇᓂᑦ, ᐱᑕᖅᐃᓇᖅᓂᖅᖢᓂᓇᓂᑦ ᑕᐃᒪᖅᑎᑦ ᐃᓕᓴᕐᕕᒃ ᐃᓄᐃᑦ ᖃᐅᔨᓚᕐᖅᑕᐅᓂᖅᑐᑦ ᑕᒪᐃᓇᓂᑦ ᐊᑐᖅᑕᕐᕕᐊᖅ ᐃᓕᓴᕐᓂᐊᖅᑐᓂᑦ ᑕᒪᐃᓇᓂᑦ ᐃᓄᐃᑦ ᖃᐅᔨᓚᐅᖅᑐᓂᑦ ᑕᒪᐃᓇᓂᑦ ᐊᑐᖅᑕᕐᕕᐊᖅ ᐃᓕᓴᕐᓂᐊᖅᑐᓂᑦ ᑕᒪᐃᓇᓂᑦ ᐱᑕᖅᑐᑎᐊᕈᒪᓂᖅᓯᑦ ᐊᐅᓚᑕᐅᓂᐊᖅᑐᑦ ᐃᓄᐃᑦ ᖃᐅᔨᓚᐅᖅᑐᓂᑦ ᑕᒪᐃᓇᓂ


ᐊᐅᓪᓗᓂᔪᓐᓇᖅ ᐃᓂᖅᑕᐅᔪᒪᓂᖅᑎᑦ
Management Objectives



ᓴᓕᓂᖅᑕᐅᓂᖅᑎᑦ ᐃᓕᓴᕐᓂᐊᓂᖅ ᑲᓇᙵᓂᖅ ᑲᑎᒪᔨᓂᖅ ᐃᒪᐃᑦ ᐅᑭᐅᖅᑕᐅᔪᒪᓂᖅᑎᑦ ᑕᐃᔭᑯᓐᓚ ᑲᑎᒪᔨᓂᖅᑎᑦ ᐃᒪᐃᑦ ᐅᑭᐅᖅᑕᐅᔪᒪᓂᖅᑎᑦ ᑕᐃᔭᑯᓐᓚ ᑲᑎᒪᔨᓂᖅᑎᑦ ᐃᒪᐃᑦ ᐅᑭᐅᖅᑕᐅᔪᒪᓂᖅᑎᑦ ᑕᐃᔭᑯᓐᓚ ᑲᑎᒪᔨᓂᖅᑎᑦ ᐃᒪᐃᑦ ᐅᑭᐅᖅᑕᐅᔪᒪᓂᖅᑎᑦ ᑕᐃᔭᑯᓐᓚ

ᓴᓕᓂᖅᑕᐅᓂᖅᑎᑦ ᓄᓇᓕᖕᒥ ᐃᓕᓴᕐᓂᐊᓂᖅ ᑲᑎᒪᔨᓂᖅᑎᑦ ᐃᒪᐃᑦ ᐅᑭᐅᖅᑕᐅᔪᒪᓂᖅᑎᑦ ᑕᐃᔭᑯᓐᓚ

ᐊᐅᓪᓕᓂᐊᓪᓗᑦ ᐃᓂᖅᑕᐅᔪᓂᖏᑦ
Management Objectives



ᐅᓴᖅᑐᑦᑎᐊᓂᖅ ᓄᓇᓕᑦ ᖃᐅᔨᒪᔨᐅᑎᑦᑎᐊᖅᑕᕐᓂᖅᓗᑦ ᐸᕐᓇᐅᑎᓴᓂᖅ ᐊᑐᑎᖅᑕᕐᓂᐊᖅᑐᓂᖅ ᖃᐅᔨᒪᑎᑦᑎᐊᖅᑕᕐᓂᖅᓗᑦ ᐊᒻᒪᓗ ᖃᐅᔨᒪᔨᐅᑎᑦᑎᐊᖅᑕᕐᓂᖅᑎᑦ ᖃᐅᔨᓴᐅᓯᓕᐊᔪᓂᖅ, ᐅᖃᓕᒫᖅᑐᓂᖅᓴᓇᓱᓗ ᐃᔪᒪᓕᑕᐅᓂᖓᓂ ᐃᓄᖅᓄᑦ ᐊᒻᒪᓗ ᐃᓄᐃᑦ ᐃᔪᒪᓕᕆᖅᓂᖓᓂ, ᖃᐅᔨᓴᖅᑕᐅᓂᐊᖅᑐᓂᖅ ᔪᓚᑦᓂᐅᑎᑕᐅᓂᐊᖅᑐᓂᖅ ᐱᓕᕆᓂᖅᓂᓂ

ᐱᔨᒥᓴᐃᖅᑕᕐᓴᓗᑎᖅ ᓴᓪᓂᕐᒥᐅᑕᓂᖅ ᐃᖅᓚᐃᓴᖅᑕᐅᓯᕐᓂᖅᓂᓂᖅ ᐊᒻᒪᓗ ᐃᔪᒥᖅᓴᐃᓴᐅᖅᑕᐅᓯᕐᓂᖅᓂᓂᖅ ᐱᓴᐊᖅᑕᖅᑐᓂᖅ ᖃᐅᔨᓴᖅᑕᐅᓂᐊᖅᑐᓂᖅ ᐊᒻᒪᓗ ᓇᐅᑦᑎᖅᑕᐅᖅᑕᕐᓂᐊᖅᑐᓂᖅ

ᖃᐅᔨᒪᔨᐅᓂᖅᓴᐅᑦᓴᓗᑎᖅ ᓄᓇᕐᓂ ᐱᓕᕆᐃᐅᔨᓇᖅᑐᓯᕐᓂᖅᓂᓂᖅ ᑎᓐᒥᐊᖅᑲᖅᓴᐃᐅᔨᑦ ᓴᓐᓂᐊᖅᑕᐅᔪᒪᖅᓯᕐᓂᖅᓂᓂ


Ensure local awareness, participation and benefits

Develop a communication plan to facilitate information and knowledge sharing, discuss concerns and input into research priorities

Encourage Salliqmiut employment and mentorship in research and monitoring

Increase local awareness of permitted activities within the MBSs

ᐊᐅᓪᓕᓂᐊᓕᒃ ᐃᓂᖅᑕᐅᔪᒪᓂᖅᑎᑦ
Management Objectives



ᐊᖏᑐᕈᓂᕐᔪᖏᑎᑦᑎᑦ ᐃᓕᒪᐃᓐᓇᕐᓂᖅ ᑭᒪᔭᖏᑐᖅ ᐅᖃᐅᔾᔨᒌᖕᖏᑦ

ᐅᔾᔨᖅᑐᑦᑎᐊᕐᓇᕐᔪᓂᖅᑎᑦ ᖃᐅᔨᓴᖅᑎᑦᑎᓂᖅ ᐊᑐᖅᑕᐅᔨᒪᔨᖅᑎᑦᑎᓂᖅ, ᐅᑭᐅᖅᑕᐅᔪᒪᓂᖅᑎᑦᑎᓂᖅ, ᒪᑐᐊᖅᑎᑦᑎᓂᖅ

ᐊᖏᑐᕈᓂᕐᔪᖏᑎᑦᑎᓂᖅ ᐃᓕᒪᐃᓐᓇᕐᓂᖅ ᑭᒪᔭᖏᑐᖅ ᐅᖃᐅᔾᔨᒌᖕᖏᑦ ᐅᖃᐅᓯᒪᔭᕐᓂᖅ ᐃᓕᓴᕐᓂᖅ ᐊᑐᖅᑕᐅᔨᒪᔨᖅᑎᑦᑎᓂᖅ

ᐊᐅᓪᓕᓂᖅᓯᓐᓂ ᖃᓄᐃᓕᓴᕐᓂᓱᐊᕐᓯᓐᓂ
Management Approaches

ᐊᐅᓪᓕᓂᖅᓯᓐᓂᒃ ᖃᓄᐃᓕᓴᕐᓂᓱᐊᖅᑐᑦ ᐅᖃᐅᓯᖃᓕᓯᐅᕙᒃ
ᖃᓄᐃᓕᓴᐅᑭᑕᐅᓂᐊᓕᓴᖅᑐᓂᒃ ᐊᒃᓯᑭᑕᐅᔨᓐᓇᖅᑐᓂᒃ ᐊᐅᓪᓕᓂᕆᓯᓐᓂᕐᒧᑦ,
ᐊᖏᓚᕆᐅᓗᑎᒍᓗ ᐊᐅᓪᓕᓂᖅᓂᕐᒧᑦ ᐱᓕᕆᐊᖏᐊᓂᔪᓚᓯᔨᑦ

- ᐱᐅᓯᑐᖅᓴᒍᑦ ᐊᐸᑎᓐᓴᓂ ᓄᓇᒥᐅᑕᓐᓂᑦ ᐊᓚᒪᓗ ᐅᓚᔨᐃᑦ ᐱᑕᖅᑎᓂᓐᓂᑦ
- ᐅᓚᔨᐃᓪᓗ ᐅᓚᔨᑎᓂᓗ ᓇᔨᓴᕆᐅᕙᐸᓐᓂᓂᓐᓂᒃ ᐊᐅᓪᓕᓂᖅᓂᖅ
- ᓇᐅᑦᑎᖅᑐᐊᖅᑕᐅᖅᑕᑕᐃᓇᖅᓂᑦ ᐊᓚᒪᓗ ᖃᐅᔨᓴᖅᑕᐅᖅᑕᓴᖅᓂᑦ
- ᐃᓄᓕᒋᓂᒃ ᖃᐅᔨᓴᒪᑦᑎᐸᓪᓕᐊᓂᖅ ᐊᓚᒪᓗ ᖃᐅᔨᓴᓕᐅᔨᕐᓴᐅᓂᖅᑐᓂᒃ ᐊᐅᓪᓕᓂᐸᓐᓂᖅ

Management Approaches outline ways to address the Management Challenges, while meeting the Management Goals

- Cultural Resources Management
- Wildlife and wildlife habitat Management
- Monitoring and Research
- Public Awareness and Information Management

ᐊᐅᓪᓂᐊᓂᖅᓯᓂᖅ ᖃᓄᐃᓕᓐᓄᑦᓯᐊᖏᓂᖅ Management Approaches

ᐱᐅᓯᑐᖃᖅᑎᒍᑦ ᐊᕐᕌᑎᓄᑦ ᓄᓇᒥᐅᑕᓂᖅᑐᑦ ᐊᓚᒐᓗ ᐅᒪᔨᐃᑦ ᐱᑕᖃᖏᓂᖅᑐᑦ

ᐅᖃᓕᖅᑐᑦᑎᐊᖃᑕᖏᓂᖅ ᓴᐳᓂᓂᐊᖅᑕᐅᓂᖅᑐᓂᓂᒃ ᐃᓄᖅᖃᖏᐅᕐᓂᐅᔪᓐᓇᖅᑐᖅᑐᑦ ᖃᑭᒋᐃᓂᖅᖃᖏᓂᖅᑐᑦ, ᐃᓄᐃᓪᓗ ᐅᖃᐅᔾᔨᑐᖅᑐᑦ, ᐃᓄᐃᓪᓗ ᓄᓇᒥᓂᒃ ᐃᒪᖏᑎᖅᑐᓂᓗ ᑕᐃᔭᐳᑐᖅᑐᑦ, ᐃᓕᓴᐅᕐᓕᓪᓕᐊᓗᑎᒃ ᖃᐅᔨᒪᔭᐅᕐᓕᓪᓕᐊᔨᓄᑦ, ᑎᑎᕋᖅᑕᐅᕐᓕᓪᓕᐊᖃᑦᑕᐃᓐᓇᓪᓗᑎᒃ ᐊᒪᓚᓗ ᓄᓇᖓᐊᓕᐅᖃᑦᑕᕋᓗᑎᒃ

Cultural Resources Management

Ensure preservation of archeological sites, Inuit language, and Inuit place names and add to knowledge through inventory and mapping projects

ᐊᐅᓪᓕᓂᐅᖅᓯᒪᓂᖅ ᖃᓄᐃᓕᓴᖅᓯᐊᕐᓯᖅᓯᒪᓂᑦ Management Approaches

ᐅᒪᔨᐃᑦᓗ ᐅᒪᔨᕐᓄᑦᓗ ᓇᔨᕕᓴᐅᕙᖅᓯᖅᓯᒪᓂᖅ ᐊᐅᓪᓕᓂᐅᖅᓯᖅ

- ᐊᓯᔾᔨᖅᑕᐅᓗᓂᓂ ᐃᓕᑕᕕᓴᐅᓂᖅᓴᓄᑦ ᑎᒃᑯᐊᖅᑕᐅᓯᒪᓂᖅᓴ ᖃᓇᑕᒥ ᐅᒪᔨᖅᓴᕐᐃᐅᓂᕐᒧᑦ, ᐱᔭᔨᑕᐅᔨᓐᓇᕐᒪᑦ ᐊᓯᔾᔨᖅᑕᐅᔨᓐᓇᕐᓂᖅᓂᓂᒃ ᑭᓪᓕᖅᓴᕐᐃᐅᓂᖅᓯᒪᓂᑦ
- ᐊᒥᓱᓴᖅᓂᒃ ᖃᓄᐃᓕᓕᐅᑕᐅᓂᐊᖅᑐᓂᒃ ᐃᑲᔨᖅᓯᐃᓗᓐᓂᑎᒃ ᐊᐅᓪᓕᑎᒃᔨᐅᑎᑲᓄᑦ ᖃᖅᓴᐃᑦ ᐊᒥᓱᓂᖅᓯᖅᓂᒃ
  - ᐱᔨᒥᓴᐃᓂᖅ ᓄᓇᓕᖅᓂ ᖃᖅᓴᓯᓯᐅᖅᑕᑕᕕᔨᓐᓂᕐᒥᒃ, ᑐᓂᐅᖅᐃᖅᑕᑕᐅᑎᐊᓐᓂᖅ ᐊᐃᑦᑐᓐᓴᒃᓗᑎᒃ ᐃᓄᖅᑎᑎᓄᓗᓄᑎᒃ ᐊᖅᓯᓓᑕᕕᔨᓐᓇᖅᑐᑕᕕᓴᐅᓂᖅ ᐊᓗᒪᓗ ᑭᓇᐅᔭᓕᐅᑕᐅᔨᓐᓇᖅᑐᑕᕕᓴᐅᓂᖅ ᐊᑐᑭᓐᓇᖅᑐᑕᕕᓴᐅᓂᖅ ᐊᑐᔨᒪᓗ ᖃᐅᔨᓴᓂᒃ ᖃᐅᔨᓴᖅᑕᐅᓐᓇᓕᓕᐅᑕᐅᓂᖅ
- ᐊᒃᑐᖅᑕᐅᓂᕐᓗᒃᑎᑦᑎᑕᐃᓕᒪᕐᐊᕐᓂᖅ ᑕᓇᐅᒃᓯᒃ ᐅᒥᐊᕐᔪᐊᕐᓂᒃ ᖃᖅᓱᑕᓯᐅᑦᓯᓕᕐᓂᖅᓴᓄᑦ ᐅᒪᔨᐃᑦ ᐱᓕᕐᓂᑦ

Wildlife and wildlife habitat Management

- Change designation to National Wildlife Area, which may involve a boundary change
- Support several strategies to manage goose population
  - Promote local harvest, sharing, sport hunting, and commercial opportunities, ongoing research
- Mitigate marine and air traffic impacts on wildlife

ᐊᐅᓪᓕᓂᖅᓯᓐᓂᖅ ᖃᓄᐃᓕᓴᕐᓯᐊᕐᓯᓐᓂᒃ
Management Approaches

ᓇᐅᑦᑎᖅᑐᑕᐊᖅᑕᐅᖃᑦᑕᐃᓐᓇᕐᓯᓐᓂᒃ ᐊᓪᓚᓗ ᖃᐅᔨᓴᖅᑕᐅᖃᑦᑕᕐᓯᓐᓂᒃ

- ᐃᑲᔪᖅᓭᐃᓂᖅ ᑲᔪᓭᓐᓂᖅᓂᑉ ᖃᐅᔨᓴᖅᑕᐅᓕᕐᓂᐊᔪᑦ ᐊᓪᓚᓗ ᓇᐅᑦᑎᖅᑐᖅᑕᐅᓴᖅᑐᖃᑦᑕᕐᓯᓐᓂᒃ, ᑎᑎᕐᖅᑕᐅᓕᕐᓂᐊᓗᑎᑦᓗ ᐃᓗᖃᖏᕐᕕᐅᓕᐅᖅᑐᑦ ᖃᑭᒪძᕕᓂᖃᖏᕐᕕᐅᔪᑦ
- ᐃᑲᔪᖅᓭᖅᑕᐅᓗᑎᑉ ᓴᓂᓕᕆᐅᑕᓗᑦ ᖃᐅᔨᓴᖅᑕᐅᕐᔪᐊᕐᓇᔪᑦ ᓭᓄᓂᓕᐅᕐᔪᐊᕐᓂᓗᑎᑉ
- ᐱᐅᓱᕈᐊᖅᑕᐅᓗᑎᑉ ᖃᐅᔨᒪᓴᓱᑦ ᐊᓕᑎᒥᑉ ᐅᒪᔪᓚᕆᐅᓕᖅᑐᑦ ᐊᒪᓗ ᐱᐅᓱᑐᖃᖅᑎᒍᑦ ᐊᓕᑎᖏᓐᓂᒃ ᓄᓇᒥᐅᑕᕐᓂᑦ ᐊᒪᓗ ᐅᒪᔪᐃᑦ ᐱᑕᖃᕐᓯᓐᓂᒃ

Monitoring and Research

- Support ongoing research and monitoring, document archaeological sites
- Support Salliqmiut research priorities
- Improve knowledge of ecological and cultural resources in the MBSs

ᐊᐅᓪᓂᓂᖅᓯᓐᓄ ᖃᓄᐃᓕᖅᑐᓂᒃᑯᓂᖅ
Management Approaches

ᐃᓄᓕᒻᓂᖅ ᖃᐅᔨᒪᑎᑦᑎᐊᓪᓕᐊᓂᖅ ᐊᓪᒪᓗ ᖃᐅᔨᒪᕗᐅᔪᕐᓂᖅᓯᓐᓄᑦ ᐊᐅᓪᓂᐊᖅᓂᖅ

- ᓴᖅᑭᑎᑦᑎᓗᑎᑦ ᑐᓴᐅᒪᕗᐅᑎᑦᑎᖃᑦᑕᕐᓂᕐᒧᑦ ᕿᕐᓇᐅᑎᐸᓴᓂᖅ ᐊᓪᒪᓗ ᐱᐃᐸᓴᖃᖅᑎᑕᐅᐊᖅᓂᕐᓂᓄᑦ ᐃᓄᐃᑦ ᑐᓴᐅᒪᑎᑦᑎᖃᑕᑭᓇᕐᓂᕐᒥᐸᑦ ᖃᐅᔨᒪᕗᖅᑭᓪᓂᕐᓂᖅᓯᓐᓄᑦ, ᖃᐅᔨᒪᕗᖅᓂᕐᓂᓪᓗ ᓄᓇᒧᖏᐅᐊᐸᑐᓄᑦ, ᐅᖃᐅᔨᔪᐅᔪᓇᖃᖅᑎᑕᕐᓗᑎᓪᓗ ᐃᔪᒪᓗᑕᐅᔪᑦ ᐃᓄᖅᓄᑦ ᐊᓪᒪᓗ ᖃᐅᔨᓴᖃᑕᐅᓂᐊᖅᑐᑐᓂᖅᓯᓐᓄᑦ ᔪᐊᓕᓐᓂᐊᐅᑎᑕᐅᓂᐊᖅᑐᑐᓂᖅᓯᓐᓄᑦ
- ᐱᔪᒥᓴᐃᓂᖅ ᐃᑲᔪᖅᔪᐃᐊᖅᓂᕐᓗᓪᓗ ᐃᓄᖅᓂᑦ ᐊᓪᒪᓗ ᐃᓄᖅᓂᑦ ᐃᓕᓴᓂᐊᖅᑎᑕᐅᔪᓂᑦ ᐱᖃᑕᐅᔪᒪᓂᖅᓂᑦ ᓄᓇᓕᖅᓂ ᐊᔪᖅᑭᐃᓂᖅᓯᓐᓄᑦ ᐊᑐᖅᓂᖃᕐᓇᖅᑎᑕᐅᓇᖅᓂᖅᓂᑦ
- ᐃᑲᔪᖅᔪᐃᐊᓂᖅ ᑲᔪᔭᖅᓇᑭᖅᑐᓂᑦ ᐊᑐᖅᑕᐅᓂᖅᓇᑭᖅᑐᓂᑦ ᐅᓪᓗᒥ ᑎᖅᑕᐊᖅᑲᐃᖅᓂᖅ ᐊᐅᓪᓂᐊᖅᓂᖅᓯᓐᓄᑦ ᐃᓗᓐᓂᐊᓂᖅ

Public Awareness and Information Management

- Develop a communication plan and provide opportunities to share information and knowledge among users, discuss concerns and research priorities
- Encourage and support Inuit and Inuit student initiatives to build local capacity
- Support sustainable tourism operations within the MBSs

ᐊᐅᓪᓕᓂᐅᖅᓯᖅ ᖃᓄᐃᓕᖅᓱᓕᐊᕐᓯᖅ
Management Approaches

ᐃᓄᓕᒻᒥᐅᖅ ᖃᐅᔨᒪᑎᑦᑎᐊᕐᓕᐊᓂᖅ ᐊᖏᒪᓗ ᖃᐅᔨᒪᓴᐅᔪᖅᓴᐅᓂᖅᓯᖅᓄᑦ ᐊᐅᓪᓕᓂᑎᐊᕐᓂᖅ

- ᑎᑎᕆᖅᑕᐅᔪᒪᓗᑎᑦ ᓄᓇᒋᑦᑕᖅᑐᕆᐊᐅᓪᐊᕐᑐᑦ, ᖃᐅᐸᐊᖅᑕᐅᖃᖅᑎᑦᑕᕆᓗᑎᒃᓗ ᐊᖏᔪᓇᔪᓰᑦ ᑲᑐᔨᖃᑎᒋᖅᓯᖅᓄᑦ ᑎᑦᔨᕆᐊᑎᒋᓄᑦ ᑕᐃᔪᑯᐊ ᓄᑕᓂᓂᑦ ᑐᔪᔪᕐᐅᑎᑦ ᑎᖅᑎᐊᖃᖅᕆᒋᑦᑐᒪᓂᖅᓯᖅᓄᑦ, ᖃᐅᔨᕆᖅᑕᐅᖃᑕᕆᓕᑎᓐᓗ ᐃᓐᓇᓂᓪᓕᐅᔪᖏᖅ
- ᒪᓪᓕᐊᖃᑕᕆᓂᖅᑎᑦ ᐱᔨᒥᕆᐃᐃᓂᖅ - ᑐᕆᐅᒪᑎᑦᑎᖃᑕᕆᓗᑎᒃ ᖃᐅᔨᒪᓴᐅᓪᓕᐊᖅᑐᓂᒃ ᒪᓪᓚᓪᓀᔨᓂᒃ ᐊᒻᒪᓗ ᒪᓪᓚᓵᓗᑐᓴᐅᐊᖅᑐᓂᒃ, ᐊᐅᓪᓕᓂᑎᓂᕐᒧᓵᓗ ᑐᖏᓚᓀᓴᐅᔨᔨᓂᒃ ᐊᒻᒪᓗ ᐱᓪᓀᐊᖏᓂᔪᒪᔨᓂᒃ

Public Awareness and Information Management

- Maintain list of camps, assess new requests inside MBSs with HTO and inspect periodically
- Compliance promotion – share information about laws and regulations, management goals and objectives

ᐊᖏᑭᑕᐅᓯᒪᔪᑦ ᐱᓕᕆᐊᖓᔪᓂᓐᓄᑦ ᑎᑭᓯᒪᖓᐊᖅᓂᓐᓄᑦᑐᓕᓗ
Authorized Activities and Access



- ᐃᓄᐃᑦ **ᐱᔨᖅᓇᐅᑎᖏᕆᐊᖏᖅᑎᑕᐅᖅᖁᓕᑦ** ᑎᑭᓯᒪᔨᖅᓇᓐᓄᑦᓂᓄ ᑎᖅᑕᐊᖏᖅᓴᑦ ᓄᓇᖅᑐᓄᑦ, ᐱᐅᓯᑐᖏᑎᒃ ᐊᑐᓂᐊᖏᑎᓪᓗᑎᑦ, ᓯᑦᓗ ᐊᖓᔪᓇᖅᓂᖅᖁᖅᑐᓂᓄ, ᓴᓇᖅᖁᐊᖓᕆᓴᒃ ᐅᓴᕋᐃᓴᐃᓂᖅᖁᖅᑐᓂᓄ, ᓄᓇᒥᑕᖏᑐᓴᖅᑐᓂᓄ (ᒪᓕᒐᖅᑎᑦ ᒪᓕᒐᓗᑎᑦ ᓄᓇᕗᓴᒥ ᓄᓇᑕᑭᑎᓄᑦ ᐊᖅᑭᑐᑎᑦ ᐊᒡᒪᓗ ᐃᓄᐃᑦ ᐊᒃᑐᖏᑕᐅᓯᒪᓂᖅᑐᓄᑦ ᐃᑲᔪᓯᐊᕐᓂᖅᑐᓄᑦ ᐊᖅᑭᑐᑎᑦ)
- ᐃᓄᐃᑦ **ᐱᔨᖅᓇᐅᑎᖏᕆᐊᖏᖅᑎᑕᐅᖅᖁᓕᑦ** ᑎᑭᓯᒪᔨᖅᓇᓐᓄᑦ ᑎᖅᑕᐊᖏᖅᓴᑦ ᓄᓇᖅᑐᓄᑦ ᐊᖅᑭᑐᑎᖅᑕᐃᔨᓂᖅᑐᓄᑦ ᐅᕝᕙᓚᐅᔪᓂᑦ ᐃᖏᓪᓗᓯᐅᑦᑎᐊᓂᖅᑐᓄᑦ

- Inuit **do not** require a permit to enter the MBSs for traditional activities, including harvesting, removal of carving stone, setting up camps *(Nunavut Agreement* and IIBA)
- Inuit **do not** require a permit to enter the MBSs when working as a hunting or fishing guide



ᐊᖏᑭᑕᐅᓕᕆᓂ ᐱᓕᕆᐊᖏᔪᓐᓇᖅᓂᖅ ᑎᑭᓕᒫᐅᔪᓐᓇᖅᓂᖏᑦᓗ
Authorized Activities and Access

- ᑕᒪᖅᓂᒃ ᐃᓄᒡᒪᓗᐅᖏᑦᑐᑦ **ᐱᔪᓐᓇᐅᑎᖅᑲᖅᑎᑕᐅᔭᑦ** ᑎᑭᓕᔪᓐᓇᖅᓂᖅ ᑎᖏᒥᖅᑲᖅᕕᖓᓄᑦ (ᓄᓇᖏᑦᓂᓄᓐᓂᑦ ᐃᒪᖅᔪᑎᖏᑦᓂᓄᓐᓂᑦ) ᑕᐃᒃᑯᐊᓗ ᐱᔪᓐᓇᐅᑎᖅᑲᖅᑲᖅᑐᑦ ᐃᓄᒡᒪᓗᐅᖏᑦᑐᑦ ᐊᑭᓐᓇᒃᑐᓂᒃᑐᒥᒃ ᐊᖏᔪᓇᓕᐊᑭᓴᓇᒃᑐᑦ, ᖃᐅᔨᓴᖅᑎᓗᑦ, ᐳᓪᓕᑎᓪᓗ ᓂᐅᖅᑕᐊᑦ

- ᑕᒪᖅᑭᒃ ᑭᓇᐅᔭᓪᓕᐅᑎᖃᖅᓂᖅᓗᑦ ᐱᓪᓚᐊᖃᖅᓂᖅᓗᓴᓪᓕᓂᑦ ᐱᓪᓚᓂᖅᓗᑦ (ᐃᓄᐃᑦ ᐊᒻᒪᓗ ᐃᓄᒡᒪᓗᐅᖏᑦᑐᑦᖃᖅ ᓇᖏᑭᓂᖅ ᐱᓪᓚᐃᖅᑯᑎᖃᖅᑐᑦ **ᐱᔪᓐᓇᐅᑎᖅᑲᖅᑎᑕᐅᔭᑦ**. ᑕᐃᒪᓇᔨᖏᑦᑐᑐᓄᑦ ᐃᓄᐃᑦ ᐊᖏᔪᓇᓕᐊᑭᓴᑦ ᐊᒻᒪᓗ ᐃᖃᓪᐊᖅᑭᒪᓕᕆᑦ

- All non-Inuit **do** require a permit to enter the MBSs (land or water) including non-Inuit sport hunters, researchers and tourists

- All commercial or business activities (Inuit and non-Inuit owned businesses) **do** require a permit. Except for Inuit hunting and fishing guides



ᐱᔨᓇᐅᑎᖅᑕᐅᔨᓇᓂᓐᓕᒃ
Permit Authorizations

- ᐱᔨᓇᐅᑎᓂᒃ ᑐᓂᔭᐅᔨᓇᖅᑐᑦ ᐃᒪᓇᐃᒃᑯᑎᒃ:
  - ᐱᓕᕆᐊᕆᔪᒪᔭᒃᓕᑦ ᓯᕐᖅᑎᕆᓗᑕᐅᓂᐊᖏᓕᑐᑐᒋᐊᕈᑎᒃ ᑎᖏᒥᐊᖅᓄᑦ, ᒪᓇᓂᖏᓇᓄᑦ, ᐅᓪᓗᖏᓇᓄᑦ, ᓇᔨᖅᐸᒃᑕᖏᓇᓄᓪᓗ ᐊᒻᒪᓗ/ᐅᕐᓂᓗᓇᓚᑦ
  - ᐱᓕᕆᐊᕆᓂᐊᖅᑕᖏᑦ ᒪᓕᑎᐊᕈᑎᒃ ᐱᑕᖅᕈᑕᐅᓂᖏᓄᓄᑦ ᑎᖏᒥᐊᖅᑲᐃᑦ ᐊᒻᒪᓗ ᐊᐅᓪᓚᑎᐅᑎᒃᓴᐃᑦ ᑕᐃᒃᑯᐊ ᕿᖏᓇᐅᑎᒃᓴᖏᑦ
- Permits may be issued if:
  - The activity is not harmful to migratory birds, eggs, nests, or habitats and/or
  - The activity is consistent with the purpose of the MBSs and their management plans



ᓱᓴᓕᐊᖅᓯᐊᖅ White-rumped Sandpiper





ᓴᐅᖅᑐᖅ Red Phalarope

ᐱᓯᓂᓇᐅᑎᖅᑭᑕᐅᓯᓂᓇᖅᓂᕐᓂᑦ
Permit Authorizations

- ᐃᓄᐃᑦ ᐱᓯᓂᓇᐅᑎᖅᑏᐊᖅᖃᑦ ᑎᑭᓕᒪᓯᓂᓇᖅᓂᖅ ᑎᓐᒥᐊᖅᖁᕕᓂᓄᑦ ᐅᖅᑐᑭᓂᓇᖅᐳᑦ ᐱᓇᓕᓴᑎᖅ ᖃᓇᑕᒥ ᐅᒪᓯᓪᓗᓵᖅᓄᑦ ᐱᓕᑎᑕᐃᓕᓴᐅᓂᓂᖅ:

ec.nupermisscf-cwspermitnu.ec@canada.ca

ᐃᖅᓴᓐᓂ ᐃᖅᓇᐃᕆᖁᕕᖅ: 867-975-4636

- ᐃᖁᓂᐅᖁᕕᖅᓄᑦ ᐊᐅᓪᓚᑎᐅᖅᑕᐅᓂᖁᑦ ᖃᑎᒪᓕᕆᓪᓕᐅᑦ ᖁᒥᖁᑭᐊᖅᑎᐅᐳᑦ ᑕᒪᐃᓂᓂᑎᐊᖅ ᐅᖅᑐᑭᑎᓂᖅ ᐱᓯᓂᓇᐅᑎᑦᑕᕐᓂᖅᑐᓂᖅ ᐊᒻᒪᓗ ᐊᑐᓪᓚᑦᓄᓕᕐᓂᖅ ᐃᖃᓗᐃᑦ ᐅᒪᓯᓪᓗᓵᖅᓄᑦ ᐃᖅᓇᐃᕆᖁᕕᖅᓄᓪᓕᐊᓂᑦ ᐊᓂᕐᑕᐅᓂᖁᕕᖅᓄᓪᓕᐊᓂᑦ ᓇᓗᓇᐃᕆᐃᓕᓴᐅᓂᖅ

- Persons who require a permit to enter the MBSs can apply for a permit from the Canadian Wildlife Service:

ec.nupermisscf-cwspermitnu.ec@canada.ca

Iqaluit office: 867-975-4636

- The Irniurviit ACMC reviews all permit applications and provides recommendations to the Canadian Wildlife Service on whether the permit should be issued or not



ᓄᓇᖅᑐᑦ ᑎᒃᓯᐊᖅᑕᐅᓯᒪᓂᖅᑐᑦ / Site Designation

- ᖃᖅᖢᐅᖅᑐᖅ ᐊᓪᒪᓗ ᐃᑲᑦᑐᐊᖅ ᐃᓐᓇᐅᔪᖅ ᐃᓕᑕᒫᓴᐅᓯᒪᖏᒪᑦ ᑎᒃᓯᐊᖅᑕᐅᓯᒪᓕᒫᑦ ᑎᖅᑎᐊᖅᖃᖅᓴᐅᓂᖅᓗᑦ ᓴᓯᐊᓂᓕᕆᔨᖅ; ᐅᓯᐊᓂᓕ, ᐅᒪᔪᐃᑦ ᐊᓪᒪᓗ ᐱᐅᓯᑐᖅᑎᒍᑦ ᐊᐳᑎᖅᑎᒍᑦ ᐊᓪᒪᓗ ᐅᒪᔪᐃᑦ ᐱᑕᖃᖅᓂᖅᑕᑦ ᐊᐳᓂᓂᐊᖅᑕᐅᓯᒪᑦᑎᐊᖅᓂᖅᖢᐅᔪᓐᓇᖅᐳᑦ ᐃᓕᓴᐅᒪᖏᑦ ᐊᓕᖅᑐᓂᒃᓴᓂ ᓄᓇᖅ ᑕᐅᓀᓐᓇ ᑲᓇᑕᒥ ᐅᒪᔪᖅᓴᖅᓴᐅᔪᑎᑐᑦ
- ᐊᐅᓪᓕᑎᐅᕐᓴᐃᑦ ᐸᖅᓇᐅᑎᖅᑦ ᐅᖅᐅᓯᖅᓕᐳᑦ ᖃᓄᐅᓕᖅᑐᑦ ᓴᓇᓕᑕᒫᓴᐅᓂᖅᑐᑦ ᒪᓕᐢᑕᐅᓯᒪᓂᖅᑐᓄᑦ ᑲᑎᒪᓂᖅᑐᑦ ᐊᓕᖅᑐᓂᒃᓴᓂ ᐊᐅᓪᓕᑎᐅᕐᓴᐃᑦ ᐱᐅᓯ


ᐊᐅᓪᓕᐊᒃᑲᐃᓕᓪ ᐸᖏᓐᓂᐅᓂᒃ ᐊᑐᓪᖅᑎᑎᓂᖅ
Management Plan Implementation

- ᐊᑐᓪᖅᑎᑕᐅᓐᓪᓚᐊᓂᖅᑐᑦ ᐊᑐᒐᒃᑲᐃᓕᓪ (ᐱᕆᒃᑲᖏᕆᐊᖅᖏᓂᔪᓂ ᑭᓇᐅᔭᐃᓪᓗᓂ)
  ᐱᔨᓴᓇᐅᑕᐅᓂᐊᖏᒪᑕ
- ᐸᖏᓇᐅᑎᓯᓂᑦ ᖃᑎᖏᑭᐊᖅᑕᐅᔨᓴᓇᖏᓂᖅ ᑕᓪᓕᒪᐅᓂᐊᖅᑐᑦ 5 ᐊᖏᕿᔪᓂᐊᖅᑐᑦ ᓇᒃᓱᖅᐸᑕ, ᑭᖑᓂᐊᒍᓪᓗ ᑕᐃᑲᖑᓐᓂ ᖁᓕᓪᓂ 10 ᐊᖏᕿᒍᐃᓪ ᓇᒃᕃᖑᓐᓇᑎᒃ



- Implemented as resources (time and money) allow
- Plan will be reviewed after 5 years, and then every 10 years after that

2025 / 2035



ᐃᖃᓗᖅᓯᒪᓂ ᐊᐅᓚᑎᕐᓗᐅᖏᓐᓄᓕᒫᖃᑦᑕᓂ ᑲᑎᒪᔨᖏᓚᐅᖅᑕᑦ ᑕᒻᒧᐊ ᑭᖑᓪᓚᐅᙵᖅ
ᑎᑎᕋᖅᓱᒪᓕᕐᓂᑦ ᐊᐅᓚᑎᑎᓕᓴᐃᑦ ᐸᖃᐅᑎᖅᓴᐃᑦ ᖃᖅᓴᐅᖅᑐᑎᒃ ᐊᓱᒪᓗ ᐃᑲᑦᑐᐊᖏᒃ
ᑎᕐᒥᐊᖅᖃᓰᕐᓂᖅ ᐊᕐᕈᖅᑕᐅᖃᓕᓚᓂᑦ ᓄᓇᕗᒻᒥ ᐅᒪᔨᓚᓱᔨᕐᓴᐊᑦ ᑲᑎᒪᔨᕐᕕᓂᑦ
The Irniurviit ACMC recommends the Final Draft Management Plans for Qaqsauqtuuq and Ikkattuaq Migratory Bird Sanctuaries for the Nunavut Wildlife Management Board's approval

ᐅᖃᖃᓗᓴᑦ ᖃᐅᔨᓕᐊᖃᓪᓚᓂ:

ᐃᖃᓕᖃᓗᓴ ᐊᐅᓚᑎᑎᓕᓴᐃᑦ ᑲᑎᒪᔨᖏᓚᓕᑦ

ᑲᒪᔨᐅᔨᖅ: ᐊᖏᓚᔨᖃᖅᑐ ᑐᖏᓕᖏᓚ ᐅᓴᐃ-ᐃᐊᑉ ᑐᓄᐊᖃ

ᖃᓚᑕᐅᓴᓴᓂᑦ ᑎᑎᖃᖅᓗᖏᓚᓂᓕ

jean-francois.dufour2@canada.ca

ᐅᖃᓕᐅᓕᒥ (867) 669-4766

Contact Information:

Irniurviit Area Co-Management Committee

c/o Vice-chairperson JF Dufour

jean-francois.dufour2@canada.ca

(867) 669-4766

Laura McKinnon

2016ᒥᑦ 2023ᒧᑦ

ᐃᓄᐃᑦ ᐊᖏᔪᖅᑕᐅᓂᐊᕐᓂᖅᐸᓂᐅᓐᓄᑦ ᐱᓕᕆᔨᑕᐅᔪᓐᓇᖅᑐᓄᑦᓗ ᐊᖑᑎᒃ

—

ᑲᓇᑕᒥ ᐅᒪᔪᖅᑲᓴᐃᑦ ᐊᓚᒪᓗ ᐅᑎᖅᑕᕐᕕᒋᑦ ᑎᖏᒥᐊᓄᑦ ᓴᓴᐅᔪᓐᓇᖅᑐᑦ ᓄᓇᓕᒫᑦ ᐅᓗᓄᐃᖅᑕᐅᔪᓂᖏᓐᓄᑦ ᑲᑐᔾᔨᓗᑎᒃ ᐅᑯᐊ ᐃᓄᐃᑦ ᓄᓇᕗᑦᑎᒋᔭᓄᑦ ᑭᓱᖅᑐᖅᑕᐅᓪᓗᑎᒃ ᓄᓇᖅᑐᑦ ᑐᖏᖅᓯᒪᔪᑦ ᐃᓄᐃᑦ ᑎᑎᖅᑕᐅᓪᓗᑎᒃ ᐊᖑᑎᒃ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑐᑦ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑐᑦ ᐊᓚᒪᓗ ᑯᐃᓐ ᐱᓕᕆᓂᖅ ᑲᓇᑕᒥ, ᑭᖑᓪᓕᖅᑐᑎᑎᓪᓗᒍ ᒥᓂᑦ ᐊᐅᓚᑦᓯᐊᕆᔭᖅᑯᓄᑦ

2016 TO 2023 INUIT IMPACT AND BENEFIT AGREEMENT FOR NATIONAL WILDLIFE AREAS AND MIGRATORY BIRD SANCTUARIES IN THE NUNAVUT SETTLEMENT AREA

—

BETWEEN
The Inuit of the Nunavut Settlement Area
represented by Nunavut Tunngavik Inc.
The Kitikmeot Inuit Association
The Kivalliq Inuit Association
The Qikiqtani Inuit Association
AND
Her Majesty the Queen in right of Canada, as
represented by the Minister of the Environment



ᖃᐅᔨᓴᖅᑎᑐᑦ ᐃᓄᐃᑦ ᑲᑎᒪᔨᖏᑦᑐᖅ
Qikiqtani Inuit Association

Environnement et Changement climatique Canada

Environment and Climate Change Canada

ᐃᓗᓕᖅᑐᑦ

2016 ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᕐᓂᖅᓯᓄᑦ ᐱᓚᒥᓕᕆᑕᐅᔨᓴᖅᑐᓄᓪᓕᒃ ᐊᖏᕈᓐᓇᑦ ᑲᓇᑕᒥ ᐅᒪᔨᖅᑎᑦᑕᓄᑦ ᑎᖏᒥᐊᓄᑦ ᓴᓴᐅᔭᓕᕐᓂᖅ ᓄᓇᕗᑦ ᓇᓗᓇᐃᖅᑕᐅᔭᓕᖏᓐᓄ **03**

ᐃᓕᖏᓐ 1 — ᑕᐃᔪᓴᖏᑦ. **05**

ᐃᓕᖏᓐ 2 — ᖃᓄᐅᑐᐃᓇᖅ ᒪᓕᒃᓴᕆᐊᓕᒥᑦ. **10**

ᐃᓕᖏᓐ 3 — ᐊᐅᓕᕐᑎᖃᑎᒋᑦ **18**

ᐃᓕᖏᓐ 4 — ᐃᓄᐃᑦ ᓄᓇᕋᑎᖏᑦ **30**

ᐃᓕᖏᓐ 5 — ᐃᓄᐃᑦ ᐱᔨᓇᕐᓂᖏᑦ ᐊᓐᒪᓗ ᐊᑐᖅᑕᐅᓴᓴᓴᖏᑦ ᑲᓇᑕᒥ ᐅᒪᔨᖅᑎᑦᑕᓄᑦ ᐊᖏᕈᓐᓇᑦ ᑲᓇᑕᒥ ᐅᒪᔨᖅᑎᑦᑕᓄᑦ ᑎᖏᒥᐊᓄᑦ ᓴᓴᐅᔭᓕᖏᓐᓄ. **36**

ᐃᓕᖏᓐ 6 — ᑐᓴᐅᒪᑎᑦᑎᓂᖅ, ᑎᑎᕋᖅᓴᒪᔨᓴᑯᑎᖏᑦ ᐊᒪᒪᓗ ᐃᓗᓐᓂᓴᑯᑎᖏᑦ. **40**

ᐃᓕᖏᓐ 7 — ᐳᓕᕋᖅᑐᑕᕆᓂᖅ ᐊᒪᒪᓗ ᐊᓴᖏᑦ ᓇᖏᒥᓂᕆᔨᐅᔭᓄᑦ ᐃᑲᔪᑎᑎᔨᑦ **46**

ᐃᓕᖏᓐ 8 — ᐃᓄᐃᑦ ᑲᓐᑐᖅᑕᐅᓴᒪᓐᑎᖅ ᐊᒪᒪᓗ ᓇᖏᒥᓂᖅᑲᖅᑐᑦ ᐱᕆᖅᑎᑎᑕᐅᓐᑐᑎᖅ **48**

ᐃᓕᖏᓐ 9 — ᐃᑕᓇᓂᐊᕐᑎᑎᓂᖅ ᐊᒪᒪᓗ ᐃᕋᖅᑲᓇᐃᔨᖅᑎᑦᑎᓂᖅ **52**

ᐃᓕᖏᓐ 10 — ᕋᖅᐅᓵᓴᕐᓂᖅ **56**

ᐃᓕᖏᓐ 11 — ᐃᓄᖓᓄᑦ ᐱᓪᒪᓚᕈᑎᐅᔨᑦ ᐊᒪᒪᓗ ᐱᕐᔪᓴᑐᖅᖏᓕᑦ **60**

ᐃᓕᖏᓐ 12 — ᓂᕐᔨᑎᖅᑎᖃᕐᑦ. **62**

ᐃᓕᖏᓐ 13 — ᐊᓴᑦᐳᐅᑎᑦ ᑕᒃᔪᓄᖓᓗ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᖏᒥᐊᓄᑦ ᑲᓇᑕᒥ ᐅᒪᔨᖅᑎᑦᑕᓄᑦ ᐊᒪᒪᓗ ᐊᖏᕈᓐᓇᑦ ᑲᓇᑕᒥ ᐅᒪᔨᖅᑎᑦᑕᓄᑦ ᐃᑲᔪᑎᑎᔨᑦ. **66**

ᐃᓕᖏᓐ 14 — ᐳᓴᖅᑐᑦ ᐃᓴᐳᓇᖅᑕᖏᑦ ᐊᒪᒪᓗ ᐊᑐᐳᓇᖅᑕᖏᑦ **71**

ᐃᓕᖏᓐ 15 — ᐊᑐᓕᖅᑎᑕᐅᔭᓕᓕᐊᔨᑦ, ᖃᐅᔨᐳᐊᓂᖅ ᐊᒪᒪᓗ ᐊᐃᔨᖅᑎᑎᔪᓐᓇᖅᑲᓇᓂᖅᓂᖅ. **74**

ᐃᓕᖏᓐ 16 — ᐊᐃᔭᐅᑎᔨᓄᑦ ᑐᐸᑕᐳᑎᑦ **79**

ᐅᐃᒍᖅ I — ᓂᖏᑎᓴᓂᖅ ᑲᓇᑕᒥ ᐅᒪᔨᖅᑎᑦᑕᓄᑦ **81**

ᐊᑎᓕᐅᖅᑐᑦ **83**

2016 ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᖅᓯᖏᓐᓄᑦ ᐱᓕᕆᓂᑕᑎᐅᔪᓐᓇᖅᑐᓄᓪᓗ ᐊᖏᑕᐅᑐᑦ ᑲᑎᒪᔨᖅᑎᒃᑯᑦ ᐊᑐᓚᓗ ᐅᑎᖅᑕᕐᕕᑦᑐᓄᑦ ᑎᖏᒪᐊᓄᑦ ᓴᓴᐅᓯᒪᓕᕐᓂᖅ ᓄᓇᕗᑦ ᐅᓪᓗᐃᐸᖅᑕᐅᓯᒪᓂᖏᓐᓄᑦ

ᐃᓕᐃᒪᓕᑦ ᑲᓇᑕᒥ ᐅᓕᔪᖅᑎᒃᑯᑦ ᓴᖅᑭᑕᐅᓯᒪᖏᓕᑕ ᐊᑕᐃᔨᑦ ***ᑲᓇᑕᒥ ᐅᓕᔪᓕᕆᓂᕐᔪᐃᑦ*** ᐱᖁᓯᖁᔪᓂᖅ ᐱᔪᓐᓇᑎᑦᓯᐊᓂ ᓯᖁᓂᕐᓂᑐᐃᓐᓇᑎᒃ; ᐊᒻᒪᓗ

ᐃᓕᐃᒪᓕᑦ ᐅᑎᖅᑕᐊᑦᑐᓄᑦ ᑎᖏᒥᐊᓄᑦ ᓴᓵᖅᑐᐅᓯᒪᓕᖏᓐᓄᑦ ᓴᓴᑭᑕᐅᓯᒪᖏᓕᑕ ᐊᑕᐃᔨᑎᑎ ***ᑎᖏᒥᐊᑦ ᐅᑎᖅᑕᐊᑦᑐᕆᐃᑦ ᐱᔪᐊᑕᐅᓂᖏᓐᖓᓄᑦ ᐱᖁᓯᖁᔪᓂᖅ*** ᐱᔪᓐᓇᑎᑦᓯᐊᓂ ᓴᓵᖅᑐᐅᓯᒪᖏᓕᑕ ᐊᑕᐃᔨᑎᑎ ᐅᑎᖅᑕᐊᑦᑐᓄᑦ ᐱᔪᓐᓇᑎᑦᓯᐊᓂ ᓴᓴᑭᑕ

ᐃᓄᐃᑦ ᐊᑲᖏᑕᐅᓂᐊᖏᓂᖅᐳᓄᑦ ᐱᓯᒪᓕᕆᑕᐅᔪᖅᑐᓄᓪᓕ ᐊᖏᕈᕆᑦ, ᐊᒻᒪᓗ ᑕᒪᖏᑲ ᖃᓄᐃᑦᑐᑦ ᐅᖏᖅᑰᒪᔪᑦ ᐊᐅᓪᓚᐃᑦ ᑲᑐᔾᔨᑦᓗᑎᒃ ᐊᑎᓕᐅᖏᖅᑰᒪᔪᑦ ᑕᒪᑐᒥᖓ ᐃᓄᐃᑦ ᐊᑲᖏᑕᐅᓂᐊᖏᓂᖅᐳᓄᑦ ᐱᓯᒪᓕᕆᑕᐅᔪᖅᑐᓄᓪᓕ ᐊᖏᕈᕆᑦ ᐊᖏᕈᔪᖅᑰᓗᑎᒃ ᐱᓪᓗᑎᑦ ᐊᒻᒪᓗ ᕿᑐᕐᖏᖅᑐᖅᑎᑦ ᑕᒪᖏᑲ ᐃᓄᐃᑦ ᓄᓇᕗᑦ ᐅᓗᓇᐃᖅᑕᐅᓂᖅᐳᓪᓚᖅ;

ᐃᓐᓇ, ᑕᐃᒪᐃᓱᒪᑦ, ᐃᓕᓴᒪᕐᕆᐅᓗᑎᒃ ᑕᐃᒪᓐᓇᐅᐅᓂᐊᖏᓕᓂᖅᑕᖏᓪᓕ, ᐊᑐᐊᖅᐃᑦ ᒪᓕᒋᐊᓪᓕᓪᓗ ᐃᓗᐊᓵᑐᑦ ᐅᕐᓂ, ᐱᓇᓕᐊᖅᑎᑎᑦᑕ ᐃᒪᐓᓇᑦ ᐊᖏᕐᕈᑎᑦ:

ᐃᓕᖏᑐ 1 — ᑕᐃᔨᓯᖏᑦ

1.1 ᑐᑭᖏᑦ ᐅᖃᖅᑳᓯᒪᓂᖏᑦᑕ ᐊᓗᐊᐃᖅᑳᓯᓪᓚᑦ *ᓄᓇᕗᒻᒥ ᓄᓇᕆᑕᑎᓄᑦ ᐊᖏᕈᑕᑎᑦ* ᐊᓯᒪᓗ ᐊᑐᖅᑳᓪᓚᑦ ᐊᔪᕐᖏᓂᒃ ᑐᑭᖏᓂᒃ ᐃᓗᐊᓂᑦ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᕐᓂᖏᓂᒃᓄᑦ ᐱᖓᓕᓯᑕᐅᔪᓂᐊᖅᑐᓄᑦᓗ ᐊᖏᕈᑕᕐᓂᑦ, ᑐᑭᖏᑦ ᐊᔪᕐᓂᑦ ᐅᖃᖅᑳᓯᒪᔪᓂ ᐃᓕᖏᑐ 1 *ᓄᓇᕗᒻᒥ ᓄᓇᕆᑕᑎᓄᑦ ᐊᖏᕈᑕᑎᓂ.*

1.2 ᕿᓯᐊᓂ ᐃᓗᓕᖏᑦ ᑎᑎᕋᖅᑳᒪᔨᕆᐊᖃᖅᑕᕐᓂ, ᐱᔪᓀᑎᑕᓗᔪ ᐅᓇ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᕐᓂᖏᓂᒃ ᐱᖓᓕᓯᑕᐅᔪᓂᐊᖅᑐᓄᑦᓗ ᐊᖏᕈᑕᕐᓂᑦ :

> “ᐊᐅᓚᕆᑎᒃᑎᑎᕐᓂᒪᑦ ᑲᑎᒪᔨᕋᓱᒃᑲᓂᕐᓂ” ᑐᑭᖅᑲᖅᑐᑦ ᐊᐅᓚᕆᑎᒃᑎᑎᕐᓂᒪᑦ ᑲᑎᒪᔨᕋᓱᒃᑲᓂᕐᓂ, ᓇᓗᓇᐃᔨᑕᑎᐊᕆᒪᔪᖅ ᐊᑦᑕᓂ;”
>
> “ᐊᑕᐊᖅ ᓄᓇᓕᓂᖅ” ᑐᑭᖅᑲᖅᑐᖅ ᓄᓇᓱᓂ ᐊᑕᐊᑦ ᐊᑕᐅᓕᖕᒪᑦ ᐅᓄᑦᑐᓄᓕᓱᓂ ᑲᓇᑕ ᐅᓘᒪᔪᖅᑳᓴᓂ ᐅᕐᓂᓐᓇᓱᓂ ᐅᑎᖅᑕᐸᑦᑐᓄᓂ ᑎᓪᓗᐊᓄᓂ ᓴᓐᓴᐅᓕᒪᓕᕆᓂᓄᑦ, ᑎᑎᕋᖅᑳᒪᔪᑦ 2-1;
>
> “ᐊᐅᓚᕆᑎᒃᑎᑎᕐᓂᒪᑦ ᑲᑎᒪᔨᕋᓱᒃᑲᓂᕐᓂ” (ᓄᓇᒥᒃ ᑲᑐᔪᖅᑎᑎᕐᑐᑦ ᑲᑎᒪᔨᕋᓱᒃᑲᓂᕐᓂᖏᓂ) ᑐᑭᖅᑲᖅᑐᖅ ᐃᓄᐃᑦ / ᓴᖅᓪᓕᑦ ᑲᑎᒪᓕᖅ ᐅᓘᒪᔪᖅᑳᓴᓂ ᐅᕐᓂᓐᓇᓱᓂ ᐅᑎᖅᑕᐸᑦᑐᓄᓂ ᑎᓪᓗᐊᓄᓂ ᓴᓐᓴᐅᓕᒪᓕᕆᓂᓄᑦ ᐊᐅᓚᕆᑎᒃᑎᑎᕐᓂᑦ ᑲᑎᒪᔨᕋᓱᒃᑲᓂᕐᓂᑦ ᑕᐃᒪᓐᓇᐅᔪᓴᕆᑎᖅᑲᖅᑎᑕᐅᔪᑦ 9.3.2, 9.3.7 ᐊᑎᓪᓗ 8.4.11 *ᓄᓇᕗᒻᒥ ᓄᓇᕆᑕᑎᓄᑦ ᐊᖏᕈᑕᑎᓂ;*
>
> *“ᑲᓇᑕᒥ ᐅᓪᓛᕐᓚᓱᓂᖓᓪᓕᒪᑦ ᐱᕋᔭᕐᔪᐊᖅᑐᖅ”* ᑐᑭᖅᑲᖅᑐᖅ *ᑲᓇᑕᒥ ᐅᓪᓛᕐᓚᓱᓂᖓᓪᓕᒪᑦ ᐱᕋᔭᕐᔪᐊᖅᑐᖅ*, ᐱᕋᔭᕐᓂᑦ ᐊᑎᓪᓗ ᓇᒡᒐᐃᑐᓐᓇᖅ ᑲᔪᓯᑎᐊᑦᖁᔭᖅᑐᖅ ᒪᒡᑐᖅᑐᖅ ᐱᓯᒪᔪᑦ ᑲᓇᑕᒥ ᐅᓪᓛᕐᓚᓱᓂᖓᓪᓕᒪᑦ;
>
> “ᓄᓇᒡᓗᓂ ᐊᖏᕈᑕᑎᓄᑦ ᐱᔪᓐᓇᐅᑎᒡᓗᓂᒃ ᑲᑎᒪᔨᕋᓱᒃᑲᓂᕐᓂ” ᑐᑭᖅᑲᖅᑐᖅ ᓄᓇᒡᓗᓂ ᐊᖏᕈᑕᑎᓄᑦ ᐱᔪᓐᓇᐅᑎᒡᓗᓂᒃ ᑲᑎᒪᔨᕋᓱᒃᑲᓂᕐᓂ, ᑐᑭᒡᐅᖅᑲᓪᓚᕐᔪᖅ ᐊᑕᐃᓂ;
>
> “ᓄᓇᕗᒻᒥ ᐊᖏᕈᑕᑎᓄᑦ ᓄᓇᒥᐅᑎᑦᓯᓂᒃ ᑲᑎᒪᔨᕋᓱᒃᑲᓂᕐᓂ” ᑐᑭᖅᑲᖅᑐᖅ ᓄᓇᕗᒻᒥ ᓄᓇᕆᑕᑎᓄᑦ ᐊᖏᕈᑕᑎᓄᑦ ᓄᓇᒥᐅᑎᑦᓯᓂᒃ ᑲᑎᒪᔨᕋᓱᒃᑲᓂᕐᓂ, ᑐᑭᒡᐅᖅᑲᓪᓚᕐᔪᖅ ᐊᑕᐃᓂ;
>
> “ᓄᓇᕗᒻᒥ ᐊᖏᕈᑕᑎᓄᑦ ᐱᔪᓐᓇᐅᑎᒡᓗᓂᒃ ᑲᑎᒪᔨᕋᓱᒃᑲᓂᕐᓂ” ᐅᕐᓂᓐᓇᓱᓂ “ᓄᓇᕗᒻᒥ ᓄᓇᕆᑕᑎᓄᑦ ᐊᖏᕈᑕᑎᓄᑦ ᓄᓇᒥᐅᑎᑦᓯᓂᒃ ᑲᑎᒪᔨᕋᓱᒃᑲᓂᕐᓂ” ᑐᑭᖅᑲᖅᑐᖅ ᑲᑎᒪᔨᕋᓱᒃᑲᓂᕐᓂ ᓴᓇᐸᐅᔨᓱᓂ ᐊᐃᑲᖅᑐᖅᑲᓂᕐᓂᖏᓂᒃ ᐊᔪᕐᖏᓂ ᐅᖃᖅᑳᓯᒪᔪᑦ ᑲᑎᒪᓗ ᐊᑐᖅᑕᐅᓂᐊᕐᓂᖏᓂᒃ ᐊᑐᖅᑕᐅᓂᐊᕐᓂᖏᓂᒃ ᐊᑎᓪᓗ ᐊᑐᖅᑕᐅᓂᐊᕐᓂᖏᓂᒃ ᐊᖏᕈᑕᕐᓂᑦ ᐃᓄᐃᑦ ᓄᓇᕗᒻᒥ ᐊᖏᕈᑕᑎᓄᑦ ᐊᖏᕈᑕᕐᓂᑦ ᐊᑎᓪᓗ ᐊᑐᖅᑕᐅᓂᐊᕐᓂᖏᓂᒃ ᐊᖏᕈᑕᕐᓂᑦ, ᐃᓕᓴᐃᑦ ᐊᑎᓪᓗ ᕿᓯᐊᓂᓕᕆᓂᑦ ᐊᑲᑐᒥᔪᑦ ᓄᓇᒥᑦ;
>
> “ᐱᖓᓐᓇᓱᑐᖃᖅᑐᖅᑲᓂᕐᓂ ᐃᓂᖏᓂ ᐱᓕᒪᓐᓇᐅᔪᑦ ᐃᓄᖕᓄᓂ” ᑐᑭᖅᑲᖅᑐᖅ ᐃᓂᒋᔭᐅᔪᑦ ᐱᓕᒪᓐᓇᐅᔪᑦ ᐃᓄᐃᑦ ᐱᐅᔪᓂᒃᑯᑦ ᐅᑯᓇᓗᓂ ᐊᑐᖅᑕᐅᓂᐊᕐᓂᖏᓂᒃ 6.4.3 ᐊᑎᓪᓗ 6.4.4;
>
> *“ᑲᓇᑕᒥ ᐅᓪᓛᕐᓚᓱᓂᖓᓪᓕᒪᑦ ᐱᕋᔭᕐᔪᐊᖅᑐᖅ”* ᑐᑭᖅᑲᖅᑐᖅ *ᑲᓇᑕᒥ ᐅᓪᓛᕐᓚᓱᓂᖓᓪᓕᒪᑦ ᐱᕋᔭᕐᔪᐊᖅᑐᖅ*, ᓇᓗᓇᐃᓕᑎᐊᑦᖁᔭᖅᑐᖅ ᖃᐅᔨᓂ;
>
> “ᑲᓇᑕᒥ ᐅᓪᓛᕐᓚᓴᕐᔪᐊᓱᐅᑦ” ᑐᑭᖅᑲᖅᑐᖅ ᑲᓇᑕᒥ ᐅᓪᓛᕐᓚᓴᕐᔪᐊᓱᐅᑦ ᑕᐃᒪᓐᓇᐅᔪᓴᕆᑎᖅᑲᖅᑎᑕᐅᔪᑦ ᑲᓇᑕᒥᑦᑎᐊᓕ ᑲᑎᒪᔨᕋᓱᒃᑲᓂᕐᓂᖏᓂᒃ ᐊᑐᖅᑕᐅᓂᐊᕐᓂᖏᓂᒃ ᐊᑎᓪᓗ ᐃᓄᖕᓄᓂ ᐅᖃᐅᓯᐊᓱᐅᑦ ᐊᑕᐃᓂ

Ċᑲძᓇᖅᑐᑦ, ᑲᒪᓕᒋᓐᓂᓪ, ᑎᒥᖅᑯᑎᒋᑦ ᐅᖅᓯᓕᕐᓂᖅᓴᓂ ᐊᐅᓚᖅᓯᓕᕐᓂᑦ ᐱᓴᕆᖅᑲᖅᑐᑦ ᑲᓇᑕᒥ ᐅᒪᓕᖅᓯ

“ᐃᓄᐃᑦ ᐅᖃᐅᓯᖏᑦ” ᑐᑭᖃᖅᑐᖅ ᐃᓄᒃᑎᑐᑦ ᐅᕝᕙᓚᐅᖅᑐᑦ ᐃᓄᐃᕐᓇᖅᑐᑦ,
ᑕᐃᒪᕐᓇᐅᔪᓕᐊᖃᖅᑎᑦᓯᓗᑎᑦ;

“ᐃᓄᐃᑦ ᑎᒥᖓᑎ” ᑐᑭᖃᖅᑐᖅ ᓄᓇᕗᑦ ᑐᓴᖏᓐᓄᒃ ᑎᒥᖏᑲ ᐊᓪᒪᓗ ᐊᕐᓇᖅᑐᖃᓕᕆᔨ
ᐃᓄᐃᑦ ᑲᑐᔾᔨᖃᑎᒌᖏᑦ;

“ᐃᓄᐃᑦ ᖃᐅᔨᓴᓕᕐᖃᖅᖏᑦ” ᑐᑭᖃᖅᑐᖅ ᐅᑯᐊ ᐱᐅᓯᑐᖃᖅᖏᑦ, ᐊᓐᓇᐅᔨᖅ ᐊᓪᒪᓗ
ᑕᒪᕐᒥᒃ ᐃᓄᐃᑦ ᐊᑐᑎᖏᑦ, ᐅᑲᐱᓇᖏᑦ, ᐊᑐᖅᓯᒪᔭᖏᑦ, ᑕᐅᑦᑐᖏᑦ ᐊᓪᒪᓗ
ᖃᐅᔨᓴᖏᑦ ᐱᓕᓗᑦ ᐊᐅᑎᑎ, ᑕᒪᐅᑯᐊᓗ ᓄᓇ, ᐃᒪᖅ, ᓂᕆᔨᑎᑦ ᐊᓪᒪᓗ ᐃᓄᐃᑦ,
ᑕᒪᐅᖏᑲ ᐃᓄᐃᑦ ᐃᓕᓴᐅᕐᓂ ᐊᐅᑎᑦᑎᓂ;

“ᐃᓄᐃᑦ ᐳᓪᕋᑦᑎᓄᑦ ᑲᒪᔨᐅᕐᓴᑦ” ᑐᑭᖃᖅᑐᖅ ᐃᓄᒃ ᐅᕝᕙᓚᐅᖅᑐᑦ ᐃᓄᐃᑦ
ᑎᒥᖅᑎᖓᑎ ᑐᓂᓯᓯᒪᓴᑦ, ᐅᕝᕙᓚᐅᖅᑐᑦ ᑐᓂᓯᓯᒪᓴᒪᓴᑦ, ᐱᖅᑎᑎᓂᒃ ᐅᕝᕙᓚᐅᖅᑐᑦ
ᐱᔨᑦᑎᕋᐅᑎᓂᒃ ᐳᓪᕋᑦᑎᓄᑦ ᐃᓄᐃᑦ ᐃᓄᓕᕐᓂᐊᕐᓂᖅ ᐊᑐᐃᓐᓇᖅᑐᑦ;

“ᐃᓄᐃᑦ ᓄᓇᖅᑎᖓᑎᑦ” ᑐᑭᖃᖅᑐᖅ ᐃᓄᐃᑦ ᓄᓇᖅᑎᖓᑎ ᓇᓗᓇᐃᖅᑕᐅᑦᑎᐊᖅᓯᒪᓴᑦ
ᐃᓕᓴᐃᖅ 1.1.1 *ᓄᓇᕗᒻᒥ ᓄᓇᕆᔨᑎᓄᑦ ᐊᖏᕈᑎᑦ*;

“ᐊᐅᓚᑦᑎᐱᓕᕐᓴᑦ ᐸᖅᓇᐅᑎᓄ” ᑐᑭᖃᖅᑐᖅ ᐊᐅᓚᑦᑎᓂᒃ ᐸᖅᓇᐅᑎᖓᑎᒃ ᑲᑎᑎᓄᑦ
ᐃᓕᓴᐃᖅ 8.4.13 ᓄᓇᕗᒻᒥ *ᓄᓇᕆᔨᑎᓄᑦ ᐊᖏᕈᑎᑦ* ᐊᒪᓗ ᐃᓕᓴᐃᖅ 3 ᒐᕐᓄᓇᓂ ᐃᓄᐃᑦ ᐊᑐᖅᑕᐅᓂᐊᖅᓂᕐᓂᓄᑦ
ᐱᓱᑦᓕᓂᕐᕕᐊᓇᖅᑐᓄᑦᓗ ᐊᖏᕈᑎᑦ;

“ᑎᖏᒥᐊᑦ ᐅᑎᖅᑕᐸᒃᑐᐃᑦ ᐱᔪᓕᐅᓂᖏᓐᓄᑦ ᐱᓂᓄᖅ” ᑐᑭᖃᖅᑐᖅ *ᑎᖏᒥᐊᑦ
ᐅᑎᖅᑕᐸᒃᑐᐃᑦ ᐱᔪᓕᐅᓂᖏᓐᓄᑦ ᐱᓂᓄᖅ*, ᐊᓪᒪᓗ ᐊᑐᖅᑕᐅᓂᐊᖅᓂᕐᓂᓄᑦ;

“ᐅᑎᖅᑕᐸᓪᓗᓄᑦ ᑎᖏᒥᐊᓄᑦ ᓴᓕᐅᑎᓄᒃᑎᑦ” ᑐᑭᖃᖅᑐᖅ ᐅᑎᖅᑕᐸᓪᓗᓄᑦ ᑎᖏᒥᐊᓄᑦ
ᓴᓕᐅᑎᓄᒃᑎᑦ;

“ᐅᑎᖅᑕᐸᓪᓗᓄᑦ ᑎᖏᒥᐊᓄᑦ ᓴᓕᐅᑎᓄᒃᑎᑦ” ᑐᑭᖃᖅᑐᖅ ᐅᑎᖅᑕᐸᓪᓗᓄᑦ ᑎᖏᒥᐊᓄᑦ
ᓴᓕᐅᑎᓄᒃᑎᑦ ᓄᓇᕗᑦ ᐊᓪᒪᓗᐃᑦᑕᐅᓯᔪᓐᓇ ᐊᒋᔪᑦ *ᑎᖏᒥᐊᑦ ᐅᑎᖅᑕᐸᒃᑐᐃᑦ
ᐱᔪᓕᐅᓂᖏᓐᓄᑦ ᐱᓂᓄᖅ*;

“ᒥᓴᕐ” ᑐᑭᖃᖅᑐᖅ ᓴᖅᓪᓕᖅᑐᒃᑯᓂᓂ ᒥᓴᕐ ᐊᖅᑎᓕᕐᓚᓱᒃᑯᑦ ᐅᕝᕙᓚᐅᖅᑐᑦ ᒥᓴᕐ
ᑎᓕᐅᕐᓂᐊᕐᓇᖅᑐᑦ, ᑭᓯᐊᓂ ᐃᓕᓴᐅ ᐅᖃᖅᖃᓯᒪᔪᓂ;

“ᑲᓂᕐᑕ ᐃᓕᓴᖃᖅᑲᐃᑦ” ᑐᑭᖃᖅᑐᖅ ᑲᓂᕐᑕ ᐃᓕᓴᖃᖅᑲᐃᑦ ᓴᖅᑭᑐᐅᓯᒪᓕᕐᓂᑦ ᓄᓇᕗᑦ
ᐊᓪᒪᓗᐃᑦᑕᐅᓯᒪᓂᖓᓐᓄᒃ ᐊᒋᔪᑦ *ᑲᓂᕐᑕ ᐃᓕᓴᓕᓚᓂᖅᓱᑦ ᐱᓂᓄᖅ*;

“ᓄᓇᕗᑦ ᓄᓇᕐᓂᑎᓄᑦ ᐊᖏᕈᑎᑦ” ᑐᑭᓯᖅᑐᑦ ᓄᓇᕗᑦ ᓄᓇᕐᓂᑎᓄᑦ ᐊᖏᕈᑎᑦ, ᐊᑦᓯᓂ ᐅᓗᓇᐃᑦᑎᐊᖅᓯᒪᔪᖅ;

“ᓄᓇᕗᑦ ᐅᓗᓇᐃᖅᑕᐅᓯᒪᓂᖏᑦ” ᑐᑭᓯᖅᑐᖅ ᓄᓇᕗᑦ ᐅᓗᓇᐃᖅᑕᐅᓯᒪᓂᖏᑦ ᐅᓗᓇᐃᑦᑎᐊᖅᓯᒪᔪᑦ 1.1.1 ᒥᖁᓇᓂ ᓄᓇᕗᑦ ᓄᓇᕐᓂᑎᓄᑦ ᐊᖏᕈᑎᑦ;

“ᓄᓇᕗᑦ ᑐᓴᖏᕕᖅ ᑎᒥᖏᑕ” ᑐᑭᓯᖅᑐᖅ ᓄᓇᕗᑦ ᑐᓴᖏᕕᖅ ᑎᒥᖏᑕ;

ᓄᓇᕗᑦ ᓄᓇᕐᓂᑎᓄᑦ ᐊᖏᕈᑎᑦ ᑐᑭᓯᖅᑐᑦ ᐊᖏᕈᑎᓴᑦ ᐊᕐᕈᒋᓂᑦ ᐃᓄᐃᑦ ᓄᓇᕗᓪᓕ ᓄᓇᖏᖅᑐᑦ ᐊᒪᓚᓗ ᑯᐃᓐᓄᑦ ᐃᓄᐃᑦ ᐊᖏᕐᕕᒃᓴᓂᑦ ᐅᑭᐅᖅᑕᕐᓂᖅ ᓄᓇᕗᑦ ᓄᓇᕐᓂᑎᓄᑦ ᐊᖏᕈᑎᑦ S.C. 1993 c. 29;

“ᑲᓇᑕᒥ ᐅᒪᔪᖅᑲᐃᑦ” ᑐᑭᓯᖅᑐᖅ ᑲᓇᑕᒥ ᐅᒪᔪᖅᑲᐃᑦ, ᖃᐅᓪᓗ ᐅᓗᓇᐃᑦᑎᐊᖅᓯᒪᔪᖅ;

“ᐱᓇᓱᐊᖅᑎᑦᑎᓪᓕ” ᑐᑭᓯᖅᑐᖅ ᓄᓇᕗᑦ ᐅᓗᓇᐃᖅᑕᐅᓯᒪᓂᖏᑦ, ᑭᖑᖅᑐᖅᑕᐅᔪᑦ ᓄᓇᕗᑦ ᑐᓴᖏᕕᖅ ᐊᒪᓚᓗ ᐱᖏᓱᑦ ᐊᐃᑦᑐᖅᓯᒪᔪᓂ ᐃᓄᐃᑦ ᑲᑎᒪᔨᖅᑎᑦᑎᓂᑦ, ᐊᒪᓚᓗ ᐅᒃᑯᓕᖅᑐᓂᖏᓐᓄᑦ ᑲᑎᒪᓇᐅᑉ ᐃᓄᐃᑦ ᐱᖏᓱᑦ ᐅᓇᓱᐊᖅᑎᑦᑎᓂᑦ, ᐊᒪᓚᓗ “ᐱᓇᓱᐊᖅᑎᑦᑎᓂᑦ” ᑐᑭᓯᖅᑐᖅ ᓇᓕᐊᑐᐃᓐᓇᖅ ᐊᑐᓂ ᖃᐅᔨᓕ, ᐊᒪᓚᓗ “ᐱᓇᓱᐊᖅᑎᑦᑎᓂᑦ” ᑐᑭᓯᖅᑐᖅ ᓇᓕᐊᑐᐃᓐᓇᖅ ᐊᑐᓂ ᖃᐅᔨᓕ ᐊᑎᖅᑲᖅᑐᑦ;

“ᐊᐃᔨᑐᖅᓯᒪᔪᑦ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑎᓂᒃᑯᑦ” ᑐᑭᓯᖅᑐᖅ ᖃᑎᖅᑎᑦᐅᑦ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑎᓂᒃᑯᑦ, ᑭᓪᓕᓯᖅᑐᖅ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑎᓂᒃᑯᑦ, ᐅᖃᓕᓴᐅᓂ ᖃᑎᖅᑕᖏᑦ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑎᓂᒃᑯᑦ, ᑎᑎᕋᖅᓯᒪᓂᖏᓐᓄᑦ ᑕᐃᒪᓐᓇᐅᔾᔨᕆᐊᖅᑎᖅᑎᓕᔪᓄ;

“ᖃᑲᐅᔾᔨᖅᑎᑎᓂ” ᑐᑭᓯᖅᑐᖅ ᐃᓄᑦ ᐱᓇᓱᐊᖅᑎᑕᖅᓯᒪᔪᖅ ᖃᑲᐅᔾᔨᖅᓯᓂᐊᖅᑐᖅ ᐃᓗᐊᓂ ᑭᓱᓕᕐᕆᐊᓂ ᐅᖓᖅᑕᐅᓪᓗᓂ ᐅᑎᖅᑕᐸᑎᓗᓄᑦ ᑎᓪᓗᐊᓄᓯᒪᔪᓄ ᓴᓇᔨᓄᑦ;

“ᐱᓱᐊᑎᒃᓴᖅ” ᑐᑭᓯᖅᑐᖅ ᐅᓴᖅᐅᓴᖅ ᐊᒪᓚᓗ ᑭᖓᔪᖅᑐᑦ ᐃᓕᓴᕐᓂᑎᓄᑦ ᓄᓇᕗᑦ;

“ᐊᐃᔨᑐᖅᓯᒪᔪᑦ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑎᓂᒃᑯᑦ” ᑐᑭᓯᖅᑐᖅ ᐊᐃᔨᑐᖅᓯᒪᔪᑦ ᓄᓇᓕᓂ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑎᓂᒃᑯᑦ, ᐅᓗᓇᐃᑦᑎᐊᖅᓯᒪᔪᖅ ᖃᐅᔨᓕ;

“ᐊᐃᔨᑐᖅᓯᒪᔪᑦ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑎᓂᒃᑯᑦ” ᑐᑭᓯᖅᑐᖅ ᐊᐃᔨᑐᖅᓯᒪᔪᑦ ᓄᓇᓖᓂ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑎᓂᒃᑯᑦ, ᐅᓗᓇᐃᑦᑎᐊᖅᓯᒪᔪᖅ ᖃᐅᔨᓕ;

“ᐅᒪᔪᓂ ᐊᕐᕌᖅᑐᒥᓂᖅᑐᑦ ᓄᖅᓯᕐᕆᓂᖅᑐᓪᓗ ᐱᖁᓯᕐᔪᐊᖅ” ᑐᑭᓯᖅᑐᖅ *ᐅᒪᔪᓂ ᐊᕐᕌᖅᑐᒥᓂᖅᑐᑦ ᓄᖅᓯᕐᕆᓂᖅᑐᓪᓗ ᐱᖁᓯᕐᔪᐊᖅ* ᐅᓗᓇᐃᑦᑎᐊᖅᓯᒪᔪᖅ ᐊᒥᓂ;

“ᓂᕆᔨᓂᓄᑦ ᐃᓱᒪᓗᑎᐅᕐᓂᖅᑐᓄᑦ ᐱᖁᓯᖅ” ᑐᑭᓯᖅᑐᖅ ᓂᕆᔨᓂᓄᑦ ᐃᓱᒪᓗᑎᐅᕐᓂᖅᑐᓄᑦ ᐱᖁᓯᖅ, ᐱᖁᓯᖅᕿᓂᖅᑐᑦ ᐊᒪᓚᓗ ᐅᓕᐊᑐᐃᓐᓇᖅ ᒪᓕᑐᖅ ᐊᑐᑎᔨᖅ;

“ᑲᑎᓪᓗᑎᑦ ᐊᖓᔪᖅᓯᒪᓂᖅᑐᑦ” ᑐᑭᓯᖅᑐᖅ ᑲᑎᓪᓗᑎᑦ ᐊᖓᔪᖅᓯᒪᓂᖅᑐᑦ, ᐊᒥᓂ ᐅᓕᐊᑐᐃᓐᓇᖅ ᒪᓕᑐᖅ ᐊᑐᑎᔨᖅ;

“ᐅᑎᖅᑐᖅ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑎᓂᒃᑯᑦ” ᑐᑭᓯᖅᑐᖅ ᐅᑎᖅᑐᖅ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑎᓂᒃᑯᑦ ᐅᓗᓇᐃᑦᑎᐊᖅᓯᒪᔪᖅ 5.1.1 *ᓄᓇᕗᑦ ᓄᓇᕐᓂᑎᓄᑦ ᐊᖏᕈᑎᑦ*;

“ᐅᓪᓚᖅᑐᑦ” ᑐᑭᓯᖅᑐᖅ ᑭᓇᐅᔾᔨᖅ, ᒥᖁᐊᓗ ᑭᓇᐅᔨᓕᕆᔨᖅᑎᑎᒃᑲᓪᓚᕆᔨᓕᕆᔨᖅ ᐃᓕᓴᕐᓂᑎᓄᑦ, ᐃᓕᓴᖅᑐᑦ ᐅᖃᓕᓴᐅᓂ ᐊᑐᖅᑐᑦ ᑲᑎᒪᔨᖅ ᐅᖃᓕᓴᐅᓂ ᐅᑎᖅᑕᐸᑎᓗᓄᑦ ᑎᓪᓗᐊᓄᓯᒪᔪᓄ, ᐅᑯᐊᖏᓐᓄᑦᑐᑦ:

(a) ᐃᓄᒃ ᐅᖃᐅᓕᖅᑐᑦ ᐃᓕᓴᐃ ᖃᑦᑐᒍᑎᓴ ᐃᖅᑲᕋᖃᑎᒋᔭᖅᑲᑦ ᐃᓅᑦ;

(b) ᑭᓇᐅᐃᖅᓇᖅ ᐊᖏᔪᓇᔪᖅᓇᐅᑎᒋᖃᖅᓯᒪᔪᖅ ᐱᔪᖅᓇᕋᑦᓂᒃ ᐊᒋᔪᑦ 5.7.34 ᒋᒃᑯᓇᓂ ᓄᓇᕗᓱᒃ ᓄᓇᒋᑲᑎᓄᑦ ᐊᖏᑲᑎᕐ;

(c) ᐃᖅᒃᑲᓇᐃᔨᕋᑎ ᐅᖃᐅᓕᖅᑐᑦ ᑲᖅᑐᕋᖃᒋᐅᓯᒪᔪᖅ ᓴᓇᑎᒋᐅᔪᖅ ᑲᓇᑎᒋᐅᔪᖅ ᑲᓇᒋᒋᐅᔪᖅ ᐅᒪᔪᓕᕆᓚᕋᔪᐊᑯᑦ ᐅᖃᐅᓕᖅᑐᑦ ᐊᓯᖏᑦ ᑲᓕᒪᒃᑯᑦ ᐱᒋᔭᑎᒋᐅᔪᖅ ᐱᓕᑐᐃ ᐃᖅᒃᑲᓇᐃᔨᓂᒃ ᐅᖃᐅᓕᖅᑐᑦ ᑲᖅᑐᕋᖃᒋᐅᓯᒪᓂᒃ; ᐅᖃᐅᓕᖅᑐᑦ

(d) ᖃᐅᐅᔨᓴᖅᑎ.

“ᓂᕆᔨᑎᖃᖅᖄᑦ ᓄᓇᑦ ᐱᒻᒪᓚᐅᐅᓕᑦ ᐃᓄᖕᓄᑦ” ᑐᑭᖃᖅᖄᖅ ᓄᓇᐃᑦ ᐱᒻᒪᓚᐅᔪᑦ ᐃᓄᖕᓄᑦ ᓂᕆᔨᑎᖃᖅᒪᑕ ᐊᒻᒪᓗ ᐊᔪᖅᑐᓂᒃ ᐱᓕᕆᑎᖃᖅᖄᑦ ᓇᓗᓇᐃᖅᓯᒪᓕᕐᓗᑎᒃ ᐊᖏᑦᑕᑎ 6.4.3 ᐊᒻᒪᓗ 6.4.4.

ᐃᓕᓐᖏᑦ 2 — ᖃᐅᔨᒋᐊᕐᓇᖅ ᒪᓕᒃᓯᒪᔨᖏᓕᒫᓂᑦ

2.1 ᑐᖓᔪᓕᕐᓂᑦ

2.1.1 ᐅᓇ ᐊᔪᐅᓇᖅᑐᑎᖅ ᐃᓕᑎᒍᓯᖓᓂᑦ ᐊᕐᑕᒋᓕᓂᑦ ᐃᓇᐃᑦ ᓄᓇᕗᒻᒥ ᐊᓚᓗᓐ ᐅᒪᔨᓕᒋᓕᑦ ᐊᕐᑎᓕᒍᖓᑦᓯᓗ ᓄᓇᕗᑦ ᓇᓗᓇᖅᑕᐅᓯᒪᓂᖅᐹᖓ ᐅᓇ ᓯᓚᑎᓕᒍᖓᑦᓯᓗ ᑭᓯᓕᒍᖓᑦᓗ ᐅᒪᔨᓕᒋᓕᑦ ᑭᓯᓕᒍᖓᑦᓗ ᐊᓚᓗᓐ ᐃᓇᑦᓯᒥᖅᓄᑦ ᐱᓯᒪᔨᖅ. *ᐃᓄᐃᑦ ᖃᐅᔨᒪᔭᑐᖅᖏᓐᓂ* ᐃᓕᑎᒍᖓᑦ ᑭᓯᐅᓕᖅᐹᓂ ᖃᐅᔨᒋᐊᕐᓇᖅᑐᑦᓯᓐᓂ, ᐅᓇᓗ ᐊᑐᓕᐊᓕᓐ ᐊᓚᓗᓐ ᑕᒪᑐᒍᓐ ᐃᓄᐃᑦ ᐱᒃᖏᓕᓂᐅᔪᓂᐊᖅᑕᖓᑦ ᐃᓕᓴᕐᓂᑦᑐᓂᓐ ᐱᖓᓱᓂ, ᐃᓕᓴᕐᓂᑦ ᐊᐅᓚᑦ ᓄᓇᕗᑦ, ᐃᒪᐃᑦ ᐊᓚᓗᓐ ᑕᓕᐅᑦ ᓄᓇᕗᑦ ᐅᓇᓗᓇᖅᑕᐅᓯᒪᓂᖅᐹ.

2.1.2 ᖃᓇᑕᒥ ᐅᒪᔨᖃᖅᖃᓕᑦ ᐊᓚᓗᓐ ᐅᑎᖅᑕᕐᑕᑐᓄᑦ ᑎᓐᓕᐊᓄᑦ ᓴᓂᖅᐅᓯᒪᓕᖅᐹ ᐱᓚᒪᓇᐅᔨᑦ ᐃᑲᔪᑎᓐᓚᒥᑦ ᓄᓇᕗᒻᒥ ᓄᓇᕐᑎᑦ ᐃᓄᐃᓐᓄᑦ ᓴᓂᐅᓄᐊᖅᑐᑦ ᓄᓇᕗᑦ ᐅᓇᓗᓇᖅᑕᐅᓯᒪᓂᖅᐹ, ᖃᓇᑕᒥ ᐊᓚᓗᓐ ᓄᓇᕐᔪᐊᖅ. ᐊᐅᑕᑎᖅᑎᑎᒍᑕᐅᓂᐊᖅᑐᑦ ᐃᓄᐃᓂᑦ ᐊᓚᓗᓐ ᖃᓇᑕᒥ ᐅᒪᔨᖃᖅᖃᓕᑦ ᓴᓂᐅᓄᐊᖅᑐᑦ ᓄᓇᕗᒻᒥ *ᓄᓇᕗᓴᖅ ᓄᓇᑕᐅᑎᓄᑦ ᐊᖏᕈᑎᑎᓂᑦ* , ᐅᓇ ᐃᓄᐃᑦ ᐊᐃᑐᖅᑕᐅᓂᐊᓂᖏᓐᓄᑦ ᐱᓕᕆᓕᕐᑕᐅᔪᓂᓇᖅᑐᓂᓐᓗ ᐊᖏᕈᑎᑎᓂ, ᐊᖏᕈᖅᑕᐅᓯᒪᔪᖅ ᐊᐅᓕᕐᑎᓴᕐᑕᐅᓂᐊᖅᑐᓄ ᕿᖃᓇᐅᑎᓂ, *ᐃᓄᐃᑦ ᖃᐅᔨᒪᔭᑐᖅᖏᓐᓂ*, ᖃᐅᔨᓴᖅᑎᑎᒪᓕᕆᓂ ᖃᐅᔨᒋᐊᕐᑎᑕᐅᑦ ᐊᓚᓗᓐ , ᑭᓯᐊᓂ ᒪᓕᕐᑎᐊᖏᕐᕿᑕ *ᓄᓇᕗᒻᒥᖅ ᓄᓇᕈᑎᑎᓄᑦ ᐊᖏᕈᑎᑎᓂ, ᑎᓐᓕᐊᓂ ᐅᑎᖅᑕᕐᕿᑐᖓᓄᑦ ᐱᔪᕈᑕᐅᓂᐊᖅᑐᓄ, ᖃᓇᑕᒥ ᐅᒪᔨᖃᖅᖃᓕᑦ ᐱᔪᓐᓇᖅᑐᖅ, ᓴᓇᕐᔪᐊᓂ ᐃᓕᓴᕐᓂᑦᑐᓂᓐ ᐱᖓᓱᓂ ᐊᓚᓗᓐ ᐊᑐᓕᐊᓕᓐ ᒪᓕᒃᓯᒪᔨᖅ ᐊᓚᓗᓐ ᐊᓕᓐᓂᐊᓕᖅᑐᑦ ᐃᓄᐃᓐᓄᑦ

2.1.9 ᐊᑐᓕᕆᑕᐅᓗᓂ ᐅᓇ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᖏᓐᓂᒃᓄᑦ ᐱᓪᓕᓕᒋᑕᐅᔨᓴᖅᓯᐅᓗᑦᓗ ᐊᖓᔪᕐᓂᐊᑦ ᐃᓄᒃᔨᓂᕐᑎᐊᖏᓗᑎᒃ, ᓴᖅᔭᓕᔪᓴᖓᓂᕐᑎᐊᖅᑐᓂᒃ, ᐊᓚᒪᓗ ᒪᓕᓴᓗᑎᒃ ᐊᖓᔪᖅᑎᑎᕐᓂᐊᖅᓯᒪᓂᑎᒃ ᑭᒃᑯᐊ ᐱᓇᓱᐊᖅᑎᕐᓂᑦ.

2.2 ᐱᓇᓱᑎᑦ

2.2.1 ᐅᑯᐊ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᖏᓐᓂᒃᓄᑦ ᐱᓪᓕᓕᒋᑕᐅᔨᓴᖅᓯᐅᓗᑦᓗ ᐊᖓᔪᖅᑎᑦ, ᐃᓕᓴᕐᓂᑦ 2-1,3-1,6-1,15-1 ᐊᓚᒪᓗ ᐅᐃᒍᖅ I, ᒪᓕᒃᑐᖅ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᖏᓐᓂᒃᓄᑦ ᐱᓪᓕᓕᐊᔪᕐᑎᐊᒃᓴᓄᑦ ᐊᖓᔪᖅᑎᑎᕐᓂᑦ ᑕᐃᒪᓐᓇᐅᔭᕆᐊᖏᖅᑎᑕᐅᑐᒃ ᐃᓕᖓᑦ 8 ᐊᓚᒪᓗ 9 ᑭᒃᑯᐊ *ᓄᓇᕗᓚᒃ ᓄᓇᕈᕐᑎᓄᑦ ᐊᖓᔪᖅᑎᑦ* ᑭᒃᑯᓄᖓ ᐊᑐᓂ ᑲᓇᑕᒥ ᐅᒪᔨᖏᑦᑐᖅ ᐊᓚᒪᓗ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᖓᒥᐊᓄᑦ ᓴᓗᖏᐅᓂᒪᓕᒃᑲ ᓇᓗᓇᐃᖅᓯᒪᔪᑦ 2-1 ᑕᕐᓂᐅᒥ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᖏᓐᓂᒃᓄᑦ ᐱᓪᓕᓕᒋᑕᐅᔨᓴᖅᓯᐅᓗᑦᓗ ᐊᖓᔪᖅᑎᑦ.

2.2.2 ᑕᕐᓂᖓᓕ ᐊᑐᑎᖃᓕᖕᒪᓂᒥ ᐅᑦᓗᒥᑦ, ᐅᐃᒍᖅ ᐊᑐᑎᖃᓄᖕᖏᑕᖅ ᓇᑦᓕᐊᖕᓇᑐᐃᓄᖅ ᑲᓇᑕᒥ ᐅᓪᓗᕈᓘᑐᑦ ᐅᕐᓂᓕᕆᓂᖅ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᖓᒥᐊᓄᑦ ᓴᓗᖏᐅᓂᒪᓕᒃᑲ ᑭᓱᓂᒥ ᓂᖓᓕᒫᓂᖅ ᐅᓪᓗᕈᓘᑐᑦ ᐅᓪᓗᕈᓘᑐᑦ, ᕕᕐᓂᐅᒃ ᐃᓄᐃᑦ ᐃᓕᓴᕐᓂᑦ ᐊᑐᓂ ᑲᓇᑕᒥ ᐊᖓᔪᖅᑎᑦ ᐱᓪᓕᓕᒋᑕᐅᔨᓴᖅᓯᐅᓗᑦᓗ ᐊᖓᔪᖅᑎᑦ.

2.2.3 ᐅᓇ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᖏᓐᓂᒃᓄᑦ ᐱᓪᓕᓕᒋᑕᐅᔨᓴᖅᓯᐅᓗᑦᓗ ᐊᖓᔪᖅᑎᑦ ᐊᑐᑎᖃᓕᖕᒪᖑᔪᖅᑐᑦ ᐊᓚᒪᓗ ᐃᓇᖓᓴᖅᑕᐅᓗᑦᓗ 2008 ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᖏᓐᓂᒃᓄᑦ ᐱᓪᓕᓕᒋᑕᐅᔨᓴᖅᓯᐅᓗᑦᓗ ᐊᖓᔪᖅᑎᑦ ᐊᑎᓕᐅᖅᑕᐅᓗᑦᓗᖅᑐᖅ ᒥᓵᖅᑐᑦ ᐊᓚᒪᓗ ᐊᖓᔪᖅᑎᑕᐅᓗᑦᓗᖅᑐᖅ ᑭᓇᐅᔭᓂᒃ ᐃᓄᐃᑦ ᐊᑐᑎᖃᓄᖕᖏᑕᖅ ᐊᑐᓂ

2.4.2 ᐅᓇ ᐊᖏᑭᐊᑕᑎᓵᖅ ᐅᕐᓂᔨᓐᓂᒃ ᐃᓕᑦᑎᐊᑕᑎᓵᖅ ᐅᐃᒍᒧᑦ ᐅᕐᓂᔨᓐᓂᒃ ᑐᓴᖏᓵᒋᓐᓇᑦ ᐱᓪᓗᒍ ᓇᓪᓕᐊᒃ ᑲᓇᕐᑕ ᐅᒪᖏᑦᑐᙳᑦ ᐅᕐᓂᔨᓐᓂᒃ ᐅᑎᖅᑕᕐᑐᓇᓐᓂᒃ ᑎᓴᒥᐊᓂᒃ ᓴᓗᐅᓕᕆᔨᒪᔭᒃ ᑎᑎᖅᑭᕆᔨᓗᑎᒃ ᐊᖏᖅᑕᐅᓕᕆᓪᓕᐊᓪᓕᐊᑦᑎᓪᓗᒍ, ᓄᓇᕗᑦ ᑐᓴᖏᔭᒃ ᑎᒥᖓ, ᐊᒻᒪᓗ ᐊᒃᑐᑎᓵᑦ ᐊᕐᕕᒃᑐᖏᓚᐅᓂᒃ ᐃᓄᓐᓂᒃ ᑐᑭᓯᓇᑎᑎᒃ.

2.5 ᐊᓯᖏᑦ ᖃᓄᑐᐃᓐᓇᖅ ᒪᓕᒃᓯᒪᔭᕆᐊᓕᕐᓂᐅᕐᖅ

2.5.1 ᐅᑯᐊ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐱᓵᓕᓕᕐᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕ ᐊᖏᕐᕈᑎᑦ ᒪᓕᓴᕐᒪᔪᑦ ᐊᖏᕐᕈᑎᑦ ᐊᖑᓪᓚᑕ ᒪᓕᒃᑕᐅᔨᓐᓇᓐᓂᒃ ᐊᖏᕐᕈᑎᑦ.

2.5.2 ᐅᓇ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐱᓵᓕᓕᕐᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕ ᐊᖏᕐᕈᑎᑦ ᐊᐅᓪᓚᑕᐅᔨᖅ ᒪᓕᒃᖏᕐᒪᔪᑦ ᓄᓇᕗᑦ ᐊᒻᒪᓗ ᑲᓇᑕ, ᐊᑐᑎᖅᑲᓪᓗᒍᒃ.

2.5.3 ᐅᓇ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐱᓵᓕᓕᕐᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕ ᐊᖏᕐᕈᑎᑦ ᐃᓪᓕᖏᓪᖏᓕᖅ ***ᓄᓇᕗᒻᒥᐅ ᓄᓇᕆᔨᕈᑎᓄᑦ ᐊᖏᕐᕈᑎᑦ***, ᐊᒻᒪᓗ ᐅᓇ ᓄᓇᕆᔨᕈᑎᓄᑦ ᐊᖏᕐᕈᑎᑦᑕᐅᖏᓪᓕᐊᓕᖅᑐᓂᒃ ᐅᕐᓂᔨᓐᓂᒃ ᐊᖏᕐᕈᑎᑦᑕᐅᖏᓪᓕᐊᓕᖅᑐᓂᒃ ᓄᓇᕗᑦ ᐱᓕᕐᓂᒃ ᐃᓄᐃᑦ 35 ***ᑲᓇᑕᐅᔪ ᐱᖁᔭᕐᔪᐊᖑᓚᓂ, 1982***.

2.5.4 ᐱᓕᕐᑲᓪᖏᓕᖅᑐᓂᒃ ᐅᕐᓂ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐱᓵᓕᓕᕐᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕ ᐊᖏᕐᕈᑎᑦ ᑐᑭᓯᐊᓱᐅᓚᓕᖏᓪᓕᐊᓪᖏᓕᖅᑐᓂᒃ ᑕᐃᒪᓗ ᐊᑐᑕᐅᓚᓕᓗᓂᓂ ᐅᕐᓂᔨᓐᓂᒃ ᐊᖏᕐᕈᑎᑦᑕᐅᖏᓪᓕᐊᓕᖅᑐᓂᒃ ᐃᓄᐃᑦ ᐱᓕᕐᓂᒃ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐃᓄᐃᑦ.

2.5.5 ᐅᓇ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐱᓵᓕᓕᕐᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕ ᐊᖏᕐᕈᑎᑦ ᐊᒻᒪᓗ ᑕᒪᕐᒥᒃ ᑐᑭᖏᕐᕈᑎᑦ ᑕᐃᔨᐅᖅᑲᑕᕐᓂᐅᕐᒪᑐᒥᒃ ᒪᓕᒃᑎᐊᕐᓗᑎᒃ ᐅᕐᓂᔨᓐᓂᒃ ᐊᑲᓇᕐᑕᐅᓚᓕᖏᓪᓕᐊᓪᖏᓕᖅᑐᓂᒃ ᐊᐅᒋᓂ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐱᓵᓕᓕᕐᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕ ᐊᖏᕐᕈᑎᑦ ***ᐊᒻᒪᓗ ᓄᓇᕗᒻᒥᐅ ᓄᓇᕆᔨᕈᑎᓄᑦ ᐊᖏᕐᕈᑎᑦ***, ᐅᑯᐊ ***ᓄᓇᕗᒻᒥᐅ ᓄᓇᕆᔨᕈᑎᓄᑦ ᐊᖏᕐᕈᑎᑦ*** ᓴᓪᓚᕐᓂᖏᓪᓕᐊᓪᖏᓕᖅᑐᓂᒃ ᒪᓕᒃᖏᕐᒪᔪᑦ ᐊᒻᒪᓗ ᐊᐅᓪᓚᑕᐅᓚᓐᓇᓗᓂ.

2.5.6 ᑐᓕᒧᐊᔪᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐅᕐᓂᔨᓐᓂᒃ ᐱᓴᖅᓯᒪᓕᑎᒃ ᑕᒪᑐᒪᖓᑦ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐱᓵᓕᓕᕐᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕ ᐊᖏᕐᕈᑎᑦ ᑐᓕᓕᕐᑕᐅᓚᐅᓂᒃ ᑕᒪᑐᒧᖓᑦ ***ᓄᓇᕗᒻᒥᐅ ᓄᓇᕆᔨᕈᑎᓄᑦ ᐊᖏᕐᕈᑎᑦ***.

2.5.7 ᐅᓇ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐱᓵᓕᓕᕐᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕ ᐊᖏᕐᕈᑎᑦ ᑕᒪᐃᓪᓗᒃ ᐊᖏᕐᕈᑎᑦᑕᐅᖏᓪᓕᐊᓕᖅᑐᓂᒃ ᐊᖏᕐᕈᑎᑦ ᐊᑦᑕᒍ 5.7.18 ***ᓄᓇᕗᒻᒥᐅ ᓄᓇᕆᔨᕈᑎᓄᑦ ᐊᖏᕐᕈᑎᑦ***, ᐊᒻᒪᓗ ᐅᓇ ᓴᖅᑭᑎᓯᒪᔨᖏᓪᓕᐊᓕᖅᑐᓂᒃ ᐊᔪᓕᖅᑎᑎᑎᓯᒪᓂᖏᑦ ᐃᓄᐃᑦ ᐊᑐᕐᓇᓕᓂᑦ ᑎᑎᖅᑭᕆᔨᓗᑎᒃ ***ᓄᓇᕗᒻᒥᐅ ᓄᓇᕆᔨᕈᑎᓄᑦ ᐊᖏᕐᕈᑎᑦ***.

2.5.8 ᐅᓇ ᐃᓴᖅᑕᐅᓯᒪᔨᖓ ᑕᒪᑐᒪᖓ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐱᓵᓕᓕᕐᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕ ᐊᖏᕐᕈᑎᑦ ᐅᓪᓚᕐᓂᖏᓪᓕᐊᓪᖏᓕᖅᑐᓂᒃ ᐃᓄᐃᑦ 8-3 ᒥᒃᑯᐃᓐᓇᑦ ***ᓄᓇᕗᒻᒥᐅ ᓄᓇᕆᔨᕈᑎᓄᑦ ᐊᖏᕐᕈᑎᑦ*** ᑐᓂᓯᒪᔨᑕᐅᖅᑎᑐᐅᓂᐊᓕᕐᓂᑐᑦ ᐅᓱᒪᓐᓂᑐᓄᓂᒃ ᐃᓕᓴᕐᓂᑐᓄᓂᒃ ᑕᒪᑐᒪᖓ ᐅᕐᓂᔨᓐᓂᒃ ᐅᓪᓚᕐᓂᖏᓪᓕᐊᓪᖏᓕᖅᑐᓂᒃ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐱᓵᓕᓕᕐᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕ ᐊᖏᕐᕈᑎᑦ.

2.5.9 ᐅᑯᐊ ᐱᖁᔭᒃ, ᐃᓕᓴᕐᓂ ᐊᒻᒪᓗ ᐅᐃᒍᑦ ᒪᓕᒃᑐᒃ ᑕᒪᑐᒪᖓ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐱᓵᓕᓕᕐᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕ ᐊᖏᕐᕈᑎᑦ ᐅᖃᖅᑐᓕᖅᑐᓂᒃ ᐊᑕᐅᓂᒃᑯᓂ ᐊᒻᒪᓗ ᑐᓂᒃᑯᑦ. ᐅᑯᐊ ᐅᖃᖅᑲᓇᐃᓐᓇᐅᑎᓂ, ᑐᓴᖅᑭᔨᑎᓂ, ᐊᒻᒪᓗ ᑐᓂᒃᑯᓪᓗᒍᓐᓇᓂ ᑕᒪᑐᒪᖓ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐱᓵᓕᓕᕐᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕ

ᐊᖏᑭᑐᑦ ᐃᑲᔪᑎᐅᓚᐅᖅᑐᑦ ᑐᑭᓯᐅᑭᑎᑦᓯᓚᑦ ᐃᓄᐃᑦ ᐊᐅᖅᑕᐅᓂᐊᓂᖏᓐᓄᑦ ᐱᓕᕆᓂᑕᐅᔪᓐᓇᖅᑐᓄᓯᓗ ᐊᖏᑭᑎᑦᓂᓐᓂ.

2.5.10 ᐅᓇ ᐃᓄᐃᑦ ᐊᐅᖅᑕᐅᓂᐊᖅᓂᖏᓐᓄᑦ ᐱᓕᕆᓂᑕᐅᔪᓐᓇᖅᑐᓄᓯᓗ ᐊᖏᑭᑐᑦ ᒪᓕᒃᑐᖅ ᑕᒪᐃᓐᓂᒃ ᐊᖏᑭᑎᓂᒃ ᐊᖓᒍᑦᓂ ᐱᓇᔪᐊᖅᑎᑦᑎᑦ ᐊᓚᒪᓗ ᐅᖅᑲᓕᑎᐊᖅᑐᓇᒪᐊᓕᓂ ᐅᖓᓯᓚᐅᑦᓂ ᑎᑎᕋᖅᑭᒪᓂᖅᑐᑦ, ᐊᔪᖏᑎᑦᑎᓴᐅᑕᐅᓂᐊᓐᖏᓂᑐᑦ ᐊᑭᓕᔪᐅᑕᐅᓯᓪᓚᖅᑭᒪᓂᖅᑐᑦ ᐅᖓᓯᓚᐅᑦᓂ ᖃᓄᐃᓕᓂᖏᓐᓄᑦ ᐊᖓᒍᑦᓂ ᐊᐅᖅᑕᐅᓂᐊᖅᓂᖏᓐᓄᑦ ᐱᓕᕆᓂᑕᐅᔪᓐᓇᖅᑐᓄᓯᓗ ᐊᖏᑭᑐᑦ ᑭᓱᐊᓂᐅ ᐅᖅᑲᖅᓱᒪᔭᐅ.

2.5.11 ᑕᐃᒪᓐᓇᐅᑎᑦᑎᓇᓱᐊᓂᖏᓐᓄᑦ ᐅᑯᐊᓗ ᓇᓗᓇᖅᓱᑎᑦᑎᓱᒪᓗᓂᑎᒃ ᑕᒪᑐᒧᖓ ᐃᓄᐃᑦ ᐊᐅᖅᑕᐅᓂᐊᖅᓂᖏᓐᓄᑦ ᐱᓕᕆᓂᑕᐅᔪᓐᓇᖅᑐᓄᓯᓗ ᐊᖏᑭᑐᑦ ᑐᑭᓯᐅᑭᑎᑦᓯᓚᑦ ᑭᓇᐅᔪᑎᑦᑎᓱᒪᓗᓂᑎᒃ ᓇᒪᒋᔭᖏᑦᓂᓐᓂᒃ ᑲᑎᕋᖅᑭᒃᑲᓐᓂᒃ ᐃᓄᐃᑦ ᐊᐅᖅᑕᐅᓂᐊᖅᓂᖏᓐᓄᑦ ᐱᓕᕆᓂᑕᐅᔪᓐᓇᖅᑐᓄᓯᓗ ᐊᖏᑭᑐᑦ ᑭᓱᐊᓂᐅ ᐅᖅᑲᖅᓱᒪᔭᐅ.

2.5.12 ᐱᑕᖅᓯᐅᔪᖅᑐᓄᑦ ᐃᓄᐃᖅᑐᑦ, ᖃᓄᓗᓇᖅᑐᑦ ᐊᓚᒪᓗ ᐅᐃᕐᕆᖅᑎᑦ ᑭᐅᔪᓂᖓ ᐃᓄᐃᑦ ᐊᐅᖅᑕᐅᓂᐊᖅᓂᖏᓐᓄᑦ ᐱᓕᕆᓂᑕᐅᔪᓐᓇᖅᑐᓄᓯᓗ ᐊᖏᑭᑐᑦ. ᐅᔪᐊ ᖃᓄᓗᓇᖅᑎᑐᖅᑐᑦ ᐊᓚᒪᓗ ᐅᐃᕐᕆᖅᑎᑦᑎᖅᑐᑦ ᐊᑐᑎᖃᓄᖅᓴᐅᓂᐊᖅᓗᑎᒃ. ᐅᔪᐊ ᐱᓇᔪᐊᖅᑎᑦᑎᓂᒃ ᐊᖏᕐᖃᑎᒃᓱᓗᑎᒃ ᐃᓄᐃᖅᑎᑐᖅᑐᑦ ᐊᑐᑎᖃᓄᖅᓴᐅᓂᐊᖅᓗᑎᑦ ᐊᑐᑎᖃᓄᖅᓴᐅᓂᐊᖅᓗᑎᑦ.

2.5.13 ᐱᑕᖃᓂᖏᓐᓄᑦ ᐅᖓᓂ ᐃᓄᐃᑦ ᐊᐅᖅᑕᐅᓂᐊᓐᓂᖏᓐᓄᑦ ᐱᓕᕆᓂᑕᐅᔪᓐᓇᖅᑐᓄᓯᓗ ᐊᖏᑭᑐᑦ ᐊᐅᖅᑕᐃᓇᓴᓐᓂᖏᓐᓄᒃ ᐅᓴᓂᐅᑎᒃ ᐅᖓᓯᓚᐅᑦᓂ ᐃᓄᐃᑦ ᐃᓕᐅᔪᓂᐅᑎᖏᓐᓄᑦ ᐊᓚᒪᓗ ᐃᑲᔪᓯᐊᖅᑕᑭᓐᓄᓕᖏᓐᓄᑦ ᐱᓕᓀᑎᑦᑎᔪᐅᓂᐊᖅᓗᓂᑦ ᓄᓇᕗᒻᒥ ᓄᓇᖃᖅᑐᖅᓯᓂ, ᐃᓄᓐᓂᑦ ᐅᖓᓯᓚᐅᑦᓂ ᐅᖓᓯᓚᐅᑦᓂ ᐅᖓᓯᓚᐅᑦᓂ

2.5.20 ᐱᑕᖃᓐᓇᕐᑕᑐᖅ ᐱᓇᓴᐅᑎᖃᑎᒋᑦ ᓄᖅᑕᕐᑕᑎᓂᖅ ᓇᑦᑕᐊᓐᓂᑐᐃᓇᖅ ᐱᒃᓴᒪᕐᓂᒃ ᐅᕙᓪᓕᐊᑦ ᐊᑐᓵᓚᖅᑐᑦ ᐊᒡᒍᑦ ᐅᒪ ᐃᓇᐃᑦ ᐊᑐᖅᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐱᓯᑦᑕᑕᐅᔪᓇᖅᑐᓇᑦᑕ ᐊᖏᕐᕈᑎᑦ ᐃᓇᖅᑎᑦ ᐅᕙᓪᓕᐊᑦ ᐅᕙᓪᓕᐊᓂᖅ ᑎᑎᐅᔪᑦ ᐅᓇᓗᐃᖅᓭᒪᔪᓂ ᐃᓇᐃᑦ ᐊᑐᖅᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐱᓯᑦᑕᑕᐅᔪᓇᖅᑐᓇᑦᑕ ᐊᖏᕐᕈᑎᑦ ᐱᒃᓴᒪᕐᓂᖏᓐᓄᑦ ᐱᓯᑦᑕᑕᐅᔪᓇᖅᑐᓇᑦᑕ ᐊᑐᖅᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐱᓯᑦᑕᑕᐅᔪᓇᖅᑐᓇᑦᑕ ᐊᖏᕐᕈᑎᑦ ᐱᒃᓴᒪ

(a) ᓇᓗᒪᓕᒍ ᐊᓗᒪᓕ ᐱᐅᓯᑐᖃᖏᓐᓄᓚᒍ ᖃᑎᖅᑕᐊᖃᑎᒌᒍᓯᓪᓚᒍ ᑐᑭᓯᑕᐊᖃᑎᒌᒍᓯᓪᓚᒍ, ᐅᑯᐊᓕ ᐱᓕᕆᓂᖏᓪᓕᓕᒫᓂ ᑐᓴᖅᑎᑐᖃᑎᒌᒍᓯᓪᓚᒍ ᑕᐅᓪᓗᖅᑎᓪᓗᒍ ᖃᓄᖅᓱᓕᐅᕙᓪᓕᐅᖏᓐᓄᑦ ᐃᓕᑐᐃᓪᓕᐃᑦ ᐊᓗᒪᓕ ᐊᓯᖏᓐᓄᑦ ᐃᓕᓴᐃᑦ;

(b) ᐃᓕᒃᑎᑐᓪᓕᐅᓚᒍᑎᒃ ᐊᓗᒪᓕ ᑐᑭᓯᓇᖅᓯᑎᑕᐅᓚᒍᑎᒃ ᐊᑐᑎᖃᖅᖄᑦ ᑎᑎᖃᖅᑦ, ᑕᐃᒪᓐᓇᕕᒃᑲᐊᖃᖅᖄᖅᓱᖅ; ᐊᓗᒪᓕ

(c) ᑭᓯᐊᓂ ᐃᓯᒪᑦᐊᓇᓯᒪᒃᐸᑦ ᐊᐅᓪᓕᑎᖃᑎᒋᐊᓂᔪᓂᑦ ᑲᑎᒪᔨᕐᓯᓪᑦ, ᑐᑭᓯᓇᖅᓯᑎᑕᐅᓚᒍᑎᒃ ᐃᓕᓴᐅᑎᓕᕐᓂᖅᓯᓂᖏᓐᓄᑦ ᑭᑭᑎᑕᐅᑕᐅᖅᓴᓪᓕᐊᔪᓂᒃ ᖃᖅᑎᓪᓚᒍᑦ ᐊᓗᒪᓕ ᐃᓕᒃᑎᑐᓂᑦ.

2.7 ᐋᖅᑭᑎᐊᖅᓚᑦᑕ ᐊᑐᕆᐊᖅᑲᖅᑕᑎᒃ

2.7.1 ᐅᑯᐊ ᐱᓇᓯᐊᖅᑎᑎᑦ ᐱᓕᕆᓂᐊᖅᑐᑦ ᑕᒪᐃᓂᒃ ᖃᓇᑐᐃᓪᓇᖅ ᐊᒃᑐᐃᓯᒪᓂᖏᓪᓄᓕᑦ ᐱᑦᓚᑦ ᑐᓐᖏᓪᓕᐊᔪᖏᓪᓄᑦ ᑕᒪᑐᒍᖓ ᐃᓕᓴᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᖅᓂᖏᓪᓄᓕᑦ ᐱᓪᓚᕆᓕᕐᓂᐅᖅᑐᓚᒍᑦᓚᓪ ᐊᖏᕐᓂᒃ. ᑐᓐᖏᓪᓕᐊᔪᖏᓪᓄᑦ ᐱᓪᓚᕆᓕᕐᓂᐅᖅᑐᓚᒍᑦᓚᓪ ᐊᖏᕐᓂᒃ.

2.8 ᑐᓴᖅᑎᑎᓂᓕᖅ

2.8.1 ᑭᓯᐊᓂ ᐊᓯᐊᔪᑦ ᐅᖅᑲᖅᑕᐅᓯᒪᓐᖏᑎᑎᓪᓚᒍ ᐊᐋᖏᑎᑕᐅᓚᒍᑎᒃ ᐅᕐᓕᔭᓐᓂᑦ ᐱᓴᕆᐊᖅᑲᖅᑎᑕᐅᓚᒍᑎᒃ, ᓇᑦᓂᐊᖓᑐᐃᓐᓇᖅ ᑐᓴᖅᑎᑎᕐᓴᐅᖅᑲᖅᓚᒍᑎᒃ, ᑐᓴᖅᑎᑎᓂᓕᓂᖅ ᐅᕐᓕᔭᓐᓂᑦ ᐱᐋᒃᓐᖏᓪᓄᓕᑦ ᐅᕐᓕᔭᓐᓂᑦ ᑐᓂᓴᐅᔪᒪᓕᓚᓕᑦ ᐱᓇᓯᐊᖅᑐᖃᑎᒌᒍᓯᓪᓚᒍᑦ ᐊᑭᒍᑦ ᑕᒪᑐᒍᖓ ᐃᓕᓐᓇ ᐊᐅᓪᓕᑎᖃᑎᒌᒍᓯᓪᓚᒍᑦ ᐱᓪᓚᕆᓕᕐᓂᐅᖅᑐᓚᒍᑦᓚᓪ ᐊᖏᕐᓂᒃ. ᐃᓕᓐᓇ ᐃᓕᓐᓇᖅᑐᑦ: a) ᓇᖏᑦᓂᖅ ᐋᑦᓚᑦᓂᑦ ᐊᐱᖓᓚᑦᓂᑦ (ᑕᐃᒪᓚᒍ ᐱᖅᐅᒍᓂ ᑕᐃᑲᓂᓂ ᑕᐃᒪ ᑐᓐᓂᑦᑯᑎᑦᓴᑦ); ᐅᕐᓕᔭᓐᓂᑦ b) ᐊᐅᑦᓭᑎᑦᓚᒍ ᑎᑎᖅᑲᑦᑯᖏᑦ ᐊᑎᓪᓕᐅᖅᑕᐅᓯᒪᓐᖏᖅᑎᖅᓚᓂ, ᐊᑦᐸᖏᓐᓇ ᐅᑎᖅᑎᑦᑎᓂᐊᐊᖏᑦ ᐱᖅᐅᒍᓂ (ᒐᑯᐊ ᓇᓚᓇᐸᑐᑦ ᐱᖅᐅᓂᖏᓪᓂᑲ ᑐᓴᖅᑎᑦᑎᓐᓄᕐᓭᑦ 7 ᐱᓇᓯᐊᕈᓯᐅᑦ ᐃᓚᐊᓂ ᑕᐃᑲᓂ ᑎᑎᖅᑲᑦᑯᖏᑦᓯᓚᒍᑦ ᐊᐅᑦᓭᑎᑕᐅᒍᓂ; ᐅᕐᓕᔭᓐᓂᑲ c) ᐊᐅᑦᓭᑎᑦᓚᓂᑦ ᓯᑲᕆᑯᑦ ᑕᖏᓐᓄᓂ ᐱᓚᑎᑦ (ᑕᐃᒪᓚᒍ ᖃᐅᔨᒪᓇᖅᓯᓂᐊᖅᑐᖅ ᐱᖅᐅᓂᖏᓪᓂᑲ ᐅᑦᓚᑦ ᐱᖅᑕᒃᓂᑎᓂ ᐊᐅᑦᓭᑎᑕᐅᓭᐅᖅᑎᑦᓚᓚᒍ ᓯᑲᕆᑯᑯᖅᑐᑦ); ᐅᕐᓕᔭᓐᓂᑦ d) ᖃᓄᖅᓱᓕᐅᕙᓪᓕᐅᖏᓐᓄᑦ, ᐅᖅᑲᓭᒪᖅᑕᐅᓯᒪᓕᓂᖏᓐᓄᑲ ᓇᓚᓇᐃᓕᖅᑐᖅ (ᑕᐃᒪᓚᒍ ᖃᐅᔨᒪᓇᖅᓯᓂᐊᖅᑐᖅ ᐅᑯᐊ ᑐᓴᖅᑎᑦᑎᓐᓄᕐᓭᑦᑎᑎᑦ ᐱᖅᐅᖅᑲᑦ ᐱᐅᓪᓕᐊᖅᑲᖅᑐᑦ).

2.8.2 ᑐᓴᖅᑎᑦᓯᐊᑎᑦ ᐊᐅᑦᓭᑎᑦᑕᐅᓂᐊᖅᑐᑦ ᐅᑯᓇᖏᓐ ᑐᖃᖅᑎᓇᑦ ᐅᖅᑲᑦᓚᒃᓇᐅᖅᑲᓚᒍᑎᖃᖅᑕᑎᒃ, ᑎᑎᕆᔨᓇᐅᓚᒍᑎᒃ ᑎᑎᖅᑲᑯᑯᓇᖏᑦ, ᐊᐅᑦᓭᑎᑦᓂᑎᖏᓇᐅᓚᒍᑎᒃ ᓯᒃᓚᓯᑯᑦ ᐅᕐᓕᔭᓐᓂᑦ ᖅᑲᓚᐅᖅᑲᑯᑦ, ᑭᓯᐊᓂ ᑕᐃᒪᓐᓇᖅᑐᑦᓚᓪᓚᒍᑎᖃᖅᑕᐅᓂᐊᖅᓂᖏᓪᓄᓕᑦ ᐱᐋᒃᓐᖏᓪᓄᓕᑦ ᐊᖏᕐᓂᒃ ᐅᕐᓕᔭᓐᓂᑦ ᐅᖅᑲᖅᑕᐅᓯᒪᓕᓂᖅ ᐱᓇᓯᐊᖅᑐᖃᑎᒌᒍᓯᓪᓚᒍᑦ ᑎᑎᖅᑲᑦᑯᖏᑦ:

ᑲᓇᑕᐅᑦ ᑲᖏᒪᑯᓂᖏᓪᓄᓕᑦ ᑐᑭᓯᑕᐅᓂ:
ᐊᐃᑲᖅᑐᒪᓕᐊᓂ ᑐᑭᒧᐊᖅᑎᖅᑲᓂ
ᐊᖏᑎᑕᓚᓂᑲᒧᖅ ᐱᓕᑎᖅᑎᑎᑦ ᑲᓇᓱᓚᒋᓂ ᐊᓚᑦ ᐅᑭᐅᖅᑕᖅᑐᖅ
ᐊᖏᑎᑕᓚᐅᑯᑦ ᑲᓇᑕᒥ
9250 49th Street, ᐃᒪᒐᓱ, ᐊᐃᑉᑕ T6B 1K5
(780) 951-8850
(780) 495-2615 (ᓯᑲᕆᑯᑦ)
david.ingstrup@canada.ca

ᓄᓇᕗᑦ ᑐᓴᓪᓗᒃ ᑎᒥᖅᓯᓄᑦ ᑐᖅᑭᒃᓴ
ᐊᓴᓪᓚᕆᒃᑐᖅ ᑎᑎᕋᖅᓇᒥᒥ
ᓄᓇᕗᑦ ᑐᓴᓪᓗᒃ ᑎᒥᖅᓯ ᑎᑎᕋᖅᑲᒃᑯᓪᓗᒃ 638
ᐃᕋᒃᓗᐃᑦ, ᓄᓇᕗᑦ X0A 0H0
(867) 975-4900
(867) 975-4949 (ᓱᒃᑲᓪᒃᑯᑦ)
jtarreak@tunngavik.com

ᕿᑎᕋᐅᑦ ᐃᓄᐃᑦ ᑲᑐᔾᕋᖅᑎᑯᓐᓂᒃᑯᓄᑦ ᑐᖅᑭᒃᓴ:
ᐊᐅᓪᓕᔪᓐᓃᑦ ᑐᑭᒧᐊᖅᑎᖅᓯ ᕿᑎᕋᐅᑦ ᐃᓄᐃᑦ ᑲᑐᔾᕋᖅᑎᑯᓐᓂᖅᓯᓕᒥᑦ
ᑎᑎᕋᖅᑲᒃᑯᓪᓗᒃ 18
ᐃᕋᒃᓗᓐᓇᕆᐊᕋᒃ, ᓄᓇᕗᑦ X0B 0C0
(867) 983-2458
(867) 983-2701 (ᓱᒃᑲᓪᒃᑯᑦ)
execdir@kitia.ca

ᑭᖑᓪᓕᕋᒃ ᐃᓄᐃᑦ ᑲᑐᔾᕋᖅᑎᑯᓐᓂᒃᑯᓄᑦ ᑐᖅᑭᒃᓴ:
ᐊᐅᓪᓕᔪᓐᓃᑦ ᑐᑭᒧᐊᖅᑎᖅᓯ ᑭᖑᓪᓕᕋᒃ ᐃᓄᐃᑦ ᑲᑐᔾᕋᖅᑎᑯᓐᓂᖅᓯᓕᒥᑦ
ᑎᑎᕋᖅᑲᒃᑯᓪᓗᒃ 340
ᑲᖏᕐᓂᓕᕋᒃ, ᓄᓇᕗᑦ X0C 0G0
(867) 645-2810
(867) 645-3855 (ᓱᒃᑲᓪᒃᑯᑦ)
gkarlik@kivalliqinuit.ca

ᕿᑭᕋᑕᓂ ᐃᓄᐃᑦ ᑲᑐᔾᕋᖅᑎᑯᓐᓂᒃᑯᓄᑦ ᑐᖅᑭᒃᓴ:
ᐊᐅᓪᓕᔪᓐᓃᑦ ᑐᑭᒧᐊᖅᑎᖅᓯ ᕿᑭᕋᖅᑕᓂ ᐃᓄᐃᑦ ᑲᑐᔾᕋᖅᑎᑯᓐᓂᖅᓯᓕᒥᑦ
ᑎᑎᕋᖅᑲᒃᑯᓪᓗᒃ 1340
ᐃᕋᒃᓗᐃᑦ, ᓄᓇᕗᑦ X0A 0H0
(867) 975-8400
(867) 979-3238 (ᓱᒃᑲᓪᒃᑯᑦ)
nbeveridge@qia.ca

2.8.3 ᑐᓴᕋᑎᔪᓕᓂ ᕋᖅᓯᓴᐅᔾᔨᒃᑯᑦ, ᐊᐅᑦᓕᓕᑎᑕᐅᓂᐊᕋᖅᑐᑦ ᐃᓄᖓᓄᑦ ᐃᕋᒃᑲᓇᐃᔾᔨᕋᖅᑐᑦ ᓇᓗᓇᐃᕋᖅᓯᒪᔪᓗᒃᑐᓄᒃ ᑕᐃᒃᑯᓂ ᑐᓴᕋᑎᔪᓕᓂᕋᖅᑕᓂᕐᓗᑎᓗᒃᑐᑦ, ᑭᓯᐊᓂ ᐱᓇᓯᐊᕋᖅᑎᑯᓐᓂᑦ ᑐᓴᕋᐅᒪᕋᖅᑎᑯᓐᓂᒃᑯᑦ.

2.8.4 ᑕᐃᒪᓗ ᓄᕋᖅᑐᒃᓯᑕ ᑎᑎᕋᖅᑕᓚᐅᒃᑯᑦ ᓇᒻᒥᓂᖅ ᐊᓱᕋᖅᓯᓗᑎᒃ, ᐅᕐᓕᔭᓐᓂᓂᖅ ᓯᒃᑲᓪᒃᑯᑦ ᐅᕐᓕᔭᓐᓂᓂᖅ ᕋᖅᓯᓴᐅᔾᔨᒃᑯᑦ.

ᑲᓇᑕᒥ ᐅᒪᔪᖏᖅᓴᓕᑦ ᐊᓚᒪᓗ ᐅᑎᖅᑕᕐᕕᑐᓂᓄᑦ ᑎᖅᒥᐊᓄᑦ ᓴᓗᐅᓕᕐᓚᕐᓴᐅ ᐅᓇ ᐃᓄᐃᑦ ᐊᐅᑐᖅᑕᐅᓂᐊᕐᓂᖅᑭᓐᓄᓄᑦ ᐱᖓᓱᓕᕐᕕᐊᔪᖅᑐᓂᓪᓗ ᐊᖏᑭᑎᒃ ᐃᑕᖅᓚᔪᑦ ᐃᓕᓴᕐᓂ 2-1 (ᑎᑎᕋᖅᓚ 2.2.1)

ᑲᓇᑕᒥ ᐅᒪᔪᖏᖅᓴᓕᑦ ᐊᓚᒪᓗ ᐅᑎᖅᑕᕐᕕᑐᓂᓄᑦ ᑎᖅᒥᐊᓄᑦ ᓴᓗᐅᓕᕐᓚᕐᓴᐅ	ᖃᓄᐅᔨᖅᑭᑦ ᓄᓇᓕᖕᓂᑦ
1. ᐊᐅᐸᐃᑦ ᑲᓇᑕᒥ ᐅᒪᔪᖏᖅᓴᓕᑦ	ᖁᑭᖅᑕᕐᕿᐊᕐᖅ
2. ᓴᓐᓂᑉᑕᑦ ᐅᑎᖅᑕᕐᕕᑐᓂᓄᑦ ᑎᖅᒥᐊᓄᑦ ᓴᓗᐅᓕᕐᓚᕐᓴᐅ	ᒥᑦᑎᒪᒋᑦᓚᐅ
3. ᐃᓚᓕᔨᖏᓂᐅ ᐅᑎᖅᑕᕐᕕᑐᓂᓄᑦ ᑎᖅᒥᐊᓄᑦ ᓴᓗᐅᓕᕐᓚᕐᓴᐅ	ᑭᓇᖏᐃᑦ
4. ᖃᐅᖅᓴᐅᖅᑐᖅ ᐅᑎᖅᑕᕐᕕᑐᓂᓄᑦ ᑎᖅᒥᐊᓄᑦ ᓴᓗᐅᓕᕐᓚᕐᓴᐅ	ᓴᓪᓕᖅ
5. ᐃᐅᓚᑐᐊᖅ ᐅᑎᖅᑕᕐᕕᑐᓂᓄᑦ ᑎᖅᒥᐊᓄᑦ ᓴᓗᐅᓕᕐᓚᕐᓴᐅ	ᓴᓪᓕᖅ
6. ᓂᓇᖏᓇᓂᓂ ᑲᓇᑕᒥ ᐅᒪᔪᖏᖅᓴᓕᑦ	ᑲᖏᕐᖅᓵᑦᓇᐱᐅᖅ
7. ᑯᐃᓴᔪᖅ ᐅᑎᖅᑕᕐᕕᑐᓂᓄᑦ ᑎᖅᒥᐊᓄᑦ ᓴᓗᐅᓕᕐᓚᕐᓴᐅ	ᐊᖅᓇᐊᑦ
8. ᓂᓴᐃᑎᖅᓴᐅᖅ ᑲᓇᑕᒥ ᐅᒪᔪᖏᖅᓴᓕᑦ	ᐊᐅᓚᐅᓪᑕᖅ
9. ᑐᐅᑕᐊᖅᓇᐅᕐᐸ ᓇᖅᓴᖓ ᑲᓇᑕᒥ ᓂᓴᐃᑎᓂᑦ ᓄᓇᖅᑯᑎᖓᑦ	ᖃᑲᐅᓚᐅᓪᑕᖅ
10. ᐊᐅᐸᑦ ᐅᑎᖅᑕᕐᕕᑐᓂᓄᑦ ᑎᖅᒥᐊᓄᑦ ᓴᓗᐅᓕᕐᓚᕐᓴᐅ	ᖃᑲᐅᓚᐅᓪᑕᖅ, ᐃᐅᐱᐊᖅᓴᖅ
11. ᖃᑲᖅᑯᓪᓗᐃᑦ ᑲᓇᑕᒥ ᐅᒪᔪᖏᖅᓴᓕᑦ	ᖁᑭᖅᑕᕐᕿᐊᕐᖅ
12. ᐃᓗᐃᓪᓕᖅ ᐅᑎᖅᑕᕐᕕᑐᓂᓄᑦ ᑎᖅᒥᐊᓄᑦ ᓴᓗᐅᓕᕐᓚᕐᓴᐅ	ᐃᖃᓗᓐᑐᑎᐊᖅ, ᐅᖅᓚᖅᑐᖅ, ᐅᒥᖓᒪᓐᑐᖅ
13. ᓚᒍᐊ ᐅᑎᖅᑕᕐᕕᑐᓂᓄᑦ ᑎᖅᒥᐊᓄᑦ ᓴᓗᐅᓕᕐᓚᕐᓴᐅ	ᖃᑲᐅᓚᐅᓪᑕᖅ

ᐃᓕᓐᓴ 3 — ᐊᐅᓕᕐᑎᕋᑎᑦᑕᓂᑦ

3.1 ᑐᑭᒋᖅᓯᒡᓗᐊᔨᓴᑎᓄᖅᓴᓂᑦ

3.1.1 ᐅᓇ ᐃᓕᓐᓴ ᐊᒋᑦᓂ ᑎᑎᕋᖅᓭᒪᓕᔪᓂ ᑐᑭᒋᕋᓯᒡᓗᐊᔨᓴᑎᓄᖅᓴᓂᖅᑐᖅ:

(a) ᐊᑐᑎᖅᓂᑎᐊᖅᑐᑐᒃ ᑲᓇᑕᒥ ᐅᒪᓕᔪᖅᓱᐃᑦ ᐊᒻᒪᓗ ᐅᑎᖅᑕᓯᒡᓗᓇᑦ ᑎᓐᓕᐊᓇᑦ ᓴᓴᐅᓭᒪᓕᐹᒃ ᐃᓇᓐᓂᑦ ᐊᐅᓕᕐᑎᕋᑎᑦᑕᓂᑦᑐᖅ ᐃᓕᓴᕐᓂᓕᕆᔨᑦ ᒪᑕᒃᑐᑐᒃᓴᓂᑦ ᐃᓇᓐᓂᑦ ᐊᐅᓕᕐᑎᕋᑎᑦᑕᓂᑦᑐᖅ ᓄᓇᕗᒻᒥᒃ *ᓄᓇᕗᒻᒥᒃ ᓄᓇᑕᐅᑎᓇᑦ ᐊᐅᑉᐳᑎᑦ*, ᐊᒻᒪᓗ ᐱᓗᐊᖅᑐᒥᒃ ᐃᓕᓐᓴ 9 ᐊᒻᒪᓗ 5 ᑕᒃᑯᓇᓂ *ᓄᓇᕗᒻᒥᒃ ᓄᓇᑕᐅᑎᓇᑦ ᐊᐅᑉᐳᑎᑦ;*

(b) ᐃᓱᒪᓕᐅᓚᓂᖅ ᐱᓕᓗᑎᑦ ᐅᑎᖅᑕᓯᒡᓗᓇᑦ ᑎᓐᓕᐊᓇᑦ ᓴᓴᐅᓭᒪᓕᐹᒃ ᐊᒻᒪᓗ ᑲᓇᑕᒥ ᐅᓇᖅᑐᖅᓴᓕᐅᕐᓂᖅ ᐊᐅᓚᑦᑎᐊᖅᑐᑦ ᐊᒻᒪᓗ ᑕᐃᒪᓐᓇᐃᑲᖅᑕᐅᓗᑎᒃ *ᐃᓇᐃᑦ ᖃᐅᔨᒪᔭᑐᖅᓂᖓᓂᑦ;* ᐊᒻᒪᓗ

(c) ᓄᓇᕗᓂᓂ ᐃᓇᐃᑦ ᐸᖅᓇᐃᓗᑎᒃ ᐊᒻᒪᓗ ᐊᐅᓕᕐᑎᓗᑎᒃ ᑲᓇᑕᒥ ᐅᓕᖅᑐᖅᓴᓕᐅᕐᓂᖅ ᐊᒻᒪᓗ ᐅᑎᖅᑕᓯᒡᓗᓇᑦ ᑎᓐᓕᐊᓇᑦ ᓴᓴᐅᓭᒪᓕᐹᒃ.

3.2 ᓄᓇᒥᒃ ᑲᑎᒪᔨᖅᑎᑎᑦᑕᓂᑦ ᑲᑎᒪᔨᕆᓭᓕᕐᓂᑦ

ᓴᖅᑭᑕᐅᓕᒪᔨᓴᑎᓂᖓᓂᑦ ᐊᒻᒪᓗ ᐱᔨᓴᑎᓂᖓᓂᑦ

3.2.1 ᐊᐅᓕᕐᑎᖅᑎᑎᑦᑕᓂᓕᓐᓂᒡᓗᑦ ᑲᑎᒪᔨᕆᓭᓕᑦ ᑲᒪᒋᔭᐅᑎᖅᑕᐅᓂᓕᓐᓂᕐᓂᓕᕆᓭᐊᖅᑐᑦ ᐊᑐᓂ ᑲᓇᑕᒥ ᐅᒪᓕᔪᖅᓱᐃᑦ ᐅᓂᒡᓘᓐᓃᑦ ᐅᑎᖅᑕᓯᒡᓗᓇᑦ ᑎᓐᓕᐊᓇᑦ ᓴᓴᐅᓭᒪᓕᐹᒃ ᐅᓂᒡᓘᓐᓃᑦ ᑲᑎᒪᔨᓂᑦ ᐊᑎᖅᑲᖅᑐᖅᓴᖅ ᐊᒥᓕᕐᓂᖅ 3-1-ᒥ.

3.2.2 ᐊᑐᓂ ᐊᐅᓕᕐᑎᖅᑎᑎᑦᑕᓂᓕᓐᓂᒡᓗᑦ ᑲᑎᒪᔨᕆᓭᓕᑦ ᐃᓕᓴᐅᓂᐊᕐᓂᐊᖅᑐᑦ ᐊᐅᓕᕐᑎᓗᑎᒃ ᐊᒋᕼᒥᔪᓄᑦ ᓄᓇᕗᔪᓄᑦ ᐊᑎᖅᑲᖅᑐᖅᓴᖅ 3-1-ᒥ, ᑭᓱᐊᓂ ᐊᕐᔨᑎᕼᐅᓂ ᐃᓗᐃᑦᓕᓂᖅ ᐅᑎᖅᑕᓯᒡᓗᓇᑦ ᑎᓐᓕᐊᓇᑦ ᓴᓴᐅᓭᒪᓕᐹᒃ, ᐅᔨᐊ ᐊᐅᓕᕐᑎᖅᑎᑎᑦᑕᓂᓕᓐᓂᒡᓗᑦ ᑲᑎᒪᔨᕆᓭᓕᑦ ᐃᓂᖅᑲᖅᑐᖅᓴᖅ ᐊᐅᓕᕐᑎᓕᓗᑎᒃ ᐃᖅᑲᓗᑐᑎᐊᖅᑎᒥ.

3.2.3 ᐱᔨᓴᑎᓂᖓᓂ ᐊᐅᓕᕐᑎᖅᑎᑎᑦᑕᓂᓕᓐᓂᒡᓗᑦ ᑲᑎᒪᔨᕆᓭᓕᑦ ᐃᓕᓴᕐᓂᐅᐅᓂᐊᖅᑐᖅ:

(a) ᐅᖅᑲᐅᔪᖅᑲᑎᑦᑕᓗᑎᒃ ᒥᓂᔅᑕᒍᑦ, ᐅᓚᓕᑲᑎᓕᓗᒍ, ᓇᐅᑲᔨᑐᐃᓇᕐᓇᖅ ᐸᖅᓇᐃᓂᖅᒥ ᐊᒻᒪᓗ ᐊᐅᓕᕐᑎᓂᖅᒥᑲ ᑕᒃᔨᓂᖅᓴᖅ ᑲᑎᒪᔨᕆᓭᓕᑦ ᑎᓐᓕᐊᓇᑦ ᓴᓴᐅᓭᒪᓕᐹᒃ ᑲᑎᒪᔨᕆᓭᓕᑦ ᑎᓐᓕᐊᓇᑦ ᓴᓴᐅᓭᒪᓕᐹᒃ, ᒪᓕᓴᓗᑎᒃ ᐃᓕᓐᓴ 3.3 ᑕᕐᔭᓂ ᐃᓇᐃᑦ ᐊᖅᑐᖅᑕᐅᓂᐊᕐᓂᕐᓂᕐᓇᓄᑦ ᐱᓵᓕᓂᕈᐅᓇᖅᑕᐅᓇᓕᓴᓗ ᐊᕐᑎᕈᑎᓂᑦ;

(b) ᑎᑎᕋᓴᓂᕈᓂ, ᐊᑦᑭᑭᐊᑦᓴᓂᕈᓂ ᐊᒻᒪᓗ ᐊᑐᓕᕆᔨᓴᓕᓯ ᑕᒃᔨᓂᖅᓴᖅ ᐊᐅᓕᕐᑎᓂᐱᑎᖅᓴᓇᓄᑦ ᐸᖅᓇᐃᓗᑎᒃ ᐃᓕᓐᓴᕈᓂ 3.5 ᑎᑭᓕᓴᒍ 3.7 ᑕᕐᔭᓂ ᐃᓇᐃᑦ ᐊᖅᑐᖅᑕᐅᓂᐊᕐᓂᕐᓂᕐᓇᓄᑦ ᐱᓵᓕᓂᕈᐅᓇᖅᑕᐅᓇᓕᓴᓗ ᐊᕐᑎᕈᑎᓂᑦ; ᐊᒻᒪᓗ

(c) ᐱᓕᓚᓂᕈᓂ ᐊᓯᓂᕈᓂ ᐊᐅᓕᕐᑎᐱᑎᓂᕆᓂᕈᓂ ᑕᒃᔨᓚᓂ ᐊᐅᓕᕐᑎᖅᑎᑎᑦᑕᓂᓕᓐᓂᒡᓗᑦ ᑎᑎᕋᖅᓭᒪᓕᐹᓂ ᐃᓇᐃᑦ ᐊᖅᑐᖅᑕᐅᓂᐊᕐᓂᕐᓂᕐᓇᓄᑦ ᐱᓵᓕᓂᕈᐅᓇᖅᑕᐅᓇᓕᓴᓗ ᐊᕐᑎᕈᑎᓂᑦ.

ᑎᒃᓯᕆᕐᕕᓂᖅᓄᑦ ᐊᒻᒪᓗ ᐊᑕᐃᖅᑑᑦ ᐃᓕᑎᐅᔪᓄᑦ

3.2.4 ᐊᑐᓂ ᐊᐅᓪᓕᕐᓴᐅᑎᒋᔭᖏᓪᓗ ᑲᑎᒪᔨᖏᓪᓕᐅᕙᓪᓕᖅᑐᑦ, ᑎᒃᓯᕆᔨᕐᓯᒪᓕᕐᑎᒃ ᐅᓯᓂᖅᓱ:

(a) ᐊᑐᑎᖏᖅᑐᖅ ᐊᐃᖅᑐᖅᓯᒪᔪᓄ ᐃᓄᐃᓄ ᑲᑐᔨᖏᓪᓕᐅᕐᓄ ᑎᒃᓯᕆᔨᕐᓴᐅᑎᒃ ᐱᖓᓱᓂᖅ (3), ᑕᒪᖏᑦ ᒎᒃᓯᓇᖅᑕ ᓄᓇᕗᒻᒥᔨᖏᑦ ᑲᕐᓚᐅᖅᓂᖅ ᓄᓇᓕ ᓄᓇᕆᔨᑎᖅᑕᐅᓂᖅ ᓄᓇᒥᐅᑕᖏᓐᓄᑦ ᓄᓇᑲᑎᒪᔨᖏᑦ;

(b) ᐅᓇ ᒥᓂᑦ ᑎᒃᓯᕆᔨᕐᓴᐅᓂ ᐱᖓᓱᓂᖅ (3), ᑕᒪᖏᑦ ᒎᒃᓯᓇᖅᑕ ᐃᓕᑎᐅᔪᓄᑦ ᖃᐅᑎᖓᖓᓂᑦ ᓄᓇᕗᖅᓂᑦ ᓄᓇᕗᒻᒥᔨᖏᑦ ᑲᑎᒪᔨᖏᓪᓕᐅᕐᓄ ᑲᑎᒪᔨᖏᓪᓕᐅᕐᓴᐅᑎᒃ ᓄᓇᑲᑎᒪᔨᖏᑦ ᓄᓇᒥᐅᑕᖏᓐᓄᑦ ᓄᓇᕗᑦ ᑲᑎᒪᔨᖏᓪᓕᐅᕐᓄ ᑲᑎᒪᔨᖏᓪᓕᐅᕐᓴᐅᑎᒃ ᓄᓇᑲᑎᒪᔨᖏᑦ. ᐅᓇ ᒥᓂᑦ ᐱᖓᓱᖓᓐᓂᖅ ᑎᒃᓯᕆᔨᕐᓴᐅᓯᒪᔪᖅᑐᖅᓄᑦ ᐃᓅᖅᑲᐅᑉᐳᖓᓂᖅ ᖃᐅᔨᓄᑦ ᐃᓕᑎᐅᔪᓄᑦ ᐅᖃᓕᒫᓂᖅ;

(c) ᑕᐃᒃᑐᓂ ᐊᐅᓪᓕᕐᓴᐅᑎᒋᔭᖏᓪᓗ ᑲᑎᒪᔨᖏᓪᓕᐅᕙᓪᓕᖅᑐᑦ ᑭᓚᖅᑐᖅᑎᖅᑕᐅᓇᖅᑐᑦ ᐊᑕᐅᓯᐅᓇᖅᑕᑐᒥᔅ ᖃᐅᑎᖓᖓᓂᖅ ᓄᓇᕗᒻᒥ, ᐅᖃᖅᑐᓯᒪᔪᖅ ᑎᑎᖅᑐᓂ 3-1, ᐃᓕᑎᐅᔪᓄᑦ ᓄᓇᓕ ᓄᓇᕆᔨᑎᖅᑕᐅᓂᖅ ᓄᓇᒥᐅᑕᖏᓐᓄᑦ ᓄᓇᑲᑎᒪᔨᖏᑦ ᐅᓇᓕᒫᓂᖅ ᓄᓇᕗᒻᒥ ᐊᓐᑕᔪᓄᑦ ᐱᓕᕆᔨᓄᑦ ᑎᒃᓯᕆᔨᕐᓴᐅᑎᒃ ᒎᒃᓯᓇᖅᑕ ᐊᐅᓪᓕᕐᓴᐅᑎᒋᔭᖏᓪᓗ ᑲᑎᒪᔨᖏᓪᓕᐅᕙᓪᓕᖅᑐᑦ ᐅᓇᓕᒫᓂᖅ ᐊᐅᓪᓕᕐᓴᐅᑎᒋᔭᖏᓪᓗ ᑲᑎᒪᔨᖏᓪᓕᐅᕙᓪᓕᖅᑐᑦ ᐅᖃᓕᒫᓂᖅ ᐊᐅᓪᓕᕐᓴᐅᑎᒋᔭᖏᓪᓗ ᑲᑎᒪᔨᖏᓪᓕᐅᕙᓪᓕᖅᑐᑦ ᐅᓇ

ᓄᓇᕗᑦᑐᓂᑦ ᐱᓕᕆᓇᐅᑎᓪᓗ ᑲᑎᒪᔨᖏᓪᓕᐊᑦ ᐱᖁᑲᓇᑐᖓᓂ ᐱᓕᕆᔨᖏᑦ ᑎᑎᖅᑲᓇᑐᖓᓂ ᐊᑐᓂᒃ ᐊᐅᓚᑦᑎᑲᑎᒃᓴᓕᒫᑦ ᑲᑎᒪᔨᖏᓪᓕᐊᑦ.

3.2.9 ᐊᐅᓚᑦᑎᑲᑎᒃᓴᓕᒫᑦ ᑲᑎᒪᔨᖏᓪᓕᐊᑦ, ᐃᓚᒋᔭᐅᓪᓗᑎᒃ ᐃᒃᓯᕐᐅᒡᓕᖅ ᐊᓯᒪᓗ ᐃᒃᓯᕐᐅᒡᓕᐅᕐᓴ ᑐᖓᓕᖓᓂᑦ, ᑎᒃᑯᐊᖅᑕᐅᓗᑎᒃ ᐃᓕᓴᕐᓂᓕᒃ.

3.2.10 ᐊᕐᓇᖅᑐᒥᔪᓂ ᐃᓄᐃᑦ ᑲᑐᔨᖏᑎᒃᓴᖓᓂ ᐅᖃᖅᖢᖓᓂ ᒥᓂᑦᑐᑦᓯᓚᖅᑐᑦ ᐊᐅᓚᑦᑎᑲᑎᒃᓴᓕᒫᑦ ᑲᑎᒪᔨᖏᓪᓕᐊᑦ ᒪᓕᒐᓗᒍ ᐊᐅᓚᑦᑎᑲᑎᒃᓴᓕᒫᑦ ᑲᑎᒪᔨᖏᓪᓕᐊᑦ ᐱᐅᔪᓚᐃᕐᓇᖅᑐᖓᓂ ᓴᖅᑭᑕᐅᔪᒥᔪᓂ ᒪᓕᒐᓘᔪᖅ 3.2.22.

3.2.11 ᑕᐃᒪᓗ ᐃᓕᓴᐅᔪᖅ ᐊᓯᖅᑕᐅᔪᓂᒃ ᐅᖃᖅᑐᓂᒃᓯᒪᓂ ᓄᖅᑲᓂ ᐊᐅᓚᑦᑎᑲᑎᒃᓴᓕᒫᑦ ᑲᑎᒪᔨᖏᓪᓕᐊᑦ, ᐅᑯᐊ ᐊᕐᓇᖅᑐᒥᔪᓂ ᐃᓄᐃᑦ ᑲᑎᒪᔨᖏᑎᒃᓴᖓᓂ ᐅᖃᖅᑐᓂᒃᓯᒪᓂ ᑎᒍᑯᓕᕐᓂᖅᑐᑦ ᐃᓕᓴᐅᔪᖅᑎᑦ, ᐊᑯᓂᐅᓗᓕᕐᓂᖓᓕᖅᑎᑦ, ᑎᒃᑯᐊᖅᑕᐅᓗᑎᒃ ᐃᓐᓇᑎᒃᓴᐊᕐᓗᓂᒃᑐᖅ ᑯᓚᖓ ᐃᓄᐃᑦ ᐊᓂᔪᓂᒃᑯᑦᑎᓂᒃᑯᑦ, ᐊᑯᓂᐅᓗᓕᕐᓂᖓᓕᖅᑎᑦ, ᑎᒃᑯᐊᖅᑕᐅᓗᑎᒃ ᐃᓐᓇᑎᒃᓴᐊᕐᓗᓂᒃᑐᖅ ᑯᓚᖓ ᐃᓄᐃᑦ ᐊᐅᓚᑦᑎᑲᑎᒃᓴᓕᒫᑦ ᑲᑎᒪᔨᖏᓪᓕᐊᑦ ᐱᐅᔪᓚᐃᕐᓇᖅᑐᖓᓂ ᑲᓇᑕᐅᔪᖅᑐᑦ ᐃᓕᓴᐅᔪᖅᑎᑦ ᓄᖅᑲᓂ ᐊᐅᓚᑦᑎᑲᑎᒃᓴᓕᒫᑦ ᑲᑎᒪᔨᖏᓪᓕᐊᑦ ᐱᐅᔪᓚᐃᕐᓇᖅᑐᖓᓂ ᓄᓇᕗᑦᑐᓂᑦ ᐱᓕᕆᓇᐅᑎᓪᓗ ᐃᓄᐃᑦ ᑲᑎᒪᔨᖏᑎᒃᓴᖓᓂ ᐅᖃᖅᑐᓂᒃᓯᒪᓂ ᑎᒍᑯᓕᕐᓂᖅᑐᑦ ᐃᓕᓴᐅᔪᖅᑎᑦ, ᐅᑯᐊ ᐃᓄᐃᑦ ᐊᐅᓚᑦᑎᑲᑎᒃᓴᓕᒫᑦ ᑲᑎᒪᔨᖏᓪᓕᐊᑦ ᐱᐅᔪᓚᐃᕐᓇᖅᑐᖓᓂ ᓄᓇᕗᑦᑐᓂᑦ ᐱᓕᕆᓇᐅᑎᓪᓗ ᐃᓄᐃᑦ ᑲᑎᒪᔨᖏᑎᒃᓴᖓᓂ ᐅᖃᖅᑐᓂᒃᓯᒪᓂ ᑎᒍᑯᓕᕐᓂᖅᑐᑦ ᐃᓕᓴᐅᔪᖅᑎᑦ, ᐊᑯᓂᐅᓗᓕᕐᓂᖓᓕᖅᑎᑦ, ᑎᒃᑯᐊᖅᑕᐅᓗᑎᒃ ᐃᓐᓇᑎᒃᓴᐊᕐᓗᓂᒃᑐᖅ ᑯᓚᖓ ᐃᓄᐃᑦ ᐊᐅᓚᑦᑎᑲᑎᒃᓴᓕᒫᑦ ᑲᑎᒪᔨᖏᓪᓕᐊᑦ ᐱᐅᔪᓚᐃᕐᓇᖅᑐᖓᓂ ᓄᓇᕗᑦᑐᓂᑦ ᐱᓕᕆᓇᐅᑎᓪᓗ ᐃᓄᐃᑦ ᑲᑎᒪᔨᖏᑎᒃᓴᖓᓂ ᐅᖃᖅᑐᓂᒃᓯᒪᓂ ᑎᒍᑯᓕᕐᓂᖅᑐᑦ ᐃᓕᓴᐅᔪᖅᑎᑦ, ᐊᑯᓂᐅᓗᓕᕐᓂᖓᓕᖅᑎᑦ, ᐊᑐᓂᒃ

ᐱᒃᓗᐅᖅᑳᑎᑕᐅᓗᑎᒃ ᑐᓴᐅᒪᒃᑭᕐᓂᖅᓯᓗ ᑐᓴᕿᑎᕆᓗᑎᒃ, ᐃᒪᒥᖅᓂᒃ ᐃᓕᓴᕿᑎᒃᑭᕐᓗᑎᒃ, ᖃᐅᔨᑐᐃᓐᓇᖅ, ᐅᒪᑭᓗᑎᓕᓕᒫᑦ.

ᐊᐅᓚᕋᑎᓂᖅᑭᑦ

3.2.16 ᐊᑐᓂ ᐊᐅᓚᕋᕿᒃᑎᑦᓂᖅᓴᑦ ᐅᑎᒪᔨᕆᔨᓚᑦ ᐅᑎᖅᓱᓗᑎᒃ ᐅᑎᒪᖃᑦᑕᕐᔪᓗᑎᒃ ᐊᖏᕐᕕᒃ ᐊᑕᐅᓯᐊᖏᓗᑎᒃ .ᐊᐅᓚᕋᕿᒃᑎᑦᓂᖅᓴᑦ ᐅᑎᒪᔨᕆᔨᓚᑦ ᐱᓇᓚᖃᑦᑕᕐᔪᓂᐊᖏᓚᐅᕐᓂ ᐅᖃᓗᓐᓄᑦᑐᑦ ᐅᑎᒪᖃᑦᑕᕐᔪᓗᑎᒃ ᐊᒻᒪᓗ ᓇᓚᐊᑐᐃᓐᓇᖅ ᐃᓯᒪᓚᖃᖅᑎᒃ ᐊᑐᑎᖃᖏᓐᓇᒃ.

3.2.17 ᐊᐅᓚᕋᕿᒃᑎᑦᓂᖅᓴᑦ ᐅᑎᒪᔨᕆᔨᓚᑦ ᐅᑎᒪᖃᑦᑕᕐᔪᓗᑎᒃ ᒪᑐᐃᖅᓱᓗᑎᒃ ᐃᓄᓚᒻᒥᓄᑦ, ᑭᓯᐊᓂ ᐊᐅᓚᕋᕿᒃᑎᑦᓂᖅᓴᑦ ᐅᑎᒪᔨᕆᔨᓚᑦ ᐅᑎᒪᔪᑐᐊᑦ ᐃᓕᓴᐅᓂᓕᑦ ᑕᐃᒪᓐᓇᖅᑐᑦ, ᑕᐃᒪᓐᓇᐅᔪᒪᓕᒫᓂᑦ ᐊᐅᓚᕋᕿᒃᑎᑦᓂᖅᓴᑦ ᐅᑎᒪᔨᕆᔨᓚᑦ ᑭᓯᐊᓂ ᐱᓕᕆᓂᖅᑎᒃ ᑕᐃᒪᓐᓇᐅᕙᓚᕐᓂᖅ.

3.2.18 ᐊᐅᓚᕋᕿᒃᑎᑦᓂᖅᓴᑦ ᐅᑎᒪᔨᕆᔨᓚᑦ ᐱᓕᕆᓚᖃᑦᑕᕐᔪᓗᑎᒃ ᐱᒃᖏᕐᕋᓂᑦ ᐃᓄᐅᑎᓂᑦ ᐅᖃᖅᑎᓕᓚᓗᑎᒃ ᐊᒻᒪᓗ ᖃᖅᑎᓗᓄᐅᕙᓚᖅᓱᓗᑎᒃ, ᐊᒻᒪᓗ ᑐᓴᕿᖅᓱᓗᑎᒃ ᐊᒻᒪᓗ ᐃᓄᐅᑎᑐᑦᓂᓚᕐᔪᖅᓱᓗᑎᒃ ᐱᔪᓚᑎᒃᑎᓂᑦ ᐃᓕᑎᐅᕙᓚᕐᓂ.

3.2.19 ᐅᓄᖅᓂᖅᑭᑦ ᐅᑎᒪᔪᓐᓇᖅᓱᓗᑎᒃ ᓇᓕᐊᑐᐃᓐᓇᖅᑲᓂ ᐊᐅᓚᕋᕿᒃᑎᑦᓂᖅᓴᑦ ᐅᑎᒪᔨᕆᔨᓚᑦ ᑎᓴᒪᐅᓚᑦ (4) ᐃᓕᑎᐅᕙᔪᑦ, ᒪᖅᑲᒃ ᒡᒃᔨᓇᖅᑐᒥᓂᑦ ᐃᓄᐃᑦ ᑲᑐᔨᕆᑎᒃᑎᑦᓂᖅᓴᑦ ᐱᔪᓚᑎᒃᑎᓂᐅᓂᓕᒫᓂᑦ ᑎᓴᔨᐊᖅᑕᐅᔨᒪᔪᑦ.

3.2.20 ᐃᓕᑎᐅᕙᔪᑦ ᑎᓴᔨᐊᖅᑭᓗᑎᒃ ᒡᒃᔨᓄᖅᓴᒃ ᐊᐅᓚᕋᕿᒃᑎᑦᓂᖅᓴᑦ ᐅᑎᒪᔨᕆᔨᓚᑦ ᐃᔨᒪᑎᕐᓄᑦ ᐱᔪᓚᑎᒃᑎᓂᑦ ᐱᔪᓐᓇᖅᑐᒥᓂᑦ ᐃᓄᐃᑦ ᑲᑐᔨᕆᑎᒃ ᐊᑕᔪ

3.2.24 ᐊᑐᓂ ᐊᐅᓪᓕᖅᑎᑦᑎᓯᒪᓂ ᑲᑎᒪᔨᖏᓪᓗ ᑐᓂᓯᔪᑎᒃ ᑐᖓᔪᖅᐅᑎᓗᒍ ᓴᖅᑭᑐᖅᑕᐅᓂᖅ ᐊᓕᒪᔪᓂ ᑲᑎᒪᔨᖏᓪᓗᒍ ᒪᑐᐊᖅᑐᓂᒃ ᐅᓇᓱᓂ ᑐᓴᕐᓯᒪᓕᕆᔨᒃᑯᑦ ᐅᓇᑲᓂ ᑐᖓᔪᖅᐅᑎᓗᒍ ᐱᖓᓱᒃᑯᑦ ᐊᑐᖅᑐᖓᓂ ᐊᐅᓚᑦᑎᓂᒃᑯᑦ ᐊᓕᒪᔪᓂ ᐃᓚᖏᑕᓗᒍ ᐊᓯᖏᓪᓗᒍ ᐊᐅᓚᑦᑎᓂᒃᑯᑦ ᐊᓕᒪᔪᓂ ᐃᓚᖏᑕ

3.2.30 ᐅᑯᐊ ᑎᑎᕎᒋᓗᐊᓂᒃ ᐃᑲᔪᖅᑎᖅᑲᓂᖅ ᐱᒋᓗᐊᖅᑐᐅᐊᖅᑐᖅ ᒐᕙᒪᖓᓂ ᑲᓇᑕᒥ ᐅᒪᔪᒋᓗᐊᕐᕕᐊᑯᑦ ᐃᒪᐠᓇ:

(a) ᖃᓄᐃᑦᐅᓇᐊᒋᖓᓄᑦ ᐃᑲᔪᕐᓗᑎᒃ ᐃᓕᖓᕆᐅᓗᑎᒃ ᐊᐅᓚᕋᓱᐃᓕ ᐊᐅᓚᕎᑎᖃᑎᒌᖓᔪᒥᑦ ᑲᑎᒪᔨᕈᓯᑦ ᑲᑎᒪᓂᖓᓄᑦ, ᐅᖃᖓᓗᓂᒃᑯᑦ ᑲᑎᒪᓂᖓᓄᑦ, ᓴᖅᑭᕎᑦ ᖃᓄᖅᑐᐃᓂᖓᓄᑦ, ᖃᓄᖓᑦᑕᐅᑎᖓᓄᑦ ᐊᓚᒪᓗ ᑐᔪᕎᒥᔪᓂᖓᓄᑦ;

(b) ᐃᑲᔪᕐᓗᑎᑦ ᐊᐅᓚᕎᑎᖃᑎᒌᖓᔪᒥᑦ ᑲᑎᒪᔨᕈᓯᑦ ᑭᓇᐅᔭᖅᑕᐃᑎᒃᓴᖓᓄᑦ ᑎᑎᕎᓗᑎᑦ ᐊᑦᑭᒃᓴᓗᑎᑦ;

(c) ᑎᑎᕎᓚᖏᓚᐊᓚᖓᓄᑦ ᐃᑲᔪᕐᓗᑎᑦ ᐊᓚᒪᓗ ᐊᑦᑭᑎᐊᖅᓴᒪᐃᓐᓇᕐᓗᑎᑦ ᐊᐅᓚᕎᑎᖃᑎᒌᖓᔪᒥᑦ ᑲᑎᒪᔨᕈᓯᑦ ᑎᑎᕎᔪᓂᖓᓄᑦ ᐊᓚᒪᓗ ᑐᔪᖅᑐᐃᓂᖓᓄᑦ; ᐊᓚᒪᓗ

(d) ᐊᓴᖓᓄᑦ ᐱᓕᕆᖓᓄᑦ ᑕᐃᒪᐃᓐᓇᑐᖏᓚᔪᑦ ᑐᑭᒧᐊᖅᑎᑕᐅᓗᑎᒃ ᐃᑲᓴᐅᑕᒥᑦ ᐊᓚᒪᓗ ᐊᖓᓕᖅᑕᐅᓴᒪᔪᑦ ᒐᕙᒪᖓᓂ ᑲᑎᒪᔨᕈᓯᑦ ᐅᒪᔪᒋᓗᐊᕐᕕᐊᑯᑦ.

3.2.31 ᐱᔪᒪᑎᓪᓗᓂᑦ ᑲᑎᒪᔨᕈᓯᑦ ᐅᒪᔪᒋᓗᐊᕐᕕᐊᑯᑦ, ᐊᓚᒪᓗ ᐊᐅᑐᑎᔪᑦ ᐊᕐᓇᖅᑐᖅᑕᐅᓴᒪᔪᑦ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑎᕐᑐᖅ ᐃᑲᔪᕐᓗᑎᒃᓴᖓᓄᑦ ᑎᓪᓗᒥᐅᕐᓂᖓᓄᑦ ᐃᖅᑲᐅᖅᑎᑎᕐᑐᖅ ᐃᑲᔪᕐᓗᑎᒃᓴᓗᑎᒃ ᐃᑲᔪᕐᓗᑎᒃ ᒐᕙᒪᖓᓂ ᐊᐅᓚᕎᑎᖃᑎᒌᖓᔪᒥᑦ ᑲᑎᒪᔨᕈᓯᑦ ᑲᑎᒪᓂᖓᓄᑦ ᐃᖅᑲᐅᖅᑎᑎᕐᑐᖅ.ᓴᓪᓕᒡᓗᐅᓴᖅᓂᖓᓪᓕᐊᓂᒃ ᐊᐅᓚᕎᑎᖃᑎᒌᖓᔪᒥᑦ ᑲᑎᒪᔨᕈᓯᑦ ᐊᐅᓚᕎᑎᖃᑎᒌᖓᔪᒥᑦ ᑲᑎᒪᔨᕈᓯᑦ ᐃᓄᐃᑦ ᐊᐅᓚᕎᑎᖃᑎᒌᖓᔪᒥᑦ ᑲᑎᒪᔨᕈᓯᑦ ᐃᓕᖓᕆᐅᓗᑎᒃᓴᖓᓄᑦ ᐊᐅᓚᕎᑎᖃᑎᒌᖓᔪᒥᑦ ᑲᑎᒪᔨᕈᓯᑦ ᐊᓯᖏᓪᓗᑎᒃ

(b) ᐊᐅᓪᓕᔪᑎᑦᓴᐃᓐᓇᓕᒃ ᓴᖅᓄᐅᑎᑦ (ᐃᓕᓐᓂᑦ 3.5 ᑎᑭᑦᓵᑐ 3.7);

(c) ᐊᕕᐊᑐᖅᓯᒪᔪᑦ ***ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑕᑎᑦᓯᓐᓂᑦ*** - ᐃᑲᔪᖅᓯᓯᑐᑦ ᐱᔪᓐᓇᐅᑎᓄᑦ ᐅᐊᑐᑎᑦ (ᐃᓕᓐᓴ 4.3);

(d) ᐋᖅᑕᐅᔪᓐᓇᖅᑐᓴ ᓴᓇᓐᓴᐊᒃᑲᖅᓱᖅ ᐅᐊᓯᓚᑦ ᑕᐃᑲᓴᑦ ᑲᓇᑕᒥ ᐅᒪᔪᖅᑐᖅᓵᐃᑦ ᐊᒪᓪᓗ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᓐᓕᐊᓄᑦ ᓴᐳᔾᔨᐅᓯᒪᕆᐊᒃ (ᐃᓕᓐᓴ 5.4);

(e) ᓄᓇᓕᕋᓱᑦ ᐊᒪᓪᓗ ᐃᒡᓗᕋᓱᑦ ᐃᓗᐊᓂ ᑲᓇᑕᒥ ᐅᒪᔪᖅᑐᖅᓵᐃᑦ ᐊᒪᓪᓗ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᓐᓕᐊᓄᑦ ᓴᐳᔾᔨᐅᓯᒪᕆᐊᒃ (ᐃᓕᓐᓴ 5.5);

(f) ᐱᖁᑎᓐᓕᑦ ᓯᓇᑐᐃᓇᐃᑦ ᐊᒪᓪᓗ ᐱᔭᐸᑎᒃᓴᑦ ᐱᓪᓚᑕᐅᔪᑦ ᐃᓄᓐᓄᑦ, ᐃᓕᑎᔨᐅᓪᓗᑎᒃ ᐃᓪᓚᑲᓪᓕᓐᓂᒋᔪᑦ, ᐃᑕᖁᓂᑕᐃᑦ ᐱᓕᕆᐊᖓᔪᑦ ᐊᒪᓪᓗ ᐃᓄᐃᑦ ᐅᖁᐅᓯᓐᓕᕆᐊᒃ ᓄᓇᐃᑦ ᐊᑎᑐᑎᓐᓕᐊᓄᑦ (ᐃᓕᓐᓂ 6.4 ᑎᑭᑦᓵᑐ 6.7);

(g) ᑲᓇᑕᒥ ᐅᓪᓚᔪᑐᖅᓵᐃᑦ ᐊᒪᓪᓗ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᓐᓕᐊᓄᑦ ᐱᔭᐸᑎᒃᓴᑦ ᖃᐅᔨᓴᖅᑎᓂᖅ (ᐃᓕᓐᓴ 10.2);

(h) ᑲᓇᑕᒥ ᐅᓪᓚᔪᓕᓗᓴᖅᓯᐊᑯᑦ ᐃᓂᓐᓴ ᐱᔭᒥᓗᑎᑦ ᐃᑕᖁᓂᑕᖁᖅᑐᑦ , ᐃᓪᓚᑲᑦᓕᓂᑕᐃᑦ ᐊᒪᓪᓗ ᓂᖁᐊᑎᑦ ᐃᓂᓐᓂᑦ ᐊᒪᓪᓗ ᐅᓪᓚᑦ ᐊᒪᓪᓗ ᐱᐅᔨᑐᖁᖅᑐᖁᖅᓴᑦ ᐃᓂᓐᓂᑦ ᐱᓪᓚᑲᐅᔪᑦ ᐃᓄᓐᓄᑦ (ᐃᓕᓐᓴ 11.3);

(i) ᐊᐅᓕᑕᐅᓂᓐᓂ ᐊᒪᓪᓗ ᐱᔭᒥᔨᐅᓂᓐᓂᑦ ᓂᖁᐊᑎᑦ ᐊᒪᓪᓗ ᓂᔪᐊᑎᑦ ᐃᓂᓐᓂᑦ ᐃᓗᐊᓂ ᑲᓇᑕᒥ ᐅᓇᓚᑦᓵᐃᑦ ᐅᐊᑐᑎᑦ ᐊᒪᓪᓗ ᐱᔭᐸᑎᒃᓴᑦ ᐃᓄᓐᓄᑦ (ᐃᓕᓐᓴ 12.2);

(j) ᓴᖇᑕᐅᓂᓐᓂᑦ, ᐊᓐᑎᓕᑎᐊᖁᓂᓐᓂᑦ, ᖃᓄᐃᓇᑕᓐᓴᓂᓐᓴᑕ ᐊᓯᔪᖁᓂᓐᓂᑦ, ᒥᓴᓂᓐᓂ ᐅᓇᓚᓱᓂᑦ ᐱᓴᖅᑕᐅᓂᓐᓂᑦ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᓐᓕᐊᓄᑦ ᓴᐳᔾᔨᐅᓯᒪᕆᐊᒃ ᐊᒪᓪᓗ ᑲᓇᑕᒥ ᐅᓪᓚᑦᓵᐃᑦ, ᓰᓪᑎᑦᓗᔪ (ᐃᓕᓐᓂ 13.3 ᐊᒪᓪᓗ 13.5); ᐊᒪᓪᓗ

(k) ᐳᓭᖅᑎᓄᑦ ᐊᑐᖅᑕᐅᓂᓐᓂ ᑲᓇᑕᒥ ᐅᓪᓚᔪᑐᖅᓵᐃᑦ ᐊᒪᓪᓗ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᓐᓕᐊᓄᑦ ᓴᐳᔾᔨᐅᓯᒪᕆᐊᒃ, ᐃᓕᑎᔨᐅᓗᑎᒃ ᐊᐅᑦᓕᑭᔾᔨᓯᒪᔪᓐᓇᖅᑐᑦ ᓄᓇᓕᖓᓄᑦ (ᐃᓕᓐᓂ 14.2 ᐊᒪᓪᓗ 14.4).

3.3.5 ᐅᓇ ᒥᓂᔾᑕ ᖃᐅᓂᓂᐊᕐᑐᖅ ᐅᓴᐅᔨᔾᔨᓂᐊᕐᑐᓂᒃ ᐊᐅᑐᑎᔨᓂᑦ ᐊᐅᓕᑦᑎᓐᓂᖅᓴᓂᑦ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᐱᓪᓚᑲᓪᓕᓐᓂᑦ ᐊᒪᓪᓗ

ᒥᓂᑦ ᐃᒪᓐᓇᐅᓂᐊᖅᑐᖅ, ᐃᓗᐊᓂ 60 ᐅᑦᓗᐃᑦ ᐱᔪᓂᐅᖅ ᐅᖅᑲᑎᕐᕕᓯᒪᓕᕆᔨᓗᓂ, ᑎᑎᕋᔨᓗᓂ ᐱᓕᕆᑎᓂᖅ ᑕᕆᔨᓇᓂᖅ. ᑕᐃᒪᓗᒃ ᑎᓂᒃ ᐃᓯᒪᓂᓂᐊᖅᑕᖏᑦ ᓇᓕᐊᑐᐃᓇᖅ ᑐᓴᖅᑕᓂ ᐱᓯᒪᓇᖏᑕᖏᑦ ᐊᐅᓚᑦᓂᖅᑎᑎᕆᔨᓂᖅ ᑲᑎᒪᔨᓂ, ᐅᓇ ᒥᓂᑦ ᐃᒪᓐᓇᐅᓂᐊᖅᑐᖅ, ᒪᓕᒐᓗᓂ ᒪᓕᒐᕋᓂᖅ ᐊᔪᖅᑎᑕᐅᓕᕆᔨᑎᑎᑉᓂᓂ ᕿᓂᕈᑎᓴᒪᐊᖅᑐᖅ, ᕿᓂᕈᑕᓪᓕᐊᑲᖅ ᑐᓴᖅᑎᑕᑎᓴᐅᑲᖅᓂᖅ ᑎᑎᕋᔨᓗᓂᕈᑦ ᐱᔪᓂᐅᖅ;

(b) ᑕᐃᒪᓗᒃ ᒥᓂᑦ ᐊᓯᕈᓯᕐᒪᑦ ᐊᐅᓚᑦᓂᖅᑎᑎᕆᔨᓂᖅ ᑲᑎᒪᔨᓂᓪᓕᖅᑐᖅ ᐅᖅᑲᔨᔭᐃᔨᔪᑎᓯᕈᓯᓂᖅ, ᓄᓇᒥᒃ ᑲᑐᔨᖅᑎᑎᕆᔨᓂᖅᑐᖅ ᑲᑎᒪᔨᓂᓪᓕᖅᑐᖅ ᐃᓕᓴᕆᑦᑎᐊᕆᔨᖅᑐᑎᓪᓗᒍ ᐅᖅᑲᔨᔭᐃᔨᔪᑎᑎᕆᔨᑲᑎᒃ, ᐊᓚᒪᓗ ᐊᔪᖅᑎᑦᑐᖅ, ᐃᓗᐊᓂ 60 ᐅᑦᓗᐃᑦ ᐱᔪᓂᐅᖅ ᒥᓂᑦᑕᐅᐳ ᐱᔨᔪᑎᖅᑎᑦ ᐊᖅᑎᕈᖅᑎᓯᒪᓄᑦ, ᑐᓂᓗᑎ ᐁᖅᑎᕆᐊᖅᓯᒪᔨᑦ ᐅᖅᑲᔨᔭᐃᔨᔪᑎᑎᕆᔨᓂᖅ;

(c) ᒥᓂᑦ ᐃᓯᒪᓕᑎᓂᐊᖅᑕᖏᑦ ᓇᓕᐊᑐᐃᓇᖅ ᐁᖅᑎᕆᐊᖅᑐᑦ ᐅᖅᑲᔨᔭᐃᔨᔪᓯᓕᑦ ᑐᓂᔪᐅᔨᑦ ᑕᕆᔨᓇᓂᖅ ᐊᐅᓚᑦᓂᖅᑎᑎᕆᔨᓂᖅ ᑲᑕᒍᑦ ᑎᑎᕋᔨᓗᓂᕈᑦ (b), ᐊᓚᒪᓗ ᐊᑐᓕᖅᑕᐅᔪᓕᑦ ᐱᔪᖁᖅᓯᔨᑎᑦ ᐃᓯᒪᓕᐊᖅᓯᒪᓕᕆᔨᑦ ᐃᓗᐊᓂ 60 ᐅᑦᓗᐃᑦ. ᒥᓂᑦ ᑎᑎᕋᔨᓗᓂ ᐱᔨᔪᑎᑎᕆᔨᓂᖅ ᕕᕕᓇᖅ ᐊᐅᓚᑦᓂᖅᑎᑎᕆᔨᓂᖅ ᑲᑎᒪᔨᓂᓪᓕᖅᑐᖅ, ᑎᑭᑎᓗᒍ ᕕᓐᓄᑦ ᒥᓂᑦ ᐊᖅᑭᖅᓯᒪᓕᐊᖅᓯᒪᔨᖅᑐᓂᖅ ᐅᖅᑲᔨᔭᐃᔨᔪᑎᑎᕆᔨᓂᖅ ᑲᑎᒪᔨᓂᓪᓕᖅᑐᖅ ᐊᐅᓚᑦᓂᖅᑎᑎᕆᔨᓂᖅ ᐁᖅᑎᕆᐊᖅᓯᒪᔨᖅᑐᖅ ᐅᖅᑲᔨᔭᐃᔨᔪᑎᑎᕆᔨᑲᖏᑦ ᐅᖅᑲᔨᔭᐃᔨᔪᑎᓂᕆᑐᖅ. ᑐᓂᓯᔪᖅᑭᑎᑐᖅ ᕕᓐᓄᑦ ᐊᐅᓚᑦᓂᖅᑎᑎᕆᔨᓂᖅ ᑲᑎᒪᔨᓂᓪᓕᖅᑐᖅ ᐁᖅᑎᕆᐊᖅᓯᒪᓕᕆᔨᖅᑐᖅ ᒥᓂᑦᑕᒍ ᐊᐅᓚᑦᓂᖅᑎᑎᕆᔨᓂᖅ ᑲᑎᒪᔨᓂᓪᓕᖅᑐᖅ, ᑎᑭᑎᓗᒍ ᐊᖅᑭᑎᓂᖅᑎᑦ ᐃᓯᒪᓕᐊᖅᓯᒪᓕᕆᔨᑦ ᐃᓗᐊᓂ 60 ᐅᑦᓗᐃᑦ. ᒥᓂᑦ ᑎᑎᕋᔨᓗᓂ ᐱᔨᔪᑎᑎᕆᔨᓂᖅ, ᑎᑭᑎᓗᒍ ᑐᑭᓯᐅᑎᓗᒍ ᐅᖅᑲᔨᔭᐃᔨᔪᑎᑎᕆᔨᓂᖅ ᑲᑎᒪᔨᓂᓪᓕᖅᑐᖅ ᐁᖅᑎᕆᐊᖅᓯᒪᓕᕆᔨᖅᑐᖅ ᐅᖅᑲᔨᔭᐃᔨᔪᑎᑎᕆᔨᑲᖏᑦ ᐅᖅᑲᔨᔭᐃᔨᔪᑎᓂᕆᑐᖅ. ᑐᓂᓯᔪᖅᑭᑎᑐᖅ ᕕᓐᓄᑦ ᐊᐅᓚᑦᓂᖅ

3.4.3 ᐱᓕᕆᓂᐊᑎᒋᔭᓂᓗᑎᒃ ᐊᖏᑭᒃᓯᓯᒪᔨᕐᓂᕆᖏᓐᓄᑦ ᐊᒻᒪᓗ ᐱᒋᓐᓇᐊᖏᓐᓄᑦ <ᖏᓐᐅᐅᑎᑦ, ᑲᓇᑕᒥ ᐅᒪᔨᒋᓐᓇᖅᖏᐊᕐᒋᑦ ᐃᓕᓴᕐᑎᐊᖏᓂᖅᓯᐊᖏᑕᖅᓯᑦ *ᐃᓇᐃᑦ ᖃᐅᔨᓴᕐᓂᖅᓯᑦ* ᑎᑎᕋᖏᓐᓄᒃᓴᓕᔨᑦ ᐊᒻᒪᓗ ᑐᵒᓂᖏᑕᐅᓴᓕᔨᑦ ᑕᒃᔪᓇᓐᖓ ᐊᐅᓪᓕᓂᕆᑎᐅᓐᓂᖏᓂᑦ ᑲᑎᒪᔨᑦᑕᐅᑎᒃ ᐱᓕᒪᓐᓇᖅᑐᑦ.

3.4.4 ᑲᓇᑕᒥ ᐅᒪᔨᓕᓚᔨᕐᓂᖏᑦ ᑐᵒᓂᖏᑎᑎᓂᐊᖏᑕᖅᓯᖓ ᐅᒃᑐᑎᑎ ᐊᖏᑭᒃᓯᓯᒪᔨᕐᓂᕆᖏᓐᓄᖅᓯᖅ ᐊᒻᒪᓗ ᐱᓕᕆᓂᐊᑎᒋᔭᓂᓗᑎᒃ <ᖏᓐᐅᐅᑎᑎ ᒥᓂᖏᑕᒪᑦ ᐊᖓᕐᖏᖅᑕᐅᓂᐊᖏᓪᓕᖅᓯᑦ. ᐅᓕᐊᑐᐃᓐᓇᖅᓯ ᓯᕐᐳᓂᕆᖏᓐᓄᑦ ᑲᑎᒪᔨᑦᑕᐅᑎᒃ ᒪᓕᓐᓂᖏᓐᓄᑦᑐᖅᓯᖅ ᐃᖓᕐᓯᓯᓕᕐᑕᐅᓂᐊᖏᓪᓕᖅᓯᖅ ᒪᓕᓐᓂᖏᓐᓄᑦ ᐊᖏᑭᒃᓯᓯᒪᔨᕐᓂᕆᖏᓐᓄᑦ ᐊᒻᒪᓗ ᐱᓕᕆᓂᐊᑎᒋᔭᓂᓗᑎᒃ <ᖏᓐᐅᐅᑎᓄᑦ.

3.4.5 ᑲᓇᑕᒥ ᐅᒪᔨᓕᓚᓂᖏᑦ ᐱᖏᓂᔪᔨᐊᖅᓯ ᐊᖏᑭᑎᐊᑐᐃᓐᓇᖅᑕᖏᑦ ᐊᖏᑭᒃᓯᓯᒪᔨᕐᓂᕆᖏᓐᓄᑦ ᐊᒻᒪᓗ ᐱᓕᕆᓂᐊᑎᒋᔭᓂᓗᑎᒃ <ᖏᓐᐅᐅᑎᑎ ᒪᓕᓚᓂᖏᓐᓄᒃ ᐱᓕᕆᓂᐊᑎᒋᔭᓂᓗᑎᒃᓴᓂᖏᓐᓄᑦ ᑎᑎᕋᖏᓐᓄᑦᑐᖅᓯᖅ ᐊᖓᕐᖏᖅᑕᐅᓂᐊᖏᓪᓕᖅᓯᖅ 3.4.1 ᑎᑭᑦᓯᔭᒍ 3.4.4-ᒧᑦ.

3.4.6 ᑭᓇᐅᔭᓄᖓᓄᑦ ᐊᑐᖅᑕᐅᓂᐊᖏᓐᓄᑦ ᐊᑐᓚᐊᖏᓄᖅᑕᑦ ᓇᓚᐅᐃᖅᓯᒪᔨᖅ ᐅᓂᓂ ᐃᓕᓐᓄᑦ ᐱᓕᕆᓂᐊᑎᒋᔭᓂᓗᑎᒃᓴᓂᖏᓐᓄᑦ ᑕᔪᓂᓐᓄᑦ ᐱᐅᓂᖏᓐᓄᑦᑐᖅᓯᖅ ᐃᓕᓴᕐᑎᐊᖏᓂᖅᓯᐊᖏᓪᓕᖅᓯᖅᓯᑦᓗ ᐊᖓᕐᖏᖅᑕᐅᓂᐊᖏᓪᓕᖅᓯᖅ ᐱᓂᓕᕆᔨᖏᑦᑐᖅᓯᖅ ᐊᑐᖅᑕᐅᓂᐊᖏᓪᓕᖅᓯᖅ ᑭᓇᐅᔭᓄᖓᓄᑦ ᐊᑐᖅᑕᐅᓂᐊᖏᓪᓕᖅᓯᖅ ᐊᖓᕐᖏᖅᑕᐅᓂᐊᖏᓪᓕᖅᓯᖅ ᐱᓂᓕᕆᔨᖏᑦᑐᖅᓯᖅ ᐊᑐ

3.5.7 ᐅძᐊ ᐊᐅᓪᓚᕐᓂᐊᕐᐳᖅᓄᑦ ᐸᖅᓇᐅᑎᑦ ᐃᓪᓚᖅᑎᓂᐊᖅᑐᑦ ᓇᓗᓇᐃᔪᕐᑎᐊᖅᓯᒪᔪᓂᒃ ᐅძᓂᖏᓐ:

(a) ᐱᔪᔾᑎᕐᒥᑦ ᑲᓇᑕᒥ ᐅᒪᔪᖅᑭᐹᑦ ᐅᕐᓂᒃᑑᑦ ᐅᑎᖅᑕᐸᖅᑐᓄᑦ ᑎᓐᓕᐊᓄᑦ ᓴᓗᐅᒪᔪᖅᓯᒪᔭᕐᓂᒃ;

(b) ᐊᐅᓪᓚᕐᔪᐸᑕᐅᔪ ᑐᖏᓐᓇᓐᒥᑦ ᐊᒻᒪᓗ ᑐᑭᓯᐅᑕᐅᑎᑕᕐᓂᓕᕆᓂᐊᖅᑐᑦ;

(c) ᓄᓇᖏᓐ ᐊᒻᒪᓗ ᐱᐅᓯᑐᖅᓇᓐᓂᑕ ᐃᓕᒪᑦᓂᓪᑲ ᖃᑲᓄᐃᓕᓐᓇᓂᖏᓐ ᐊᒻᒪᓗ ᐃᓗᓕᓐᒥᑦ ᐃᓗᐊᓂ ᑕᑦძᐊ ᑲᓇᑕᒥ ᐅᒪᔪᖅᑭᐹᑦ ᐅᖅᑕᐸᖅᑐᓄᑦ ᑎᓐᓕᐊᓄᑦ ᓴᓗᐅᒪᔪᖅᓯᒪᔭᕐᓂᒃ ᐊᐅᓚᵅᓂᖏᓐᓂᑦ;

(d) ᐊᑐᐊᖏᐃᑦ ᐅძᐊᓗ ᑐᑭᒧᐊᔪᑕᐅᓂᐊᖅᑐᑦ ᐊᐅᓪᓚᕐᓂᐱᔾᔪᑕᐅᔾᔨᓄᑦ ᑕᑦძᓂᖏᓐ ᑲᓇᑕᒥ ᓂᔾᔪᐊᑎᓄᑦ ᑲᓇᑕᒥ ᐅᒪᔪᖅᑭᐹᑦ ᐅᖅᑕᐸᖅᑐᓄᑦ ᑎᓐᓕᐊᓄᑦ ᓴᓗᐅᒪᔪᖅᓯᒪᔭᕐᓂᒃ;

(e) ᐅᓪᓗᖏᓐᓂ ᐊᑐᓕᖅᑎᑕᐅᕆᓚᖅᓴᖏᓐᓂ ᐊᐅᓪᓚᕐᓂᐱᔾᔪᑕᐅᔾᔨᓄᑦ ᐸᖅᑕᐸᖅᑐᖏᓐᓂᑦ ᖃᓄᐃᓕᐅᕕᖅᑕᐅᑎᑕᕐᓂᒃ ᑭᓯᓂᖏᓐᓂᑦ; ᐊᒻᒪᓗ

(f) ᑭᓄᖏᓐᓂ ᓇᓗᓇᐃᔪᖅᓯᒪᔪᓕᒃ ᐊᖏᑭᑎᑎᓂ 14.2.2 ᐊᒻᒪᓗ 14.4.1.

3.5.8 ᑕᐃᒃᑐᓂ ᑲᓇᑕᒥ ᐅᒪᔪᖅᑭᐹᑦ ᐅᕐᓂᒃᑑᑦ ᐅᑎᖅᑕᐸᖅᑐᓄᑦ ᑎᓐᓕᐊᓄᑦ ᓴᓗᐅᒪᔪᖅᓯᒪᔭᕐᓂᒃ ᐃᓗᐊᓄᑦ ᓄᓇᖅᑯᐊᖏᓐᓂᑦ, ᐊᐅᓪᓚᕐᓂᐱᔾᔪᑕᐅᔾᔨᓄᑦ ᐸᖅᓇᓐᓂᑦ ᒪᑕᕐᓂᐊᕐᓂᐊᖅᓯᒪᔭᕐᓂᒃ ᐊᒻᒪᓗ ᐅᖅᑲᖅᓯᒪᔭᓗᑎᒃ ᓇᓕᓕᐊᖏᓐᓇᑐᓂᓗᐃᑦᓇᓱᒃ ᐱᔾᔪᑕᐅᔾᔨᓂᑦ ᓴᖅᓂᑐᑦ ᐃᓗᐊᓂ ᐃᓄᐃᑦ ᓄᓇᖅᑯᐊᖏᓐᓂᑦ.

3.5.9 ᑕᐃᒃᑐᓂ ᐅᓇ ᐊᐅᓪᓚᕐᓂᐱᔾᔪᑕᐅᔾᔨᖅ ᐸᖅᑕᐅᑎ ᐱᑕᖅᑲᓱᖅᑐᖅ ᑕᑦძᐊᓄᖓᓐ ᑲᓇᑕᒥ ᐅᒪᔪᖅᑭᐹᑦ ᐅᖅᑕᐸᖅᑐᓄᑦ ᐅᕐᓂᒃᑑᑦ ᐊᐅᓪᓚᕐᓂᐱᔾᔪᑕᐅᔾᔨᓄᑦ ᑎᓐᓕᐊᓄᑦ ᓴᓗᐅᒪᔪᖅᓯᒪᔭᕐᓂᒃ, ᐅძᐊ ᐊᐅᓪᓚᕐᓂᖅᓱᑎᑲᓂᖅᓯᓕᕐᒪᑦ ᑲᑎᒪᔨᖏᑦᑑᒃ ᖃᑭᖅᑭᐅᓂᐊᖅᑕᕐᓂᑦ ᐅძᐊ ᐊᐅᓪᓚᕐᓂᐱᔾᔪᑕᐅᔾᔨᓄᑦ ᐸᖅᓇᓐᓂᑦ ᐃᒪᓐᓇᕐᓂᖅᑐᖅᓯᒪᔭᕐᓂᒃ, ᒪᑕᓚᑕᑦ ᐃᑕᖓᒃ 3.6, ᐊᑐᓕᖅᑕᐅᖅᑯᓯᐅᔭᕐᓂᒃ ᐋᖅᑭᑎᐊᖅᓗᓂᒃ ᐅᖅᑕᐸᓐᓂᑦ ᐃᓐᖏᑦᓗᓂᒃ ᐅᓇ ᐸᖅᑕᐅᑎ ᐅᓴᒪᑦᑐᒪᑦ.

3.6 ᐊᐅᓪᓚᕐᓂᐱᔾᔪᑕᐅᔾᔨᓄᑦ ᐸᖅᑕᐅᑎᑦ ᐊᖏᑭᖅᑕᒪᔨᐊᖅᑎᓂᖓᓐᓄᓪᓗ

3.6.1 ᐊᐅᓪᓚᕐᓂᖅᑲᑎᕐᓂᖓᓐᖏᑦᑐᒥᑦ ᑲᑎᒪᔨᖏᑦᑑᒃ ᐊᑐᓕᖅᑕᐅᔾᔨᓕᕐᓂᒃᑑᑦ ᐱᔭᕆᓂᐊᖅᓯᒪᔪᓂ ᐊᐅᓪᓚᕐᓂᐱᔾᔪᑕᐅᔾᔨᓄᑦ ᐸᖅᑕᐅᑎᑦ ᑕᑦძᓂᖏᓐ ᓄᓇᕗᒻᒥ ᐅᒪᔪᖅᑭᐹᑦᑐᓂᓗᐃᑦᓇᓱᐊᖅᑐᓂᖅ ᐊᖏᑭᖅᑕᒪᔨᐊᖅᑎᓂᖓᓐᓄᓪᓗ ᑲᑎᒪᔨᖏᑦᑑᒃ ᐊᐅᓪᓚᕐᓂᐱᔾᔪᑕᐅᔾᔨᓄᑦ ᐸᖅᑕᐅᑎᑦ ᐅძᐊᓗ ᐊᐅᓪᓚᕐᓂᐱᔾᔪᑕᐅᔾᔨᓄᑦ ᓄᓇᕗᒻᒥ ᐅᒪᔪᖅᑭᐹᑦᑐᓂᓗᐃᑦᓇᓱᐊᖅᑐᓂᖅ. *ᓄᓇᕗᒻᒥᒃ ᓄᓇᕕᑕᐅᓂᕐᓂᒃ ᐊᖏᑭᖅᑕᒪᔨᐊᖅᑎᓂᖓᓐᓄᓪᓗ ᐅᒪᔪᖅᑭᐹᑦ* 5.2.34 (c) ᐊᒻᒪᓗ 5.3.16.

3.6.2 ᐅძᐊ ᐊᐅᓪᓚᕐᓂᖅᑲᑎᕐᓂᖓᓐᖏᑦᑐᒥᑦ ᑲᑎᒪᔨᖏᑦᑑᒃ ᑐᓴᕆᓂᐊᖅᑐᑦ ᐊᔪᖅᑐᖅᑭᐹᑦ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑲᑎᕐᓂᖓᓐᖏᑦᑑᒃ ᐊᐅᓪᓚᕐᓂᐱᔾᔪᑕᐅᔾᔨᓄᑦ ᐸᖅᑕᐅᑎᑦ ᐅძᐊᓗ ᐊᐅᓪᓚᕐᓂᐱᔾᔪᑕᐅᔾᔨᓄᑦ ᓄᓇᕗᒻᒥ ᐅᒪᔪᖅᑭᐹᑦᑐᓂᓗᐃᑦᓇᓱᐊᖅᑐᓂᖅ.

3.6.3 ᑕᐃᒪᓗ, ᒪᑕᓚᑐᒃ ᐃᓕᓴᕐᐅᓚᐅᕐᓂᐊᔪᒃᑑᑦ ᐅᖅᑲᖅᓯᒪᔪᓂ ᐊᖏᑭᖅᑕᒥ 5.2.34(c) ᐊᒻᒪᓗ 5.3.17 ᑎᑭᓪᓚᑐ 5.3.23 *ᓄᓇᕗᒻᒥᒃ ᓄᓇᕕᑕᐅᓂᕐᓂᒃ* ᐊᖏᑭᖅᑕᒪᔨᐊᖅᑎᓂᖓᓐᓄᓪᓗ, ᐅძᐊ ᓄᓇᕗᒻᒥ ᐅᒪᔪᖅᑭᐹᑦᑐᓂᓗᐃᑦᓇᓱᐊᖅᑐᓂᖅ ᐅᕐᓂᒃᑑᑦ ᒥᓴᑕ ᐊᒃᑲᖅᐸᕐᓂᑦ, ᑕᒪᔾᒥᑦ ᐅᕐᓂᒃᑑᑦ ᐃᓕᓴᕐᐅᓚᐅᕐᓂᐊᔪᒃᑑᑦ ᐱᔭᕆᓂᐊᖅᓯᒪᔪᓂ ᐊᐅᓪᓚᕐᓂᐱᔾᔪᑕᐅᔾᔨᓄᑦ ᐸᖅᑕᐅᑎᑦ ᐊᒻᒪᓗ ᐸᖅᑕᐅᑎᓄ ᐅᑎᖅᑐᓂ ᑲᑎᒪᔨᖏᑦᑑᒃ ᐊᐅᓪᓚᕐᓂᐱᔾᔪᑕᐅᔾᔨᓄᑦ ᓄᓇᕗᒻᒥ ᐅᒪᔪᖅᑭᐹᑦᑐᓂᓗᐃᑦᓇᓱᐊᖅᑐᓂᖅ ᐃᓕᓴᕐᐅᓚᐅᕐᓂᐊᔪᒃᑑᑦ ᐱᔭᕆᓂᐊᖅᓯᒪᔪᓂ ᐊᐅᓪᓚᕐᓂᐱᔾᔪᑕᐅᔾᔨᓄᑦ ᐸᖅᑕᐅᑎᑦ ᐊᒻᒪᓗ ᑐᓴᕆᓂᐊᖅᑐᑦ ᓄᓇᕗᒻᒥ ᐅᒪᔪᖅᑭᐹᑦᑐᓂᓗᐃᑦᓇᓱᐊᖅᑐᓂᖅ.

3.6.4 ᒪᓕᒐᔪᒍ 8.4.13 *ᓄᓇᕗᑦᑐᓄᑦ ᐊᖏᕈᑎᓄᑦ*, ᐅᓴᒪᒋᔭᐅᔨᑦ ᐊᐅᓚᓂᕆᔨᐸᑎᒃᓴᓄᑦ ᕿᖅᓇᐅᑎᑦ ᒪᓕᒃᖁᓂᐊᖅᑐᒃ ᐊᑐᓂᖅᑯᔭᐅᔨᓂᒃ ᐊᐅᓚᓂᑎᖃᑎᒋᖓᓂᖅᓯᑦ ᑲᑎᒪᔨᕆᓚᑦ, ᐊᑐᖅᓗᑎᑦ ᐊᑐᓂᖅᑯᔭᐅᔨᑦ ᐱᔨᒪᔨᓄᑦ ᐊᓯᒪᓂᖅᓄᑦ ᐅᖃᐅᔾᔨᒌᑦ ᑎᒥᐅᔨᓄᑦ.

3.6.5 ᑕᐃᒪᓗ ᒥᔅᓯ ᐊᖏᕐᕋᔪ ᐊᐅᓚᓂᑎᔨᐸᑎᒃᓴᓄᑦ ᕿᖅᓇᐅᑎᑦ, ᒥᔅᓯ ᐱᑎᐊᖅᑎᖓᔪᐊᖅᑕᖏᑲ ᑲᔪᓯᕆᓇᐊᓕᑦ ᑕᒃᑯᐊ ᐊᑐᖅᑎᒃᖅᑐᑦ ᐊᑐᓕᖅᑕᐅᑎᐊᓕᑦ ᕿᖅᓇᐅᑎᑦ.

3.6.6 ᐅᑯᐊ ᐊᐅᓚᓂᖅᑎᑦᓯᓂᖅᓯᑦ ᑲᑎᒪᔨᕆᓕᑦ ᑐᓂᓯᓂᐊᖅᑐᑦ ᑕᕐᕉᓚᓂ ᐊᖏᕐᕋᐅᔨᓯᓪᓚᕐᒥᒃ ᐊᐅᓚᓂᑎᔨᐸᑎᒃᓴᓄᑦ ᕿᖅᓇᐅᑎᑦ ᑕᒃᑯᓇᖏᓪ ᓄᓇᕗᑦ ᑐᓴᕐᕋᓚᒃ ᑎᒥᖏᓪ ᐊᓱᒪᓗ ᐊᒃᑐᑎᔨᑦ ᐊᕐᓚᑐᖅᑭᔪᓕᑦ ᐃᓄᐃᑦ ᑲᑐᔾᕋᖅᑭᑎᒃᓴᑦ.

3.7 ᐋᖅᑭᒪᐊᑭᑎᔨᑦ ᐊᐅᓚᓂᑎᔨᐸᑎᒃᓴᓄᑦ ᕿᖅᓇᐅᑎᓄᑦ

3.7.1 ᑌᓐᒪᒃᑯᑦ, ᑎᓕᕆᐅᓪᓚᔨᑦ ᐃᓄᐃᑦ ᐊᐅᓚᓂᓪᓗᒋᑦ, ᐅᖃᓕᓵᔪᓂᒃ ᑭᓇᑐᐃᔪᓇᖅ ᐱᓕᕆᓚᒪᔨᖅ ᐊᑲᖅᑕᐅᔨᔨᑦ ᑕᒪᑐᒍᓐᓇᑦ ᐊᐅᓚᓂᑎᔨᐸᑎᒃᓴᓄᑦ ᕿᖅᓇᐅᑎᑦ ᐋᖅᑭᒪᐊᑭᑎᒃᓴᓪᓚᕆᓇᖅᑐᑦ ᐊᐅᓚᓂᑎᔨᐸᑎᒃᓴᓄᑦ ᕿᖅᓇᐅᑎᑦ ᑐᓴᕐᕋᓚᒃ ᐊᐅᓚᓂᑎᖅᑯᑎᒃᖅᑐᓂᖅᓯᑦ ᐅᖃᐅᔾᔨᒌᑦ.

3.7.2 ᐅᑯᐊ ᐊᐅᓚᓂᖅᑎᑦᓯᓂᖅᓯᑦ ᐅᖃᒪᒃᑯᐊᖅᑎᑦᓯᓂᖅᓯᑦ ᐃᓕᓴᕐᓂᐊᖅᑎᑦᓯᓂᖅᑐᓂᑦ ᑐᓴᕐᕋᓚᒃᑯᑦ ᐋᖅᑭᒪᐊᑭᑎᒃᓴᓪᓚᕆᓇᖅᑐᓂᑦ ᐋᖅᑭᒪᐊᑭᑎᒃᓴᓪᓚᕆᓇᖅᑐᓂᒃ ᐊᐅᓚᓂᑎᔨᐸᑎᒃᓴᓄᑦ ᕿᖅᓇᐅᑎᑦ ᒪᓕᒐᓂᐊᖅᑐᒃ ᐱᓕᕐᓂᐊᓂᕐᓂᒃᓴᓂᒃ ᑎᑎᕋᖅᑭᒪᔨᖅ ᐃᓕᓴᕐᓂᐊᖅᑐᑦ 3.5 ᐊᓚᒪᓗ 3.6.

ᐊᐅᓪᓗᖅᑎᑎᕆᔨᓂᕐᒧᑦ ᑲᑎᒪᔨᖏᓪᓗᒃ ᑎᑎᕋᖅ 3-1 (ᐊᖏᕐᕈᑎᓂ 3.2.1 ᐊᓚᒪᓗ 3.2.2)

ᑲᓇᑕᒥ ᐅᒪᔪᖅᑭᒃᑲᓂᕐ ᐊᓚᒪᓗ ᐅᑎᖅᑕᐸᖅᑐᓗᓂᕐ ᑎᖏᒥᐊᓂᕐ ᓴᓗᐅᓕᒪᓐᓇᒃ	ᖃᓄᕆᔨᖏᑦ ᓄᓇᓕᒃ	ᐊᐅᓪᓗᖅᑎᑎᕆᔨᓂᕐᒧᑦ ᑲᑎᒪᔨᖏᓪᓗᒃ
1. ᐊᒃᐸᐃᑦ ᑲᓇᑕᒥ ᐅᒪᔪᖅᑭᒃᑲᓂᕐ ᐊᓚᒪᓗ ᖃᖅᑯᓂᓗᐃᑦ ᑲᓇᑕᒥ ᐅᒪᔪᖅᑭᒃᑲᓂᕐ	ᖃᓄᖅᑕᖅᓴᐊᖅ	ᓯᓚᓐᓕᓕᒃ
2. ᓴᓐᓂᕐᑦ ᐅᑎᖅᑕᐸᖅᑐᓗᓂᕐ ᑎᖏᒥᐊᓂᕐ ᓴᓗᐅᓕᒪᓐᓇᒃ	ᒥᖅᑎᒪᒃᑕᓐᓄᒃ	ᐊᓯᖏᓐᓄᓯᖏᓐᓇᖅ
3. ᐃᓯᓕᓐᓴᖅᓂᒃ ᐅᑎᖅᑕᐸᖅᑐᓗᓂᕐ ᑎᖏᒥᐊᓂᕐ ᓴᓗᐅᓕᒪᓐᓇᒃ	ᑭᖏᓐᓄᐃᑦ	ᐃᓯᓕᓴᓂ
4. ᖃᖅᖏᓴᐅᖅᑐᑉ ᐅᑎᖅᑕᖅᐸᖅᑐᓗᓂᕐ ᓴᓗᐅᓕᒪᓐᓇᒃ ᐊᓚᒪᓗ ᐃᒃᓯᑐᐊᖅ ᐅᑎᖅᑕᖅᐸᖅᑐᓗᓂᕐ ᑎᖏᒥᐊᓂᕐ ᓴᓗᐅᓕᒪᓐᓇᒃ	ᓴᓕᓪᓕᒃ	ᐃᖃᓂᐅᖃᐃᑦ
5. ᓂᖏᖅᓴᓂ ᑲᓇᑕᒥ ᐅᒪᔪᖅᑭᒃᑲᓂᕐ	ᑲᖏᖅᓵᔨᓕᐱᒃ	ᓂᖏᖅᓴᓂᖅ
6. ᑯᐃᔪᒃ ᐅᑎᖅᑕᐸᖅᑐᓗᓂᕐ ᑎᖏᒥᐊᓂᕐ ᓴᓗᐅᓕᒪᓐᓇᒃ	ᐊᖃᕕᐊᑦ	ᓂᓗᕕᐊᓕᒃ
7. ᓂᖅᔪᑎᖅᑭᕕᒃ ᑲᓇᑕᒥ ᐅᒪᔪᖅᑭᒃᑲᓂᕐ	ᐊᐅᓯᐃᑦᑐᖅ	ᓂᖅᔪᑎᖅᑭᕕᒃ
8. ᑐᒃᑐᓕᐊᖅᕕᐅᑉ ᓇᖅᓴᖅᓴᑌ ᑲᓇᑕᒥ ᐅᒪᔪᖅᑭᒃᑲᓂᕐ ᐊᓚᒪᓗ ᐊᒃᐸᑦ ᐅᑎᖅᑕᐸᖅᑐᓗᓂᕐ ᑎᖏᒥᐊᓂᕐ ᓴᓗᐅᓕᒪᓐᓇᒃ	ᖃᖅᐅᓯᐃᑦᑐᖅ	ᓯᓗᒃᖆᐃᑦ
9. ᐃᓗᐃᑦᓕᓐᓄᒃ ᐅᑎᖅᑕᐸᖅᑐᓗᓂᕐ ᑎᖏᒥᐊᓂᕐ ᓴᓗᐅᓕᒪᓐᓇᒃ	ᐃᖃᓗᒃᑐᑦᑎᐊᖅ, ᐅᖅᓯᓄᖅᑐᖅ, ᐅᒥᖓᓕᖅᑐᖅ	ᐊHᐃᐊᒃ

ᐃᓕᓐᓴ 4 — ᐃᓄᐃᑦ ᓄᓇᕗᑎᓐᓂᑦ

4.1 ᑐᑭᒃᓯᑦᓕᐊᔨᐸᑎᒃᓴᑦ

4.1.1 ᐅᖏᓂ ᐃᓕᓐᓴᓂ ᐊᑕᓂ ᑎᑎᕆᒃᓭᒪᔪᑦ ᑐᑭᒃᓯᑦᓕᐊᔨᐸᑎᒃᓴᓐᓂᑦ:

(a) ᐊᑦᑭᑦᑎᐊᖅᓭᒪᑏᓐᓇᓗᑦᓂ ᓄᓇᒥᐅᑕᐃᑦ ᐱᔨᔪᑎᒃᓴᐃᑦ ᐊᖓᓂᖅᓇᖅᑐᑦ ᐃᓄᐃᑦ ᓄᓇᕗᑎᓐᓂᑦᓂ ᐅᑯᐊᓗ ᐅᑎᖅᑕᐅᔪᓐᓇᖅᑐᓂᒃ ᐊᓚᒪᓗ ᑲᑎᒪᔨᓂᒃ ᐱᖓᓱᐅᓭᒪᓕᖅ ᐊᓚᒪᓗ ᑲᑎᒪᔨᓂᒃ ᐃᒪᓕᕆᔨᒃᑯᑦ;

(b) ᐃᓕᓴᕆᓗᓂᑦ ᐊᑐᓂ ᐃᓂᓐᓂᑦ ᐊᓚᒪᓗ ᐱᖅᓴᓐᓂᑦ ᐊᕐᓇᑦᑐᖅᓭᒪᔪᑦ ᐃᓄᐃᑦ ᑲᑐᔪᖅᑎᑎᒋᓐᓂᑦ ᐊᓚᒪᓗ ᒥᓵᑦ ᐊᐅᓪᓚᐅᐅᓂᖓᓂᓄᑦ ᐃᓄᐃᑦ ᓄᓇᕗᑎᓐᓂᑦ ᐃᓗᐊᓂᑦ ᐅᑎᖅᑕᐅᔪᓐᓇᖅᑐᓂᒃ ᑎᓐᒥᐊᓄᑦ ᐱᖓᓱᐅᓭᒪᓕᖅ ᐊᓚᒪᓗ ᑲᑎᒪᔨᓂᒃ ᐃᒪᓕᕆᔨᒃᑯᑦ;

(c) ᓴᓚᒪᑐᑐᓄᑦ ᐊᓚᒪᓗ ᐊᔪᕈᒥᒃ ᐊᑦᑭᒃᑲᑦᑕᖅᓗᑎᓄᑦ ᐊᐃᖏᐅᑎᑎᕆᔭᓄᑦ ᐊᖏᒡᑕᓂᑦ ᐊᐅᓚᑦᑐᖅᓭᒪᔪᑦ ᐃᓄᐃᑦ ᑲᑐᔪᖅᑎᑎᒋᓐᓂᑦ ᐊᓚᒪᓗ ᑲᑎᒪᔨᓂᒃ ᐃᒪᓕᕆᔨᒃᑯᑦ ᐊᓚᒪᓗ ᐅᑎᖅᑕᐅᔪᓐᓇᖅᑐᓂᒃ ᐃᓄᐃᑦ ᓄᓇᕗᑎᓐᓂᖅᓂᑦ ᐃᓗᐊᓂᑦ ᐱᖓᓱᐅᓭᒪᓕᖅ ᐊᓚᒪᓗ ᐅᑎᖅᑕᐅᔪᓐᓇᖅᑐᓂᒃ ᐱᖓᓱᐅᓭᒪᓕᖅ ᐊᓚᒪᓗ ᑲᑎᒪᔨᓂᒃ ᐃᒪᓕᕆᔨᒃᑯᑦ;

(d) ᑐᓴᐅᒪᑎᑕᐅᖏᑦᓗᑎᓄᒃ ᐃᓄᐃᑦ ᑕᐃᑲᓂ ᑲᑎᒪᔨᓂᒃ ᐃᒪᓕᕆᓚᔪᓄᑦ ᐱᓕᕆᑎᕐᓂᑦ ᐱᓇᓱᐊᖏᑦᑐᑦ, ᐃᖅᑲᓇᐃᓕᖏᑦᑐᑦ ᐊᓚᒪᓗ ᑲᒪᒋᔭᖏᓂᒃ ᐃᓱᔪᓂᐊᖏᑦᑐᓕᕆᓂᑦ ᐃᓄᐃᑦ ᓄᓇᕗᑎᓐᓂᖓᓄᑦ; ᐊᓚᒪᓗ

(e) ᐃᓱᖏᓕᖅᑲᑎᐊᖏᖅᓗᑎᓄᒃ ᓇᓂᑐᓇᖅ ᑲᑎᒪᔨᓂᒃ ᐃᒪᓕᕆᔨᒃᑯᑦ ᐊᓚᒪᓗ ᐅᑎᖅᑕᐅᔪᓐᓇᖅᑐᓂᒃ ᑎᓐᒥᐊᓄᑦ ᓴᓚᒪᓗ ᑲᑎᒪᔨᓂᒃ ᑕᒪᐃᓐᓇᖅᑐᑦ ᐃᓄᐃᑦ ᓄᓇᕗᑎᓐᓂᖓᓄᑦ.

4.2 ᑐᑭᓐᓂᑦ

4.2.1 ᐱᔨᔪᑎᑎᑦᓗᒍ ᐅᓇ ᐃᓕᓐᓴᓂ:

(a) “ᐊᕐᓇᑦᑐᖅᓭᒪᔪᑦ ᐃᓄᐃᑦ ᑲᑐᔪᖅᑎᑎᒋᓐᓂᑦ - ᐃᑲᔪᖅᓭᓕᒃᑯᑦ ᐱᔪᓐᓇᐅᑎᓄᑦ ᐅᖃᓕᓐᓂᖓᓄᑦ” ᑐᑭᖅᑲᖅᑐᖅ ᐅᓂᐊ ᐅᖃᓕᑎᑦ ᑕᑕᑎᖅᓭᒪᔪᑦ ᑭᒃᓂᓂᐊᖅᑐᑦ, ᐱᓇᓵᑦᓂᑦ ᐅᕐᓚᓕᓐᓂᒃ ᐃᑲᔪᖅᑕᐅᔭᓗᑎᓄᒃ ᑎᑎᕆᔭᓗᑎᓄᒃ ᑭᒃᓂᐊ ᐊᕐᓇᑦᑐᖅᓭᒪᔪᑦ ᐃᓄᐃᑦ ᑲᑐᔪᖅᑎᑎᒋᓐᓂᑦ, ᑭᒃᓂᐊᓂᒃ ᑲᑎᒪᔨᓂᒃ ᐅᕐᓚᓕᓐᓂᒃ ᐅᑎᖅᑕᐅᔪᓐᓇᖅᑐᓂᒃ ᐱᖓᓱᐅᓭᒪᓕᖅ ᐱᔪᓐᓇᐅᑎᑦ ᐱᓕᕆᓚᐅᖅᓯᒪᔪᑦ ᐃᓄᐃᑦ ᓄᓇᕗᑎᓐᓂᖓᓄᑦ ᐃᓗᐊᓂᑦ ᐱᖓᓱᐅᓭᒪᓕᖅ ᐊᓚᒪᓗ ᐅᑎᖅᑕᐅᔪᓐᓇᖅᑐᓂᒃ ᐱᖓᓱᐅᓭᒪᓕᖅ ᐊᓚᒪᓗ ᑲᑎᒪᔨᓂᒃ ᐃᒪᓕᕆᔨᒃᑯᑦ ᑕᐃᑲᓂᓗ, ᒪᓕᒃᑐᑦ 9.3.3 ᐅᕐᓚᓕᓐᓂᖓᓄᑦ 9.3.4 *ᓄᓇᕘᒥᒃ ᓄᓇᑕᐱᑎᓄᑦ ᐊᖏᓪᓗᑎᑦ*, ᖃᑲᓇᐃᓕᖏᑦᑕᐅᔪᓐᓇᖅᑐᖅ ᐱᔪᓐᓇᐅᑎᖃᑎᖃᖅᑎᓕᓐᓇᓗᑎᒍ ᐊᕐᑕᒍ *ᑲᑎᒪᔨᓂᒃ ᐃᒪᓕᕆᓚᐅᖅᓴᒍ ᐱᖃᓐᓇᖅᑐᓄᑦ* ᐅᕐᓚᓕᓐᓂᖓᓄᑦ *ᑎᑎᕆᐊᖅᑐᑦ; ᐱᖃᓐᓇᖅᑐᓄᑦ* ᐊᖏᓪᓗᑎᑦ ᐅᑎᖅᑕᐅᔪᓐᓇᖅᑐᓂᒃ ᐱᔪᓐᓇᐅᑎᓄᑦ ᐊᓚᒪᓗ

(b) “ᐊᕐᓇᑦᑐᖅᓭᒪᔪᑦ ᐃᓄᐃᑦ ᑲᑐᔪᖅᑎᑎᒋᓐᓂᑦ - ᐃᑲᔪᖅᓭᓕᒃᑯᑦ ᐱᔪᓐᓇᐅᑎᓄᑦ ᐅᖃᓕᓐᓂᖓᓄᑦ” ᑐᑭᖅᑲᖅᑐᖅ ᐅᓇ ᐃᓄᐃᑦ ᑲᑐᔪᖅᑎᑎᒋᓐᓂᑦ - ᐃᑲᔪᖅᓭᓕᒃᑯᑦ ᐱᔪᓐᓇᐅᑎᓄᑦ ᐅᖃᓕᓐᓂᑦ ᐊᖏᓪᓗᑎᓕᓐᓇᖅᑐᓂᒃ ᑎᑎᕆᔭᓗᑎᓄᒃ.

4.3 ᐊᑐᖅᓗᑎᒃ ᐃᓄᐃᑦ ᓄᓇᕗᑎᖏᓐᓂᑦ ᓄᓇᒥᒃ ᐅᑎᖅᑕᕐᕕᑐᓄᑦ ᑎᖏᒥᐊᓄᑦ ᓴᓗᕆᐅᓯᒪᔪᒃ ᐊᓪᓚᓗ ᑲᓇᑕᒥ ᐅᒪᔨᖅᑭᖅᓴᕐ

ᐃᓕᓴᐅᓕᖅᑲᖅᓴᖅ ᐱᑦᓗᑎᑦ ᖃᐅᔨᓴᖅᓯᖅ

4.3.1 ᒥᓂᑦ ᐊᖏᑎᖅᓕᓗᓂ ᐊᕕᒃᑐᖅᓕᒫᑦ ᐃᓄᐃᑦ ᑲᑐᔪᖅᑎᑎᒋᖏᑎᑦ -ᐃᑲᔫᖅᓕᕆᑐᑦ ᐱᔨᓄᓇᐅᑎᓄᑦ ᐅᒃᑐᑎᑎᖏᓄᑦ ᑕᐃᑲᑐᓂ, ᓇᖏᒥᓂ ᐅᖅᑲᐅᒪᕗᖅᑐᓗᒍ, ᐱᓴᓕᑎᐊᖅᑐᓗᓂ, ᑐᒃᓕᕆᖅᑐᓗᓂ ᖃᓄᐃᑦᑐᕈᒪᕗᑎᓄᑦ ᒪᓕᓕᑎᐊᖅᑐᓗᓂ ᒡᒃᔅᓴᓕᖅᑐᒍ ***ᑲᑦᓗᑎᒃ ᐅᒪᔨᓕᓕᓄᑦᓯᒍ ᐱᖅᔅᖁᔨᐊᖅᖁᖅ, ᑎᖏᒥᓂ ᐅᑎᖅᑕᕐᕕᑐᐃᓐᓂ ᐱᔨᔪᑕᐅᓂᖏᓐᓄᓄᑦ ᐱᖅᔅᖁᖅ*** ᐊᓪᓚᓗ ᓂᖅᔪᑎᓄᑦ ᐃᓕᓴᐅᓕᖅᑲᖅᓴᖅᓂᑦ - ᐃᑲᔫᖅᓕᕆᑐᑦ ᐱᔨᓄᓇᐅᑎᓄᑦ ᐅᒃᑐᑎᑎᖏᓄᑦ, ᒥᓂᑦ ᐃᓕᓴᐅᓕᖅᑲᖅᓴᖅᓂᑦ - ᐃᑲᔫᖅᓕᕆᑐᑦ ᐱᔨᓄᓇᐅᑎᓄᑦ ᐅᒃᑐᑎᑎᖏᓄᑦ, ᒥᓂᑦ ᐃᓕᓴᐅᓕᖅᑲᖅᓴᖅᓂᑦ ᐱᖅᔅᖁᖅ, ᐊᑐᑎᖅᑎᖏᖅᑕᖅᓯᒪᓂᑦ ᐃᓄᐃᑦ ᑲᑐᔪᖅᑎᑎᒋᖏᑎᑦ - ᐃᑲᔫᖅᓕᕆᑐᑦ ᐱᔨᓄᓇᐅᑎᓄᑦ ᐅᒃᑐᑎᑎᖏᓄᑦ, ᒥᓂᑦ ᐃᓕᓴᐅᓕᖅᑲᖅᓴᖅᓂᑦ ᐱᖅᔅᖁᖅ, ᐊᑐᑎᖅᑎᖏᖅᑕᖅᓯᒪᓂᑦ ᐃᓄᐃᑦ ᑲᑐᔪᖅᑎᑎᒋᖏᑎᑦ - ᐃᑲᔫᖅᓕᕆᑐᑦ ᐱᔨᓄᓇᐅᑎᓄᑦ ᐅᒃᑐᑎᑎᖏᓄᑦ, ᒥᓂᑦ ᐃᓕᓴᐅᓕᖅᑲᖅᓴᖅᓂᑦ ᐱᖅᔅᖁᖅ, ᐊᑐᑎᖅᑎᖏᖅᑕᖅᓯᒪᓂᑦ ᐃᓄᐃᑦ ᑲᑐᔪᖅᑎᑎᒋᖏᑎᑦ - ᐃᑲᔫᖅᓕᕆᑐᑦ ᐱᔨᓄᓇᐅᑎᓄᑦ ᐅᒃᑐᑎᑎᖏᓄᑦ, ᒥᓂᑦ ᐃᓕᓴᐅᓕᖅᑲᖅᓴᖅᓂᑦ

(a) ᐊᒃᑐᑭᓇᖅᑕᖏᑐᓂᒃ ᖃᓄᐃᑦᑕᐅᖅᑕᐅᔨᓕᕆᖅ ᓄᓇᕗᒻᒥ ᑎᑭᐸᒃᑐᓄᑦ ᑎᖏᒥᐊᓄᑦ ᐊᓕᓗᓂ ᐃᓄᖏᑐᓄᑦ, ᓄᓇᕗᒻᒥ ᓂᔪᐸᑎᑦ ᐊᓕᓗᓂ ᐃᓄᖏᑐᓂ, ᓄᖓᒍᕆᖅᑐᑦ ᐅᓗᓴᐊᓇᖅᑐᒥᕆᕕᖅᑐᑦ ᐅᕝᕙᓘᓐᓃᑦ ᓯᕆᖅᑕᐅᑎᐊᖅᑐᑦ ᓂᔪᐸᑦᑐᑦ ᐊᑕᒍᑦ ᓂᔪᐸᑎᓄᑦ ᐃᓕᓴᐅᓕᖅᑲᖅᓴᖅᓂᑦ ᐱᖅᔅᖁᖅ, ᐅᕝᕙᓘᓐᓃᑦ ᑕᒃᓴᓕᖅ ᐅᒪᔨᓕᕆᑐᑦ ᐃᓄᐃᑦ, ᐅᑯᐊ ᐃᓕᑎᕐᓂᓗᑎᒃ:

(i) ᓄᓇᕗᒦᖅ ᐊᓚᓗᓗ ᖃᓄᖅ ᐊᒃᑎᑎᓂᒃ ᖃᓄᐃᑦᑕᐅᖅᑕᐅᔨᕕᖅ;

(ii) ᓇᓂᓐᓂᒃ ᐊᓚᓗᓗ ᓄᓇᓐᖓᐊᑎᒍᑦ ᓇᓗᓇᐃᔅᓗᒍ ᖃᓄᐃᑦᑕᐅᖅᑕᕐᕕᖅᑐᖅ;

(iii) ᖃᓐᖓᑕᐅᓴᖅᓂᒃ ᐊᖅᑦᒍᒥᒥ ᖃᓄᐃᑦᑕᐅᖅᑕᖅᑕᐅᔨᕕᖅ;

(iv) ᑕᑭᓂᒃ ᖃᓄᐃᑦᑕᐅᖅᑕᖅᑕᐅᔨᕕᖅ;

(v) ᖃᓄᐅᓚᕆᐅᔨᑦ ᐊᒃᑐᕋᖅᑕᖏᑦᑐᑦ ᖃᓄᐃᑦᑕᐅᖅᑕᖅᑕᐅᔨᕕᐸᑦ; ᐊᓚᓗᓗ

(vi) ᖃᓄᐅᓚᕆᐅᔨᑦ ᐅᓄᖅᓂᖏᑦᑐᑦ ᐊᓚᓗᓗ ᐃᓂᖏᑦᑐᑦ ᐃᓂᑐᖅᖃᓄᖅᓯᓄᑦ ᐅᑎᖅᑎᑦᑎᐊᓗᑦᑐᑦ;

(b) ᐃᓅᓯᑦᓴᓂᓐᓂᒃ, ᐱᐅᓯᑐᖅᑕᓴᓂᓐᓂᒃ ᐊᓚᓗᓗ ᑭᓇᐅᔨᖅᑎᑦ ᐱᓕᕆᑦᑕᓴᓂᑦᑎᔨᕈᑎᐅᔨᑦ ᐱᓚᒪᓴᐅᔨᑦ ᖃᓄᐃᑦᑕᐅᑭᐅᑦᑕᐅᔨᓕᒥᑦ ᐃᓅᓐᓄᑦ;

(c) ᐅᑯᐊ ᐊᕕᒃᑐᖅᓕᕆᑐᑦ ᐃᓄᐃᑦ ᑲᑐᔪᖅᑎᑎᒋᖏᑎᑦ ᐊᓚᓗᓗ ᐊᑕᔨᑦ ᓄᓇᓕᓪᓕᑦ ᑐᑭᑕᐅᐅᑎᑎᓕᓪᓕᓴᐅᑎᖅᑕᖏᑦᑐᑦ ᐃᑲᔪᖅᑐᐊᖅᑐᓗᒍᑎᑦ ᖃᓄᐃᑦᑕᐅᖅᑕᖅᑕᐅᔨᕕᑦ;

(d) ᖃᓄᐅᑐᐃᓐᓇᖅ ᑭᓇᐅᔨᕈᓴᓂᐅᑎᓗᒍᓚᐊᖅᐸᑦ ᐊᓚᓗᓗ ᐊᑐᑎᓴᐅᒃ ᖃᓄᐃᑦᑕᐅᓐᓚᔨᐊᑕᐅᔨᕕᐃᓐᓇᖅᑐᐅᖅᑕᐅᒃ ᖃᓄᐃᑦᑕᐅᖅᑕᖅᑕᐅᔨᕕᖅ ᐃᓗᐊᓂ ᐅᑎᖅᑕᐅᐸᑦᑐᓄᑦ ᑎᖏᒥᐊᓄᑦ ᓴᓗᕆᐅᓯᒪᔪᒃ ᐊᓚᓗᓗ ᐃᓕᓴᐅᓕᖅᑲᖅᓴᖅᓂᑦ;

(e) ᐃᓕᐊᑐᐃᓐᓇᖅ ᐄᓐᓄᑎᑎᑎᐅᔨᕕᖅ ᐅᓂᒃᑲᓐᓄ ᐊᑕᒍᑦ 4.3.8 ᐊᓚᓗᓗ ᐃᓕᐊᑐᐃᓐᓇᖅ ᐊᕕᒃᑐᖅᓕᕆᑐᑦ ᐃᓄᐃᑦ ᑲᑐᔪᖅᑎᑎᒋᖏᑎᑦ ᐊᓐᓄᓕᕆᖅᑲᒃᑐᑎᑦ - ᒥᓂᑦ ᐅᖅᑲᖅᑎᑎᒍᓕᑎᑎᓄᑦᑕᐅᑎᖅᑕᖏᑦᑐᑦ ᐊᑕᒍᑦ 4.3.9;

(f) ᐊᓯᖏᑦᑕᐅᕕᖅ ᐱᓕᒃᓴᑦ ᑕᐃᒪᓗᓗ ᒥᓂᑦ, ᐱᓕᕆᑦᑎᐊᖅᓯᑎᑦ, ᐃᓯᓚᑦᑎᐊᖅᓯᑎᑦ ᐊᑐᑎᖅᑲᖅᑐᑦ; ᐊᓚᓗᓗ

(g) ᖃᓄᖅ ᖃᓄᐃᑦᑕᐅᖅᑕᖅᑕᐅᔨᕕᖅ ᐃᓅᖅᑲᑎᑦᑎᔨᓇᖅᓯᓚᐅᑦ ᐃᓗᐊᓂ ᐅᑎᖅᑕᕐᕕᑐᓄᑦ ᑎᖏᒥᐊᓄᑦ ᓴᓗᕆᐅᓯᒪᔪᒃ ᐊᓚᓗᓗ ᑲᑐᔪᖅᑎᑎᒋᖏᑎᑦ, ᐱᓕᓗᓇ ᖃᓄᐃᑦᑲᖅᓴᖅᑎᑦ ᐅᖅᑲᖅᑎᑎᖅᑕᐅᔨᕕᖅ.

ᐃᓱᒪᓕᐅᕆᓚᐅᖅᓯᒪᔪᖅ

4.3.2 ᐅᓇ ᐊᑐᑎᓴᖅ ᐊᐅᓪᓚᑎᕆᐊᖃᖅᓯᓐᓂᖓᓪᓗᒍ ᑲᑎᒪᔨᖏᓪᓗ ᐅᓴᕆᔨᖅᑲᒋᑦᑕᓐᓄᑦ ᐊᕐᕌᖅᑕᐅᓴᒪᓕᕆᔨᒃᑯᑦ ᐃᓚᐅᑦ ᑲᔪᓯᑎᑦᑎᓐᓄᑦ ᐊᕐᕌᖅᑕᐅᓴᒪᓕᕆᔨᒃᑯᑦ ᐃᓚᐅᑦ ᑲᔪᓯᑎᑦᑎᓐᓄᑦ - ᐃᑲᔪᖅᓱᐅᑎᑦ ᐱᓕᕆᔨᐅᑎᓚᑦ, ᐃᓪᓕᖅᓴᐅᓚᑎᑲᖅ ᓚᓪᒋᑕᐅᔪᓕᕆᓪᓕᐊᓪᓚᖅ ᐊᓪᓚᑎᑲᖅᑕᐅᓴᒪᓕᕆᔨᒃᑯᑦ ᐃᓚᐅᑦ ᑲᔪᓯᑎᑦᑎᓐᓄᑦ - ᐃᑲᔪᖅᓱᐅᑎᑦ ᐱᓕᕆᐊᐅᑎᓚᑦ.

4.3.3 ᐃᓱᒪᓕᐊᖓᔪᓪᓕᐊᓪᓚᑐ ᐅᓴᖅᑲᓱᒪᓕᕆᔨᒃᑯᑦ 4.3.4-ᒥ ᑎᑭᑦᓵᔭᒍ 4.3.13-ᔪᑦ, ᐊᑐᖅᓱᓕᑎᒃ ᖃᐅᔨᓴᕐᓂᒃᑲᒥᒃ ᑎᑎᕋᖅᑲᒪᓕᕆᔨᖅ 4.3.1-ᒥ ᐊᑐᑎᓂᐊᖅᑐᖅ ᐊᕐᕌᖅᑕᐅᓴᒪᓕᕆᔨᒃᑯᑦ ᐃᓚᐅᑦ ᑲᔪᓯᑎᑦᑎᓐᓄᑦ - ᐃᑲᔪᖅᓱᐅᑎᑦ ᐱᓕᕆᐊᐅᑎᓚᑦ ᐅᖅᑐᑎᓂᓚ. ᐱᖓᓕᓕᐊᓂᖅᓯᒪᓕᕆᔨᖅ 3.3.6-ᒥ ᐊᓪᓚᑦ 3.3.7-ᒥ ᐊᑐᑎᓂᐊᓐᖓᓕᑐᑦ ᑕᑯᔨᓚᖓᓐᓂ ᐊᐅᓪᓚᑎᕆᐊᖃᖅᓯᓐᓂᖓᓪᓗᒍ ᑲᑎᒪᔨᖏᓪᓗᑕ ᐅᖅᑐᑎᓂᐊᖅᑐᖅ ᐱᓕᕆᔨᒃᑯᑦ ᐃᓚᐅᑦ ᑲᔪᓯᑎᑦᑎᓐᓄᑦ - ᐃᑲᔪᖅᓱᐅᑎᑦ ᐱᓕᕆᐊᐅᑎᓚᑦ ᐅᖅᑐᑎᓂ.

4.3.4 ᐅᓇ ᐊᕐᕌᖅᑕᐅᓴᒪᓕᕆᔨᒃᑯᑦ ᐃᓚᐅᑦ ᑲᔪᓯᑎᑦᑎᓐᓄᑦ - ᐃᑲᔪᖅᓱᐅᑎᑦ ᐱᓕᕆᐊᐅᑎᓚᑦ ᐅᖅᑐᑎᓂ, ᐱᓴᑎᑎᖅᑕᒍ ᓇᓕᐊᖅᓂᑐᐃᓐᓇᖅ ᐃᑲᔪᒋᑎᖅᓂᒃ ᐅᖅᑐᑎᓂᖅ, ᐅᖅᑲᐅᓱᖅᑐᓂᐊᖅᑐᑦ ᖃᐅᔨᓴᕐᓂᒃᑲᒥᒃ ᐊᕐᕌᖅᑕᐅᓴᒪᓕᕆᔨᒃᑯᑦ ᐃᓚᐅᑦ ᑲᔪᓯᑎᑦᑎᓐᓄᑦ - ᐃᑲᔪᖅᓱᐅᑎᑦ ᐱᓕᕆᐊᐅᑎᓚᑦ ᐅᖅᑐᑎᓂ ᐱᓱᓪᓕᐊᓂᖅᑐᑦ ᐃᓚᐅᑦ ᑲᔪᓯᑎᑦᑎᓐᓄᑦ ᐃᓚᐅᑦ ᑲᔪᓯᑎᑦᑎᓐᓄᑦ ᐅᖅᑐᑎᓂ ᐱᓱᓪᓕᐊᓂᖅᑐ

ᐊᑯᓂᐅᓗᐊᓂᕐᒪᑐᑐᒃ, ᓂᕆᐊᖏᓐᓄᓗᒃ ᐊᓐᕈᕐᖏᑎᒍᑉᑐᒃ ᐃᓐᓴᕐᖏᑎᒋᔭᖏᑎᒋᓂᐅᓂᒃᑯᓂ ᐊᒻᒪᓗ ᐃᓐᓴᕐᖏᑎᒋᔭᖏᑎᓯᓪᓕᐊᓂᖏᖏᓂᒃᑯᓂ, ᐊᒻᒪᓗ ᐃᓐᓴᕐᖏᑎᒋᔪᒍᐅᔪᓂᒃ ᐅᓂᒃᑲᓐᓂᒃ ᑐᓐᓴᕐᖏᑎᒋᔭᐅᓂᒃᑐᓂᓗ ᐊᓐᓴᕐᕿᑲᓐᓂᒃᑐᓂᓗ ᐊᐃᓐᑐᕐᖏᓕᕆᔨᓂ ᐃᓚᐅᓂ ᑲᑐᔾᔨᖏᓐᓄᑦᑐᓂᒃ ᐊᒻᒪᓗ ᒥᓂᑕᒧᑦ.

4.3.9 ᑕᐃᒪᓗ, ᐃᓕᓴᕐᓂᐊᓕᐅᖅᑕᒍ ᐃᓐᓴᕐᖏᑎᒋᔭᖏᑎᓂᑎᔪᖅ ᐅᓂᒃᑲᖏᓐ ᐊᒋᔪᑦ ᑎᑎᕐᓂᒃᑯᓂᖏᑎᒋᔭᖏᑎᓂᑎᔪᖅ 4.3.8, ᐅᑯᐊ ᐃᓄᐃᑦ ᐊᐃᐸᑐᖅᓯᒪᔪᑦ ᑲᑐᔾᔨᖏᑎᒋᔭᖏᑦ ᐊᑐᕈᒪᔭᖏᓐᖏᑦ ᐱᓕᕆᐊᖓᓂᖏᓐᓄᑦ ᑲᔪᓯᑎᑦᑕᒍ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖏᑎᒋᔭᖏᑦ - ᐃᑲᔪᖅᓯᕗᑦ ᐱᔪᓐᓇᐅᑎᓄᑦ ᐅᐸᑐᑎᑦ, ᑐᓴᕈᐅᑎᖏᓐᓇᖅᑐᓐᓇᖅᑐᖅ ᐅᐸᑐᑎᖓ ᑕᑦᓯᒧᖓ ᐊᓐᓴᕐᕿᑲᓐᓂᒃᑯᓂᖏᓐᓄᑦ ᐊᐃᐊᖅᑐᓂᒃ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖏᑎᒋᔭᖏᑦ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖏᑎᒋᔭᖏᑎᓂᑎᔪᖅ ᒥᓂᑕᒧᑦ, ᑕᐃᒪᓗ, ᑕᒪᐃᓐᓂᒃ ᓇᒪᒋᔭᐅᓗᑎᒃ ᖃᐅᖓᑲᓂᒃ, ᐅᖃᖅᖃᑎᒋᔭᖏᑎᓗᓂᒃ ᐅᐸᑐᑕᐅᔪᖅ, ᑲᑎᖅᑎᑎᒋᔭᖏᑎᓗᓂᒃ ᐅᐸᑐᑎᑦ ᐊᑐᕐᓂᒃᑯᓂᖏᓐᓄᑦ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖏᑎᒋᔭᖏᑦ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖏᑎᒋᔭᖏᑎᓂᑎᔪᖅ

ᐊᓚᓗ ᑐᓂᔪᐅᓗᑎᒃ ᒪᑦᔨᐅᓗᒃ ᐊᐃᒃᑐᖅᓯᒪᔨᓂ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑎᒋᔭᖅᓂ ᐅᕙᓪᓕᐊᓂ ᐊᐅᓪᓚᖅᑎᑎᒋᓐᓂᕐᒧᑦ ᑲᑎᒪᔨᖏᓪᓗᒍ.

4.5 ᐃᓄᐃᑦ ᓄᓇᕗᑎᖏᓪᓕ ᐊᖏᕈᑎᑎᓂᖅᖏᓪᓕᑐᑦ 9-3

4.5.1 ᐃᓄᐃᑦ ᓄᓇᕗᑎᖏᓪᓕ ᐅᑯᐊᓗ ᐱᔨᒪᔨᐅᓕᕆᔨᑦᑐᑦ ᑐᓴᕐᕋᖅᑕᐅᓴᔨᒥᒃ ᑲᓇᑕᒥ ᐅᒪᔨᔭᕐᓂᖅᓱᑦ ᐅᖓᓯᒃᑐᐊᕐᓂᖅᓄᑦ ᐅᖓᖅᑎᑦᓯᓄᑦ ᑎᓪᓗᐊᓄᑦ ᓴᓇᔨᐅᓴᒪᓕᕐᒃ ᐅᕙᓪᓕᐊᓂᓂ ᑲᑎᒪᔨᖏᓪᓕ ᐅᕙᓪᓕᐊᓂᓂ ᐅᖓᓯᒃᑐᐊᕐᓂᖅᓄᑦ ᑎᓪᓗᐊᓄᑦ ᓴᓇᔨᐅᓴᒪᓕᕐᒃ ᐊᓪᓚᓗ ᐅᓪᓗᐊᓂᖅᓯᒪᓕᕐᓂᑐᑦ 9.3 *ᓄᓇᕗᑦᑐᒃ ᓄᓇᕕᑦᑎᓄᑦ ᐊᖏᕈᑎᑎᓂ* ᑭᓇᐅᔨᒪᓂᖅ ᐊᖏᕈᑎᑎᓂᖅᓄᑦ ᐊᐃᑐᑎᓪᓗᒍ ᐊᐃᒃᑐᖅᓯᒪᔨᓂ ᐊᖏᕈᑎᑎᓂᖅᓄᑦ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑎᒋᓐᓂᕐᒧᑦ ᐃᓄᐃᑦ ᓄᓇᕗᑎᖏᓪᓕ ᐊᓯᖏᓪᓕᐊᓂᖅᓄᑦ ᐊᐃᒃᑐᖅᓯᒪᔨᓂ ᐊᐃᑐᑎᓪᓗᒍ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑎᒋᓐᓂᕐᒧᑦ, ᐃᓕᓴᐅᑎᖅᓯᒪᔨᓂ ᐃᓚᖓᓂ ᐊᖏᕈᑎᑎᓂᖅᓄᑦ ᐃᓄᐃᑦ ᓄᓇᕗᑎᖏᓪᓕ ᐊᐃᒃᑐᖅᓯᒪᔨᓂ ᐃᓕᓴᐅᑎᖅᓯᒪᔨᓂ ᐃᓄᐃᑦ ᓄᓇᕗᑎᖏᓪᓕ ᐊᖏᕈᑎᑎᓂᖅᓄᑦ ᐃᓄᐃᑦ ᓄᓇᕗᑎᖏᓪᓕ ᐊᐃᒃᑐᖅᓯᒪᔨᓂ

4.6.4 ᑲᓇᑕᒥ ᐅᒪᔨᓕᓚᕐᓂᔨᐊᕐᑯᑦ ᑐᓂᔅᓯᒪᔨᓂᖅ ᑎᒍᕐᓯᓂᖅᑐᑦ ᑕᒪᐃᓐᓄᑦ ᑲᓇᑕᒥ ᐅᒪᔨᓕᓚᔨᓄᑦ ᐱᓕᕐᑎᕐᑎᑦ ᐱᓇᓯᐊᕐᑎᓄᑦ, ᐃᖅᑲᓇᐃᔨᕐᑎᓄᑦ ᐊᒻᒪᓗ ᑲᓐᑐᖅᑕᐅᓯᒪᔨᓄᑦ, ᖁᑲᐅᒪᕐᑎᐊᕐᓂᐊᕐᒪᑕ ᑕᒪᑲᑯᐊ ᐃᓄᐃᑦ ᒪᓕᕐᓂᐊᕐᑯᓐᓂᑦ ᐊᖏᕐᑎᑐᑎ 4.6.3-ᒥᑲ.

4.7 ᐃᓯᖅᑕᐅᔨᓇᖅᑐᑦ ᐃᓄᐃᑦ ᓄᓇᕐᑯᑎᖏᑦ

4.7.1 ᐅᑯᐊ ᒪᓕᒐᐃᑦ ᐃᓕᖏᓂᒃ 4.3 ᐊᑐᑎᓂᐅᔪᖅᑐᑦ ᐱᔨᓇᐅᑎᑕᕐᓯᑎᓄᑦ ᑕᑲᑯᓄᖏᖅ ᑲᓇᑕᒥ ᐅᒪᔨᓕᓚᔨᓂᕐᓂᔨᐊᕐᑯᑦ ᐃᑲᔨᖅᑕᐅᓗᑎᒃ ᑕᑲᑯᓇᖏᖅ ᐊᕐᓚᖅᑐᖅᑐᑦ ᐃᓄᐃᑦ ᑲᑐᑦᓇᖅᑎᑎᒋᑦ ᐊᒻᒪᓗ ᐃᓯᕐᓄᓇᕐᓂᖏᖅ ᓇᓂᑐᐃᓇᖅ ᑲᑎᒪᔨᖏᑦ ᐅᕙᓪᓕᐊᓂᖅ ᐅᑎᖅᑕᐸᑐᓄᑦ ᑎᓕᕐᒥᐊᓄᑦ ᓴᓱᖏᐅᕙᓚᐃᒃ ᑭᖑᓪᓕᓄᓕ, ᐅᕙᓪᓕᐊᓂᓕᑦ ᑕᑯᔪᑎᒃ ᓇᐸᖅᑕᐅᔨᓚᐅᖅᑐᑦ ᐅᕙᓪᓕᐊᓂᓕᑦ ᑭᒧᑐᖅ ᐅᖃᓕᓕᒫᑦ ᐅᕙᓪᓕᐊᓂᓕᑦ.

ᐃᓕᓐᓄ 5 — ᐃᓄᐃᑦ ᐱᓱᓇᓂᓕᕆᑦ ᐊᓐᓚᓗ ᐊᑐᖅᑕᐅᓴᓱᓕᕆᑦ ᑲᓇᑕᒥ ᐅᓚᔨᖅᑎᓂᑦ ᐊᓚᓗ ᐅᑎᖅᑕᕐᑐᓄᑦ ᑎᓪᓗᐊᓄᑦ ᓴᓲᐅᓕᒪᓂᖅ

5.1 ᑐᑭᒃᖅᑕᕐᓕᐊᔪᑎᒃᓴᖅ

5.1.1 ᐅᕙᓂ ᐃᓕᓐᓄᓂ ᐊᒋᓂ ᑎᑎᕋᖅᓯᒪᔨᑦ ᑐᑭᒃᖅᑕᕐᓕᐊᔪᑎᒃᓴᓐᓂᑦ:

(a) ᐃᓕᓴᕆᓗᑐᑦ ᐃᓄᐃᑦ ᐱᔨᓇᓕᕋᓂᖅᑎᒃᓴᓐᓂᑦ ᐊᖏᓇᓯᒡᓇᖅᓯᒪᑎᒃ ᐃᓗᐊᓂ ᑲᓇᑕᒥ ᑲᑎᒪᔨᖅᑎᑦᑎᓂᑦ ᑎᓪᓗᐊᓄᑦ ᐅᑎᖅᑕᕐᑐᓄᑦ ᐊᑐᓚᓗᒃ, ᐅᕋᖅᓯᒪᔨᑦ *ᓄᓇᕗᒻᒥ ᓄᓇᕈᑭᑎᓄᑦ ᐊᖏᑭᑎᖏᓐᓂᒥ;*

(b) ᐅᕋᐅᓯᓗᓕᑐᑦ ᐱᔨᓇᐅᑎᕈᑭᑎᓄᑦ ᑕᐃᒪᓐᓇᐅᕐᓗᖅᓴᕆᕋᓂᖅ ᐊᐅᓪᓚᑭᐱᓕᒪᓂᐊᖅᑐᒃ ᐊᐅᓪᓚᑭᐱᓕᒪᓂᐊᖅᑐᒃᑐᑦ ᐊᖏᓇᓯᒡᓇᖅᓯᒪᓂᑦ ᐃᓗᐊᓂ ᑲᓇᑕᒥ ᐅᑭᒃᖅᑕᕐᓕᐊᔪᑎᒃᓴᓐᓂᑦ ᑎᓪᓗᐊᓄᑦ ᐅᑎᖅᑕᕐᑐᓄᑦ ᐊᑐᓚᓗᒃ, ᐅᕋᖅᓯᒪᔨᑦ *ᓄᓇᕗᒻᒥ ᓄᓇᕈᑭᑎᓄᑦ ᐊᖏᑭᑎᖏᓐᓂᒥ;*

(c) ᐱᔨᓇᓕᓗᑎᒃ ᐃᓄᐃᑦ ᐱᔨᓇᓕᓂᕆᔪᑦ ᐅᑲᑯᓯᐱᑲᓐᓂᑐᒃ ᓴᓇᓐᓯᐊᐱᑲᓕᒪᓂᑦ ᑕᐃᑲᓐᓂᑦ ᑲᑎᒪᔨᖅᑎᑦᑎᓂᑦ ᐃᓗᐊᓂ ᑲᓇᑕᒥ ᐅᑎᖅᑕᕐᑐᓄᑦ ᐅᑎᖅᑕᕐᑐᓄᑦ ᑎᓪᓗᐊᓄᑦ ᐊᑐᓚᓗᒃ, ᐅᕋᖅᓯᒪᔨᑦ *ᓄᓇᕗᒻᒥ ᓄᓇᕈᑭᑎᓄᑦ ᐊᖏᑭᑎᖏᓐᓂᒥ;*

(d) ᑐᓂᓯᒪᓕᑐᑦ ᐃᓄᐃᑦ ᐊᐳᑭᐱᓯᓇᐅᑎᓗᑎᒃ ᓄᓇᓕᕋᓵᒃᓴᒃᓐᓂᒃ ᐃᓗᓐᓯᓐᓂ ᑲᓇᑕᒥ ᐅᓚᔨᖅᑎᑦᑎᓂᑦ ᑲᑎᒪᔨᖅᑎᑦᑎᓂᑦ ᐅᑎᖅᑕᕐᑐᓄᑦ ᐊᑐᓚᓗᒃ, ᐅᕋᖅᓯᒪᔨᑦ *ᓄᓇᕗᒻᒥ ᓄᓇᕈᑭᑎᓄᑦ ᐊᖏᑭᑎᖏᓐᓂᒥ.*

5.2 ᐃᓄᐃᑦ ᐊᑐᕈᓇᖅᑕᖏᓐᓄᑦ ᐊᑎᒪᓗ ᐃᓱᕈᓇᖅᓴᓕᑐᑦ ᑲᓇᑕᒥ ᓅᓚᔨᖅᑎᑦᑎᓂᑦ ᐊᑎᒪᓗ ᐅᑎᖅᑕᕐᑐᓄᑦ ᑎᓪᓗᐊᓄᑦ ᓴᓲᐅᓕᒪᓂᖅ

5.2.1 ᐅᕋᖅᓯᒪᔨᕋᖅ ᐃᓕᓐᓄᑦ 5 ᐊᑎᒪᓗ ᒪᓕᒋᐊᓕᓵ 5.7.18 *ᓄᓇᕗᒻᒥ ᓄᓇᕈᑭᑎᓄᑦ ᐊᖏᑭᑎᖏᓐᓂ* ᐃᓄᐃᑦ ᐃᓱᒪᖅᓯᖅᑐᑦ ᐊᑎᒪᓗ ᐃᓱᖅᑕᐃᓕᓴᖅᓄᖅᓯᑦᑐᑦ ᐊᖏᓇᓯᒡᓇᖅᑐᓂᒃᑯᑦ ᑕᒪᕐᒥ ᓄᓇᕗᒻᒥ ᐃᓗᐊᓂ ᑲᓇᑕᒥ ᐅᑎᖅᑕᕐᑐᓄᑦ ᑎᓪᓗᐊᓄᑦ ᓴᓲᐅᓕᒪᓂᖅ.

5.2.2 ᐅᕋᖅᓯᒪᔨᑦ ᐊᑎᒪᓗ ᒪᓕᒋᐊᓕᓂ ᑕᕐᔪᓐᓇ ᐃᓕᓐᓄ 5 *ᓄᓇᕗᒻᒥ ᓄᓇᕈᑭᑎᓄᑦ ᐊᖏᑭᑎᖏᓐᓂᒥ*, ᐃᓕᑎᒃᐅᓪᓗᑐᒃ ᓇᓕᐊᑐᐃᓇᓕᖅ ᐊᑲᑐᑎᑦ ᐊᑐᓇᐊᖅᓴᖅᓯᑎᑎᓂᑎᑎᔨᐊᕐᓕᐅᔨᑦ ᓴᓱᑭᐅᐅᔨᒪᔨᑦ ᐃᓕᓐᓄᓂᒥ ᑕᑯᓐᓇᓕᒫᑦ ᓄᓇᕗᒻᒥ ᐃᓱᒪᖅᓯᖅᑐᑦ ᐅᓇ ᐃᓄᒃ ᐅᕝᕙᓘᓐᓃᑦᑐᒃᑯᑦ ᑎᑎᕋᐅᓕᒪᓂᖅᑐᒃ ᐊᑐᓚᓗᒃ ᖃᓄᐃᓕᓐᓂᒃ ᐊᑐᔪᓐᓇᖅᑐᑦ ᐊᑎᒪᓗ ᐊᑐᖅᑕᐅᓴᓱᓕᕆᑦ ᐊᑐᔪᓐᓇᖅᑐᑦ ᐃᓕᓐᓄᑦ 5.7.34 *ᓄᓇᕗᒻᒥ ᓄᓇᕈᑭᑎᓄᑦ ᐊᖏᑭᑎᖏᓐᓂᒥ* ᐃᓚᖓᓄᐅᔨᓇᖅᑐᓴᖅ, ᐃᓗᐊᓂ ᑲᓇᑕᒥ ᐃᓗᐊᓂ ᑲᑎᒪᔨᖅᑎᑦᑎᓂᑦ ᐅᑎᖅᑕᕐᑐᓄᑦ ᑎᓪᓗᐊᓄᑦ ᐊᑐᓚᓗᒃ, ᐊᐅᓪᓚᑭᐱᓕᒪᓂᐊᖅᑐᒃ ᓂᕆᔨᐅᓂᒃ ᐊᑎᒪᓗ ᖃᓄᐃᓕᓐᓂᒃᑯᑦᑎᓂᑦ ᐊᐅᓪᓚᑭᐱᓕᒪᓂᐊᖅᑐᒃᑐᑦ ᓯᐃᓴᕆᓂᖅᑐᓂᒃᑯᑦᑎᓂᒃ, ᐱᔨᓇᐅᑎᖅᓂᖅᑐᓂᒃᑯᑦᑎᓂᒃ ᐅᕙᓘᓐᓃᑦᑐᒃᑯᑦ ᐃᓕᓴᕆᓂᖅᑐᓂᒃᑯᑦ ᐊᑐᓚᓗᒃ *ᓄᓇᕗᒻᒥ ᐃᓗᐊᓕᓵᓂᕐᒧᑦ ᐱᖃᓐᓇᕐᓂᖅ* ᐅᕙᓘᓐᓃᑦᑐᒃ *ᑎᑎᕋᐅᓕᒪᓂᖅᑐᒃᑐᑦ ᐱᖃᓐᓇᕐᓂᖅ, ᑎᓪᓗᐊᓄᑦ ᐅᑎᖅᑕᕐᑐᓄᑦ ᐊᑐᓚᓗᒃ ᐱᓪᓚᕆᓂᖅ* ᐃᓄᐃᑦ ᐱᔨᓇᐅᑎᖅᑕᕐᑐᓄᑦ ᐊᑐᓚᓗᒃ.

5.2.3 ᑐᑭᓯᓇᓕᑎᑲᓐᓂᔪᓂᕐᒪᑦ, ᑎᑭᓕᑐᒍ ᐅᑯᐊ ᒪᓕᒋᐊᓕᔪᓕᑎᒃ *ᓄᓇᕗᒻᒥ ᓄᓇᕈᑭᑎᓄᑦ ᐊᖏᑭᑎᖏᓐᓂᑦ*, ᐅᕙᓘᓐᓃᑦᑐᒃ *ᐱᔨᓇᓕᕋᓂᖅᑎᑦᑎᓂᑦ ᐱᖃᓐᓇᕐᓂᖅ*, *ᑎᑎᕋᐅᓕᒪᓂᖅᑐᒃᑐᑦ ᐱᖃᓐᓇᕐᓂᖅ, ᐃᓗᐊᓂ ᑲᓇᑕᒥ ᐅᑎᖅᑕᕐᑐᓄᑦ ᑎᓪᓗᐊᓄᑦ ᐊᑐᓚᓗᒃ*, ᐅᕙᓘᓐᓃᑦᑐᒃ *ᓄᓇᕗᒻᒥ ᓄᓇᕈᑭᑎᓄᑦ ᐊᖏᑭᑎᖏᓐᓂᒥ* ᐃᓄᐃᑦ ᐱᔨᓇᐅᑎᖅᑕᕐᑐᓄᑦ ᐊᑐᓚᓗᒃ *ᐱᔨᓇᓕᕋᓂᖅᑎᑦᑎᓂᑦ ᐱᖃᓐᓇᕐᓂᖅ, ᑎᓪᓗᐊᓄᑦ ᐅᑎᖅᑕᕐᑐᓄᑦ ᐊᑐᓚᓗᒃ ᐱᓪᓚᕆᓂᖅ* ᐃᓄᐃᑦ ᐱᔨᓇᐅᑎᖅᑕᕐᑐᓄᑦ ᐊᑐᓚᓗᒃ *ᐱᔨᓇᓕᕋᓂᖅᑎᑦᑎᓂᑦ ᐱᖃᓐᓇᕐᓂᖅ* ᐃᓄᐃᑦ ᐱᔨᓇᐅᑎᖅᑕᕐᑐᓄᑦ ᐊᑐᓚᓗᒃ.

5.2.4 ᓄᓇᕗᑦ ᑐᓴᖏᓐᓄᐊᕐ ᑎᒥᖅᓯ ᑎᑎᕋᔨᓕᑦ ᐃᓄᐊᑎᑐᑦ ᐅᕙᑦᓴᓕᕆᔨᑦ ᐊᓚᒪᓗ ᖃᓄᐃᓗᐊᖅᑐᑦ ᕿᑐᐊᓕᕐᓂᐊᕐᓇᖅᑐᑦ ᐃᓄᐃᑦ ᐊᓯᐅᓕᕐᓴᓐᓇᖅᑐᑦ ᐊᓚᒪᓗ ᑕᒪᐃᓄᓯᖅ ᖃᓄᐃᓕᒥᐅᓴᖅᑐᑦ ᐱᔨᓴᓂᐊᕐᓇᖅᑐᑦ ᐃᓗᐊᒥᑦ ᐅᑎᖅᑕᕐᓂᑐᑦ ᑎᓕᒥᓄᑦ ᓴᓐᖏᓕᕐᓂᖅ. ᑕᐃᑲᓂᓗ, ᑲᓇᑕᒥ ᐅᒪᔨᓕᓪᓕᐊᑎᑦ ᐊᓚᒪᓗ ᓄᓇᕗᑦ ᑐᓴᖏᓐᓄᐊᕐ ᓄᓇᖃᖅᑐᓕᕐᓂᐊᕐᓇᖅᑐᑦ. ᐃᓚᒪᖃᓕᓐᓄᑦ ᑕᒪᒃᑯᓐᓄᑦ ᑲᑎᒪᔨᓕᓪᓕᐊᑎᑦ ᐊᓚᒪᓗ ᓄᓇᕗᑦ ᑐᓴᖏᓐᓄᐊᕐ ᐅᑎᖅᑕᕐᓂᑐᑦ ᖃᓄᐃᓕᒥᐅᓴᖅᑐᑦ. ᐃᓚᒪᖃᓕᓐᓄᑦ ᑕᒪᒃᑯᓐᓄᑦ ᓄᓇᖃᖅᑐᓕᕐᓂᐊᕐᓇᖅᑐᑦ, ᑲᑎᒪᔨᓕᓪᓕᐊᑎᑦ ᐊᑭᓕᓐᓄᑦ ᐃᓗᐊᒥᑦᑐᓂᓕᕐᓂᐊᕐᓇᖅᑐᑦ ᐊᓚᒪᓗ ᓄᓇᕗᑦ ᑐᓴᖏᓐᓄᐊᕐ ᓄᓇᖃᖅᑐᓕᕐᓂᐊᕐᓇᖅᑐᑦ ᑐᓴᖏᓐᓄᐊᕐ ᐃᓗᐊᒥᑦᑐᓂᓕᕐᓂᐊᕐᓇᖅᑐᑦ ᐊᓚᒪᓗ ᑐᓴᖏᓐᓄᐊᕐ ᓄᓇᖃᖅᑐ

ᓄᓇᕗᑦ ᓄᓇᕐᔪᐊᑎᖅᑐᓂᒃ ᐃᓗᐊᓂᒃ ᐅᓇᕐᓂᒃ ᑲᑎᒪᓗ ᐅᑎᖅᑕᕐᕕᒋᓄᑦ ᑎᒥᐅᔪᓄᑦ ᓴᓇᕐᓂᖅᓯᒪᔭᖅ.

5.4.2 ᐃᓅᓴ ᐊᑐᐸᓇᖅᑕᖏᑦ ᐱᓕᓇᕐᓂᖅᓂᖅᓂᒃ ᐅᖅᑲᖅᓯᒪᔨᑦ ᐊᖏᑎᑕᑎᑦ 5.4.1, ᐊᑐᐸᓇᕐᓗᒍ ᐄᓴᐃᓕᓇᕐᓂᑎᑦ ᐊᓪᓚᓗ ᐊᓴᖅᑐᑦᓗᑎᑦ ᓴᓇᓇᒃᑯᓕᖅᑐᑦ ᓴᖅᑭᑎᓂᖅᑐᒃ ᓴᓇᕐᓂᑦ ᐃᓂᖏᓇᓂᖅ, ᐃᓕᑎᒥᐅᓂᒃᑐᒃ ᐊᑦᑭᑎᐊᖅᑯᓯᒪᐃᓇᕐᓂᒍ ᖃᐅᔨᓴᖅᑎᑦᓯᒪᓂᖅ ᓴᓇᖅᑐᒃ ᐊᓪᓚᓗ ᓴᓇᖅᑐᑦ ᐃᓂᖏᓇᓂᖅ, ᐃᓕᑎᒥᐅᓂᒃᑐᒃ ᐊᑦᑭᑎᐊᖅᑯᓯᒪᐃᓇᕐᓂᒍ ᖃᐅᔨᓴᖅᑎᑦᓯᒪᓂᖅ ᐊᑐᐸᓇᕐᓗᒍ.

5.4.3 ᐊᖏᔪᑎᑦᑕ ᐊᐅᓪᓚᕆᓇᖅᑎᑎᕆᔨᒪᓯᑦᓴᑦ ᑲᑎᒪᔨᕿᓪᓚᕆᑐᒃ ᑕᒃᑯᓂᖓ ᑲᓇᖅ ᐅᓪᓗᒥᐅᑕᕐᓂᖅᓯᐊᑯᓂ, ᐃᓄᐃᑦ ᐊᑐᖅᑯᑦᑕ ᐱᓕᕆᓂᖅᓯᒪᔪᓂ ᐊᒡᓂᑎᑦ 5.4.1 ᐊᑎᒪᓗ 5.4.2 ᐊᑎᒪᓗ, ᑕᐃᒪᓐᓇᐅᔪᓂ ᐃᓄᐃᑦ ᓄᓇᕐᓂᖅᑐᓂ ᐃᓗᐊᓂᒃ ᐅᓇᕐᓂᖅ ᑲᑎᒪᔨᕿᓪᓚᕆᑐᒃ ᐅᑎᖅᑕᕐᕕᒋᓄᑦ ᑎᒥᐅᔪᓄᑦ ᓴᓇᕐᓂᖅᓯᒪᔭᖅ, ᐊᑐᓂᑦᓂ ᐊᕐᓇᖅᑐᖅᑕᐃᓕᓇᕐᓂᑦ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑦᓯᒪᓂᖅ, ᓴᓇᓇᒃᑯᓕᖅᑐᑦ ᐊᑦᑭᑎᐊᖅᑯᓯᒪᐃᓇᕐᓂᒍ ᐅᖅᑲᖅᓯᒪᔨᑦ ᐊᓪᓚᓗ ᐊᑦᑭᑎᐊᖅᑯᓯᒪᐃᓇᕐᓂᒍ ᓴᓇᓇᒃᑯᓕᖅᑐᑦ ᐱᓕᕆᓂᖅᓯᒪᔭᖅ ᑲᑎᒪᔨᕿᓪᓚᕆᑐᒃ ᐅᑎᖅᑕᕐᕕᒋᓄᑦ ᑎᒥᐅᔪᓄᑦ ᓴᓇᕐᓂᖅᓯᒪᔭᖅ.

5.4.4 ᓚᕐᓗᑎᒃ ᐊᖏᔪᕕᑎᓂ 5.4.1 ᐊᑎᒪᓗ 5.4.2, ᐊᐅᓪᓚᕆᑐᒃᑯᓕᖅᑐᑦ ᐸᑐᐅᑎᑦ ᓄᓗᐅᐃᖅᑯᓯᒪᓗᑎᒃ ᑭᓯᓂᖏᓐᓂᒃ ᑲᓇᐃᖅᑯᓯᒪᔭᖅ ᐅᖅᑰᓯᐊᑯᓂ ᐃᓗᐊᓂᒃ ᐅᓇᕐᓂᖅ ᐃᓗᐊᓂᒃ ᐅᑎᖅᑕᕐᕕᒋᓄᑦ ᑎᒥᐅᔪᓄᑦ ᑲᑎᒪᔨᕿᓪᓚᕆᑐᒃ, ᐊᑐᓂᑦᓂ ᐊᐃᓐᓇᓂᖅ ᐊᐃᓐᓇᖅᑐᖅᑕᐃᓕᓇᕐᓂᑦ ᐃᓄᐃᑦ ᒃᑐᔾᔨᖅᑎᑦᓯᒪᓂᖅ, ᑲᑎᒪᔨᕿᓪᓚᕆᑐᒃ ᐊᑦᑭᑎᐊᖅᑯᓯᒪᐃᓇᕐᓂᒍ ᐃᓕᓴᕐᓂᖅᓯᒪᔭᖅ ᐊᑐᐸᓇᕐᓗᒍ ᐊᑦᑭᑎᐊᖅᑯᓯᒪᐃᓇᕐᓂᒍ ᑲᓇᓇᒃᑯᓕᖅᑐᒃ ᐅᖅᑰᓯᐊᑯᓂ ᐊᐃᓐᓇᖅᑐᖅᑕᐃᓕᓇᕐᓂᑦ ᐊᑦᑭᑎᐊᖅᑯᓯᒪᐃᓇᕐᓂᒍ ᐃᓄᐃᑦ ᐊᑐᐸᓇᕐᓗᒍ ᐊᑦᑭᑎᐊᖅᑯᓯᒪᐃᓇᕐᓂᒍ ᑲᓇᓇᒃᑯᓕᖅᑐᒃ ᐃᓕᓴᕐᓂᖅᓯᒪᔭᖅ ᐊᑦᑭᑎᐊᖅᑯᓯᒪᐃᓇᕐᓂᒍ ᐃᓄᐃᑦ ᐊᑐᐸᓇᕐᓗᒍ

5.5.5 ᑕᐃᒪᓗ ᑲᓇᑕᒥ ᐅᒪᔨᓕᓚᕐᔪᐊᑯᑦ ᖃᐅᔨᒪᓕᕆᑎᑲ ᐅᑯᐊ ᐳᓕᕋᑦ ᐳᓯᕈᒪᔪᑎᑲ ᓄᓇᓕᕋᓯᓄᑦ, ᐱᖃᑎᑎᕐᐅᑐᓗᑎᑲ ᐳᓯᕋᑦ ᐅᖃᐅᓱᐅᑐᓗᓂ ᖃᑭᒥᖅᑕᐊᖃᑎᖃᖅᑐᓗᑎᑲ ᐊᐅᑐᒍᔨᑦ ᐊᐅᓪᓕᕋᑎᖃᑎᒋᔭᐅᓂᒧᑦ ᑲᑎᒪᔨᕋᓯᓯᑦ. ᐅᑯᐊ ᐊᐅᓪᓕᕋᑎᖃᑎᒋᔭᐅᓂᒧᑦ ᑲᑎᒪᔨᕋᓯᓯᑦ ᐅᖃᐅᔨᔪᓱᒪᓂᐊᖅᑐᑦ ᐳᓯᖅᑎᓄᑦ ᐅᓪᒪᒪᕋᖏᓐᓇᑦᑐᑦ ᐳᓯᓗᐊᖅᓴᓂᖅ.

ᐃᓕᓐᓄ 6 — ᑐᓴᐅᒪᑎᑎᓂᖅ, ᑎᑎᕆᖏᒪᔪᕐᓯᑎᓐᓂᑦ ᐊᓚᒪᓗ ᐃᓗᓪᕐᓯᑎᓐᓂᑦ

6.1 ᑐᑭᒃᖢᑦᓂᐊᕐᔪᑎᒃᓴᑦ

6.1.1 ᐅᖏᓂ ᐃᓕᓐᓄᓂ ᐊᒋᓂ ᑎᑎᕆᖏᒪᔪᑦ ᑐᑭᒃᖢᑦᓂᐊᕐᔪᑎᒃᓴᓐᓂᑦ:

(a) ᑎᑎᕆᖏᒪᔪᑦ ᐃᑕᕐᓂᑕᑦ, ᐃᓕᓴᕐᔪᖏᑕᒃ ᐅᓂᒃᑲᖅ, ᐊᓚᒪᓗ ᐅᓂᒃᑲᖅᑐᐊᑦ ᐃᓚᒪᑲᑦᓕᓂᑕᐃᑦ ᑎᑎᕆᖏᒪᔪᖅ ᑕᒪᒃᔨᓂᐅᓐᓄ ᑲᓇᖑᓕᖅᑕᐃᑦ ᐊᓚᒪᓗ ᐅᑎᖅᑕᖢᑦᑐᓄᑦ ᑎᓐᓕᐊᓄᑦ ᓴᐳᒃᓴᐅᓕᕐᓂᖅ;

(b) ᓇᓗᓇᐃᕐᓗᑎᑦ ᐱᐅᔪᑐᖅᑲᖅᑕᐃᑦ ᐃᓂᓐᓂᑦ ᐱᓚᒪᑕᐅᔪᑦ ᐃᓄᓐᓄᑦ ᐊᓚᒪᓗ ᓂᕐᔪᑎᖅᑲᖅᓵᐅᔪᑦ ᐱᓚᒪᑕᐅᔪᑦ ᐃᓄᓐᓄᑦ;

(c) ᓴᖏᑦᑎᓗᑎᒃ ᑐᑭᔪᓇᖅᔪᑎᑦᑎᔪᓚᕐᓂᒃ ᑎᑎᕆᖏᒪᔪᓚᕐᓂᒃ ᐃᑲᔪᑎᓂᐊᕐᕆᑐᖅᒃ ᐅᓪᓚᕆᑎᓄᑦ ᐅᑯᐊᓗ ᖃᓄᐃᓄᓐᓚᖅᑐᑦᑐᒃ ᑕᓯᑯᓄᑦᑐᑦ ᑕᒪᒃᑯᐊᓂᖅ ᐊᑐᓚᖅᑐᑦ ᐃᓄᓐᓄᑦ ᑎᓐᓕᐊᓄᑦ ᑎᑎᕆᖏᒪᔪᑦ ᐊᓚᒪᓗ ᑎᓐᓕᐊᓄᑦ ᑎᑎᕆᖏᒪᔪᑦ ᐊᑐᓚᖅᑐᑦ ᐊᓚᒪᓗ ᐃᓚᒪᑲᑦᓕᓂᑕᐃ

6.4 ᐱᕐᔪᐊᑎᖅᑐᒃ ᐱᓕᕆᔨᒃᑲᓪᓚᕆᐊᓕᒃ ᐃᓄᖅᓄᑦ: ᑕᒪᐃᓐᓄᑦ ᒪᓕᖅᑕᐅᕐᓚᕆᓕᒃ

6.4.1 ᓄᓇᕗᑦ ᑐᓐᖓᕐᓂᒃ ᑎᒥᖓ ᑲᓐᑐᖏᓪᓗᐊᖅᑐᑦ ᑎᑎᕋᖅᑕᐅᓚᑎᑉᑲ ᐱᕐᔪᐊᑎᖅᑎᑦ ᐱᓕᕆᔨᒃᑲᓪᓚᕆᐊᓕᒃ ᐃᓄᖅᓄᑦ (“ᐅᔪᐊ ᐱᕐᔪᐊᑎᖅᑎᑦ”), ᐊᒋᓐᓂ ᑎᑎᕋᖅᑲᓕᕐᓂ ᐊᖏᕐᕋᖅᑕᐅᖏᓕᕐᓂ ᑲᓐᑐᖏᖅᑕᐅᖏᓕᕐᓄᑦ ᐱᐅᖏᑦ ᐊᒻᒪᓗ ᐱᓕᕆᓐᓂᕐᖏᑦ, ᑕᒪᐃᓐᓄᑦ ᐅᕙᓪᓕᐊᓂᖅ ᐃᓕᓴᖅᑐᑦ ᐊᒻᒪᓗ ᐅᑎᖅᑕᕐᕕᒋᔭᖅᑐᑦ ᑎᓯᒥᐊᓄᑦ ᓴᓇᓕᐅᕋᖅᑕᐅᓚᑎᑉᑲ ᓴᓇᔨᐅᔪᓕᐊᑉᑲ 6-1. ᐅᔪᐊ ᐱᕐᔪᐊᑎᖅ ᐃᓐᓇᕐᓴᓂᑕᐅᓂᐊᖅᑐᑦ ᐊᒻᒪᓗ ᐱᔨᓴᖅᑕᐅᓚᑎᑉᑲ ᐊᑐᓕᖅᑎᑕᐅᔨᓴᓕᐊᖅᑐᖅᓂᑦᑐᖅ ᐸᖅᓇᐅᑎᓂ ᒪᓕᒐᓗᑎᑉᑲ 15.3.1.

6.4.2 ᐱᕐᔪᐊᑎᖅᑎᑦ ᐱᕐᔪᐊᑎᖅᑭᑎᑉᓕᑎᑦ ᐅᔪᐊ:

(a) ᐃᑲᔪᖏᓕᑦ ᐱᓕᕐᓂᐊᑎᖅᑎᓂᖅᓱᖅ ᐊᐅᓪᓂᖅᐱᕐᔪᐊᑎᖅᓴᓕᑦ ᐸᖅᓇᐅᑎᖅᑦ;

(b) ᐃᑲᔪᖏᓕᑦ ᐱᓕᕐᓂᐊᑎᖅᑕᐅᓂᖅᑎᑦ ᑐᑭᓯᓇᖅᓯᑎᖅᓂᖅᑎᓂᖅᓱᖅ ᐃᓄᖅᑎᑐᓪᓚᕆᓂᖅᓱᖅ ᑭᓯᑐᐃᓐᓇᐅᓂᖅ ᐊᓕᓇᖅᓯᒪᔪᑦ ᐃᓪᓕᖅᑐᓂ 6.8;

(c) ᑎᑎᖅᑦ ᑐᓴᐅᒪᓂᖅᑎᔪᑦ ᐱᐅᓯᑐᖅᓴᓕᑦ ᐱᓕᕆᐅᔪᓂᖅ ᐃᓄᖅᓄᑦ; ᐊᒻᒪᓗ

(d) ᐃᑲᔪᖏᓕᑦ ᐱᓕᕐᓂᐊᑎᖅᑕᐅᓂᖅᑎᑎᑦ ᐃᓄᖅᑎᑐᑦ ᓄᓇᕗᑦ ᐊᑎᖏᓕᑦ ᒐᖅᔨᓄᖓ, ᑲᓇᑕᒥ ᐅᓕᔪᖅᑲᓇᑦ, ᐅᑎᖅᑕᐸᓪᓕᓄᑦ ᑎᖏᒥᐊᓄᑦ ᓴᓇᔨᐅᔪᓕᐊᖅᑐᖅ ᐊᒻᒪᓗ ᓄᓇᕗᑦᓴᐅᔨᔪᑦ ᐃᓗᐊᒥᑐᖅ.

6.4.3 ᐱᕐᔪᐊᑎᖅᑎᑦ ᐅᔪᐊ ᐃᓕᓐᓴᐅᓂᐊᖅᑐᑦ ᑎᑭᐅᔪᓚᑎᑦᑎᓂᑦ ᐅᓪᓚᑎᓐᓴᖏᓐᓄᑦ ᐃᓄᐃᑦ ᐱᓇᔪᐊᖅᑎᑲᑎᑦᓯᑦ:

(a) ᐃᓄᐃᑦ ᐅᖅᑲᐅᔪᑐᖅᑎᑦ ᐱᓕᕆᐊᖓᔪᑦ ᐊᒻᒪᓗ ᐃᓕᖅᓂᕐᓕᖅᑐᖅᑐᑦ ᐱᓕᕆᐊᖓᔪᑦ ᐅᓚᐅᐃᖅᓯᒪᔪᑦ ᐃᓕᓐᓂ 6.5 ᐊᒻᒪᓗ 6.6 ᐊᒋᓂᓕ;

(b) ᐅᓚᐅᐃᖅᓯᒪᔪᑦ ᐊᒻᒪᓗ ᑭᓯᓂᖓᑦ ᓂᕐᔨᑎᖅᑐᖅᑐᑦ ᓄᓇᕗᑦ ᐱᓕᕆᐅᔪᑦ ᐃᓄᖅᓄᑦ ᐊᒻᒪᓗ ᐱᐅᓯᑐᖅᑐᖅᓴᖅᑐᑦ ᐃᓂᖓᑦ ᐱᓕᕆᐅᔪᑦ ᐃᓄᖅᓄᑦ; ᐊᒻᒪᓗ

(c) ᐅᓚᐅᐃᖅᓯᒪᔪᑦ ᒐᖅᔨᐊᖅᓱᓇᖅᑐᑦ ᐃᓄᖅᑎᑐᑦ ᐊᑎᖏᓕᑦ ᖃᓇᑕᒥ ᐅᓕᔪᖅᑲᓇᑦ, ᐅᑎᖅᑕᐸᓪᓕᓄᑦ ᓄᓇᕗᑦ ᑲᓐᖓᕐᕕᒋᔭᖅᑐᖅ ᐊᒻᒪᓗ ᐊᑐᑎᖅᑐᑦ ᓄᓇᕗᑦ ᐃᓚᐅᓂ, ᐅᖓᓂ ᐅᖅᑲᖅᓯᒪᔪᑦ 6.7.1 ᐊᒋᓂᓕ.

6.4.4 ᓂᕐᔨᑎᖅᑐᑦ ᓄᓇᕗᑦ ᐱᓕᕆᐅᔪᑦ ᐃᓄᖅᓄᑦ ᐊᒻᒪᓗ ᐱᐅᓯᑐᖅᑐᖅᓴᖅᑐᑦ ᐃᓂᖓᑦ ᐱᓕᕆᐅᔪᑦ ᐃᓄᖅᓄᑦ ᐅᓚᐅᐃᖅᑕᐅᓚᑎᑉᑲ ᑕᒪᓐᓇ ᐅᓂᒃᑲᑐᐊᖓᑎᒍᑦ ᐊᐱᖅᓯᖅᑕᐅᓚᑎᑉᑲ ᓂᐱᑦᑕᐅᖅᑕᐅᓚᑎᑉᑲ, ᕿᒥᕐᔪᐊᖅᑕᐅᓚᑎᑉᑲ ᐃᓐᓇᐅᔭᕋ ᐱᑎᖅᑕᐅᑦᑐᖅ, ᐅᕐᓂᒍᓐᓇᖅᑐᑦ ᕿᒥᕐᔪᐊᖅᑕᐅᓚᑎᑉᑲ ᐃᓐᓇᐅᔭᕋ ᐊᒻᒪᓗ ᐃᓄᖅᓄᑦ ᐱᓕᕆᐅᔪᑦ ᐃᓄᖅᓄᑦ ᐅᓂᒃᑲᑐᐊᖓᑎᒍᑦ, ᐊᐅᑎᓕᓐᓂᖅᑐᑦ ᐅᓂᒃᑲᑐᐊᖓᑎᒍᑦ (ᓄᓇᕗᑦ), ᐱᐅᓯᑐᖅᑕᐅᓚᑎᑉᑲ, ᐊᒻᒪᓗ ᐃᓄᖅᓄᑦ ᐱᐃᓕᖅᑐᖅᑐᖅᓴᖅᑐᑦ ᓄᓇᕗᑦ. ᑕᐃᑲᓂ ᐅᓄᐃᓴᖅᓯᒪᔭᕐᓂᒃᑐᑦ, ᓂᕐᔨᑎᖅᑎᖏᓐᓄᑦ ᓄᓇᕗᑦ ᐱᓕᕆᐅᔪᑦ ᐃᓄᖅᓄᑦ ᐅᓂᒃᑲᑐᖅᑐᑦ ᐃᓕᖅᑎᔭᓂᓂᑦ ᓄᓇᕗᑦ ᐊᒻᒪᓗ ᑭᓯᓂᖓᑦ ᓂᕐᔨᑎᖅᑐᖅ ᐊᒻᒪᓗ ᐱᓕᕆᐅᔨᔪᑎᖅᑐᑦ ᐃᓄᖅᓄᑦ.

6.4.5 ᐱᕐᔪᐊᑎᖅᑐᑦ ᐅᓂᒃᑲᖅᑐᐊᑦ, ᐃᕐᓂᕐᔨᑎᖅᑎᑦ ᐊᒻᒪᓗ ᓂᕐᔨᑎᓂᖅ ᖃᐅᔨᓚᑎᓂᑎᒐᖏᓕᕐᓂ ᐊᑐᖅᑕᐅᓚᑎᑉᑲ ᐱᓇᔪᐊᑎᓂᖅ, ᐊᒻᒪᓗ ᐅᓂᒃᑲᖅᑐᐊᑦ ᐊᒻᒪᓗ ᐃᑕᕐᓂᕐᑕᓕᕆᔨᐊᕐᓕᖅᑐᑦ ᐊᔨᔪᖓᓕᒫᓂᖅ ᐱᓕᕆᓐᓂᕐᖏᓐᓄᑦ ᐊᑕᖅᓯᒪᔪᖅᑐᑦ ᐃᓕᓴᖅᑐᑦ.

6.4.6 ᐱᓕᕆᔭᕐᓕᐊᓕᓂᕐᓂᐊᖅᓯᓚᑎᑦ ᐱᓇᔪᐊᑎᖅᑎᑦ, ᓄᓇᕗᑦ ᑐᓐᖓᕐᓂᒃ ᑎᑎᕋᖅᑕᐅᓚᑎᑉᑲ ᓴᓇᔨᐅᔪᓕᐊᖅᑐᖅᑐᑦ ᐊᒻᒪᓗ ᐃᓄᖅᓄᑦ ᐱᓇᔪᐊᑎᖅᑎᑲᑎᑦᓯᑦ, ᐊᒋᓐᓂᖅᑎᖅᑐᑦ, ᒐᖅᔨᓄᖓ ᐊᔭᕆᐊᓕᖅᑐᖅ (ᓄᓇᕗᑦ), ᐱᐅᓯᑐᖅᑕᐅᓚᑎᑉᑲ, ᐊᒻᒪᓗ ᐃᓄᖅᓄᑦ ᐱᓇᔪᐊᑎᖅᑎᑲᑎᑦᓯᑦ, ᐊᒋᓐᓂᖅᑎᖅᑐᑦ ᐊᐅᓪᓂᖅᐱᕐᔪᐊᑎᖅᑐᖅᓴᖅᑐᑦ ᐊᒻᒪᓗ ᐃᓄᖅᓄᑦ ᐱᓇᔪᐊᑎᖅᑎᑲᑎᑦᓯᑦ

ᑲᑎᒪᓕᕆᓪᓕᐅᑦ ᐊᓚᒐᓗ ᓄᓇᕗᖅᓂ ᐃᓄᐃᑦ. ᑕᒪᕐᒪᑲ ᐱᖏᓂᖅᑎᑦ ᓴᓇᔨᐅᔪᑦ ᒪᓕᖅᑎᐊᖏᓐᓂᑲᑎᑲ ᑕᐃᒪᓐᓇᖅᑐᑦᓴᓂᐅᑉᑐᐊᔪᓂᖅᑐᑦ *ᓄᓇᕗᒻᒥ ᐃᑕᖏᑕᓕᕆᓂᖅ ᐊᓚᒐᓗ ᐅᔾᔨᖓᖏᓂᖅᓯᒪᔪᓂᖅᑐᑦ ᐃᓂᖓᓄᑦ ᐱᖏᓂᕆᓕᕆᓂ* ᐊᓚᒐᓗ ᐃᓕᖅᓱ 33 *ᓄᓇᕗᒻᒥᖅ ᓄᓇᕆᑕᑎᓄᑦ ᐊᖓᑕᑕᑎᑦ*, ᐊᑐᑎᖏᑎᓪᓗᑎᑦ.

6.4.7 ᑎᑭᐅᑎᓯᒪᓗᐊᔪ ᐊᑐᑎᖏᑕᖅᓴᓂᖅᓱᒍᑦ, ᓄᓇᕗᑦ ᑐᖅᖁᓕᕐᑲᑦ ᑎᑎᑦ ᐊᑐᐃᖅᓴᐅᓕᓂᖏᑎᓪᓗᑎᑦ ᐱᖏᑎᑦ ᐊᖅᑐᑎᖏᑕᖏᑐᓂ ᐃᓂᖓᖅᑐᑦ ᐊᒡᔪᓄᑦ ᓄᓇᕗᖅᓄᑦ.

6.4.8 ᓄᓇᕗᑦ ᑐᖅᖁᓕᕐᑲᑦ ᑎᑎᑦ ᐱᓕᓴᑎᐊᖏᓂᕐᓂᖏᖅᑐᑦ ᐊᑐᐃᖅᓴᐅᓕᓂᖏᑎᑦᓂᑎᓯᒪᔪᓗᑎᑲᑎᑲ ᐊᐅᓕᖏᑎᖏᑦᑐᖅᓴᓂᖅᑐᑦ ᑲᑎᕕᖅᑕᕐᓂᖅ ᓄᑖᖅᑐᑦ ᐃᓕᓴᐅᑎᓪᓗᑎᑦ ᑭᖑᓂᖓᓄᑦ ᐱᖏᓂᖅᑐᑦ ᐊᐅᓕᖏᑎᖏᑦᑐᖅᓴᓂᖅᑐᑦ ᐊᐅᓕᖏᑎᖏᑦᑐᖅᓴᓂᖅᑐᑦ <ᖏᓄᐅᑎᑦ.

6.5 ᐅᓂᒃᑐᑦᑕᕆᐃᑦ ᐃᓚᒪᑦᓕᓂᑕᓂᒃ ᐱᓕᕆᐊᖓᔪᖅ

6.5.1 ᐅᑯᐊ ᐅᓂᒃᑐᑦᑕᕆᑐᕆᔪᑦ ᐃᓕᖏᑦᓗᑎᒃ ᐊᐱᒃᓱᐃᓗᑎᒃ ᖃᐅᐅᔨᖏᓗᑎᒃ ᖃᐅᐅᔨᒪᔭᖓᓂᒃᓂᓂᒃ ᐃᓂᓇᑦᑕᕆᐃᑦ ᐊᓚᒐᓗ ᖁᑎᖏᑦᑕᐊᖏᓗᑎᑦ ᐊᑐᐃᖅᓴᐅᓕᔪᑦ ᑎᑎᕐᓗᑎᑦᓂᑎᓯᒪᔪᓗᑎᑲᑎᑲ ᐃᓕᓐᓂᐅᓗᑎᑦ ᑎᑎᕐᓗᑎᑦᓂᑎᓯᒪᔪᓗᑎᑲᑎᑲ 6-1 ᑲᓇᑕᒥ ᐅᒪᔪᕆᑲᓴᐃᑦ ᐊᓚᒐᓗ ᐅᑎᖅᑕᕐᕕᒃᓂᑦ ᑎᓐᓕᐊᓂᑦ ᓴᓗᐅᐅᓕᓂᖅᑐᑦ. ᐊᑐᑎᖏᑕᖅᑐᑦ ᐊᐱᖅᓱᑕᐅᔪᑦ ᖃᐅᐅᔨᓴᐅᑎᑦ ᑭᓱᑕᓚᓂᖓᓄᑦ ᐃᓕᖏᑦᓗᑎᒃ, ᖃᓄᐃᑦᐃᓐᓇᐅᓕᖏᑎᖏᑦᑐᖅᓴᓂᖅᑐᑦ ᐃᓂᓇᑦᑕᕆᐃᑦ ᐃᓱᕐᑕᓚᓂᖓᓄᑦ ᐃᓕᓐᓂᐅᓗᑎᑦ:

(a) ᐃᓚᒪᑦᓕᓐᓂᖅ ᐃᓕᑭᔪᓱᓂᖓᓄᑦ ᐃᓄᐃᑦ, ᓂᔪᐊᑎᑦ ᓄᓇᖅᓴᓂ, ᐊᓚᒐᓗ ᐊᖏᐅᓱᔪᓱᓂᖓᓄᑦ ᐊᓚᒐᓗ ᓇᓂᓐᓂᖓᓄᑦ;

(b) ᐅᓂᒃᑐᖅᑐᐊᑦ ᐊᓚᒐᓗ ᐅᓂᒃᑐᑦ;

(c) ᓄᓇᕗᓐᖏᖅᑕᑦᓕᐊᓱᑎᐊᖅᓯᒍᑎᒃ; ᐊᓚᒐᓗ

(d) ᐃᓂᖓᓄᑦ ᐊᓚᒐᓗ ᑭᓱᓂᖓᓄᑦ ᐱᐅᓱᑐᖅᑎᖅᑎᕐᓂᐅᕗᑦ ᐃᓂᖓᓄᑦ ᐱᓚᒪᐅᕗᑦ ᐃᓄᖓᓄᑦ ᐊᓚᒐᓗ ᓂᔪᐊᑎᖅᑎᕐᓂᐅᕗᑦ ᓄᓇᐃᑦ ᐱᓚᒪᐅᕗᑦ ᐃᓄᖓᓄᑦ.

6.5.2 ᐱᓕᕆᔨᒪᔭᑦ ᐃᓂᖓᓄᑦᑐᖅᑎᑦ ᐊᑕᔨᒥᑦ ᓄᓇᕗᖅᓂᓕ ᐊᑐᑎᖅᑎᖅᑎᖅᑐᑦ ᑭᖅᑯᓄᖅᓱ ᐃᓄᐃᑦ ᐱᓇᓱᐊᖅᑎᑦᓂᑎᓯᒪᓕᖅᑐᓂᒃ ᐳᓴᕐᓗᑎᒥ ᐃᑕᖏᑕᓕᕆᓂᖅᑐᑦᓴᓂᐅᑉᑐᐊᔪᓂᖅᑐᑦ ᐃᓂᖓᓄᑦ ᐊᓚᒐᓗ ᑐᓂᓗᑎᒥ ᐃᓚᖏᑕᕐᓂᖅᑐᑦ ᐃᓂᓇᑦᑕᕆᐃᑦ ᐃᓱᕐᑕᓚᓂᖓᓄᑦ ᐊᓚᒐᓗ ᑐᖓᔪᖅᑐᑦᓴᓂᐅᑉᑐᐊᔪᓂᖅᑐᑦ ᐅᓂᒃᑐᖅᑐᐊᑦ ᐊᓚᒐᓗ ᐃᓄᐃᑦ ᐱᓇᓱᐊᖅᑎᑦᓂᑎᓯᒪᓕᖅᑐᓂᒃ.

6.5.3 ᓄᓇᕗᑦ ᑐᖅᖁᓕᕐᑲᑦ ᑎᑎᑦ ᑲᓇᑐᖅᑲᓱᒪᓗᑎᑲᑎᑲ ᐊᑐᓂ ᑐᓴᕐᓂᕐᓂᔨᑦ ᑎᑎᕐᓗᑎᑦᓂᑎᓯᒪᓗᑎᑲᑎᑲ ᐊᐱᖅᓱᑎᑦ ᖃᐅᐅᔨᓴᐅᑎᑦ ᐅᖃᐅᔨᓴᐅᓂᖅᑐᑦ ᐃᓚᒪᑦᓕᓐᓂᐅᓂᓴᓂᑦ ᐱᓕᕆᔨᒪᔭᑦ, ᐊᐅᓱᐊᖅᑎᑲᑎᑲ ᐅᓂᒃᑐᖅᑐᐊᑦ ᐊᓚᒐᓗ ᐃᓄᐃᑦ ᑐᓴᕐᓂᕐᓂᔨᑦ ᐊᓚᒐᓗ ᐃᓄᐃᑦ ᐱᓇᓱᐊᖅᑎᑦᓂᑎᓯᒪᓕᖅᑐᓂᒃ.

6.5.4 ᐱᓕᕆᐊᑎᑕᐅᑎᓪᓗᑎᑦ ᐊᓚᒐᓗ ᑐᓂᐅᖅᑎᑕᐅᓗᑎᒃ ᐅᖃᐅᔨᑐᖅᑎᑦ ᐃᓚᒪᑦᓕᓐᓂᑕᖅᑐᑦ ᐊᑕᒍᑦ ᐅᒪᔪᓕᖅᑐᑦ ᐃᓕᓐᓂᐅᓗᑎᑦ ᐊᓚᒐᓗ ᐃᓄᐃᑦ ᑐᓴᕐᓂᕐᓂᔨᑦ ᐃᓚᒪᑦᓕᓐᓂᑕᖅᑐᑦ ᐅᖃᐅᔨᑐᖅᑎᑦ ᐱᓕᕆᔨᒪᔭᑦ ᐃᓄᐃᑦ ᐱᓇᓱᐊᖅᑎᑦᓂᑎᓯᒪᓕᖅᑐᓂᒃ ᐊᑐᐃᖅᓴᐅᓕᓂᖏᑎᖏᑦᑐᖅᓴᓂᖅᑐᑦ ᐊᓚᒐᓗ ᐃᓄᐃᑦ ᐱᓇᓱᐊᖅᑎᑦᓂᑎᓯᒪᓗᑎᑲᑎᑲ

ᓄᓇᐃᑦ, ᓇᓗᓇᐃᖅᑕᐅᓂᒪᔪᑦ ᖃᑎᖅᑕᐋᖅᑎᑎᕐᓴᓗᑎᒃ ᐃᓕᓴᐅᔭᖅᓯᓚᐅᕙᓪᓕᐊᑦ, ᐋᐅᓪᓚᖅᑎᑎᕐᓴᐅᓕᖅᓯᓂᖅ ᑲᑎᒪᔨᖏᓪᓂ ᐋᒪᓗ ᓄᓇᓕᖅᓱᓂ ᐃᓄᐃᑦ. ᒥᐅᓇᐅᔪᖅ ᐊᑐᖅᑕᐅᔪᑦ ᐱᑕᖅᖃᐅᔪᑦ ᐃᓐᓇᐃᒪᓕ ᐋᖅᑎᓕᖅ ᐊᐅᓚᑦᑐᑕᐅᔪᑦ ᖃᐅᔨᓴᖅᑕᐅᔪᑦ ᐃᓕᓴᐅᔭᖅᓯᓚᐅᕙᓪᓕᐊᑦ, ᐊᐅᓚᑦᑐᑕᐅᔪᑦ ᐃᓕᓴᐅᔭᖅᓯᓚᐅᕙᓪᓕᐊᑦ ᑲᑎᒪᔨᖏᓪᓂ ᐃᓕᓴᐅᔭᖅᓯᓚᐅᕙᓪᓕᐊᑦ ᐃᓚᖃᑎᒌᖅᑐᑦ

6.8 ᑐᑭᑐᖏᓐᓂᒃ ᑎᑎᕆᔨᒪᔪᓂ

6.8.1 ᓄᓇᕗᑦ ᑐᓴᐃᓚᖅ ᑎᒥᖓ ᓴᖅᑭᑎᑎᓂᐅᕐᓂᖅᑐᑦ ᐅᖃᖅᑎᖅᓯᒪᔪᓂ ᑲᑎᒪᔨᖅᑎᑦ ᐱᖓᓱᐃᓐᓇᖅᑐᑦ ᑐᑭᓯᐅᑉ ᐱᓕᕆᔨᒃᑯᑦ ᑐᑭᓯᐅᑉ ᐱᓕᕆᔨᒃᑯᑦ, ᓴᖅᑭᑦᓴᖅᑐᑦᓴᖅᑎᑦ, ᐅᖃᓕᒫᓂᖅ ᐱᓕᕆᔨᒃᑯᑦ, ᐊᓯᔪᓐᓇᖅᑐᑦ ᑐᑭᓯᐅᑉ ᐱᓕᕆᔨᒃᑯᑦ ᐱᒻᒪᕆᐅᑉ ᐃᓚᖓ ᐊᓯᔪᓐᓇᖅᑐᑦ ᐱᓕᕆᔨᒃᑯᑦ ᐊᑐᖅᑐᑦ ᐊᓯᔪᓐᓇᖅᑐᑦ ᐱᒻᒪᕆᐅᑉ ᐃᓚᖓ

(d) ᐊᓯᖏᑦ ᐊᐢᑐᑎᓯᑦ ᑐᓴᖅᑕᐅᓯᒪᓕᕆᖅᑐᑦ ᖃᐅᔨᓴᕆᐅᑎᐊᖅᑎᓪᓗᐊᖅᓯᒪᒃᑕ ᑲᓇᑕᒥ ᐅᒪᓯᕐᒥᖅᓂᒃ ᐅᕙᓪᓕᐊᑉᐸᑦ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᖏᒥᐊᓄᑦ ᓴᐳᒪᐅᓄᑦ ᐊᓚᒪᓐᓇ ᐊᕐᓇᖅᑐᖅᓯᒪᓂᖏᑦ ᑐᓴᕐᑎᑕᐅᔨᓴᓪᓚᕆᑦ, ᐅᕙᓪᓕᐊᓂᑉᐸᑦ ᐃᓄᐃᑦ ᓇᓗᓇᖅᑐᖅᑐᑦ.

6.9 ᐊᑐᖅᓂᖅ ᓄᓇᕗᒻᒥᑦ ᓇᐸᐸᖅᑕᐅᔨᓴᓂ

6.9.1 ᑕᐃᒪᓐᓇ ᑲᓇᑕᒥ ᐅᒪᓴᓕᓚᕐᔪᐊᕈᑦ ᓴᖅᑭᑦᓯᑎᐃᓇᖅᓴᖅᑎᑦ ᐱᒃᓗᑎᑦ ᑲᓇᑕᒥ ᐅᒪᓴᖅᑐᖅᓂᒃ ᐊᓯᖏᑦ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᖏᒥᐊᓄᑦ ᓴᐳᒪᐅᓄᑦ, ᐊᑐᑎᖅᑐᒥᐅᖅᑐᑦ ᐅᓪᓗᒥᑦ ᐊᓚᒪᓐᓇ ᑕᐃᒪᓐᓇᐃᑐᓗ ᐊᐸᑦᑕᐅᔨᓴᓂᖏᑦ ᐊᑕᓴᓂ ᓄᓇᕗᒻᒥᑦ.

ᑭᓇᐅᔨᒪᓂᖅ ᐊᓚᒪᓐᓇ ᐃᓕᓴᕐᓂᖅ ᑲᑎᑎᖅᑕᐅᔨᓴᓂ ᐊᖏᑕᐅᓂᖅ 6-1 (ᒪᓕᑐᑦ 6.4.1, 6.5.1, 6.6.1 ᐊᓚᒪᓐᓇ 6.8.4)

ᑲᓇᑕᒥ ᐅᒪᓴᖅᑐᖅᓂᒃ ᐊᓚᒪᓐᓇ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᖏᒥᐊᓄᑦ ᓴᐳᒪᐅᓄᑦ	ᖃᐅᔨᒪᖅᓯᒪᓂᖏᑦ ᓄᓇᓕᓂᑦ
1. ᐊᐳᐸᐃᑦ ᑲᓇᑕᒥ ᐅᒪᓴᖅᑐᖅᓂᒃ	ᖃᑭᖅᑕᖅᓴᕐᕕᖅ
2. ᓴᓐᓂᑦ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᖏᒥᐊᓄᑦ ᓴᐳᒪᐅᔨᓴᓕᓴᐅᖅ	ᒥᑦᑎᒪᒪᑕᓚᐅᖅ
3. ᐃᓱᓚᐅᓴᖅᓂᐅᖅ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᖏᒥᐊᓄᑦ ᓴᐳᒪᐅᔨᓴᓕᓴᐅᖅ	ᑭᓇᖏᐃᑦ
4. ᖃᐅᖅᓴᐅᖅᑐᑲᖅ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᖏᒥᐊᓄᑦ ᓴᐳᒪᐅᔨᓴᓕᓴᐅᖅ ᑎᖏᒥᐊᓄᑦᑎᐹᐸᐅᑐᓄᑦ ᓴᐳᓇᓂᕐᓴᖅ	ᓴᓪᓕᓂᑦ
5. ᐃᐅᓱᑐᐊᖅᑐ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᖏᒥᐊᓄᑦ ᓴᐳᒪᐅᔨᓴᓕᓴᐅᖅ	ᓴᓪᓕᓂᑦ
6. ᓂᓇᑎᓇᑐᓂ ᑲᓇᑕᒥ ᐅᒪᓴᖅᑐᖅᓂᒃ	ᑲᓇᑎᖅᓵᐃᒪᐃᓇᐅᖅ
7. ᑯᐃᓴᓛᖅ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᖏᒥᐊᓄᑦ ᓴᐳᒪᐅᔨᓴᓕᓴᐅᖅ	ᐊᖅᓚᐊᑦ
8. ᓂᓴᐊᑎᖅᑐᖅᓚᐅᖅ ᑲᓇᑕᒥ ᐅᒪᓴᖅᑐᖅᓂᒃ	ᐊᐅᓱᐃᐅᑦᑐᖅ
9. ᑐᖅᑐᒥᐊᖅᓚᐅᐸᑦ ᓇᖅᓴᓇᑲ ᑲᓇᑕᒥ ᐅᒪᓴᖅᑐᖅᓂᒃ ᓄᓇᖅᑯᑎᓂᖏᑦ	ᖃᑲᐅᓱᐃᐅᑦᑐᖅ
10. ᐊᐳᑦ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᖏᒥᐊᓄᑦ ᓴᐳᒪᐅᔨᓴᓕᓴᐅᖅ	ᖃᑲᐅᓱᐃᐅᑦᑐᖅ, ᐃᐳᐱᐊᖅᓴᖅ
11. ᖃᑲᖅᑯᓪᓗᐃᑦ ᑲᓇᑕᒥ ᐅᒪᓴᖅᑐᖅᓂᒃ	ᖃᑭᖅᑕᖅᓴᕐᕕᖅ
12. ᓱᒧᐊ ᐅᑎᖅᑕᐸᑦᑐᓄᑦ ᑎᖏᒥᐊᓄᑦ ᓴᐳᒪᐅᔨᓴᓕᓴᐅᖅ	ᐃᖅᑲᓪᓕᑦᑐᖅᓴᖅ, ᐅᖅᓱᖅᓕᖅᓴᖅ, ᐅᒥᖓᒪᓕᖅᓴᖅ

ᐃᓕᓐᓴ 7 — ᐊᓕᕆᔨᑐᑎᓂᖅ ᐊᒻᒪᓗ ᐊᓯᖏᑦ ᓇᖏᑕᓂᕆᔨᒃᓴᓂ ᐃᑲᔪᑎᔪᑦ

7.1 ᑐᑭᒃᓯᒪᓕᕆᔨᓂᑎᑦ

7.1.1 ᐅᕐᓂ ᐃᓕᖏᓄ ᐊᒋᓂ ᑎᑎᕆᔨᓯᒪᔪᑦ ᑐᑭᒃᓯᒪᓕᕆᔨᑎᑦ:

(a) ᖃᐅᔨᓴᒪᐅᕆᑎᐊᓕᕆᓐᓇᑎᑦ ᑭᓴᐅᔭᓕᐅᑭᓐᓇᖅᑐᓂᖅᐳᑦ ᐃᒪᒥᓂᖃᓯᑭᓐᓇᖅᑎᑦ ᐊᒻᒪᓗ ᐱᐅᔨᑐᖃᖅᑐᒃᓴᑎᒃ ᐊᒻᒪᓗ ᐃᓅᔨᖏᑦᑎᐊᖅᓴᑎᒃ ᐃᓅᐃᑦ;

(b) ᐱᓯᒪᓕᐊᑎᕆᑎᓇᑎᒃ ᓅᓚᒃᑐᑐᓂ ᓴᓕᖅᐅᔪᓐᓇᖅᑐᑦ ᐊᒻᒪᓗ ᐊᑎᖏᑦ ᑭᓇᐅᔭᓕᐅᑕᐅᔪᓐᓇᖅᑐᑦ ᐃᓗᐊᓂ ᐊᒻᒪᓗ ᓴᓇᔨᒥᒃ ᐅᒪᖅᓯᓐᓂᑦ ᑎᓐᓕᐊᓂ ᓴᓐᖅᓴᐅᔨᓚᒃ;

(c) ᐱᓴᓇᐅᖅᑲᑎᑦᓴᓂᑦ ᐊᒻᒪᓗ ᑲᔪᔨᔪᒥᒃ ᐃᑲᔪᖅᑕᐅᕐᓚᖅᑎᖅᑐᑎᒃ ᐃᓄᖓᓄᑦ ᐃᓗᐊᓂ ᐊᑕᔪᑦ ᓄᓇᓕᒫᑦ;

(d) ᐊᔪᖓᑎᑦᓴᓂᑦ ᐃᓄᐃᑦ ᓴᖅᑭᑎᔨᓴᑎᒃ ᐊᒻᒪᓗ ᐁᖅᑕᑎᐊᖅᔨᒪᐅᓇᖅᓴᐃᒍ ᐅᓪᓗᕆᑕᐅᓂᖅᓴᒧᑦ ᓇᖏᑦᑐᖅᑲᑎᕆᔨᖅᑕᖅ ᐊᒻᒪᓗ ᐃᑕᐅᓂᐊᖅᓴᐃᒍ ᐃᖅᑲᓇᐃᕐᓂᐊᖅᓴᐃᒍ;

(e) ᐅᓄᖅᓯᒋᐊᖅᓴᓂᑦ ᐅᓪᓗᕆᑕᐅᓂᖅᓗᑦ ᐱᓯᒪᔪᑦ ᐱᔭᑎᑎᕐᕈᑎᑦ ᐊᒻᒪᓗ ᓄᓇᑕᖅᓂ ᐅᓪᓗᕆᑕᐅᓚᑦᑐᖅᑲᑎᐊᑕᖅᓴᓂᐅ ᐊᑕᔪᑦ ᓄᓇᓕᒫᑦ;

(f) ᐅᓄᖅᓯᒋᐊᖅᓴᓂᑦ ᒥᖅᔪᑦ ᓄᓇᒥᒃ ᓯᑎᑦᑎᐅᖅᑭᑦᑐᑦ ᐅᓪᓗᕆᑕᐅᓂᖅᓴᖅ ᐊᐃᑐᑎᔪᑦ ᑲᓇᑕᑦ ᐃᒪᔪᖅᑎᓴᑦ ᐊᒻᒪᓗ ᐅᑎᖅᑕᑦᑐᓄᑦ ᑎᓐᓕᐊᓄᑦ ᓴᓐᖅᓴᐅᔨᓚᒃᑎᒃ ᐃᓄᖓᓄᑦ; ᐊᒻᒪᓗ

(g) ᐃᑲᔪᑎᔪᑦ ᑲᔪᓯᐅᓕᐅᓇᖅᑐᓂᒃ ᐱᓕᕆᔨᐊᑎᑎᓂᖅᓴᖅ ᓄᓇᕐᓂ ᓇᓗᓇᐃᖅᑕᐅᓯᒪᓂᖓ.

7.2 ᐃᓄᐃᑦ ᐅᓪᓗᕆᑕᐅᓚᑦ ᑭᓇᐅᔭᖅᑎᖓᑦ

7.2.1 ᓄᓇᕐᓂ ᑐᐃᖓᓕᒃ ᑎᒥᖓ ᐊᐅᓪᓚᕆᔨᐅᓗᑎᒃ ᐃᓄᐃᑦ ᐅᓪᓗᕆᑕᐅᓚᑦ ᑭᓇᐅᔭᖅᑎᖓᐃᓂᒃ ᐊᖏᕐᓴᕐᓪᓚᕆᐊᑎᐃᓂᐊᕐᒪᑦ ᐃᓗᓪᓕᖅᑐᐃᓇᖅᓴᓂᐅ ᐊᐅᑦᑕᐃᓂ ᐃᓄᐃᑦ ᐅᓪᓗᕆᑕᐅᓚᑦ ᐊᑕᔪᓂ ᓄᓇᕐᓂ ᐊᒻᒪᓗ ᐱᐅᓪᓚᕆᐊᑎᐃᓂᒃ ᐊᐃᑎᖃᖅᑐᑐᓂ ᓄᓇᕐᓂ ᐅᓪᓗᕆᑕᐅᓂᖅᓴᖅ ᐱᔭᑎᑎᕐᕈᑎᑦ. ᒪᓕᒐᐊᓕ 15.2.1-ᒥᒃ, ᑲᓇᑕᒥ ᐅᒪᔨᓕᓚᔨᖅᐊᑯᑦ ᑐᓂᔪᓚᓂᐊᖅᑐᑦ ᑲᓇᑕᒥ ᑕᒪᑯᓂᖓ ᐃᓄᐃᑦ ᐊᐅᑦᑕᐃᓂᐊᖅᓴᓂᐅᓄᑦ ᐱᐅᓪᓚᕆᐊᑎᐃᓂᐊᕐᒪᑦ ᐊᑐᕐᓴᐃᓂ ᐊᑐᕐᓴᐃᓂ ᑭᓇᐅᔭᖅᑎᖓᑦ ᖃᓄᖅᑭᑦᑐᖅᑲᑎᐊᑕᖅᓴᓂᐅ ᐱᐅᓪᓚᕆᐊᑎᐃᓂᐊᕐᒪᑦ 5-ᒥ ᐊᖏᕐᕈᑎᓂ 15-1 ᑭᒃᑯᓄᖓ ᐃᓄᐃᑦ ᐅᓪᓗᕆᑕᐅᓚᑦ ᑭᓇᐅᔭᖅᑎᖓᓂᐅᓂᖅ.

7.2.2 ᑭᓇᐅᔭᖅᑎᐃᑦ ᑐᓂᐅᕆᔨᐅᑕᐅᓚᑦ ᐅᑯᓄᖓ ᐊᒻᒪᓗ ᐊᔨᔨᑎᓂᑦ ᐊᑐᖅᓂᐊᖅᑕᖏᑦ:

(a) ᐱᓴᓇᐅᖅᑲᑎᕐᕈᑎᓂᖅ ᐃᓄᖓᓂ ᐅᓪᓗᕆᑕᐅᓚᐅᓂᒃ ᐅᑯᓂᖓ, ᐊᓯᖏᑐᓂᓗ:

(i) ᑎᑎᕐᓂᖅ ᑭᓇᐅᔭᖅᑲᓂᑦ ᑐᒃᓯᕐᕆᐅᑎᓂᒃ;

(ii) ᑎᑎᕐᓂᖅ ᓇᖏᑦᓂᑯᑎᖓᓂᑦ ᐁᖅᑭᒃᓯᒪᓚᖅᑲᓂᒃ ᐁᖅᑭᒃᓯᒪᓕᐅᑎᖏᑐᓂᓗ;

(iii) ᑎᑎᕋᖏᓪᓗᑎᑦ ᓴᓪᓕᔪᓐᓇᖅᑐᖏᓪᓗᒍ ᐊᑐᖅᑕᐅᓂᐊᖅᑐᑦ ᐅᑯᐊᓗ ᑎᑭᐅᑎᓂᒃᓴᖅ ᐱᓕᕆᔨᖅᑐᑦ ᐊᓂᒍᓐᓇᖅᑐᑦ ᐊᑐᖅᑐᑦ ᐊᒻᒪᓗᓐ ᓄᓇᕗᒻᒥᐅᑏᑦ;

(iv) ᐱᓕᕆᓚᐅᑎᓂᓗᑎᒃ ᐃᑲᔪᖏᑦᑕᖅᑕᑦ ᐊᓪᓚᖅᑎᑦ ᐱᔪᓂᑎᕋᐅᑎᓂᒃ ᐅᑯᐊᓗ ᑎᑭᐅᑎᓂᒃᓴᖅ ᖃᓄᐃᓕᓚᐅᖏᓪᓗᒍ ᓴᖅᑭᑐᓂᒃ ᐊᒻᒪᓗᓐ ᖃᓄᖅᑐᑦ ᑭᖑᓪᓕᖅᑐᖅ;

(v) ᓂᐅᕐᕕᓂᖅᓴᖅ ᐊᓪᓚᖅᑎᓂ ᐊᑐᖅᑕᐅᓂᐊᖅᑐᓂᒃ ᑲᑎᖓᔪᓂ ᐊᒻᒪᓗᓐ ᐊᓕᖅᑎᓂᒃ ᐱᔪᓂᑎᕋᐅᑎᓂᒃ;

(vi) ᐋᕐᑭᓂᐊᓕᒫᓂᖅᓯᓚᑦ ᓇᖕᒥᓂᓕᖓᔪᓈᑎᑭᓂᑎᓂᒃ ᐊᒻᒪᓗᓐ ᑭᓇᐅᔭᓕᐊᓕᖓᑎᓂᒃ; ᐊᒻᒪᓗᓐ

(vii) ᑐᓂᓕᓂᖅ ᑐᑭᓕᐅᑎᖓᑦᓂᒃ ᓄᓇᒥᐅᑕᐃᑦ ᐊᒻᒪᓗᓐ ᐱᐅᓕᑐᖅᓄᓕᑦ ᐱᔪᐊᑎᖅᑦ;

(b) ᐃᓕᓴᐅᕐᓂᑎᓂᓗᑎᒃ ᐱᓕᕆᔨᒪᔪᑦ ᐃᓄᖕᓂ ᐅᑭᐅᑐᑎᓕᕆᓂᐅᕐᓂᒃ ᐅᑭᐅᑐᑎᓂᓪᓗᑎᑦ ᐱᓕᕆᑦᑐᕐᓂᑦ ᑐᓂᓕᓂᑦᑐᓂᒃ ᐅᑭᐅᑐᑎᓂᓪᓗᑎᑦ ᐱᓕᕆᔨᒪᔪᑦ ᐃᓄᖕᓂ ᐊᒻᒪᓗᓐ ᐱᓕᕆᔨᖅᑕᐅᓂᐊᖅᑐᓂᒃ;

(c) ᑭᖑᓂᓚᐅᒃ ᓄᓇᕗᖕᓂ ᐱᔪᓂᑎᕋᐅᑎᓂᒃ ᐅᑭᐅᑎᓄᓂ, ᐊᒻᒪᓗᓐ, ᐱᓚᐊᖅᑐᒥᒃ, ᐅᑭᐅᑎᓂ ᐅᒥᐊᖅᓯᐊᓂᑦᑕᓂ;

(d) ᐊᑭᓕᖅᑐᑦ ᓇᖕᒥᓂᔨᑎᖅᑎᓂᖅᓯᓐᓂᓄᓂ ᐊᒻᒪᓗᓐ ᓇᓕᓕᐅᔨᒪᓂᒃ ᐊᑭᖕᓂ ᐊᒻᒪᓗᓐ ᑕᐃᒪᐃᓕᑐᓄᓂ ᐊᑭᓕᖅᑐᑐᒃᑯᑎᕐᓂᖅᓯᓕᕆᓕᓕᖅ; ᐊᒻᒪᓗᓐ

(e) ᐱᓕᕆᓚᐅᑎᓂᓗᑎᒃ ᐅᑭᐅᑐᑎᓕᕆᓂᖅᓴᖅ ᐋᕐᑭᐳᒪᓴᐊᑎᖅᓴᓂᒃ, ᐅᑭᐅᑐᑎᓄᓂ ᐊᑐᖅᑕᐅᓂᐊᖅᑐᑦ ᐊᒻᒪᓗᓐ ᓂᐅᕐᕕᓂᖅᓴᖅᑕᐅᓂᐊᖅᑐᑦ <ᖃᓄᐅᑎᑦ, ᑕᐃᒃᓂᓂ ᐱᓕᕆᔨᓚᐅᑎᑦ ᐊᑕᔪᑦ ᓄᓇᓕᖕᓂ.

7.2.3 ᐅᑯᐊ ᑭᓇᐅᔭᖅᓴᐃᑦ ᐊᑐᐃᓐᓇᐅᓂᐊᖅᑐᑦ ᑕᖅᑯᓄᖓ ᐊᒻᒪᓗᓐ ᐃᓕᓴᐅᑕᐅᓚᐅᑎᓂᒃ ᐃᓄᐃᑦ ᐅᑭᐅᑐᑎᓕᕆᓚᐅᓂᒃ ᐊᑐᖅᑕᐅᓂᐊᖅᑎᓚᐅᑎᒃ ᒪᑕᒃᓚᐅᑦ ᓇᓚᐅᐃᖅᓯᒪᔪᑦ 7.2.2 ᐊᒻᒪᓗᓐ ᒪᑕᒃᓚᐅᑎᒃ ᐅᑯᓂᖓ ᐃᒪᐃᑕᖓᔪᖏᓪᓗᒍ ᐊᑐᖅᑕᐅᓂᐊᖅᑎᓚᐅᑎᒃ:

(a) ᖃᓄᐅᔨᒪᔭᐅᑎᐊᑕᑎᑕᓂᓚᐅᑎᒃ ᐅᕐᓀᑦᓂᓄᖅᑐᑦ ᑐᑭᓯᓇᖅᓯᒪᔨᓚᐅᑦ ᐱᐅᓯᑐᖅᓴᓂᐅᑦ ᐊᓂᒍᓐᓇᖅᑐᑦ ᑕᖅᑯᓄᖓ ᐅᒃᑯᒪᔪᓄᑦ ᐅᕐᓀᑦᓂᓄᖅᑐᑦ ᐅᑭᐅᑐᑎᓕᕆᓂᐅᕐᓂᒃ ᐊᑐᖅᑕᐅᓂᐊᖅᑐᑦ ᐅᑭᐅᑐᑎᓕᕆᓂᐅᕐᓂᒃ ᐃᓄᐃᑦ ᐅᑭᐅᑐᑎᓕᕆᓚᐅᑎᒃ

ᐃᓕᓐᓴ 8 — ᐃᓄᐃᑦ ᑲᓇᑕᖏᑦᑐᓂᒃ ᐊᓚᒐᓗ ᓇᓗᒋᓂᖅᓯᑦ ᐱᕆᖅᑎᑕᐅᓗᑕᒃ

—

8.1 ᑐᑭᖏᖅᓯᒪᓕᕆᔨᐸᑎᒃᖏᒪᑦ

8.1.1 ᐅᕐᓂᑕ ᐃᓕᓐᓄᓂᑕ ᐊᒋᑕ ᑎᑎᕋᖏᓪᓗᒪᔪᑦ ᑐᑭᖏᖅᓯᒪᓕᕆᔨᐸᑎᒃᖏᒪᖕᓐᓂᑦ:

(a) ᖃᐅᐅᓚᓴᐅᓕᓐᑎᑦᓯᑐᑦ ᐃᓄᐃᑦ ᐃᓕᐅᑎᑦᓯᑐᑎ ᖃᐅᓇᑕᒥ ᐅᒪᓚᕆᔨᕐᓇᐸᑎᑦᑕ ᒪᕐᒪᒃᑯᓂ ᑲᓇᑕᖏᑦᑐᓂᒃ ᓄᓇᕗᑦ ᐅᓪᓗᐃᖏᑕᐅᓪᓚᕐᓂᒪ, ᒪᓕᓐᑎᐊᖅᓯᓂᐅᑎ ᐃᓕᓐᓄ 24 ᑕᒃᑯᓂᐅᑎ *ᓄᓇᕗᒻᒥᒃ ᓄᓇᑕᑭᑎᓄᑦ ᐊᖏᕐᑭᑎᑦ;* ᐊᓚᒐᓗ

(b) ᐱᐅᓯᕐᓕᓐᓂᑦᓯᑐᑦ ᐃᓗᓕᓕᑭᐅᔪᖏᓐᓂᑦ ᐃᓄᐃᑦ ᑲᓇᑕᖏᑦᑐᓂᒃ ᑐᓂᔨᐅᓯᒪᓕᕆᓂᒃ ᒪᕐᒪᒃᑯᑦ ᑲᓇᑕᖏᓐᓂᑦ.

8.2 ᑐᑭᖏᑦ

8.2.1 ᐱᔪᓐᓇᕐᓯᑐᑦ ᑲᔪᓯᒪᖏᓐ ᐃᓕᖏᓄᑦ:

(a) “ᓄᓇᖅᑲᖅᑲᖅᓯᒪᔪᑦ ᓇᓗᒋᓂᔅᓴᐅᓕᓕᒃ” ᑐᑭᖃᖅᑐᒃ ᓄᓇᖅᑲᖅᑲᖅᓯᒪᔪᑦ ᓇᓗᒋᓂᔅᓴᐅᓕᓕᑦ ᐅᓇᓗ ᑐᑭᖏᑦ ᐅᖃᖅᑐᑦᓯᒪᓕᕆᔨᐸᑎᒃᑐᒃ ᒪᕐᒪᒃᑯᓂ ᓄᓇᕗᑦ ᑲᓇᑕᖏᑦᑐᓂᒃ ᑲᑎᕆᔨᕐᓂᐊᕆᐊᖅᑐᑦ ᓄᓇᖅᑲᖅᑲᖅᓯᒪᔪᑦ ᐅᓇᓗᒋᓂᔅᓴᐅᓕᓕᑦ ᐅᓇᓗᒋᓂᔅᓴᐅᓕᓕᒃᑯᓂᒃ ᓄᓇᖅᑲᖅᑲᖅᓯᒪᔪᑦ ᐅᓇᓗᒋᓂᔅᓴᐅᓕᓕᑦ ᐅᓇᓗᒋᓂᔅᓴᐅᓕᓕᒃᑯᓂᒃ (ᑲᑎᑕᐅᓯᒪᔪᑦ ᑲᓇᑕᖏᑦᑐᓂᖅᑐᑦ ᐊᑐᐊᒃᓴᖅ ᑐᐊᖏᑎᑎᓂᑐᑦ 1996-6), ᑲᑎᑭᒋᐊᖅᑕᖏᓪᓗᑎᒃ ᐃᓐᓇᐸᒃᓯᑦ, ᐊᓚᒐᓗ ᐃᓕᖏᖅᑎᑐᒃ “ᐃᓄᐃᑦ ᐅᓇᓗᑎᒃᖅᑐᑦ” ᑐᑭᖓ ᐅᓪᓗᐃᖅᑲᓚᖅᓯᒪᔪᖅ ᐃᓕᓐᓄ 24 ᑕᒃᑯᓂᐅᑎ *ᓄᓇᕗᒻᒥᒃ ᓄᓇᑕᑭᑎᓄᑦ ᐊᖏᕐᑭᑎᑦ.*

(b) “ᐅᓇᒥᓂᔪᓂᑎᒃᖅᑐᓄᑦ ᐱᙵᖅᑎᑦᑎᕐᓂᖅ ᐅᕐᓚᓴᐅᓂᐅ ᑭᓇᐅᔨᓕᒥᐅᓂᔪᓐᓇᖅᑐᑦ” ᑐᑭᖃᖅᑐᖅ, ᒪᓕᒃᑕᐅᔨᓚᐸᓕᒃ (c), ᑲᓇᑐᒃ ᐊᑐᑭᓇᖅᑐᖅ ᐊᖏᕐᕋᒧᑦ ᐅᖓᑕᓂᓴᐅᓂ, ᐅᕐᓚᓴᐅᓂ ᑕᓇᓕᑕᐃᓇᖅ ᑲᓇᑐᒃ ᐊᔨᖓ ᑐᓂᔨᐅᓯᖅ ᐅᖓᑕᓂᓂ ᐊᖏᕐᕋᒧᑦ:

(i) ᒪᓕᒃᑐᖅ ᑐᖓᔪᓂᓇᓂ 60 ᐃᓂᒃ ᓴᓇᔪᓇᓂᖓᑕ ᐊᖏᕐᕋᒧᒥ; ᐅᕐᓚᓴᐅᓂᓂ

(ii) ᑲᓇᑐᖓ ᐊᑭᑐᑎᒋᓂᓂ $20,000 ᐊᖏᕐᕋᒧᑕᒪᓐᓇᑦ; ᐊᓚᒐᓗ

(iii) ᐊᓯᖏᑦ ᑲᓇᑐᒃ ᐅᕐᓚᓴᐅᓂᓂ ᑲᓇᑐᑦ ᑕᐃᒪᓗ ᑲᓇᑕᖏᑦᑐᓂᖅ ᐊᖏᕐᕋᒧᑦ ᐊᓚᒐᓗ ᐊᐅᖓᑎᔪᑦ ᑎᒍᔨᓕᒪᔪᑦ ᐃᓄᐃᑦ ᐊᐅᓪᓕᔪᓐᓇᑦ ᐊᖏᕐᕿᑲᑎᑦᑎᓐᓂᒃ ᐃᓕᓐᓄᓂᑦ, ᑎᑎᕿᑲᑎᒧᑦ, ᓇᓗᒋᓂᔅᓴᐅᓕᓕᑦ ᐱᐃᕿᑲᑎᑕᐅᔪᑦ;

(c) ᓇᓗᒋᓂᔅᓴᐅᓕᓕᑦ ᐱᐃᕿᑲᑎᑦᑎᕐᓂᖅ ᐅᕐᓚᓴᐅᓂᓂ ᑭᓇᐅᔨᓕᒥᐅᓂᔪᓐᓇᖅᑐᑦ ᐅᒥᖓ ᐃᓕᖏᑲᓐᖏᓪᓕᑦᑐᑦ:

(i) ᑲᓇᑐᒃ ᐱᓴᓵᔪᔭ ᐃᔭᐊᓂ ᐅᑕᐅᔨᖅ 12-ᑕᖅᑭᒥ, ᐱᓕᓴᓂᖓ ᑎᑭᐅᑎᔨᓕᒥᐅᓂᔪᓐᓇᖅᑐᑦ ᐅᖓᑕᓂᓂ ᐊᖏᕐᕋᒧᑦ ᐱᔪᓐᓇᕐᓯᑐᑦ ᑭᖓᔪᓂᖅᓯᒪᓂᒪ, ᐅᓚᒪᓂᖓᔪᓂᐊᖏᓚᑕ ᓯᓚᐅᓇᓂᖓᑐᑎᑦᑕᑦ ᓄᓇᒥ ᑕᐃᒪᓇᐅᑦᑐᓚᕆᔨᖏᑦᑐᑦ;

(ii) ᐃᖅᑲᓇᐃᔭᖅᑎᑕᐅᓪᓚᕐᓂᑎᒃ ᑲᓇᑐᑦ; ᐅᕐᓚᓴᐅᓂᓂ

(iii) ᓄᓇᓗ ᐊᓂᔪᓕᑦ ᑲᔪᓯᖏᖅ ᐅᕙᓪᓕᐊᔪᖅ ᑲᔪᓯᖏᑦ ᐅᑯᐊ ᑲᓇᑕᒥ ᐃᒪᓕᕆᔨᖏᑦᑐᐊᑯᑦ ᐊᒻᒪᓗ ᐊᖏᑐᐃᓐᓇᑦ ᑎᓕᕆᔨᓂᖅᑐᑦ ᐃᓇᐃᑦ ᐊᐅᓚᓂᓂᕆᔨᓕᕆᔨᓕᕆᑦ ᐊᓕᕐᓂᖅᑎᑦᓂᖅᓯᓚᖅᑐᑦ ᐃᓕᓴᕐᓂᖅᑯ, ᑎᑎᖅᑲᑎᒍᑦ, ᐅᑯᐊ ᓇᖅᑎᓂᖅᓱᖅᑐᓂ ᐱᕕᖅᑲᑎᑦᑎᓂᐅᕐᓇᖅᓯᑦᑐᓂᑦᑐᓂᖅᑐᑦ ᐃᓕᓴᕐᓂᖅᑯ;

(d) “ᑲᓇᑕᖏᑦ” ᑐᑭᖅᑲᖅᑐᖅ ᑕᒪᖅᑎᑦ ᑲᓇᑕᖏᑦ, ᑭᓯᐊᓂ ᑲᓇᑕᖏᑦ ᓴᖅᓚᒪᓂᕐᒥᓂ ᐃᖅᑲᖅᓯᐅᑉᐳᖅᑲᐃᑦᑐᖅᑐᓂ ᐃᓕᓴᕐᓂᖅᑯ ᐃᓕᓴᕐᒥ 23 ᒡᒃᑯᓚᓐᓂ ***ᓄᓇᕗᒻᒥᐅ ᓄᓇᕆᑎᑎᓄᑦ ᐊᖏᕐᑕᑎᓄᑦ***, ᑐᓂᓴᐅᔪᑦ ᑲᓇᑕᒥ ᐅᒪᔪᓕᕆᔨᓕᕆᑦ ᐱᔪᓐᑎᑦᓯᑐᑦ ᑲᓇᑕᒥ ᐅᒪᔪᖏᖑᓂ ᐅᖓᖅᑕᕐᓂᖅᑐᑦ ᑎᑭᓕᓄᑦ ᓴᓇᔨᓴᕐᓂᖅᑐᑦ, ᐃᓕᑦᑎᓴᐅᓂᓗᒃ, ᖃᓄᐃᔨᓴᖅ ᐱᔪᓴᖅᑐᑦ:

(i) ᑲᓇᑕᖏᑦ ᑲᔪᓯᔪᓴᒪᓕᒫᑦ, ᐱᓴᓯᒪᔪᑦ, ᐱᓕᕐᓂᐊᑎᕐᑎᓂᖅ, ᓂᐅᕕᖏᓂᖅ, ᐃᓗᓐᓂᐅᔪᖅ, ᐃᓅᓯᖏᓐᓄᑎᓂᖅ, ᐱᐅᓯᖏᓐᓄᑎᓂᖅ, ᐃᓚᖅᑎᐊᑎᓂᓂᖅ, ᓄᓇᕗᑦᑐᓯᓂᖅ, ᐱᖓᓱᒪᔪᓂᖅ ᐱᐅᓯᖏᓐᓄᑎᓂᖅ, ᐃᓅᓯᖏᓐᓄᑎᓂᖅ, ᐃᓗᓐᓂᐅᔪᖅ ᐃᓗᓐᓂᐅᔪᖅ ᐃᓗᓐᓂᐅᔪᖅ

8.3.5 ᑲᓇᑕᒥ ᐅᒪᔨᓕᕆᔨᖅᐳᑎᑦ ᐃᓱᒪᑎᑎᐊᖅᓗᓂᑎᑦ ᐃᓕᒧᔪᖅᑐᑦ ᑲᐟᔪᓂᑎᐊᖅᑐᑦ ᐃᓇᐃᑦ ᑕᐃᑲᓂᓗ ᐃᓕᒧᔪᑦ ᐱᓇᓱᖅᑎᖅᑐᑦ ᒪᓕᒃᓗᓂᑎᑦ ᑕᐃᒪᓐᓇᐅᔾᔨᖅᑎᖅᑎᑕᐅᔨᑦ ᑭᓱᓕᕆᔨᐅᓐᓇᖅᓂᓐᓇᓂᑦ ᓇᓐᒥᓂᔪᑎᑎᓗᓂᑎᑦ ᐊᖏᖅᑲᓱᒪᔨᖅᓴᓂᑦ ᓇᓇᖅᑲᖅᑲᖅᓱᒪᔨᓇᓂᑦ ᓇᖅᒥᓂᖅᑕᓗᓄ.

8.3.6 ᐅᓇ ᐃᓕᖅᓴ ᒪᓕᑎᐊᖅᑲᖅᑐᖅ ᓇᓇᑦᓴ-ᐊᖅᑭᖅᑎᑎᑦᔪᑎᑦ-ᐱᓱᒪᔨᑦ ᑭᓱᓕᕆᔨᐅᓐᓇᖅᑭᖅᓂᓂᑦ ᓇᖅᒥᓂᔪᑎᑎᓗᓂᑎᑦ ᐊᑐᐊᐱᖅ, ᑕᐃᒪᓗ ᐊᑐᓕᖅᑕᐅᑐᐊᖅᐸᑦ ᒪᓕᓪᓗᓄ ᓇᓇᑦᓴ ᑐᓴᖅᓴᖅ ᑎᒥᖅᓴ-ᑐᕐᒪᓚᕆᔪᑎᖅᓴᓂ ᐱᓪᓗᒍ ᐊᑭᓕᐅᑎᑦ ᐃᖅᑲᖅᑐᐃᐱᖅᔪᖅᑎᑕᐅᓂᖅ ᐊᓚᒍᓗ ᓇᓕᓕᐅᓂ ᐊᑐᓕᖅᑕᐅᑐᐊᖅᑲᓚᐊᖅᑲᖅᑐᓂᖅ ᐱᓪᓗᒍ ᓇᓕᓕᐅᓂ ᐊᑐᓕᖅᑕᐅᑐᐊᖅᑲᓚᐊᖅᑲᖅᑐᓂᖅ ᐊᓚᒍᓗ

ᐊᓯᖏᑦ ᑎᑎᕆᖃᓯᒪᔨᑦ ᑕᒃᑯᓇᓂ ᑲᓇᑕᒥ ᐅᒪᔨᓪᓗᕆᔨᐊᕐᑯᑦ ᐱᓯᒪᔭᖏᑦ ᐊᑐᑎᕆᒃᖁᑦ ᑭᓇᐅᔭᑦᐅᑭᐅᑎᓯᓐᓄᑦᓗᓂᑦ ᑕᒃᑯᓂᔪᑦ ᐱᕕᕆᕆᑎᓂᑎᓂᖅ ᐅᖅᓯᔪᓐᓇᓂ ᑭᓇᐅᔭᑦᐅᑭᐅᑎᓯᓐᓄᑦᓗᓂᑦ ᐱᕕᕆᕆᑎᓂᑎᓂᖅ ᐅᖅᓯᔪᓐᓇᓂ ᑭᓇᐅᔭᑦᐅᑭᐅᑎᓯᓐᓄᑦᓗᓂᑦ.

8.4.8 ᑕᐃᒪᓗ ᑎᓕᕆᐅᓯᒪᔨᑦ ᐃᓄᐃᑦ ᐊᐅᓪᓚᕆᔪᓂᑦ ᑐᓂᔪᐊᑎᒥᓂᑦ ᑭᒪᔭᑦ ᑎᑎᕆᓂ ᐊᑐᓕᕆᖃᖅᑐᑦ ᐱᓕᑦᑐᓂᑦ ᑲᑎᒪᔨᓴᖅᑎᑦᑎᓂᒃ ᑕᐃᒪᓕᕆᔨᒃᑯᐊᑎᖃᖅᑐᑦ, ᑲᓇᑕᒥ ᐅᒪᔨᓪᓗᕆᔨᐊᕐᑯᑦ ᑲᓐᑐᖅᓯᓂᐊᖃᖅᑐᑦ ᑕᒃᑯᓂᒃ ᑎᓕᕆᐅᓯᒪᔨᑦ ᐃᓄᐃᑦ ᐊᐅᓪᓚᕆᔪᓂᑦ ᐱᓕᑦᑐᓂᑦ ᐅᖅᓯᔪᓐᓇᓂ ᐱᕕᕆᕆᑎᓂᑎᓂᖅ ᐅᖅᓯᔪᓐᓇᓂ ᑭᓇᐅᔭᑦᐅᑭᐅᑎᓯᓐᓄᑦᓗᓂᑦ.

8.5 ᐊᖏᖁᒍᒥᓵ ᖃᒥᖏᑭᐊᖅᑕᐅᔪᓐᓇᑎᒃ

8.5.1 ᐊᖏᖁᒍᒥᓵ ᖃᒥᖏᑭᐊᖅᑕᐅᖅᑲᑕᖏᓐᓇᑎᒃ ᒪᑕᒃᓴᐃᒍ ᐃᑕᓇᒃ 15.4 ᑕᒃᑯᓇᓂ ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᖏᓂᒃᓂᓕᓂ ᐱᔪᓐᓀᑦᑕᐅᔪᓐᓇᖅᑐᓄᑦᓐᓇ ᐊᓇᑎᑭᑐᑦ, ᑲᓇᑕᒥ ᐅᒪᔨᓪᓗᕆᔨᐊᕐᑯᑦ ᐅᓂᒃᑲᓕᐊᓚᖅᑲᑕᖏᓐᓇᑎᒍᓐᓇ ᑕᒃᑯᓇᓂᒃ ᐱᓇᓱᐊᖅᑎᑦᑎᓂᒃ ᑕᒪᖏᑦ ᐱᕕᕆᓚᐅᖅᑎᑦ ᒪᑕᑐᒃ ᐱᓕᕆᔨᒃᑯᐊᑎᖃᖅᑐᑦ ᐃᓄᐃᑦ ᐅᓇ ᐃᓕᓐᓂ, ᐃᓕᓐᓂ 24 ᑕᒃᑯᓇᓂ ᓄᓇᕗᒻᒥᒃ ᓄᓇᕈᑎᒋᓄᑦ ᐊᓇᑎᑭᑐᑎᑦ ᐊᓚᒪᓗ ᑭᓱᓕᕆᓪᓚᕆᔪᓐᓇᖅᑐᓄᑦᓐᓇ ᓄᓇᖅᑐᖅᑲᓐᓂᑎᑦ ᐊᑦᑭᒃᓱᒪᓕᕐᓂᒃᖁᓕᖅᑐᑦ ᓄᓇᖅᑐᖅᑲᓐᓂᑎᓂᖅ, ᐃᓕᑭᒃᐅᓕᓐᓂᑎᒃ ᐊᑐᑎᖃᖅᑐᑦ ᖁᑉᑭᒃᓱᒪᓕᕐᓂᒃᖁᓕᖅᑐᑦ ᑲᓐᑐᖅᓯᓂᐊᖃᖅᑐᑦ ᓄᓇᖅᑐᖅᑲᓐᓂᑎᓂᖅ ᐅᖅᓯᔪᓐᓇᓂ ᐃᓕᑭᒃᐅᓕᓐᓂᑎᒃ ᐊᑐᑎᖃᖅᑐᑦ ᖁᑉᑭᒃᓱᒪᓕᕐᓂᒃᖁᓕᖅᑐᑦ ᐊᑐᑎᖃᖅᑐᑦ ᐱᓕᑦᑐᓂᑦ ᐃᓄᐃᑦ ᐅᓇ ᐃᓕᓐᓂ, ᐃᓕᓐᓂ 24 ᑕᒃᑯᓇᓂ ᓄᓇᕗᒻᒥᒃ ᓄᓇᕈᑎᒋᓄᑦ ᐊᓇᑎᑭᑐᑎᑦ ᐊᓚᒪᓗ ᐱᓕᕆᔨᒃᑯᐊᑎᖃᖅᑐ

ᐃᓕᓐᓴ 9 — ᐃᓕᓐᓂᐊᕐᓂᑎᓂᖅ ᐊᒻᒪᓗ ᐃᖅᑲᓇᐃᔭᖏᑎᓂᑎᓂᖅ

9.1 ᑐᑭᒋᖅᓯᓪᓚᐊᔨᓱᑎᐅᒃᖦᓪ

9.1.1 ᐅᕐᓄᓂ ᐃᓕᓐᓴᑐᓂ ᐊᒋᓂ ᑎᑎᕆᖅᓯᒪᔪᓪ ᑐᑭᒋᖅᓯᓪᓚᐊᔨᓱᑎᐅᒃᖦᓪᕈᓪ:

(a) ᖃᓇᑎᕆᓗᑎᐅᒃ, ᐱᔾᓀᐅᖅᖦᖅᑎᕆᓗᑎᐅᒃ ᐊᒻᒪᓗ ᐃᖅᑲᓇᐃᔭᖏᑎᕆᓗᑎᐅᒃ ᐃᓄᖓᓂᒃ ᖃ>ᓐᓂᐊᖏᖅᓂᓂ-ᐱᓯᒪᔪᓪ ᐃᖅᑲᓇᐃᓴᒃᖦᐃᓪ;

(b) ᐊᑐᖅᑕᐅᓗᑎᐅᒃ ᑲᓇᑕᒥ ᐅᒪᔪᓚᓀᔪᕐᐊᐊᑯᓪ ᐃᖅᑲᓇᐃᔭᓚᖏᑎᕆᑎᓂᔨᓱᓯᖏᓪᕈᓪ, ᖃᓇᑎᕆᑎᔨᓱᓯᖏᓪᕈᓪ ᐊᒻᒪᓗ ᐃᖅᑲᓇᐃᔭᖏᑎᓄᓪ ᐱᓚᓂᔨᓱᓯᖏᓪᕈᓪ ᐊᑐᖅᑭᓱᖏᓪᕈᓪ ᐊᒻᒪᓗ ᖁᑲᐃᑦᓂᖅᓚᖏᖅᑕᖅᑕᖅᑕᖅᓂᑯᓪᓕ;

(c) ᐅᓄᔪᓕᐊᖅᓗᑎᓪᓕ ᐱᐃᖅᑲᖏᑎᕆᑎᔨᓱᑎᐅᒃᖦᓪᕈᓪ ᐃᓄᖓᓂᒃ ᐱᓚᓂᔨᓱᓴᖅᑕᖓᓄᓄ ᑲᑎᒪᔨᓴᖅᑐᓂᒃ ᐃᓄᖓᓂᒃ ᐱᓚᓂᔨᓱᓴᖅᑐᓂᒃ ᐃᓕᓐᓴ 23 ᑭᒃᑯᓄᓪᓕ ᓄᓇᕗᑦ ᓄᓇᕗᒻᒥᒃ ᓄᓇᕆᔨᑎᓄᓪᓕ ᐊᖏᕐᕕᑎᓂ.

9.2 ᐃᓄᖓᓂᒃ ᐱᔾᓀᐅᖅᖦᖅᑎᕆᑎᕆᔨᑕᐅᔨᖅᖦᖅ

9.2.1 ᑲᓇᑕᒥ ᐅᒪᔪᓚᓀᔪᕐᐊᐊᑯᓪᓕ ᑲᔨᓱᔨᒥᒃ ᒪᑕᖅᑕᑕᓪᓗᑎᕐᓂ ᓄᓇᕗᑦ ᑲᒪᕐᓂᒃ ᑎᑎᕆᖅᓯᒪᔪᓪ ᐊᒻᒪᓗ ᑐᐱᒪᑎᕆᑎᕆᔨᑕᐅᓴᖅᑎᑦᓂᑎᓂᒃ ᐊᓯᔪᒪᔪᓪ ᐃᖅᑲᓇᐃᔭᓴᖅᑕᐅᓂᒃ ᖃᓄᖓᓂᒃ ᐃᖅᑲᓇᐃᔭᖏᑎᓄᓪᓕ, ᐃᓕᓴᐅᓂᒃ ᐃᓄᖓᓂᒃ ᐃᖅᑲᓇᐃᔭᖏᑎᓄᓪᓕ ᐊᒻᒪᓗ ᓇᖕᒥᓂᔨᓱᓯᖏᓪᕈᓪ ᐃᓄᖓᓂᒃ ᐱᔾᓀᐅᖅᖦᖅᑎᕆᑎᕆᔨᑕᐅᔨᖅᖦᖅᑎᓄᓪᓕ.

9.2.2 ᑲᓇᑕᒥ ᐅᒪᔪᓚᓀᔪᕐᐊᐊᑯᓪᓕ ᐃᖅᑲᓇᐃᔭᖏᑎᕆᑎᓂᔨᓱᓯᖏᓪᕈᓪ, ᐱᔾᓀᐅᖅᖦᖅᑎᕆᑎᓂᔨᓱᓯᖏᓪᕈᓪ ᐊᒻᒪᓗ ᐃᓕᓐᓴᐃᔪᖅᑎᖏᖅᑎᕆᑎᓂᔨᓱᓯᖏᓪᕈᓪ, ᖃᓄᖓᓂᒃ ᐃᖅᑲᓇᐃᔭᖏᑎᓄᓪᓕ ᐃᓄᖓᓂᒃ ᐃᓕᓐᓴᐃᔪᖅᑎᖏᖅᑎᕆᑎᓂᔨᓱᓯᖏᓪᕈᓪ, ᐊᒻᒪᓗ ᐃᓕᓐᓴᐃᔪᖅᑎᖏᖅᑎᕆᑎᓂᔨᓱᓯᖏᓪᕈᓪ ᐃᓄᖓᓂᒃ ᐃᓕᓐᓴᐃᔪᖅᑎᖏᖅᑎᕆᑎᓂᔨᓱᓯᖏᓪᕈᓪ ᐃᓄᖓᓂᒃ ᐱᔾᓀᐅᖅᖦᖅᑎᕆᑎᕆᔨᑕᐅᔨᖅᖦᖅᑎᓄᓪᓕ

9.3 ᐊᒥᐃᓐᓇᕆᐊᓐᖏᑦᑐᑦ ᐊᓚᒐᓂ ᐃᓱᒪᓕᕆᓂ ᐃᖅᑲᓇᐃᔨᖅᑎᑕᐅᓂᖅ ᐊᓚᒐᓂ ᐃᖅᑲᓇᐃᔨᒪᓂᑎᑎᓴᒥᐅᔪᓂ

9.3.1 ᑲᓇᑕᒥ ᐅᒪᔪᒪᓇᓴᕐᔪᐊᑯᑦ ᖃᐅᒥᖅᑕᐊᖅᑲᑎᑕᒍᑦ ᐱᒪᓪᓕᖅᑲᑎᑕᒍᑦ ᓄᓇᕗᑦ ᑐᓴᖅᑲᓚᐅᑦ ᐊᓚᒐᓂ ᐃᖅᑲᓇᐃᔨᒪᒥᑎᑦᑲᑎᑎᓴᒥᐅᔪᓂ ᐊᒥᐃᓐᓇᕆᐊᓐᖏᑦᑐᑕᓄᑦ ᐊᓚᒐᓂ ᐃᓱᒪᓕᒫᓄᑦ ᐃᖅᑲᓇᐃᔨᖅᓴᓄᑦ ᓄᓇᕗᒻᒥ:

(a) ᐊᑐᐊᖏᐃᑦ ᐊᓚᒐᓂ ᐱᐅᓱᖅᓴᒪᓪᓕᐊᓱᒪᓕᓴᓂ ᑎᑎᕋᖅᑕᓪᓕᓴᓂ ᐃᖅᑲᓇᐃᔨᖅᓴᐃᑦ, ᐃᓕᑎᕐᐅᑦᑕᓗᑎᒃ ᐃᖅᑲᓇᐃᔨᖅᓴᓂᒃ ᓴᕿᔨᕐᑎᑎᓂᖅᓂᖅ;

(b) ᐃᖅᑲᓇᐃᔨᒪᓂᑎᑎᓂᕐᑕᐊᓂᕐᒧᑦ ᐃᒪᓐᓇᐃᓕᓐᖏᓚᕐᔪᐊᑎᖅᑐᑦ ᐊᓚᒐᓂ ᐊᔪᖏᑦᑕᓪᓕᐊᕋᖅᑕᖏᑦ, ᐃᓕᑎᕐᐅᑦᑕᓗᑎᒃ ᖃᓄᐃᓕᐊᑎᒥᓇᕐᓂᖏᓐᖓ ᐊᕐᑕᐅᔪᖅ ᐅᓯᒪᓕᐊᑦ; ᐊᓚᒐᓂ

(c) ᐃᖅᑲᓇᐃᔨᖅᓴᐃᑦ ᑭᓱᑕᓪᓗᐊᓂᖏᓐᓂᖅ ᐊᓚᒐᓂ ᐅᓂᖃᖅᓭᒪᓕᒫᑦ ᐊᔪᖏᑦᑕᓪᓕᐊᕋᖅᑕᖏᑦᑐᖅ.

9.3.2 ᑲᓇᑕᒥ ᐅᒪᔪᒪᓇᓴᕐᔪᐊᑯᑦ ᐃᓕᑕᐅᑎᓭᒪᓗᐊᕋᖅᑕᖏᑦᑐᑦ ᐅᑯᐊ ᐊᑕᑦᓂ ᑎᑎᕐᓂᖅᓭᒪᓕᒫᑦ ᖃᑭᓂᓭᓪᓗᐊᕋᖅᑕᖏᑦᓴᓂ ᐊᓚᒐᓂ ᐃᖅᑲᓇᐃᔨᖅᓴᒥᐅ ᑭᓱᑕᓪᓗᐊᓂᖏᓐᖓᓄᑦ ᐊᐅᑦᑕᓪᓕᐊᕋᖅᑕᖏᑦᑐᓄᑦ ᐊᓚᒐᓂ ᐃᓱᒪᓕᕐᓂᖅᑲᖅᓯᒪᓕᒫᑦ ᐃᖅᑲᓇᐃᔨᖅᓴᐃᑦ ᓄᓇᕗᒻᒥ ᐃᓄᐃᑦ ᐃᓄᓇᕗᑦᓄᓇᕗᑦ:

(a) ᖃᐅᔨᓴᓂᑦ ᐃᓄᐃᑦ ᐱᐅᓱᖅᓴᒪᓕᒫᑦ, ᐃᓄᐃᑦ ᐊᑐᐊᖏᐃᑦ ᐊᓚᒐᓂ ᓄᓇᕗᑦ ᐊᔪᖏᑦᑕᓪᓕᐊᕋᖅᑕᖏᑦᑐᑦ;

(b) ᐃᓄᐃᑦ ᐊᑐᐊᖏᐃᑦ ᐊᓚᒐᓂ ᓄᓇᕗᑦ ᐊᔪᖏᑦᑕᓪᓕᐊᕋᖅᑕᖏᑦᑐᑦ;

(c) ᐃᓄᐃᑦ ᐊᑐᐊᖏᐃᑦ ᐊᓚᒐᓂ ᓄᓇᕗᑦ ᐊᔪᖏᑦᑕᓪᓕᐊᕋᖅᑕᖏᑦᑐᑦ;

(d) ᐃᓄᐃᑦ ᐊᑐᐊᖏᐃᑦ ᐊᓚᒐᓂ ᓄᓇᕗᑦ ᐊᔪᖏᑦᑕᓪᓕᐊᕋᖅᑕᖏᑦᑐᑦ;

(e) ᐃᓄᐃᑦ ᐊᑐᐊᖏᐃᑦ ᐊᓚᒐᓂ ᓄᓇᕗᑦ ᐊᔪᖏᑦᑕᓪᓕᐊᕋᖅᑕᖏᑦᑐᑦ; ᐊᓚᒐᓂ

(f) ᐃᓄᐃᑦ ᐊᑐᐊᖏᐃᑦ ᐊᓚᒐᓂ ᓄᓇᕗᑦ ᐊᔪᖏᑦᑕᓪᓕᐊᕋᖅᑕᖏᑦᑐᑦ.

9.3.3 ᑲᓇᑕᒥ ᐅᒪᔪᒪᓇᓴᕐᔪᐊᑯᑦ ᐱᑎᐊᖏᑦᑕᓗᑎᒃ ᖃᐃᐊᖏᑦᑕᓗᑎᒃ ᐃᓄᐃᑦ ᐊᑐᐊᖏᐃᑦ ᐊᓚᒐᓂ ᐃᓱᒪᓕᒫᑦ ᐊᓚᒐᓂ ᐃᖅᑲᓇᐃᔨᖅᓴᐃᑦ ᓄᓇᕗᑦ ᐊᔪᖏᑦᑕᓪᓕᐊᕋᖅᑕᖏᑦᑐᑦ ᐃᓄᐃᑦ ᐊᑐᐊᖏᐃᑦ ᐊᓚᒐᓂ ᐃᓱᒪᓕᕐᓂᖅᑲᖅᓯᒪᓕᒫᑦ

9.4 ᐊᒥᐃᓐᓇᕆᐊᓐᖏᑎᑐᒃ ᐊᓚᒐᓂᒃ ᐃᓱᒪᓕᓕᒋᑦ ᐃᖅᑲᓇᐃᔨᖅᓴᑦ ᐊᓚᒐᓂᒃ ᓴᓇᔨᓂᒃ ᐱᖓᓱᖅᓴᖅᓯᖅ

9.4.1 ᑕᐃᒪᓐᓇᐅᓱᒪᖓᒪᑦ ᓇᒥᐊᑐᐃᓐᓇᖅ ᐃᖅᑲᓇᐃᔪᒥᕐᓂᑎᓂᒋᖅᓯᒥ ᑭᖑᔪᓐᖓ ᐊᒥᐃᓐᓇᔨᓐᖏᑎᑐᒃᓴᓂᒃ ᐊᓚᒐᓂᒃ ᐃᓱᒪᒥᕐᕕᒃᓴᓂᒃ ᐃᖅᑲᓇᐃᔨᖅᓴᓂᒃ, ᐊᓚᒐᓂᒃ ᑎᒥᐅᒥᕐᓂᑐᒃ ᐱᓪᓕᒥᑎᐊᖅᑎᑐᒃ ᐃᓇᐃᓂᒃ, ᐃᖅᑲᓇᐃᔪᖅᑐᑦ ᐅᔾᔨᖅᑐᕐᓂᒃ ᐃᒪᓐᓇᐅᓂᒋᖅᑐᑦ, ᑲᑐᑎᓂᒃ ᐃᖅᑲᓇᐃᔪᖅᑐᖅ, ᐱᓪᓕᒥᑎᐊᕐᓂᑎᓂᓗᑎᖅ ᐃᒥᓐᓂᐊᕐᓂᒋᖅᑐᑦ ᐃᖅᑲᓇᐃᔨᓕᕆᓂᐊᖅᑕᖅᓴᓂᒃ ᐊᓚᒐᓂᒃ ᐱᖓᓱᖅᓴᖅᑎᑐᒋᖅᑕᐅᑦ <ᖁᓇᐅᑎ.

9.4.2 ᐅᑯᐊ ᐃᒥᐠᓂᐊᕐᓂᒋᖅᑕᑦ ᐃᖅᑲᓇᐃᔨᓕᕆᓂᐊᕐᓂᒋᖅᑕᕐᓗᓂᒃ ᐊᓚᒐᓂᒃ ᐱᖓᓱᖅᓴᖅᑎᑐᒋᖅᑕᐅᑦ <ᖁᓇᐅᑎ ᑕᐃᒪᓐᓇᐃᔪᓕᒥᓂᒃ ᐊᑕᔪᑦ 9.4.1 ᐃᓕᖅᑲᖅᓗᑎᒃ ᐅᑯᓂᖓᓱᒃᓗ, ᐊᓱᖏᑦᑐᓂᓗ:

(a) ᑭᓱᓂᖏᑦᑐᒃ ᐃᖅᑲᓇᐃᔪᖅᑐᑦ ᑭᓪᓕᐅᔨᓂ ᐊᓚᒐᓂᒃ ᐊᑯᓂᐅᔨᕈ ᐃᓕᐠᓂᐊᕐᓂᒋᖅᑕᑦ ᐃᖅᑲᓇᐃᔨᓕᕆᓂᐊᕐᓂᒋᖅᑕᕐᓗᓂᒃ ᑎᑭᐅᑎᐅᓱᖏᓂᒋᖅᑕᓂ; ᐊᓚᒐᓂᒃ

(b) ᐅᖅᑐᖅᑭᒪᔪᖅ ᓴᓇᓱᒪᓪᓗᓂ ᐃᑲᔪᖓᓗᑦ ᐃᖅᑲᓇᐃᔪᖅᑎᑦ ᑎᑭᐅᑎᓂᒋᖅᑐᑦ ᐃᓕᐠᓂᐊᕐᓂᒋᖅᑕᑦ ᐃᖅᑲᓇᐃᔨᓕᕆᓂᐊᕐᓂᒋᖅᑕᕐᓗᓂᒃ ᑎᑭᐅᑎᓂᒋᖅᑕᓂ.

9.4.3 ᐃᓕᐠᓂᐊᕐᓂᒋᖅᑕᑦ ᐃᖅᑲᓇᐃᔨᓕᕆᓂᐊᕐᓂᒋᖅᑕᕐᓗᓂᒃ ᐊᓚᒐᓂᒃ ᐱᖓᓱᖅᓴᖅᑎᑐᒋᖅᑕᐅᑦ <ᖁᓇᐅᑎ ᐊᒥᐃᓐᓇᔨᓐᖏᑎᑐᒃᓴᓂᒃ ᐃᖅᑲᓇᐃᔪᖅᑎᓇᓂᒃ ᐊᓚᒐᓂᒃ ᐃᓱᒪᓕᓴᖅᑎᓇᓂᒃ ᐃᖅᑲᓇᐃᔪᖅᑐᖅᑕᑦ ᐊᒥᐅᓱᑦᒧᑦ ᐊᖏᔪᒧᑦ ᐅᕙᓪᓕᓐᓂᑦ ᐊᑯᓂᐅᔨᓂᖅᑎᑦᑎᓂᒃ ᐅᑯᓂᖓᓱᒃᓗ ᐃᓕᖅᑲᖅᓗᑎᒃ ᐅᑯᓂᖓᓱᖅᓴᖅᑎᑐᒋᖅᑕᐅᑦ ᐊᓚᒐᓂᒃ ᐱᖓᓱᖅᓴᖅᑎᑐᒋᖅᑕᕐᓗᓂᒃ ᐅᑯᓂᖓᓱᒃᓗ ᐃᓕᖅᑲᖅᓗᑎᒃ:

(a) ᓇᓪᓕᐊᓐᓂᑦ ᐱᖓᓱᖅᓴᖅᑎᑐᒋᖅᑕᕐᓗᒃ ᐱᐅᓱᕙᓪᓕᓐᓂᐊᕐᓂᒋᖅᑕᓂ ᑲᓇᓕᓚᓴᓐᓇᓂᖏᑦᑐᒃ ᑕᐃᒪᓐᓇᐅᓴᖅᑎᖏᓐᓂᖏᑦᑐᒃ ᐃᖅᑲᓇᐃᔪᖅᑎᑎᒃᓴᑦ, ᐃᓚᒐᓂᒃ

(b) ᐃᓇᐅᑎᑦ ᐅᖅᑐᖅᑭᒪᔪᖅ ᖃᐅᓪᓚᐃᑎᑦ ᐅᖃᖓᑕᖅᑲᓇᑎᒃ ᐃᓕᐠᓂᐊᕐᓂᑎᒃ, ᑕᐃᒃᓂᒥ ᐱᖓᓱᖅᑲᓇᑎᑦ ᐃᒃᓚᖅᓂᐊᕐᓂᑎᑦ ᐃᖅᑲᓇᐃᔪᖅᑎᑎᑦ ᑲᓇᓴᖅᑭᓇᓂᒃ, ᐊᓚᒐᓂᒃ ᑕᐃᒃᓂᒥ ᐅᖃᖓᑕᖅᑲᓇᖅ ᐊᑐᖅᑕᐅᓚᖅ ᐱᖓᓱᖅᑲᓇᖅᓴᐅᓇᓂᑎᓂᒥ; ᐊᓚᒐᓂᒃ

(c) ᐊᓱᖅᑭᓐᓂᒃ ᐱᓴᒪᓕᓚᒥᒃ ᐱᖓᓱᖅᑲᓇᖅᓴᐅᓇᖅᑐᓂᒃ ᐱᐃᓴᖅᑎᑎᑎᒍᑎᒃ, ᐅᑯᐊᑐᓗ ᐃᓕᒋᑐᑎᑦ ᐃᖅᑲᓇᐃᔪᖅᑐᓗᑎᓗᑐ ᐱᖓᓱᖅᑲᓇᓂᖅ, ᑲᓱᓱᑎᑦᑎᓐᓂ ᐃᓕᐠᓂᐊᕐᓂᑎᓂᒋᖅᑐᑦ ᐊᓚᒐᓂᒃ ᐃᓕᐠᓂᐊᕐᓂᐃᖅᓯᒪᓂᖅ ᓄᖅᑲᖅᓱᓕᒋᐅᖅᑐᓂᒃ.

9.4.4 ᐅᑯᐊ ᐱᖓᓱᖅᓴᖅᑎᑐᒋᖅᑕᐅᑦ ᐱᐃᓴᖅᑎᑎᒍᓕᒃ ᑐᓂᓱᓱᓕᑎᒃ ᐅᓴᖅᑲᓱᓕᓕᒃ ᐊᒥᐃᓐᓇᔨᓐᖏᑎᑐᒃᓴᓂᒃ 9.4.3 ᐊᓴᑭᖅᓱᓕᓕ ᐊᑐᑎᖅᑲᖅᓗᑎᒃ ᐃᖅᑲᓇᐃᔪᖅᑎᑎᒃ ᐱᓕᒪᓐᓇᖅᑐᖅ, ᐃᓕᓴᕐᕕᑐᓗᑎᒃ:

(a) ᑎᑭᐅᑎᓗᑐ ᐊᑐᖅᑕᐅᑦ, ᐃᓕᐠᓂᐊᕐᑎᑕᐅᑎᓗᑎᒃ ᐃᓇᐃᑎᑦ ᐅᓴᖅᑲᓕᑦᓗᑎᒃ;

(b) ᑐᓂᓱᓕᑎᑦ ᐃᓕᐠᓂᐊᕐᓂᑎᓴᓕᓂᐊᕐᓂᒋᖅᑕᕐᓕᐠᓂᖅ ᐊᑐᓂ ᐱᓕᕆᑎᓂᑎᕐᔪᑎᓂᒋᖅᑐᒃ; ᐊᓚᒐᓂᒃ

(c) ᑐᓂᓗᑎᑦ ᐱᖓᓱᖅᑲᑎᒃᓗᑎᑦ ᓄᓇᕗᓪᓕᒥ, ᑕᐃᒃᓂᓗ ᑕᓪᒃᑯᐊ ᐱᖓᓱᖅᑲᓇᖅᓴᐅᓇᖅᑕᑦ ᐊᑐᐃᓐᓇᐅᓗᑎᒃ ᐱᐅᓂᓂᒋᖅᑐᒃ.

9.4.5 ᑭᓇᐅᔨᖅᑲᑦ ᐊᑐᕆᐊᓂᒋᖅᑕᑦ ᐊᑐᓂᓯᐊᓂᑎᑦ ᐅᓗᓇᐃᖅᓱᓕᕐᓂ ᐃᓕᓐᓄ 9.3 ᐊᓚᒐᓂᒃ 9.4 ᐱᖃᐅᓂᓯᓐᖏᑎᑐᒃᓴᓂᒃ ᐃᓇᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᓯᓂᖓᓐᓄᓇᓕᒃ ᐱᓕᒪᓐᓇᕕᐅᓐᓇᖅᑐᓇᓕᓂᓗ ᐊᓐᖏᑭᒃᑯᑦ ᐊᑐᓕᖅᑎᑕᐅᓂᖅᓯᓕᕐᓂᓕᒃ ᑭᓇᐅᔨᖅᑲᐃᐃ.

9.5 ᐊᕿᑦᔪᒍ ᐃᓕᓴᐅᓂ ᐃᓕᓐᓂᕐᒧᑦᑎᓂᒃᑯᓐᓂᕐᒧᖅᑐᑦ

9.5.1 ᐅᑯᐊ ᐱᓇᓱᐊᖅᑎᑎᕐᑐᑦ ᐊᓐᓴᖅᑎᑎᕐᑐᖅᑐᑦ ᖃᑎᖅᑕᐊᕿᓗᓕᑦ ᐊᓯᒪᓗ ᐃᓱᒪᑦᑎᐊᑲᓐᓂᕐᒧᑦ ᓴᖅᑭᑕᐅᔨᓐᓇᕐᓂᖅᑐᑦ ᐊᕿᑦᔪᒍ ᐃᓕᓴᐅᓂ ᒥᔅᓯᐊᓂᑦ ᓴᓇᔨᐅᓱᒪᓕᕆᔨ 6-ᓴᐅᓕ ᐊᕿᑦᔪᒍᓂ ᖃᑎᖅᑕᐊᕿᖅᑕᐅᓗᓂ ᐅᖃᖅᑎᑦᓯᓐᓇᕐᓂᖅ ᓄᓇᕗᒻᒥᔨᓂᒃ ᓴᓇᔨᐅᓱᒪᓕᕆᔨ ᑲᑎᕋᐅᖅᑕᐅᓗᓂ ᐅᖃᖅᑎᑦᓯᓐᓇᕐᓂᖅᑐᑦ ᐊᒻᒪᓕᒃᑯᑦ ᐊᑕᐃᓂ ᐃᓕᓴᐅᓂ 15.4 ᐅᓪᓚᓂ ᐃᓄᐃᑦ ᐊᐅᑐᖅᑕᐅᓂᐊᕐᓂᖓᓄᑦ ᐱᓪᓚᑦᑕᑕᐅᔨᓐᓇᕐᓂᖅᑐᓗᓂᒃ ᐊᕿᑦᔪᒍ ᐃᓕᓴᐅᓂ

ᐃᓕᓐᓴ 10 — ᖃᐅᔨᓴᖅᓯᖅ

10.1 ᑐᑭᒃᖢᑦᓕᐊᔨᓂᐅᕐᓴᖅ

10.1.1 ᐅᓇ ᐃᓕᓐᓴ ᐊᒋᓂ ᑎᑎᕆᖅᓯᒪᔪᑦ ᑐᑭᒃᖢᑦᓕᐊᔨᓂᐅᕐᓴᖅᓗᓐᓂᑦ:

(a) ᐅᓄᖅᓯᑎᐊᕆᓗᑎᑦ ᐃᓄᐃᑦ ᐃᓕᐅᔪ ᓂᕆᔨᑎᓂᒃ ᐊᓪᓚᓗ ᓂᕆᔨᑎᑦ ᐃᓂᒃᑎᓵᓂᒃ ᖃᐅᔨᓴᖅᑐᓂ ᐊᓪᓚᓗ ᐱᖏᓇᐅᖅᓴᕐᔪᓂ;

(b) ᑎᓕᐅᓇᔪᑦ ᐊᓪᓚᓗ ᐱᖏᓇᐊᑭᓕᓇᔪᑦ ᐃᓄᐃᑦ ᑲᔪᓯᑎᐊᕆᔭᓇᑎᑦ ᓴᓪᓂᒋᐊᕆᓂᒃ ᐊᓪᓚᓗ ᖃᐅᔨᓴᖅᑎᒋᔭᖅᑎᓗᑎᑦ ᐊᓪᓚᓗ ᐃᓕᓴᐅᓇᐃᖏᖅᓯᐊᕆᖅᑕᑦ ᐃᓄᖅᑲᓇᐃᖏᓇᑎᓂᖅᓯᐊᕆᖅᑕᒥᓐᓄᑦ;

(c) ᐊᐅᓚᑎᖅᑎᑎᑦ ᖃᐅᔨᓴᐃᓂᕆᑲᒃ ᐱᔪᓐᓇᐅᑎᑕᖅᖢᑦᓕᐊᓂᓐᓄᑦᑕ ᐱᓕᑦᓚᐊᓂᓐᓄᑦ ᑕᒃᑯᓇᓐᓄᑦ ᑲᑎᒪᔨᖅᓯᓐᓂ ᑲᑎᒪᔨᖅᓯᓐᓂ ᐊᐅᓚᑎᖅᑎᑎᓐᓄᑦ ᐊᐅᓚᑎᖅᑎᑎᓐᓂᒃ ᐊᓪᓚᓗ ᐊᑐᑎᖅᖢᖅᑐᑦ ᐊᐅᓚᑎᖅᑎᑎᓐᓂᒃ ᐊᓪᓚᓗ ᐃᓄᐃᑦ ᐅᓂᒃᑳᖅᑐᑦ ᐊᓪᓚᓗ ᐊᑐᑎᖅᖢᖅᑐᑦ ᐃᓄᐃᑦ ᐅᓂᒃᑳᖅᑐᑦ ᐊᓪᓚᓗ

(d) ᐃᓕᓴᐅᑎᒃᓯᑦ ᐃᓄᐃᑦ ᖃᐅᔨᓴᕐᖃᖅᖢᑦ ᐊᓪᓚᓗ/ᐅᕙᓪᓕᔪᓐᓇᑦ ᐃᓄᐃᑦ ᐅᓂᒃᑲᖅᑭᑦ ᐃᓘᓚᒃᑯᓕᓂᓂᒃ ᖃᐅᔨᓴᖅᓂᕐᒧᑦ, ᑕᐃᒃᓂ ᐊᑐᑎᖃᖅᑎᓯᓚᑦ; ᐊᓪᓚᓗ

(e) ᐱᖅᑎᖅᖅᓚᑎᒃ ᐊᐅᑯᓕᔪᓂᓴᓕᒪᓕᐅᑐᒃ ᑕᐃᒃᓂ ᑕᐃᓚᵒᓇᐅᔨᵒᓇᖅᖅᑕᐅᑎᑯᓕᓚᒍ ᐊᐅᓕᕆᖅᑎᑎᐅᓐᓂᖅᓕᒧᑦ ᑲᑎᓚᔪᖅᓭᑯᑦ ᖃᐃᒃᓂ ᖃᐅᔨᓴᖅᑕᓇᓕᑦ ᐱᔨᵒᓇᐅᑎᓴᖅᓗᓂᵒᓂᒃ ᐱᓕᕆᓐᓕᐊᑎᑯᓕᑦ.

10.3 ᐃᓇᐃᑦ ᓇᓇᒥ ᖃᐅᔨᓴᖅᑐᓂ ᐃᒃᔨᖅᑎᑦ

10.3.1 ᑭᓯᐊᓂ ᐊᓐᕆᖅᑲᑎᐅᓐᓯᓚᑦᓴᑦᓂᖅ ᑕᒃᑯᐊᓗ ᐊᑐᑎᖅᑲᖅᑐᑦ ᐊᐃᒃᑐᑦᖅᓯᓚᔨᑦ ᐃᓇᐃᑦ ᑲᑐᔨᖅᑎᑎᐅᓐᓂᕆᑦ, ᑲᓇᒥᒥ ᐅᒪᔨᔨᓕᓚᔪᔨᖅᔨᐊᑯᑦ (ᐃᓕᓐᓴᐅᑐᓴᑎᒃ ᑕᒪᖅᒥᒃ ᐃᓅᖅᑲᓇᐃᖅᑲᐅᑎᑦ ᐊᓪᓚᓗ ᑲᓐᑐᑦᑕᐅᓯᓚᔨᑦ) ᐃᒪᓐᓇᐅᓂᐊᖅᑐᑦ, ᐊᑐᓂ ᓇᓇᒥ ᐊᖅᑦᔪᑦ ᐃᓕᓐᓄᑦ:

(a) ᐃᓅᖅᑲᐃᑦᕈᑎᓴᑎᒃ ᐅᓇᓀᓚᐊᖅᓴᓕᑦ ᐃᓇᐃᑦ ᖃᐅᔨᓴᖅᑐᓂ ᐃᑲᔨᕈᑎᓂᒃ ᐅᑯᐊᓴ ᐊᑐᓂ ᑲᑲᓇᒥᒥ ᐅᒪᔨᔨᓕᓚᔪᔨᖅᔨᐊᑯᑦ ᓇᓇᓂ ᐱᓕᓚᓚᖏᓕᑦ ᐃᓂᖅᑎᑎᕈᓐᓇᓐᓂᒃ, ᐊᓪᓚᓗ ᖃᓇᐃᓕᓂᐅᓴᕆᓴᖅᑎᑎᓇᓕᑦ ᐊᑕᐅᔪᖅᑭᒃ ᐃᑲᔨᔨᖅᑎᑎᓴᑎᒃ ᐊᑐᓂ ᖃᐅᔨᓴᖅᑕᖏᓂᓂ; ᐊᓪᓚᓗ

(b) ᑐᓂᓴᓕᑦ ᓇᓇᒥ ᐱᓴᓚᐅᖅᓴᑎᓐᓴᓕᑦ ᐃᓅᖅᑲᓇᐃᖏᑎᓐᓂᒃ ᒪᓕᓪᓴᓗᒍ 10.3.1(a), ᑕᒪᑲᑯᐊᓗ ᐅᖅᑲᐅᔪᑦ ᐊᓪᓚᓗ ᑎᑎᕋᖅᔪᓚᔨᑦ ᖃᐅᔨᓴᖅᑎᑐᐊᖅᓴᓕᑦ ᐊᑐᓂ ᐃᓅᑉ ᐱᓕᓚᓂᖏᓐᓄ ᑕᐃᒃᓂ ᐃᓕᓴᑎᓇᓗᒍ ᐃᓅᖅᑲᐃᑦᕈᓚᖏᓐᓄ.

10.3.2 ᑲᓇᑕᒥ ᐅᒪᔨᓕᓚᔪᔨᐊᑯᑦ ᓴᓂᖅᓀᐃᔪᒪᓂᐊᖅᑐᑦ ᐃᓇᐃᑦ ᐊᑲᖅᑕᐅᓂᐊᕐᓂᖏᓐᓄᓇᑦ ᐱᓀᑦᓕᕐᑕᐅᔨᓄᖅᑐᓇᓕᑦᓴ ᐊᖏᕐᕕᑦ ᑎᑎᕋᖅᔪᓚᔨᖅ 8 15-1 ᑭᓇᐅᔭᖅᓴᐃᑦ ᐊᑭᖏᑦ ᐃᓇᐃᑦ ᖃᐅᔨᓴᖅᑐᓂ ᐃᑲᔨᕈᑎᑎᒍᓂᑦ ᐱᓕᕆᓐᓕᐊᑎᑯᓕᖅᖅ ᐅᓴᓇᐃᖅᔪᓚᔨᖅ 10.3.1. ᑕᒪᖅᑭᒃ ᐱᔭᐅᔨᖅ ᑭᓇᐅᔭᖅᓴᕐᑎᑎᒍᔨᓂᒃ ᐊᔪᖏᓐᓂᒃ ᑲᑦᑯᓇᒥᒃᑕ ᐃᓇᐃᑦ ᐊᑲᖅᑕᐅᓂᐊᕐᓂᖏᓐᓄᓇᑦ ᐱᓀᑦᓕᕐᑕᐅᔨᓄᖅᑐᓇᓕᑦᓴ ᐊᑐᓂ ᐃᓕᓴᑎᓇᓗᒍ ᐊᖏᓕᓚᒍᓂ ᖃᐅᔨᓴᖅᑎᑎᒍᓂᑦ ᐃᓇᐃᑦ ᐊᑲᖅᑕᐅᓂᐊᕐᓂᖏᓐᓄᓇᑦ ᐱᓀᑦᓕᕐᑕᐅᔨᓄᖅᑐᓇᓕᑦᓴ ᐊᑐᓂ ᐃᓕᓴᑎᓇᓗᒍ ᐊᖏᓕᓚᒍᓂ ᖃᐅᔨᓴᖅᑎᑎᒍᓂ

10.4 ᐃᓄᐃᑦ ᖃᐅᔨᓴᖅᑎᑦ

10.4.1 ᐱᖁᔭᓚᑎᓪᓗᑎᑦ ᐃᓄᐃᑦ ᖃᐅᔨᓴᖅᑎᑦ, ᑲᓇᑕᒥ ᐅᓚᔪᓕᓐᓄᔪᐊᕆᔪᑦ ᑐᓂᓯᓂᐊᖅᑐᑦ ᖃᐅᔨᓴᖅᑎᑎᒃ ᐅᔪᓂᖅ:

(a) ᐃᓱᐳᓇᔭᓗᑎᑦ ᐃᓄᓕᒻᒥᓄᑦ ᑐᓴᖅᑕᐅᓚᐊᕈᑎᒃ ᖃᓄᐃᙱᑦᑐᑦ ᐅᔪᐊᓗ ᐊᑐᐃᓇᐅᔭᔪᑦ ᐃᓗᐊᓂᑎ ᑲᑎᒪᔨ ᑲᑎᑕᐅᔭᓕᕆᔨᑦ ᑲᑎᒪᓗ ᐅᔪᐊ ᐃᑲᔪᐳᓇᐅᓕᑦ ᖃᐅᔨᓴᖅᑎᓂᒃ ᐱᓕᓚᓗᑎᒃ ᖃᐅᔨᓴᖅᓯᐊᖅᑐᖅᑕᑎᒃ, ᐃᓕᒥᕆᔭᒃᓗᑎᒃ ᐊᑐᑎᖅᑐᖅᑐᑦ ᖃᐅᔨᓴᕈᑎᓄᑦ ᐅᓂᒃᑐᑦ, ᑐᒃᓱᕈᐅᑎᑦ, ᑲᑎᕐᕕᐅᔪᓄᑦ ᖃᓄᐃᑲᖅᑐᑦ ᐊᑲᓕᑦ ᑐᑭᓯᒪᔪᑦ;

(b) ᐃᓕᓐᓂᓪᓚᒃ ᐃᓱᐳᓇᔭᓗᑎᒃ ᒎᒃᓯᓚᓐᓇ ᑲᓇᑕᒥ ᐅᓚᔪᓕᓐᓄᔪᐊᕆᔪᑦ ᐃᒃᓗᔭᓯᑎᓐᓂᓪᓚᓄᑦ ᐊᑲᓕᑦ ᓴᓇᕕᒻᒥᓐᓂᓪᓚᓄᑦ, ᐊᑐᐃᓇᐅᔭᔪᑦ, ᑕᐃᑲᓂᓂᑎᑦ ᐊᑲᓕᑦ ᐊᑐᐃᓇᐅᔭᔪᑦ ᐃᓄᐃᑲᖅᑕᐅᔭᕆᓚᖅᑐᓂᒃ ᐃᓄᐃᑲᖅᑕᐅᔭᔪᖅ ᖃᐅᔨᓴᖅᑕᐅᔭᐅᓯᑎᐊᖅᖢᑦᑕᐊᑎᓗᓗᑎᒃ ᖃᐅᔨᓴᖅᑕᐅᔭᔪᖅ;

(c) ᐱᓕᓚᑦᓴᖅᑕᖓᓯᑎᑦ ᑕᒪᕐᕉᒃ ᑲᑎᑕᐅᓇᓕᑦ ᐅᓚᔪᓕᓐᓄᔪᐊᕆᔪᑦ ᐱᓯᒪᔪᑦ ᖃᐅᔨᓴᖅᑎᑦ ᖃᐅᔨᓴᖅᑕᖓᓯᓂᓕᑦ; ᐊᑲᓕᑦ

(d) ᓇᒪᒃᑎᓪᓗᔪᓄ ᐊᑲᓕᑦ ᐊᑐᐃᓇᐅᔭᑎᓪᓗᔪᓄ, ᐃᓱᓪᓗᔪᓄ, ᐊᑭᓐᓇᖅᑐᓪᓗᑎᒃ ᑕᒃᑯᐊ ᖃᐅᔨᓴᖅᑎᑦ, ᖃᖓᑕᒋᔭᕐᓗᑎᒃ, ᓄᓇᓕᖅᑎᓂᒃ ᐅᖅᓯᓇᓐᓂᑦ ᐅᑎᐊᓂᒃᑐᑦ ᖃᓄᐃᙱᑦᑐᑦ ᐃᓕᓴᕐᓂᖅᑐᑦ ᐃᓚᖓᑎᓪᓗᔪᓄ ᐅᖅᓯᓇᓐᓂᑦ ᐅᑎᖅᑕᐸᓪᓕᐊᓂᒃ ᐅᖅᓯᓇᓐᓂᑦ ᐅᑎᖅᑕᐸᓪᓕᐊᓂᒃ ᐊᑐᐃᓇᐅᔭᔪᑦ ᐱᓚᒋᑎᓪᓗᔪᓄ ᐊᐅᓚᑎᓂᓪᓚᓄᑦ ᐊᑲᓕᑦ ᐱᓚᒋᑎᓪᓗᓂᓪᓚᓄᑦ ᖃᐅᔨᓴᖅᑕᖓᓂᒃ ᐃᓗᐊᖓᓂᓪᓚᓄᑦ ᐅᑎᖅᑕᐸᓪᓕᐊᓂᒃ ᐃᓄᐃᑲᖅᑕᐅᔭᔪᑦ ᐱᓚᒋᑎᓪᓗᔪᓄ ᐅᖅᓯᓇᓐᓂᑦ ᐅᑎᖅᑕᐸᓪᓕᐊᓂᒃ ᐱᓚᒋᑎᓪᓗᓂᓪᓚᓄᑦ ᐅᑎᖅᑕᐸᓪᓕᐊᓂᒃ ᐃᓄᐃᑲᖅᑕᐅᔭᔪᑦ ᐃᓗᐊᖓ

10.6 ᐊᑐᓕᕆᔨᐅᓪᓗᒃ ᐅᕙᓪᓚᓂᓕᓂᒃ ᐊᐅᓪᓚᑦᑎᐊᓕᖅᓴᖅᑕᐊᓚᑦ ᓄᓇᓄᑦ

10.6.1 ᐊᐅᓚᓪᑎᔨᐱᕆᔭᓄᑦ ᐸᖅᓇᐅᑎᑦ ᓇᓗᓇᐃᖅᓭᓕᒫᖅᑐᑦ, ᐊᑐᑎᖅᑲᖅᓂᖏᓐᓄᑦ:

(a) ᓄᓇᐃᑦ ᐊᓖᒪᓗ/ᐅᕙᓚᓂᓂᒃ ᖃᓄᐃᓕᓂᖏᑎᓪᓗᑎᑦ ᑕᐃbᓂ ᐊᑐᖅᓄᑦᑐᐊᖅᓂ ᐅᕙᓚᓂᓂᒃ ᑕᐃᒪᓇᐅᔭᕆᐊᖅᑲᖅᑐᑦ ᐅᓄᐊ ᖃᐅᐅᔨᖅᑎᑦ ᒪᓕᒃᑎᖅᑲᓂᐊᖅᓯᓗᑎᒃ ᐃᓄᓐᓂ ᐊᐅᓪᓚᑦᑕᑎᕆᐅᓯᒪᓗᑎᒃ; ᐊᖓᒪᓗ

(b) ᐊᑐᓪᓄᓕᐅᔨᑦ ᐊᔪᖏᑦᑕᕆᐊᖅᑲᖅᑕᖏᓐᓂᓂᒃ ᐊᐅᓪᓚᑦᑕᐅᖅᑲᖅᑕᓂᐊᖅᓯᓗᑎᓗᓂ, ᐃᓚᒋᔭᐅᓗᑎᒃ ᐊᑐᓪᓄᓕᐅᔨᑦ ᐱᓪᓗᐊ ᐊᑭᓕᓂᖅᑕᖅᑕᑕᐃᓚᒋᔭᐱᑎᖅᓱᓗᒃ ᓇᓗᓇᐃᖅᓭᓕᒫᖅᑐᑦ ᐊᐅᓚᓪᑎᔨᐱᕆᔭᓄᑦ ᐸᖅᓇᐅᑎᑦ.

10.6.2 ᐅᑯᐊ ᐊᐅᓚᑦᑎᖅᑎᑉᓴᓂᓐᒧᑦ ᑲᑎᒪᔨᖏᓪᓕ ᖃᑎᒌᖓᔪᖅᑎᖅᑲᓐᓂᒃ ᐊᑐᑎᔨᓂᒃ ᓇᓗᓀᓂᓂ ᐊᖏᔪᓂᑐᐃᓐᓇᑯᑦ ᓇᓗᐃᑦ ᐃᓗᐊᓕᐊᖅᓯᒪᔨᑦ ᐃᓗᐊᓂ ᐊᐅᓚᓪᑎᔨᐱᕆᔭᓄᑦ ᐸᖅᓇᐅᑎᑦ ᒪᓕᒃᓗᒍ 10.6.1(a).

10.6.3 ᑕᐃᒪᓗᒍ, ᐊᖏᑕᐅᔨᓴᖅᓯᒪᓕᕐᓂᐅᓂ ᐊᐅᓚᑦᑎᔨᐱᕆᔭᓄᑦ ᐸᖅᓇᐅᑎᑦ, ᐊᑐᑎᖅᑲᖅᑐᑦ ᓇᓗᒥᒃ ᑲᑐᔭᖅᑲᑎᑦᑕᖅᑐᑦ ᑲᑎᒪᔨᖏᓪᓕ ᑐᓴᖅᑎᑕᐅᓂᐊᖅᑐᑦ ᓇᓗᒥ ᐊᑐᖅᑲᓂᐊᖅᑕᖏᓐᓂᓂᒃ ᐊᔪᖏᑦᑕᑎᐊᖅᑐᑦ ᐊᐅᓚᓪᑎᔨᐱᕆᔭᓂᐊᖅᓯᒪᓕᖅᑐᑦ ᐅᕙᓚᓂᓂᒃ ᑕᐃᒪᓇᐅᔭᕆᐊᖅᑲᖅᑐᑦ, ᐃᓕᓴᕐᓂᑎᑦᑐᓗᒍ ᖃᖅᑲᓇᐅᔭᖅᑐᖅᓯᒪᔨᖅᑐᑦ ᐊᖏᔪᓂᑐᐃᓐᓇᑯᑦ ᐊᑭᑭᓕᒫᖅᑐᑦ ᓇᓗᒥᒃ ᑲᑐᔭᖅᑲᑎᑦᑕᖅᑐᑦ ᐃᓕᓴᕐᓂᑎᑦᑐᓗᒍ ᑎᑎᕋᖅᓴᕐᓂᖅ ᐃᓕᓴᕐᓂᑎᑦᑐᓗᒍ 3.6 ᐊᑐᑎᓂᐊᖅᑐᑦ.

10.6.4 ᑲᓇᑕᒥ ᐅᒪᔨᓕᑦᓚᔨᓄᑦ ᐱᓕᕆᑎᖅᑎᑦ ᐃᓕᓕᐅᑎᓴᒪᓗᑎᑦ ᑕᒪᑲᔪᓄᖓ ᐱᔨᓄᐊᓕᐅᑎᑦ ᐊᖓᒪᓗ ᓇᓕᐊᑐᐃᓄᐊᓐᓇᖅ ᐊᐅᓪᓚᑦᑕᐅᖅᑲᖅᑐᑦ ᐱᓴᒪᔨᑦ ᐱᔨᓄᐊᓐᓇᖅᑎᓄᑦ ᑕᐃᒪᓇᐅᔭᕆᐊᖅᑲᖅᑕᖏᓐᓂᒃ ᐊᖓᒪᓗ ᐃᓕᓕᐅᑎᓴᒪᓗᑎᒃ ᐱᔨᓄᐊᓐᓇᖅᑎᓄᑦ ᓇᓕᐊᑐᐃᓄᐊᓐᓇᖅ ᐊᐅᓪᓚᑦᑕᐅᖅᑲᖅᑕᓂᑐᑦ ᐊᑐᓕᖅᑕᐅᖅᑲᓐᓂᖏᓐᓄᑦ ᓇᓗᓇᐃᖅᓭᓕᒫᖅᑐᑦ ᐃᓗᐊᓕᐊᖅᓯᒪᔨᑦ ᐃᓗᐊᓂ ᐊᐅᓚᓪᑎᔨᐱᕆᔭᓄᑦ ᐸᖅᓇᐅᑎᓂᓐ.

ᐃᓕᖅ 11 — ᐃᓄᖅᓄᑦ ᐱᒻᒪᓗᑎᑐᐃᔨᑦ ᐊᒻᒪᓗ ᐱᖏᓪᓗᕆᖅᑎᑦ

11.1 ᑐᑭᒃᓯᑦᑕᐊᔨᐱᑎᒃᓴᑦ

11.1.1 ᐅᖏᓂ ᐃᓕᖅᓴᓂ ᐊᑕᓂ ᑎᑎᕋᒃᓱᒪᔨᑦ ᑐᑭᒃᓯᑦᑕᐊᔨᐱᑎᒃᓴᖅᑎᑦ:

(a) ᓴᐳᒻᒥᓗᑎᑦ ᐊᒻᒪᓗ ᓄᖅᔪᑕᐅᑕᐃᓕᑕᑎᑦᓗᑎᑦ ᐃᑕᖏᑕᕋᒃᖁᑐᑦ ᓄᓇᒥ ᐃᓂᖅᑎ, ᐃᑕᖏᑕᐃᒃ, ᐊᒻᒪᓗ ᐃᓄᖅᓄᑦ ᐃᓂᕆᐅᕆᔨᑦ ᐱᒻᒪᓕᕆᔨᑦ ᐃᓄᖅᓄᑦ; ᐊᒻᒪᓗ

(b) ᒪᓕᕐᑎᐊᖏᓗᐊᖏᓐᓇᑕ ᑕᒪᖏᒃ ᖃᓄᐃᓕᓕᕆᕆᔨᑦ ᐃᓗᖅᑎᓂᒃ ᑐᓴᕐᓂᒃᑎᑦ ᓄᓇᖏᑎᖅᑎᑦ ᐊᒻᒪᓗ ᑎᖅᕿᐊᓄᑦ ᑎᑭᖅᑐᐃᓄᑦ ᓴᐳᖏᑕᐊᖏᓐᓄᑦ ᑕᐃᒪᓂᓄᐅᓕᖏᑐᐃᓄᑦ ᑕᒥᔪᑎᒍᑦ ᐊᒻᒪᓗ ᐃᓕᖅᓴᓂ 33 ᑭᒃᑯᓄᓂ *ᓄᓇᕗᒻᒥᒃ ᓄᓇᑕᕆᑎᓄᑦ ᐊᖏᕆᕆᑎᑦ.*

11.2 ᑐᑭᖅᑎᑦ

11.2.1 ᐱᔪᐊᑎᑦᓗᑎᑦ ᕆᕐᖅᓱᒃ ᐃᓕᖅᓴᓄᑦ:

(a) “ᐃᑦᑕᕋᑦᑕ ᐱᕋᐊᑎᑦ” ᑕᐃᒪᓐᓇᓕᑕᐃᓐᓇᖅ ᑐᑭᓕᕆᖅᓱᒪᔨᖅ ᑕᐃᒃᑐᒥ *ᓄᓇᕗᒻᒥ ᐃᑕᕋᑦᑕᓕᕐᓂᖅ ᐊᒻᒪᓗ ᐅᓴᕋᖅᓯᒪᔨᓂᒃ ᐃᒥᖅᑎᓄᑦ ᐱᕋᐊᑎᑦᑦ;*

(b) “ᐃᑕᕋᑦᑕᑦ ᑭᓱᓂᖅᑎᑦ” ᑕᐃᒪᓐᓇᓕᑕᐃᓐᓇᖅ ᑐᑭᓕᕆᖅᓱᒪᔨᖅ ᑕᐃᒃᑐᒥ *ᓄᓇᕗᒻᒥᒃ ᓄᓇᑕᕆᑎᓄᑦ ᐊᖏᕆᕆᑎᑦ;*

(c) “ᐃᑕᕋᑦᑕᑦ ᓄᓇᒥ ᐃᓂᖅᑎᑦ” ᑐᑭᖅᑲᖅᑐᖅ ᐃᑦᑎᑦ ᓇᓗᓇᐃᖅᓱᒪᔨᑦ ᓇᓕᐊᖏᑐᐃᓐᓇᖅ ᐱᕋᐊᑎᑦᑕᒥ ᐅᖏᓪᓗᑎᑦ *ᓄᓇᕗᒻᒥᒃ ᓄᓇᑕᕆᑎᓄᑦ ᐊᖏᕆᕆᑎᑦ;* ᐊᒻᒪᓗ

(d) “*ᐱᕋᐊᑎᑦ*” ᑐᑭᖅᑲᖅᑐᑦ *ᓄᓇᕗᒻᒥ ᐃᑕᕋᑦᑕᓕᕐᓂᖅ ᐊᒻᒪᓗ ᐅᓴᕋᖅᓯᒪᔨᓂᒃ ᐃᓂᖅᑎᓄᑦ ᐱᕋᐊᑎᑦᑦ.*

11.3 ᐱᐅᓯᑦᑲᖅᑲᖅᓇᐅᔨᑦ ᐃᓂᖅᑎᑦ ᐱᒻᒪᓗᐅᔨᑦ ᐃᓄᖅᓄᑦ

11.3.1 ᐱᓇᓯᐊᖅᑲᑎᑦᑦ ᐃᓕᓴᖅᓯᒪᔪᑦ ᐅᓇ ᒥᓂᑦ ᐃᓇᓯᓕᓗᓴᐅᑯᓄᑦ ᐊᒻᒪᓗ ᐃᓄᐃᑦ ᐱᕋᐊᑎᑦᑕᖅᑑᑎᑦ ᐊᔪᖅᑎᑦᑎᐊᖅᑐᑦ ᓇᐅᑦᓯᓂᑐᐊᖅᑐᑦ ᐊᒻᒪᓗ ᓄᓇᖏᑎᖅᑎᑦ ᐳᓴᕐᓂᒃᑎᑦᑦ ᐊᒻᒪᓗ ᓄᓇᖏᑎᖅᑎᑦ, ᐃᑕᖏᑕᓂᒃ ᐃᑕᖏᑕᐃᒃ ᖃᐅᔨᓴᓕᑎᒋᔪᑦ, ᐊᒻᒪᓗ ᐃᑕᖏᑕᐃᒃ ᐃᓄᖅᓄᑦ ᐊᒻᒪᓗ ᐅᓴᕋᖅᓯᒪᔨᑦ ᐃᓄᖅᓄᑦ ᐊᑎᓂᑎᓗᑎᒃ ᐊᒻᒪᓗ ᐃᓄᖅᓄᑦ ᐊᑎᓂᑎᓗᑎᒃ ᑕᒪᐃᓐᓇ ᐱᕋᐊᑎᑦᑦ ᐃᓄᖅᓄᑦ ᐊᒻᒪᓗ ᐃᓄᖅᓄᑦ ᐊᑎᓂᑎᓗᑎᒃ

11.3.3 ᐅᓇ ᒥᓂᑦ ᖃᐅᔨᓴᖅᓂᐊᖏᑐᖅ ᐅᑯᐊ ᑕᒪᖅᑭᒃ ᑲᓇᑕᒥ ᐃᒪᓕᒫᓴᕐᕕᐊᑯᑦ ᐃᖏᑲᓇᐃᔨᕐᑎᖓᑦ ᐊᓯᒪᓕ ᑲᓐᑐᖁᑕᐅᓗᒍᑦ ᓴᖅᑭᑐᖅᑐᑦ, ᐊᓯᒪᓕ ᐱᒫᒥᓂᐊᖏᑐᖅ ᐊᑐᑎᖃᖅᑐᓂᖅ ᐱᑎᐊᖅᑐᓂ ᐅᑯᐊ ᐱᕐᓇᐃᓇᑎᒋᑐᑦ ᐊᒋᔪᑦ *ᑲᓇᑕᐅᑦ ᓂᕐᔪᑎᖓᓐᓇᓂ ᐱᖁᓯᕐᕕᖏᖅ* ᐊᓯᒪᓕ *ᑎᓐᓯᐊᓇᑦ ᑎᑭᓂᖅᑐᓂ ᑲᒪᒋᔭᐅᓂᖅ ᐊᖏᖅᑐᓂ* ᑐᓴᖏᑦᑐᑎᓪᓗᒍ ᐱᖁᓯᕐᕕᖏᖅ ᑐᓴᖅᑎᑕᐅᔪᓐᓇᑎᑐᖅ ᐱᖁᓯᕐᕕᖏᓂᖅ ᐊᓯᒪᓕ ᒪᑕᑭᖅᑕᕐᓂᖓᑦ ᑕᒪᓐᓇᖅᑐᑦ.

11.3.4 ᑕᐃᒪᓗ, ᐊᑯᓂᐅᔪᒥ ᓄᓇᒥ ᓴᓇᔨᐅᓂᖓ, ᐃᖏᑲᓇᐃᔨᕐᑎᖓᑦ ᐊᓯᒪᓕ ᑲᓐᑐᖁᑕᐅᔪᓐᓇᑦ ᓇᓂᔪᒍᑎᑲ ᐊᖏᑲᓴᔪᒍᑎᑲ ᐃᑕᖏᓂᕆᓪᓕ ᐃᓂᖓᑦ, ᐃᑕᖏᓂᑕᓂᑲ ᖃᑲᐅᔨᑕᑎᒋᔭᖅᐅᔪᑦ, ᐊᓯᒪᓕ ᐃᑕᖏᓂᑕᑦ ᐱᖏᖁᖃᑲᐅᔨᓴᖅᑐᓂᖅ ᐊᑐᔪᑦ ᐃᓕᓂᖅ 6.4 ᑎᑭᑦ 6.6 ᐅᕐᓂ ᐃᓇᐃᑦ ᐊᑐᖏᑕᐅᓂᐊᖏᖅᑐᓂᖅ ᐱᕐᓕᓕᕐᒋᐅᔨᓇᖅᑐᓇᓕᓪᓗ ᐊᖅᑭᒋᑦ, ᐊᔪᓕᓕᔨᓕᑎᑦ, ᐊᓯᒪᓕ ᓇᓕᓇᖅᑐᑦ ᓇᓂᖅ ᑲᒪᑎᕐᓇᒪᖅᓂᖅ ᐃᓂᖅᑭᑦᓂ, ᖃᐅᔨᓴᖅᑎᑕᐅᔪᓐᓇᑦ ᐅᕐᕕᓇᓐᓂᑦ ᐃᑕᖏᓂᑕᓂᑦ ᐱᖁᓯᕐᕕᖏᓂᑦ, ᐊᓯᒪᓕ ᑐᓴᖅᑎᑕᐅᔪᓐᓇᑦ ᑕᒪᑐᖅᓱᒪ ᐱᐅᔪᑐᖅᑕᓚᕐᓇᐃᓇᓂᑦ, ᐃᓇᐃᑦ ᐱᖁᓯᕐᕕᖏᓂᑦ ᐅᑭᔪᖅᑐᑦ ᐱᓕᕐᓂᖅ ᐃᓕᓴᕐᕕᖏᓂᑦ ᐃᓇᐃᑦ ᐱᖁᓯᕐᕕᖏᓂᑦ ᐊᑐᔪ

ᐃᓕᓐᓴ 12 — ᓂᕆᑎᖅᓴᕐ

12.1 ᑐᑭᒃᓯᕐᓗᐊᓂᖅᓴᕐ

12.1.1 ᐅᕙᓂ ᐃᓕᓐᓴᓄᓂ ᐊᑦᓂ ᑎᑎᕆᖅᓯᒪᔪᓕ ᑐᑭᒃᓯᕐᓗᐊᓂᖅᓴᖅᓯᓐᓂᕐ:

(a) ᓴᓗᒥᖅᓕ ᓂᕆᔪᑎᓄᓕ ᓄᓐᔪᑕᐃᓕᓗᑎᑕᑎᓂᔨᔪᑕᐅᔪᓕ ᐊᒻᒪᓗ ᓂᕆᔪᑎᓕ ᐃᓂᓐᓂᕐ; ᐊᒻᒪᓗ

(b) ᐊᑐᑎᔨᓂᒃ ᐊᑭᓕᖅᑕᐅᓗᑎᒃ ᐃᓄᐃᓕ ᑐᐊᐃᓴᕆᓐᖃᖅᑐᓄᓕ, ᐱᖓᓗᐊᖅᑭᔨᓄᓕ ᐅᕝᕙᓂᓐᓂᑦᓂᕐ ᑐᓴᕐᓗᐊᖅᑐᕆᕐᖃᖅᑐᓂᕐ ᓇᓐᓂᐊᕐᓂᖅ ᑎᓇᓕᒫᓂ, ᐊᒃᓱᕈᓯᕐᓗᒍ ᐅᕝᕙᓂᓐᓂᑦᓂᕐ ᑎᑎᕆᖅᓯᒪᔪᓕ ᐅᑎᖅᑕᕐᓂᑐᐃᓐᓇᓕᒫᒃ.

12.2 ᓂᕆᔪᑎᓄᓕ ᐊᐅᓪᓕᖅᑎᔨᕐᓗᐊᓂᖅ

12.2.1 ᓄᓇᕗᓂ ᑐᓴᖏᓐᓄᒃ ᑎᒥᖓ ᑐᓂᓯᓂᐊᕐᓂᖅᑐᖅ ᓄᓇᕗᓪᓕ ᐅᓪᓗᕆᓚᐅᔨᔪᐊᕐᑯᓄᓕᓂ, ᑲᓇᑕᒥ ᐅᓪᓗᕆᓚᐅᔨᓴᔨᔪᐊᕐᑯᓄᓕ ᐊᒻᒪᓗ ᐊᓯᖏᓂ ᐊᑐᑎᓴᕐᓂᖅᑐᖅ ᑎᑎᕐᐅᔨᓄᓕ ᓂᕆᔪᑎᓴᕐᓂᖅᑐᖅᓂᒃ ᕿᑐᐅᓪᓚᕆᓴᕐᓂᖅᑐᖅ ᐱᓕᓚᐅᔨᓂᒃ ᐃᓄᖅᓱᓕᓂ ᐅᑯᐊᓗ ᐱᖓᐅᓯᓚᕐᓂᖅ ᒪᓕᒐᕐᓂᖅ ᐃᓕᓐᓂ 6 ᐱᖓᑯᖏᓂᕐ.

12.2.2 ᐅᖓᑲᐅᑎᖓᐅᓯᒪᓗᑎᒃ ᐊᑐᑎᓴᕐᓂᖅᑐᖅ ᐊᐅᓪᓕᖅᑎᓂᖅᑎᑲᖏᓐᓂᔪᓕ ᑲᑎᒪᔨᖅᓭᓕ, ᐅᑯᐊ ᑲᑭᑕᒥ ᐅᓪᓗᕆᓚᐅᔨᔪᐊᕐᑯᓄᓕᓂ ᓇᓗᓇᕐᑐᖅᓯᓗᓂᒃᑯᓐᓂᔪᓕ ᐱᓕᓚᐅᔨᓄᓕ ᐃᓄᐃᓕ ᑕᒪᐃᓐᓇᕐᓂᑎᔪᓕ ᑲᑎᒪᔨᖅᓭᓕ ᐅᕝᕙᓂᓐᓂᑦᓂᕐ ᑎᓐᓂᒃᑕᕐᓂᑐᐃᓐᓇᓕᒫᒃ ᓄᓇᓕᖕᒥᐊᓐᓂᕐᓂᑎᔪᓕ ᐅᑯᐊ <ᐃᓕᓴᐅᔪᓕ ᐊᒻᒪᓗ ᑐᓂᐅᖅᑲᐅᑕᐅᔨᓴᓇᓐᓂᖅᑐᓂᕐ (<ᐃᓕᓴᐅᔪᓗᑎᒃ ᐅᕝᕙᓂᓐᓂᓂᕐ ᖃᓄᐊᖅᑲᐅᑯᑦ) ᐊᑐᔪᖅᑭᒪᓕᕆᓂᕐᓂᖅᓯᓂᕐ ᐱᑎᐊᖅᑲᓐᓇᑐᓂᕐ ᐅᑦᓗᓐᓄᑦ, ᑕᐃᒪᓗ ᖃᐅᔨᓴᒪᓂᕐᓂᔨᓂᕐ ᑕᒪᐃᓐᓇᕐᓂᖅᓯᒪᓂᕐ ᐃᓕᓐᓂ 5 ᒡᒃᑯᓇᓂᑦ ᓄᓇᕗᓪᓕᖅᑐᒃ ᓄᓇᕗᑦᑕᑎᓄᓕᓂ ᐊᖏᕐᑕᑎᓂᕐ. ᐃᓱᒪᓕᕆᓂᑦᓗᔪ, ᑲᓇᑕᒥ ᐅᓪᓗᕆᓚᐅᔨᔪᐊᕐᑯᓄᓕᓂ ᐃᓪᓕᖅᑐᓂᐅᔪᐊᕐᓂᖅᓯᓗᓂᒃᑯᓐᓂᖅ:

(a) ᐊᑦᑎᒃᓱᓱᒪᑎᓕᓪᓗᑎ ᕿᓚᐊᓕᐃᑐᓴᖅᑐᖅ ᓂᕆᔪᑎᓂᖅ ᐊᒻᒪᓗ ᓂᕆᔪᑎᓂ ᐃᓂᓐᓂᕐᓂ, ᐊᒻᒪᓗ ᖃᐅᔨᓴᓕᖅᐅᓕᕆᓂᑦᓂᕐᓂᐊᖅᑭᓪᓗᑎᒃ ᓴᓇᓱᒪᓂᕐᓂᖅᓯᒪᓂᕐ, ᓂᕆᔪᑎᓂ;

(b) ᖃᓄᐃᓕᓐᓄᕆᓂᖅ ᐱᓕᕆᓂᖅᐊᖏᓚᓂᑎᓂ, ᒪᓕᓂᑎᐊᖅᑭᓗᑎᒃ ᒥᓂᔾᔪᐅᕙᓂ ᓄᓇᕗᑦᑕᕐᓂᖅᓯᓂᕐ ᐊᐅᔪᓂ ᒡᒃᑯᓇᓂᖅ *ᑲᓇᕗᓂ ᐅᓪᓗᕆᓚᐅᔨᔪᐊᕐᑯᓄᓕ, ᑎᓐᓂᐊᓂ ᐅᑎᖅᑕᕐᓂᑐᐃᓐᓇᓕᒫᓂ ᐱᔨᔪᐊᕐᑕᐅᓂᕐᓂᖅᓯᓂᕐᓂᖅ ᐱᖓᑯᖏᓂᕐ*, ᐅᕝᕙᓂᓐᓂᑦᓂᕐ ᓂᕆᔪᑎᓄᓕ ᐃᓄᐃᓕᓐᓄᕆᓂᖅᓴᕐᓂᖅᓯᓂᕐᓂᖅ ᐱᖓᑯᖏᓂᕐ, ᐊᑐᑎᖅᑲᐅᑎᒃᓴᓂᕐᓂᖅ ᐱᐅᔪᑐᖅᑲᓂᕐ ᐱᓐᓂᐊᖅᑭᔨᓄᓕ ᐱᓕᕆᓂᖅᐊᖏᓚᓂᑎᓂ ᐱᓕᕆᓂᖅᐊᖏᓚᓂᑎᓂᓪᓗ ᓇᓕᓚᐃᐅᓪᓗᖅ *ᐃᓄᐃᓕ ᖃᐅᔨᓴᖅᑐᕆᕐᖃᖅᑐᓂᕐ* ᑎᑎᕆᖅᓯᒪᔪᓕ ᐊᒻᒪᓗ ᑐᓂᓯᓂᐊᕐᓂᖅᑐᖅᓂᒃ ᐃᓄᐃᓕ ᑕᒪᐃᓐᓇᕐᓂᖅᓯᒪᓂᕐ ᐊᓯᖏᓂᕐ ᖃᐅᔨᓴᓕᖅᐅᓕᕆᓂᑦᓂᕐ ᐊᒻᒪᓗ ᐊᓯᖏᓂᕐ ᖃᐅᔨᓴᓕᖅᐅᓕᕆᓂᑦᓂᕐ ᐱᓕᕆᓂᖅᐊᖏᓚᓂᑎᓂ;

(c) ᖃᐅᔨᓴᕐᑎᐊᖅᓯᓗᑎᑦ ᐃᓄᓕᒻᒥᑦ ᐃᓱᒪᓕᕐᓂᖅᑎᑦ ᓕᕆᔨᒃᑯᑦ ᓴᓪᓕᐊᖅᓯᓂᖅ ᐅᖅᓯᒪᔭᕐᓂᖅ ᐊᐅᓚᓕᑎᓂᖅ ᓂᕆᔨᐸᑎᓂ ᐊᐅᓚᓕᑎᓂᖅ, ᐅᖅᓯᒪᔭᓂᓂ ᐊᓱᒪᓚᑦ ᓂᕆᔨᐸᑎᓂ ᐊᐅᓚᓕᑎᓂᖅᓗᑦ ᑎᒥᐅᔨᑦ ᓄᓇᕗᑦᐸᖅᑐᑦ; ᐊᒻᒪᓗ

(d) ᐊᑐᑎᖃᖅᑎᖅᓯᓗᒍ, ᖃᐅᒥᖅᑕᐊᖅᐸᑎᓗᑎᑦ ᐃᓄᐃᑦ ᐊᐅᓚᕆᓴᖅᑎᑦ.

12.2.4 ᒪᓕᕆᐊᖅᑲᖅᑐᖅ ᐊᖏᕈᑎᑦ 4.6.1 ᐊᐅᓚᕆᑎᔨᐸᖅᓴᓂ ᐱᕆᐊᓚᐅᓂᖏᓂᖏᓂ ᐃᓱᐱᕆᑎᐊᖅᓂᑦ ᐃᓄᐃᑦ ᓄᓇᕗᑦᐸᖅᑎᑦ ᒪᓕᖅᓴᓂᖅ ᐃᓚᖅ 12.2.3, ᐅᔪᐊ ᑲᓇᑕᒥ ᐅᒪᔨᓚᓱᓂᔨᐊᔪᑦ ᑐᓴᕆᑎᕆᓂᐊᖅᑐᑦ ᐊᑐᑎᖃᖅᑐᓄᑦ ᐊᖁᖅᑐᖅᓯᒪᔨᑦ ᐃᓄᐃᑦ ᑲᑐᔭᖅᑎᑎᒋᓂᖅ. ᑲᓇᑕᒥ ᐅᒪᔨᓚᓱᓂᔨᐊᔪᑦ ᐅᓂᒃᑲᓚᐅᑐᓄᖅ ᐊᐅᓚᓕᑎᓂᖅᐸᑎᑎᕐᓂᖅ ᖃᓄᖅᑲᐅᒪᔨᓗᒃᑐᖅ ᐊᐅᓚᓕᑎᓂᖅᐸᑎᑎᕐᓂᖅ ᖃᖅᑐᖅᓯᒪᔨᑦ ᐃᓄᐃᑦ ᓄᓇᕗᑦᐸᖅᑎᑦᑕ ᒪᓕᖅᓴᓂ ᐃᓚᖅ 12.5.1.

12.2.5 ᐊᐅᓚᕆᐸᖅᑲᓄᑦ ᐸᖁᐅᑎᑦ ᓇᓴᐅᐃᖅᓯᓂᖅᑐᖅᑎᑦ ᖁᑲᓄᖅ, ᐱᑕᖅᑲᑭᓂ, ᐅᐸᑐᖁᑲᓂᖅᓴᓂᕆᐊᖅᑐᖅᑎᑦ, ᐊᒻᒪᓗ ᖁᑲᒪᔪᑦ ᑕᒃᑯᓄᑦ ᒪᓕᐹᑕᐅᓕᕆᓚᐅᖅᓂᑦ 12.2.1 ᑎᑭᓕᓴᐃ 12.2.4 ᓴᓇᒥᓴᑎᑦ ᐊᒻᒪᓗ ᓄᓯᒋᐅᑕᐃᓕᓚᑎᓴᓕᑎᑦ ᓂᕆᔨᑎᑦ ᐊᒻᒪᓗ ᓂᕆᔨᑎᑦ ᐃᓂᖅᑎᑦ ᐊᑕᔨᑦ ᑕᒃᑯᓄᑦ ᐅᓂᒃᑲᓚᐅᑐᓄᖅ ᐃᑎᒪᓂᖅᑐᓂᖅ ᑎᑎᕋᐅᓚᐅᑐᖅ, ᐃᓕᑎᔭᐅᓚᑎᑐᖅ ᓂᕆᔨᑎᖃᖅᑐᑦ ᐊᓯᒪᓕᕆᐅᔨᑦ ᐃᓄᐃᑦ ᓄᓇᕗᑦᐸᖅᑎᑦ ᐃᓂᖅᑎᑦ.

12.2.6 ᑐᑭᓯᓇᑎᐊᑲᓂᓂᐅᐊᖅᓯᑦ, ᐅᔪᐊ ᐊᐅᓚᕆᓂᖅᑎᑎᒋᓂᖅᓯᑦ ᑲᑎᒪᔨᕋᓱᒃᑦ ᐅᓂᖅᑲᔨᓯᒪᓂᐊᖅᑐᖅᑎᑦ ᒥᓂᕐᒥᑲ ᑕᒪᐃᓂᓂᖅ ᓂᖅᑲᐃᓕᓕᒫᐊᓂᖅ ᑕᒪᑲᔪᓂᖅ ᐊᐅᓚᕆᑎᔨᐸᖅᑐᓂᖅᓯᑦ ᐊᒻᒪᓗ ᓂᕆᔨᑎᑦ ᐃᓂᖅᑐᓂᖅ ᐃᓗᖅᑐᓂᖅ ᑲᑎᒪᔨᑦ ᐅᓂᒃᑲᓚᐅᑐᓄᖅ ᐅᑎᖅᑕᕐᕕᒃᓴᓂᖅ ᑎᖅᑎᓕᕐᓂᖅ ᐅᑎᖅᑕᕐᕕᒃᓴᓂᖅ ᑎᖅᑎᓕᕐᓂᖅ ᐃᓗᖅᑐᓂᖅ ᐅᑎᖅᑕᕐᕕᒃᓴᓂᖅ ᑎᖅᑎᓕᕐᓂᖅ ᐃᓗᖅᑐᓂᖅ ᐅᑎᖅᑕᕐᕕᒃᓴᓂᖅ ᑎᖅᑎᓕᕐᓂᖅ ᐊᒻᒪᓗ ᑭᒃᑯᐊ ᐃᓗᐊᓂᑐᑦ ᑲᑎᒪᔨᑦ ᐅᓂᒃᑲᓚᐅᑐᓄᖅ ᐅᑎᖅᑕᕐᕕᒃᓴᓂᖅ ᑎᖅᑎᓕᕐᓂᖅ ᐃᓗᖅᑐᓂᖅ ᐅᑎᖅᑕᕐᕕᒃᓴᓂᖅ ᑎᖅᑎᓕᕐᓂᖅ ᑲᑎᒪᔨᑦ ᐃᓄᐃᑦ ᑭᒃᑯᐊ ᑲᑎᒪᔨᑦ ᐊᖏᔪᓂᖅᑐᑦ ᐊᐅᓚᕆᓴᖅᑎᑎᒋᓂᖅ ᐊᖏᔪᓂᖅᑐᑦ ᐃᓄᐃᑦ ᓄᓇᕗᑦᐸᖅᑎᑦ ᐊᐅᓚᕆᑎᔨᐸᖅᑐᓂᖅᓯᑦ

ᖃᐅᔨᓴᓂᐊᕐᓗᓂᑐᑎᒃ ᖃᐅᔨᓴᖅ ᐊᓯᖏᑐᓂᓂ ᐱᓚᕆᓐᖏᓐᓇᑦ ᑭᓇᐅᔭᖅᑕᕐᓂᑎᑦᑕᐅᔪᓐᓇᕐᓂᑐᑎᒃ. ᑕᐃᒪᓕᕐᓂ, ᒪᓕᒐᓴ ᖃᐅᔨᓴᓇᓂᖏᓐᓇᑦ, ᓄᓇᕗᑦ ᑐᓴᖏᓐᓇᐃᒪᒃ ᑎᑎᖏᓐ ᐱᔭᕆᖃᖅᓯᓯᒪᓂᖅᑐᑦ ᐃᒪᓐᓇ (i) ᐱᓕᕆᔨᐤᓂᓗᑎᒃ ᑭᓇᐅᔭᖅᑕᑎᑦᑕᐅᓇᔭᓂᖏᓐᓂᐤᓂᒃ ᐊᓯᒥᓐᓂᒃ ᓂᕆᐅᓇᐤᓯᑐᑐᖅ ᐊᒻᒪᓕᓗ (ii) ᐊᑭᓕᐅᑕᐅᔪᖅ ᐊᑭᓕᐅᑕᐅᔭᕆᕆᐊᓕᓕᒃ, ᑐᓴᖅᑎᑎᓂᐅᐊᖅᑐᖅ ᑕᒃᑯᓄᖓ ᑲᓇᑕᑦ ᐅᒪᔨᓕᕆᔨᓄᑦ ᐱᔭᕆᑎᖅᑎᑦ. ᐱᔪᑎᒃ ᓄᓇᕗᑦ ᑐᐤᖓᓇᖅ ᑎᑎᖓ ᑐᓴᖅᑎᑦᑎᐱᓗᒃᓴᓂᒃ ᐅᕐᓀᑑᐤᓂᑦ ᑐᓴᖅᑎᓪᓗᑦ ᓄᓇᕗᑦᓯᓄᑦ ᓄᓇᕗᓐᓄᑦ ᐊᖏᔪᓚᐅᑐᑦᓯᓄᑦ ᖃᐅᔨᖅᑐᑎᓂᒃ ᓄᓇᕗᓐᓄᑦ, ᑲᓇᑕᑦ ᐅᒪᔨᓕᕆᔨᓄᑦ ᐊᑭᓕᐅᑕᐅᔪᖅ ᑕᒃᑯᓄᖓ ᑲᓇᑕᑦ ᐅᒪᔨᓕᕆᔨᓄᑦ ᐊᑭᓕᐅᑕᐅᔪᖅ ᐊᑭᓕᐅᑕᐅᔭᕆᕆᐊᓕᓕᒃ ᐊᑭᓕᐅᑕᐅᔭᕆᕆᐊᓕᓕᒃ ᑕᒃᑯᓄᖓ ᑲᓇᑕᑦ ᐅᒪᔨᓕᕆᔨᓄᑦ ᐊᑭᓕᐅᑕᐅᔪᖅ

12.4 ᐃᑭᑕᐅᖏᑦᑕᓗᓂᒃ ᐊᑐᑎᖏᖅᑐᑦ ᓂᖅᐸᑎᐃᓐᓂᑦ

12.4.1 ᓇᓕᐊᑐᐃᓇᖅ ᐊᑐᑎᖏᖅᑐᑦ ᓂᖅᐸᑎᐃᓐᓂᐅᓕᐅᑦ ᑐᖅᑯᑕᐅᓯᒪᔪᖅ ᑐᐊᐃᖅᓇᖅᑐᒃᑯᑦ, ᐱᕆᔭᖅᓂᒃᑯᑦ, ᐅᕙᓪᓕᐊᓪᓚᑦ ᐱᖅᑭᓗᐊᖏᓂᒃᑯᑦ ᐃᑭᑕᐅᓂᐊᖏᖅᑐᒃ ᒪᑕᒃᓗᐊ 5.6.55 ᒎᒃᑯᓇᓂ *ᓄᓇᕗᒻᒥᒃ ᓄᓇᕈᑎᑎᓄᑦ ᐊᖏᕈᑎᑎᑦ.*

12.5 ᐅᖏᐅᓯᖏᓗᑎᑦ

12.5.1 ᐊᖏᕐᕌᒥ ᖃᑎᖏᑭᐊᖏᖅᑕᐅᔪᑦ ᐱᓕᕆᔨᐅᔪᑦ ᒪᓕᒃᓴᐊᓗ ᐃᓕᖓ 15.4 ᐅᓪᓚᓂ ᐃᓄᐃᑦ ᐊᒃᑐᖏᖅᑕᐅᓂᐊᖏᓂᖅᐳᓂᓪᓕᑦ ᐱᓪᓕᓕᓕᑕᐅᔪᓪᓚᕆᓄᑦᓗ ᐊᖏᕈᑎᑎᑦ,ᑲᓇᑕᒥ ᐅᒪᔪᓕᕆᔨᖏᔪᐊᑦ ᐅᖏᐅᓯᖏᓗᑎᑦ ᒎᒃᑯᓄᖓ ᑕᒪᐃᓪᓚᑦ ᐅᒃᑐᑕᐅᔪᑦ ᒪᓕᒃᓴᐊᓗ ᐅᓪᓚᓂ ᐃᓕᖓᓂ. ᑲᓇᑕᒥ ᐅᒪᔪᓕᕆᔨᖏᔪᐊᑦ ᐅᖏᖅᑮᖏᓗᑎᒃ ᐅᓂᒃᑲᓕᐅᖏᓐᓇᑎᒃ ᒎᒃᑯᓄᖓ ᐊᐅᓕᕐᑎᖅᑎᑦᓂᓪᓕᒪᑦ ᑲᑎᒪᔨᖏᓪᓕᐊᑦ, ᐊᒃᑐᑎᔪᑦ ᓄᓇᕗᒻᒥᒃ ᐊᓚᖅᑐᖅᑯᓕᕆᓂᖅᑐᒃᑯᑦ, ᐊᒃᑐᑎᔪᑦ ᐊᓚᒃᑐᖅᓯᒪᔪᓂ ᐅᒪᔪᓕᕆᔨᒃᑯᑦ, ᓄᓇᕗᒻᒥ ᐅᒪᔪᓕᕆᔨᖏᔪᒃᑯᑦ ᐊᓱᒪᓗ ᐊᓯᖏᕐᓂ ᑎᒥᐅᔪᑦ ᐱᖁᓇᖅᑐᖅᑯᑦ ᓂᖅᐸᑎᓄᑦ.

ᐃᓕᓐᓴ 13 — ᐊᓯᔾᔨᐅᑎᑦ ᑕᕆᔨᓴᓐᓴ ᐅᑎᖅᑕᕐᕕᑦᑐᓄᑦ ᑎᓐᓗᐊᓄᑦ ᓴᓱᐅᓯᒪᓂᖅ ᐊᒻᒪᓗ ᑲᓇᑕᒥ ᐅᒪᔾᔪᖅᓴᖅᑐᑦ

—

13.1 ᑐᑭᖅᕕᑦᓕᐊᔪᐊᑎᖅᓴᑦ

13.1.1 ᐅᓂᒃ ᐃᓕᓐᓄᓂ ᐊᒋᓂ ᑎᑎᕋᖅᓯᒪᔾᔪᑦ ᑐᑭᖅᕕᑦᓕᐊᔪᐊᑎᖅᓴᖅᑎᓐᓇᑦ:

(a) ᓴᕋᑭᑎᓗᑎᒃ ᐊᓯᔾᔨᕕᖏᓐᓄᖅᑐᒃ ᐊᒻᒪᓗ ᐊᑐᑎᖅᑭᑎᐊᖅᑐᒥᒃ ᐱᔪᔪᑎᖅᓴᓂᒃ ᖃᐅᔨᓴᓯᑎᐊᕋᓂᖅ ᓴᕋᑭᑎᓂᐊᕋᖅᑐᓄᑦ, ᐊᓐᓯᑕᒥᐊᕋᓂᐊᕋᖅᑐᓄᑦ ᐊᒻᒪᓗ ᖃᐅᔨᓴᓯᑎᐊᖅᑐᓄᑦ ᐊᓯᔾᔨᕕᓂᐊᕋᖅᑐᓄᑦ;

(b) ᑐᓂᓗᑕᑎ ᐃᓄᐃᑦ ᐊᒻᒪᓗ ᑌᓐᒪᒃᑯᑦ ᑐᓴᐅᒪᔭᕆᐊᕋᖅᑕᕐᓯᓂᒃ ᐱᔪᔪᑎᖅᓴᖅᑎᓐᓇᑦ ᐱᑕᖅᑲᖅᑐᑦ ᑐᐅᔪᓕᕐᓂᐊᕋᖅᑐᓄᑦ, ᐊᓐᓯᑕᒥᐊᕋᓂᐊᕋᖅᑐᓄᑦ ᐊᒻᒪᓗ ᖃᐅᔨᓴᓯᑎᐊᖅᑐᓄᑦ ᐊᓯᔾᔨᓴᓂᐊᕋᖅᑐᓄᑦ, ᐊᓐᓯᑕᒥᐊᕋᓂᐊᕋᖅᑐᓄᑦ ᐊᒻᒪᓗ ᖃᐅᔨᓴᓯᑎᐊᖅᑐᓄᑦ ᐊᓯᔾᔨᓴᓂᐊᕋᖅᑐᓄᑦ ᐃᓯᒪᓕᐊᓕᐊᑲᖅᑐᑦ;

(c) ᐊᑐᖅᑕᐅᑎᑦᑎᓗᑎᒃ ᐅᑯᐊ ᓴᕋᑭᑎᓂᐊᕋᖅᑐᓄᑦ, ᐊᓐᓯᑕᒥᐊᕋᓂᐊᕋᖅᑐᓄᑦ, ᖃᐅᔨᓴᓯᑎᐊᖅᑐᓄᑦ ᐊᓯᔾᔨᓴᓂᐊᕋᖅᑐᓄᑦ, ᐊᑎᕆᔨᓯᐊᕋᖅᑐᑦ ᐊᒻᒪᓗ ᐋᖅᑕᐅᔪᑦ ᐱᓕᓓᓴᐅᔭᓐᓗᑎᒃ ᑭᓯᐊᓂ ᖃᑎᖅᑭᐊᖅᑎᑕᐅᓴᓗᑎᒃ ᐊᒻᒪᓗ, ᐊᔪᖑᓐᓂᑐᐊᖅᓐᓇᑦ, ᐃᑲᔪᖅᑕᐅᓗᑎᒃ ᖃᐅᓕᑐᓂ ᐊᑕᔪᓂ ᓄᓇᓕᒻᒥ, ᐊᒻᒪᓗ ᐃᓄᐃᑦ ᐱᔪᒪᓕᓗᑕᑦ ᐊᒻᒪᓗ ᐱᔪᑕᐅᔪᓄᑦ ᐃᓯᖅᑕᐅᔪᓴᐅᓕᖅᓗᑎᒃ ᐃᓄᐃᑦ ᓄᓇᔪᑎᖏᑦ ᐃᓯᒪᓕᐊᓕᐊᑲᖅᓯᒪᓕᖅᑎᓗᑎᒃ ᐅᑯᐊ ᐃᓐᓯᕐᓂᐊᕋᖅᑐᓄᑦ;

(d) ᐊᑐᖅᑕᐅᑎᑦᑎᓗᑎᒃ ᐅᑯᐊ ᐊᖅᑐᖅᑕᐅᔪᑦ ᐊᒻᒪᓗ ᐃᑲᔫᓯᔪᖅᑎᓗᑎᒃ ᐃᓄᐃᑦ ᐅᖅᑲᓯᐅᓗᑎᒃ ᑕᐃᑲᓂ ᓴᕋᑭᑎᓂᐊᕋᖅᑐᓄᑦ, ᐊᓐᓯᑕᒥᐊᕋᓂᐊᕋᖅᑐᓄᑦ, ᖃᐅᔨᓴᓯᑎᐊᖅᑐᓄᑦ ᐊᓯᔾᔨᓴᓂᐊᕋᖅᑐᓄᑦ, ᐊᑐᖅᑕᐅᔪᑦ ᐊᒻᒪᓗ ᐋᖅᑕᐅᔪᑦ ᐅᑎᖅᑕᕐᕕᑦᑐᓄᑦ ᓄᓇᔪᑎᖏᑦ ᐊᒻᒪᓗ ᐃᓕᓴᕐᕕᖅᑎᑦ; ᐊᒻᒪᓗ

(e) ᑐᓂᓗᑕᑎ ᐊᑎᑕᐅᔪᑦᑎᓗᑎᒃ ᐊᒻᒪᓗ ᐋᖅᑕᐅᔪᓗᑎᒃ ᐅᑎᖅᑕᕐᕕᑦᑐᓄᑦ ᓄᓇᔪᑎᖏᑦ ᓴᕋᑭᑎᓂᐊᕋᖅᑐᓄᑦ ᐊᒻᒪᓗ ᐃᓕᓴᕐᕕᖅᑎᑦ ᐊᒻᒪᓗ ᐊᐅᓴᖅᑎᑦᑎᓗᑎᒃ ᑕᐅᖅᓯᓗᑎᒃ ᐃᓄᐃᑦ ᓄᓇᔪᑎᖏᑦ ᒪᓄᓇᖅᑐᒃ ᐃᓗᐊᓂᕐᑐᑦ ᐅᑎᖅᑕᕐᕕᑦᑐᓄᑦ ᓄᓇᔪᑎᖏᑦ ᓴᕋᑭᑎᓂᐊᕋᖅᑐᓄᑦ ᐊᒻᒪᓗ ᐃᓕᓴᕐᕕᖅᑎᑦ.

13.2 ᑐᑭᓯᒪᓂ

13.2.1 ᐅᓂᒃ ᐃᓕᓐᓄᓂ ᐊᒋᓂ ᑎᑎᕋᖅᓯᒪᔾᔪᑦ ᑐᑭᖅᕕᑦᓕᐊᔪᐊᑎᖅᓴᖅᑎᓐᓇᑦ:

(a) “ᖃᐅᔨᓴᓯᑎᐊᖅᑎᓂ ᑲᑎᒪᔨᓂ” ᑐᑭᖃᖅᑐᖅ ᑲᑎᒪᔨᓂ ᑌᓐᒪᒃᑯᑦ ᐃᓄᐃᑦ ᐱᔪᓐᓂᒃ ᑭᑑᑎᓂᖅᑐᑦ ᓴᕋᑭᑎᓂᐊᕋᖅᑐᓄᑦ ᒪᓕᓕᖅᑐᑦ ᓴᕋᑭᑎᓂᐊᕋᖅᑐᓄᑦ ᐊᒻᒪᓗ ᐃᓄᐃᑦ 13.4;

(b) ”ᐋᖅᑕᐅᔪᑦ” ᑐᑭᖃᖅᑐᖅ ᐋᖅᑕᐅᔪᑦ ᐊᓐᓯᑕᒥᓂ ᑲᑎᒪᔨᓂ ᐅᖅᑲᓯᐅᓗᑎᒃ ᐅᑎᖅᑕᕐᕕᑦᑐᓄᑦ ᓄᓇᔪᑎᖏᑦ ᐊᒻᒪᓗ ᐃᓕᓴᕐᕕᖅᑎᑦ;

(c) “ᓄᓇᕗᓂᖓ 1 ᖃᐅᔨᓴᓯᑎᐊᖅᑕᐅᔪᑦ” ᑐᑭᖃᖅᑐᖅ ᐱᔪᓐᓂᒃ ᖃᐅᔨᓴᐅᓯᑎᐊᖅᓗᑎᒃ ᓴᕋᑭᑎᓂᐊᕋᖅᑐᓄᑦ ᐊᒻᒪᓗ ᖃᐅᔨᓴᓯᑎᐊᖅᑐᒥᒃ ᐃᓄᐃᑦ ᐱᔪᓐᓂᒃ ᐃᓗᐊᓂᕐᑐᑦ;

(d) “ᓄᓇᕗᑦ 2 ᖃᐅᔨᓴᓂᐊᕐᕕᒃᓴᖅ” ᑐᓂᖅᑐᒃ ᐱᓕᕆᔨᒃ ᖃᐅᔨᓴᖅᑎᑦᓂᐊᖅᓯᒃᑐᑦ, ᒪᓕᒐᔪᒍ ᓄᓇᕗᑦ 1 ᖃᐅᔨᓴᓂᐊᕐᕕᒃᓴᖅ ᐊᓚᒪᓗ ᐃᓕᓴᕐᔨᓐᓂᑎᒃ ᓄᓇᒥ ᐃᓕᓴᕆᓴᓂᐊᖅᑐᑦ ᐊᓚᒪᓗ ᖃᐅᔨᓴᖅᑎᑦᓂᐊᖅᓯᒃᑐᑦ;

(e) “ᒥᖅᑕᑎᓪᓗᒍ” ᑐᓂᖅᑐᒃᑐᒃ ᓇᓕᐊᑐᐃᓇᖅ ᐊᓂᔨᖅᑐᑦ ᑭᖅᓕᕐᓂ ᐃᓗᕐᓂᑎᓇ ᑲᑎᒪᔨᖅ ᐅᓇᓕᓗ ᐊᑐᐃᑦᓯᒃᑐᑦ ᒥᖅᑕᑎᓪᓗᓂ ᐃᓗᕐᓂᑎᓇ ᑲᑎᒪᔨᖅ ᐅᓇᓕᓗ ᐊᑐᐃᑦᓯᒃᑐᑦ ᑲᑎᒪᔨᓂᖅᓯᒃ ᐅᓇᓕᓗ ᐊᑐᐃᑦᓯᒃᑐᑦ ᐃᓗᕐᓂᑎᓇ ᑲᑎᒪᔨᖅ ᐅᓇᓕᓗ ᐊᑐᐃᑦᓯᒃᑐᑦ ᐃᓗᕐᓂᑎᓇ ᑲᑎᒪᔨᖅ;

(f) “ᐱᓕᕆᔨᒃᑲ ᖃᐅᔨᓴᐅᔨᓴᓂᐊᖅᓯᒃᑐᑦ” ᑐᓂᖅᑐᒃᑐᒃ ᖃᐅᔨᓴᓂᐊᕐᓂᖅᓯᒃ ᐱᓕᕆᔨᒃᑲᖅᓯᒃ ᐱᓕᕐᓇᖅᑐᒃᑐᑦ ᓄᓇᕗᑦ, ᑕᐅᑦᑐᖅᓯᒃ ᖃᐅᔨᓴᐅᔨᓴᓂᐊᕐᓂᖅᓯᒃ ᖃᐅᔨᓴᖅᓯᐅᓪᓚᑐᑦ ᑕᕆᔨᒃᑲᖅᓯᒃ ᖃᐅᔨᓴᓂᐊᖅᑎᑦᓯᒃ ᐃᓚᖓᑕᐅᓪᓚᑐᑦ ᐊᓚᒪᓗ ᖃᐅᔨᓴᖅᑎᑦᓂᐊᖅᓯᒃᑐᑦ ᐃᓗᕐᓂᑎᓇ ᑲᑎᒪᔨᖅ ᑕᕆᔨᒃᑲᖅᓯᒃ ᐊᓚᒪᓗ ᐃᓗᕐᓂᑎᓇ ᑲᑎᒪᔨᖅ ᐃᓗᕐᓂᑎᓇ ᑲᑎᒪᔨᖅ ᑕᕆᔨᒃᑲᖅᓯᒃ ᐊᓚᒪᓗ ᐃᓗᕐᓂᑎᓇ ᑲᑎᒪᔨᖅ

ᑲᓇᒪᒃᑯᑦ ᑎᒥᖅᑯᓂᒋᑦ ᐱᓕᕆᔨᒪᔨᑦ ᑕᒪᔭᒥᖏᓪ ᐱᔪᐊᑎᒃᖁᑦ ᖃᐅᔨᖏᑦᐅᓇᓂᕆᐊᖅᓯᒪᑎᒃ ᑭᒃᑯᓂᓪ ᐱᖃᑦᒪᔨᓂᑦ ᓴᖅᑭᑎᓂᐊᖅᓯᒪᓂᖅ, ᐊᓂᒍᓕᕆᓂᐊᖅᓯᒪᓂᖅ, ᐅᖃᓕᓴᐅᓂᒃ ᖃᓄᐃᓇᓕᒫᓂᖏᓪ ᐊᑐᕐᓴᖅᑐᑦ.

13.4.2 ᒪᓕᒪᓐᓇᐃ ᖃᐅᔨᓴᖅᑎᑎᕆᔭᐅᓪᓚᑦ 13.4.1, ᑲᓇᑕᒥ ᐅᒪᔨᓕᓕᖅᓴᔨᐊᑯᑦ ᐱᖓᓱᖅᑯᓂᐊᖅᑐᑦ ᐊᓚᒪᓂ ᐃᓕᓴᕐᕕᑕᐅᓂᒃ ᖃᐅᔨᖏᑦᐅᓇᓂᕆᐊᖅᑎᑦ ᑲᑎᒪᔨᑦ. ᐊᑐᐅ ᖃᐅᔨᖏᑦᐅᓇᓂᕆᐊᖅᑎᑦ ᑲᑎᒪᔨᑦ ᐱᖓᓱᖅᑯᓂᐊᖅᑐᑦ ᖃᒪᐃᓕᐅᑎᒃ ᐱᔪᐊᑎᒃᖁᑦ ᖃᐅᔨᖏᑦᐅᓇᓂᕆᐊᖅᓯᒪᑎᒃ ᐱᔪᐊᑎᒃᖁᑦ, ᐊᓂᒍᓕᕆᓂᐊᖅᓯᒪᓂᖅ, ᐅᖃᓕᓴᐅᓂᒃ ᖃᐅᔨᖏᑦᐅᓇᓂᕆᐊᖅᓯᒪᓂᖅ ᐊᑐᕐᓴᖅᑐᑦ. ᐊᑐᐅ ᖃᐅᔨᓴᖅᑎᑎᕆᔭᐅᓪᓚᑦ ᑲᑎᒪᔨᑦ ᐃᒪᐃᓐᓇᐅᑎᑦᑐᑦ, ᐃᓚᖏᑕᐅᓪᓚᖅ:

(a) ᖃᐅᔨᓴᖅᑎᑎᕆᔭᐅᓪᓚᑦ ᓄᓇᖓᓂ ᖃᓄᐃᓇᓕᒫᓂᖓ ᓇᓪᓕᐊᑐᐃᓐᓇᖅ ᐱᔪᐊᑎᒃᖁᑦ ᖃᐅᔨᖏᑦᐅᓇᓂᕆᐊᖅᑐᑦ;

(b) ᖃᐅᔨᓴᖅᑎᑎᕆᔭᐅᓪᓚᑕᑦ ᖃᓄᐃᓕᑎᒋᔭᐅᑦ ᐊᖏᑕᓂᖏᑦ ᖃᐅᔨᔪᖅᑕᐅᔪᑦ ᓇᓪᓕᐊᑐᐃᓐᓇᖅ ᐱᔪᐊᑎᒃᖁᑦ ᖃᐅᔨᖏᑦᐅᓇᓂᕆᐊᖅᓯᒪᔪᑦ ᑕᐃᒪᐃᓐᓇᐅᓇᓕᒋᐊᑦ, ᐃᓕᑎᒃᐅᓪᓚᑕᑦ ᑕᒪᕐᒥ ᓴᓪᕿᓂᖓᓐᓂ 1 ᐊᓚᒪᓚᑦ ᓴᓪᕿᓂᖓᓐᓂ 2 ᖃᐅᔨᖏᑦᐅᓇᓂᕆᐊᖅᑦᑎᓪᓗᐊᓂᖓᓐᓂᑦ, ᐊᓚᒪᓚᑦ ᖃᐅᓄᖅ ᑭᓇᐅᔨᓪᓗᐅᑕᐅᔪᓐᓇᓐᓂᖓᓂᒃ ᐅᖃᓕᓴᐅᓂᒃ ᐃᓐᓇᓴᓪᓚᑦ - ᑭᓇᐅᔨᑎᒍᑦ ᐱᓕᕆᓪᓚᖅᓯᒪᓂᖅᑐᑦ ᖃᐅᔨᔪᖅᑕᐅᓚᑐᑦ ᑕᐃᒪᐃᓐᓇᐅᔨᓚᕆᐊᖃᑎᒃ ᖃᐅᔨᖏᑦᐅᓇᓂᕆᐊᖅᓯᒪᓂᖅ ᖃᐅᔨᖏᑦᐅᓇᓂᕆᐊᖅᓯᒪᖅᓯᖅ;

(c) ᐱᓕᕆᓪᓚᐊᑎᑕᐅᓚᑐᑦ ᑎᑎᕋᖅᑕᐅᓚᑐᑦ ᔭᓇᓂᓚᓂᐊᖅᑕᖏᓐᓄᓚᑦ ᑮᓇᐅᑎᑦ ᐱᓕᕆᔨᑦ ᐱᔪᓱᖅᑕᐅᔪᒍᑎᒃ ᓇᓪᓕᐊᑐᐃᓐᓇᖅ ᐱᔪᐊᑎᒃᖁᑦ ᖃᐅᔨᖏᑦᐅᓇᓂᕆᐊᖅᓯᒪᖅᓯᖅ;

(d) ᓂᕈᐊᕐᓗᑎᑦ ᒪᓕᒪᓐᓇᐃ ᑎᒥᖅᑯᓂᒋᑦ ᐅᖃᓕᓴᐅᓂᒃ ᑲᓐᓂᑲᖅᑕᐅᓚᑐᑦ ᐱᑕᓚᐅᓚᑐᑦ ᑕᒪᕐᒥ ᓴᓪᕿᓂᖓᓐᓂ 1 ᐊᓚᒪᓚᑦ ᓴᓪᕿᓂᖓᓐᓂ 2 ᖃᐅᔨᓴᖅᑎᑎᕆᔭᐅᓪᓚᑦ ᑕᐃᒪᐃᓐᓇᐅᔨᓚᕆᐊᖃᑎᒃ ᐱᔪᐊᑎᒃᖁᑦ, ᐃᓕᑎᒃᐅᓪᓚᑕᑦ ᐃᓅᖏᓪᓕᐊᓂᕆᐊᖅᓯᒪᓂᖅ ᐃᓅᓴᓂᖅᑎᑎᕆᔭᐅᓪᓚᑦ ᐊᓚᒪᓚᑦ ᐱᔪᐊᑎᒃᖁᑦ ᐃᓅᓴᓂᖅᑎᑎᕆᔭᐅᓪᓚᑦ ᐊᓚᒪᓚᑦ ᖃᐅᔨᖏᑦᐅᓇᓂᕆᐊᖅᓯᒪᓂᖅ ᐃᓅᓴᓂᖅᑎᑎᕆᔭᐅᓪᓚᑦ ᐱᔪᐊᑎᒃᖁᑦ ᓄᓇᓕᖅᑐᓂ ᓄᓇᓕᖅᑐᓂ ᐱᓕᕆᔨᑦ ᐃᓅᓴᓂᖅᑎᑎᕆᔭᐅᓪᓚᑦ ᐱᔪᐊᑎᒃᖁ

13.4.4 ᑕᐃᒪᓗ ᑲᓇᑕᒥ ᐅᒪᔨᓕᓚᖅᓴᔨᐊᓂᕐᑯᑦ ᐃᓱᒪᓕᐅᖅᑕᑦ ᐃᓕᐅᕕᖏᐅᑉᓯᓚᑎᒃ ᑕᒪᑐᒪᑦ ᐱᔨᕈᑎᒃᓴᖅ ᖃᐅᔨᔪᐅᓇᓱᓕᕆᔨᑎᑦ ᓴᖅᑭᑎᓂᐊᕐᓯᖅᑐᖅ ᓄᓇᖕᓂᓄ ᑐᓴᓱᖅᑕᐅᓱᒪᔨᓄᑦ ᓴᓇᑭᑎᓂᕐᓯᖅᑐᖅᑐᑦ, ᐊᖏᕐᓂᓕᕆᐊᓯᐊᓯᖅᑐᑦ ᐅᖏᖓᔪᓂᑦ ᖃᓄᐃᓕᓖᖏᓂᕐᓂᑦ ᐊᓱᓕᔪᖅᑐᑦ, ᑲᔨᓱᒃᖏᓂᑐᖅ ᑐᓴᓱᖅᑕᓂᖓ ᒡᒃᑯᓄᖓ ᓄᓇᕗᒻᒥ ᐅᒪᔨᓕᓚᖅᓴᔨᐊᓂᕐᑯᓄᑦ.

13.4.5 ᐃᓄᐃᑦ ᐊᑐᖅᑕᐅᓂᐊᓯᓂᖏᓐᓄᑦ ᐱᖓᓕᓕᐊᔨᓂᕐᓚᓄᑦ ᐊᖏᕐᕈᐃᑦ ᐊᑐᓕᖅᑎᑕᐅᖓᓕᓕᐊᔨᓄᑦ ᑭᓇᐅᔭᖅᓴᐃᑦ ᐃᒪᐧᐃᓇᐅᓂᐊᖏᓐᓂᑐᑦ, ᑭᓱᐊᓂ ᐊᖏᖅᑕᐅᔪᒪᔪᑎᒃ ᒡᒃᑯᓇᖕᓂᒃ ᑭᓇᐅᔭᖅᓴᓂᒃ ᐊᑭᖏᓐᓄᑦ ᓀᓕᓕᐊᔨᓂᕐᓚᓄᑦ ᐃᓕᖅᑐᓂ ᐱᖓᓱᒃᑐᖅ ᐊᑐᖅᑕᓂᖓ ᐊᑐᖅᑕᐅᓂᐊᖏᓐᓂᑐᑦ ᐱᖓᓱᒃᑐᖅ ᐊᓕᐅᓂᐊᖏᓐᓂᑐᑦ ᐊᑐᖅᑕᐅᓂᐊᓯᓯᖅᑐᑦ.

13.4.6 ᐃᓕᖏᓐ 13.4 ᐊᑐᑎᓴᓇᖏᑐᑦ ᓂᐅᖏᓗᓂ, ᖃᖅᑯᓕᓪᓗᐃᑦ ᐅᖓᓚᓇᓂ ᐊᑐᐸᐃᑦ ᑲᓇᑕᒥ ᐃᓕᔪᖅᑯᓴᓕᑦ.

13.5 ᐋᓴᖅᑕᐅᓂᖏᓂᑦ ᐊᒻᒪᓗᓪ ᒥᖏᓪᓗᐅᐊᖅᑕᐅᓂᖏᓂᑦ

13.5.1 ᑕᐃᒪᓗᓪ ᓄᓇᕗᑦ ᑐᑎᖏᖓ ᐅᖓᓚᓇᓂᓂᑦ ᐊᑐᑎᖅᑯᖅᑐᑦ ᐊᕐᓚᑲᖅᓯᒪᔪᑦ ᐃᓄᐃᑦ ᑲᑐᔪᖅᑎᑦᓯᖏᓂᑦ ᐅᑯᐱᓯᓴᑐᑦ ᐅᑯᐊᑦ ᑕᒪᖅᑎᒃ ᐅᖓᓚᓇᓂᓂᑦ ᐃᓕᖏᓂᑦ ᒡᒃᑯᓇᖕᓂᒃ ᐅᑎᖅᑕᐅᓂᐊᖏᓐᓂᑐᑦ ᐃᓄᐃᑦ ᓄᓇᕗᑎᖏᓂᑦ ᐃᑐᐊᑦ, ᐃᑲᔪᑎᑐᐊᖏᓐᓂᒃᑎᑐᑦ ᐅᑐᖅᑐᐃᑐᑐᓂᑦ ᐃᓕᖏᓂᑦ ᐊᒻᒪᓗᓪ ᐃᓄᐃᑦ ᑲᑐᔪᖅᑎᑦᓯᖏᓂᑦ ᐃᓄᐃᑦ ᑲᑎᒪᔨᖅ ᐊᑐᖅᑕᓂᖓ ᐅᑎᖅᑕᐅᓂᐊᖏᓐᓂᑐᑦ ᑎᒥᐅᑦ ᐊᑐᖅᑕᐅᓂᐊᓯᓯᖅᑐᑦ ᐃᓄᐃ

13.5.5 ⊃Pᓯᓇᑦᑎᐊᖏᓐᓂᕐᐊᖅᓯᑦ, ᐋᓴᖅᑕᐅᓂᖓᓗᓄᑦ ᐅᕐᓂᐅᓂᖓᓗᓄᑦ ᐱᓕᕆᔨᖅᑕᐅᓂᖓᓗᓄᑦ ᐋᓴᐃᖅᐸᑕ ᒪᑦᔨᑐᒃᑯᐊᑎᒃ ᐅᒥᖓ ᐃᓕᖓ ᐊᔪᖅᑎᑎᓂᐊᕐᓂᑦᑐᑦ ᑲᑎᒪᔨᖅᑕᐅᓂᖓᓗᓄᑦ ᑲᖏᖅᑎᖏᓐᓂᐊᕐᓂᖅᓴᖅ ᐱᖓᓱᖅᑕᐅᓂᖓᓗᓄᑦ ᐊᔪᖅᑎᑎᓂᐊᕐᓂᖅᓴᖅ ᐱᖓᓱᖅᑕᐅᓂᖓᓗᓄᑦ ᐱᐅᓯᖅᑕᐅᓂᖓᓗᓄᑦ ᐊᔪᖅᑎᑎᓂᐊᕐᓂᖅᓴᖅ ᐱᖓᓱᖅᑕᐅᓂᖓᓗᓄᑦ ᐱᐅᓯ

ᐃᓕᓐᓴ 14 — ᓴᓪᓕᖅᑐᑦ ᐃᓱᑭᓐᓇᖅᑕᖏᑦ ᐊᓚᒐᓂ ᐊᑐᑭᓐᓇᖅᑕᖏᑦ

14.1 ᑐᑭᒋᖅᓯᒃᓕᐊᔨᐱᑎᒃᓴᑦ

14.1.1 ᐅᓂᒃ ᐃᓕᓐᓴᓄᒃ ᐊᒋᒃ ᑎᑎᕆᖅᓱᓕᔪᑦ ᑐᑭᒋᖅᓯᒃᓕᐊᔨᐱᑎᒃᓴᖏᑦ:

(a) ᓴᓪᓕᖅᑎᑦ ᐊᑐᑭᓐᓇᖅᑕᖏᑦ ᒪᓕᒃᑎᐊᕐᓗᑎᒃ ᐃᓇᐃᑦ ᐊᑐᑭᓐᓇᖅᑕᖏᓐᓂᒃ ᐊᓚᒐᓂ ᑲᖏᐅᓴᐊᕆᓇᖅᑐᓂᓯᖅᑕᖏᑦ ᑲᓇᖅᑕᒥ ᐅᒪᔨᑲᖅᓴᐃᓐ ᐅᖃᓕᒫᒃᑯᑦ ᐅᓂᖅᑕᐸᕐᓗᓇᑦ ᑎᓐᓕᐊᓇᑦ ᓴᓵᖏᐅᓴᐊᕆᓚᐅᒃ;

(b) ᑕᐃᑲᓂᒃ ᐊᑐᑎᖅᑲᖅᑎᑦᓗᔪ, ᐱᖏᓕᐊᑭᓚᓕᓗᑎᑦ ᐊᐅᑦᓪᓗᔭᓂᐅᑐᑦ ᐃᓇᓐᓂᑦ;

(c) ᑐᓴᕐᑎᓗᑎᑦ ᓴᓪᓕᖅᑎᑦ ᐃᓇᐃᑦ ᐱᔪᓇᕐᓂᖅᓂᓂᒃ, ᐊᑐᑭᓐᓇᖅᑕᖏᓂᒃ ᐊᓚᒐᓂ ᑲᖏᐅᓴᐊᕆᓚᐅᑎᑦᓗᑎᑦ ᑲᑎᒪᓂᖅᑲᖅᑎᑦ ᐅᒪᔨᑲᖅᓴᐃᓐ ᐅᖃᓕᒫᒃᑯᑦ ᑎᓐᓕᐊᓇᑦ ᓴᓵᖏᐅᓴᐊᕆᓚᐅᒃ; ᐊᓚᒐᓂ

(d) ᐊᐅᐅᓐᖓᓕᓴᐅᓇᖅᑕᐃᓕᓚᓗᑎᒃ ᐊᓚᒐᓂ ᒥᑭᒃᑎᑦᓗᑎᑦ ᐊᐅᐅᓐᖓᓕᓴᐅᓇᔪᑦ ᐊᐅᒋᓂᑦ >ᓕᕆᑎᑦ ᐊᓚᒐᓂ ᓂᕆᔪᑎᑦ.

14.2 ᐃᓇᐃᑦ ᐊᑐᑭᓐᓇᖅᑕᖏᑦ ᐊᓚᒐᓂ ᐅᖃᐅᓕᐊᑎᖅᓱᓕᓗᑎᑦ

14.2.1 ᐅᑯᐊ ᐊᐅᓪᓕᕆᑎᖅᑎᑯᓂᖅᓴᓪᓕᔭᑦ ᐅᑎᓚᓕᕆᓪᑦ ᐅᖃᐅᓕᓴᓂᐅᔪᖅᑐᑦ ᒥᓂᔾᔪᑎᒥᒃ, ᐊᓚᒐᓂ ᐊᓱᖏᓐᓂᓂᒃ ᒥᓂᔾᔪᑎᓂᒃ ᐊᓚᒐᓂ ᐊᐅᓪᓕᔪᓱᓂᒃ ᐅᓐᓴᒪᔪᑦ ᑕᐅᒃᑯᓂᖓᓐᖓ ᐅᓪᓕᕆᑎᖅᑎᑯᓂᖅᓴᓪᓕᔭᑦ ᐅᖃᐅᓕᓴᓂᐅᔪᖅᑐᑦ ᐃᓇᐃᑦ ᐅᒪᔨᑲᖅᓴᐃᓐ ᐅᓂᖅᑕᐸᕐᓗᓇᑦ ᑎᓐᓕᐊᓇᑦ ᓴᓵᖏᐅᓴᐊᕆᓚᐅᒃ, ᐃᓕᒥᓴᐅᓕᓗᑎᒃ ᖃᐅᔨᑎᑎᖅᑎᓂᖅᓴᖏᑦ, ᓄᖅᑎᕕᓕᒥ ᐊᓚᒐᓂ ᒪᓕᒥᐊᖅᑲᖅᑐᑦ ᓴᓪᓕᖅᑎᑦ ᓴᓵᖏᐅᓴᐊᕆᓚᐅᒃ ᐊᓚᒐᓂ ᐊᓱᐅᑦᑎᐊᓕᐊᔪᓐᓇᒋᑦ ᒪᓕᒥᐊᖅᑲᖅᑐᑦ ᓴᓪᓕᖅᑎᑦ ᐊᑐᑭᓐᓇᖅᑕᖏᑦ ᐊᓚᒐᓂ ᐅᖃᐅᓕᓴᓂᐅᔪᖅᑐᑦ ᐃᓇᐃᑦ ᐊᑐᑭᓐᓇᖅᑕᖏᓐᓂᒃ ᐊᓚᒐᓂ ᐅᖃᐅᓕᓴᓂᐅᔪᖅᑐᑦ ᐃᓕᓴᕐᓂᐊᓕᖅᑐᑦ ᐃᓇᐃᑦ ᐅᖃᐅᓕᓴᓂᐅᔪᖅᑐᑦ ᐊᓚᒐᓂ ᑎᓐᓕᐊᓇᑦ ᐅᖃᐅᓕᓴᓂᐅᔪᖅᑐᑦ ᐊᓚᒐᓂ ᑎᓐᓕᐊᓇᑦ ᐅᑭᐅᖅᑕᖅᑐᑦ. ᑕᐃᑲᓂᒃ ᐅᓇᓕᒃ ᐃᓕᓴᕐᓂᐊᓕᖅᑐᑦ ᐅᐊᖅᑐᑦᓗᑎᑦ ᐊᓚᒐᓂ ᑎᓐᓕᐊᓇᑦ ᐅᖃᐅᓕᓴᓂᐅᔪᖅᑐᑦ ᐃᓇᐃᑦ ᐊᑐᑭᓐᓇᖅᑕᖏᓐᓂᒃ ᐊᓚᒐᓂ ᑎᓐᓕᐊᓇᑦ ᐅᑭᐅᖅᑕᖅᑐᑦ ᐃᓇᐃᑦ ᐅᖃᐅᓕᓴᓂᐅᔪᖅᑐᑦ ᐊᓚᒐᓂ ᑎᓐᓕᐊᓇᑦ ᐅᑭᐅᖅᑕᖅᑐᑦ ᐃᓕᓴᕐᓂᐊᓕᖅᑐᑦ ᐃᓇᐃᑦ ᐅᖃᐅᓕᓴᓂᐅᔪᖅᑐᑦ ᐊᓚᒐᓂ ᑎᓐᓕᐊᓇᑦ ᐅᑭᐅᖅᑕᖅᑐᑦ ᐃᓇᐃᑦ ᓄᖅᑎᕕᓕᒥ ᐅᑭᐅᖅᑕᖅᑐᑦ ᐃᓇᐃᑦ ᐅᖃᐅᓕᓴᓂᐅᔪᖅᑐᑦ ᐊᓚᒐᓂ ᑎᓐᓕᐊᓇᑦ

(d) ᖃᑉᓱᓕᕆᔨᒃᑲᓪᓚᕆᐊᓕᖅ ᐊᕐᓇᖅᑐᑦ ᐊᑐᖅᑕᐅᓂᖅ ᐊᑐᖅᑕᐅᓂᖅ ᐊᑐᖅᑕᐅᓂᖅ ᐊᕐᓇᖅᑐᑦ ᐊᑐᖅᑕᐅᓂᖅ ᐊᑐᖅᑕᐅᓂᖅ ᐊᑐᖅᑕᐅᓂᖅ

14.4 ᐊᑐᓕᖁᐅᔾᔪᑎ ᐅᕐᓂᒍᓐᓂᑎ ᑕᐃᒪᐢᓇᐅᑦᓴᕆᐊᖅᑐᖅᑐᑎ ᐊᐅᑦᓭᓐᔨᓕᕆᐅᔾᔪᓂᒃ ᓄᓇᒃ

14.4.1 ᐊᐅᓭᓯᐊᕐᑲᓇᑦ <ᖁᓇᐅᑎᑦ ᓇᓗᓇᐃᖅᓯᒪᓂᐊᖅᑐᑎ, ᐊᑐᑎᖅᑭᑎᒃ:

(a) ᓄᓇᐃᑦ ᐊᓪᒪᓗ/ᐅᕐᓂᒍᓐᓂᑎ ᖃᓄᓇᐃᓕᖅᑲᓪᓚᔪᑦ ᑕᐃᒃᑐᓂ ᐊᑐᓕᖁᐅᔾᔪᓕᖅᑎᑦ ᐅᕐᓂᒍᓐᓂᑎ ᑕᐃᒪᐢᓇᐅᑦᓴᕆᐊᖅᑐᖅᑐᑎ ᐅᑯᐊ >ᓪᓚᖅᑐᑦ ᐱᖅᑎᖅᖁᓪᓗᑎᒃ ᐊᐅᓭᓯᐊᕐᑲᓇᖅᑐᒃᓴᓕᐅᔾᔪᓂᒃ ᐃᓄᖅᓂᒃ; ᐊᓪᒪᓗ

(b) ᐊᐅᓭᓯᐊᕐᑲᓇᑦ ᐊᔪᖅᑎᑕᓚᐊᖅᑲᖅᑕᖅᑎᑎ ᒎᒃᑯᓄᖅᓱ ᐊᐅᓭᓯᐊᕐᑲᓇᖅᑐᒃᓴᓕᐅᔾᔪᓂᓄᑦ, ᐃᓕᑎᕆᐅᓭᓗᑎᒃ ᐊᐅᓭᓯᐊᕐᑲᓇᑦ ᐱᓪᓗᒍ ᐊᑭᑎᓴᑭᐅᓇᖅᑐᑎ ᐊᐅᓭᓯᐊᕐᑲᓇᖅᑐᒃᓴᓕᐅᔾᔪᓂᒃ <ᖁᓇᐅᑎᑦ ᒪᑕᕐᓗᒍ 14.4.1(a).

14.4.3 ᑕᐃᒪᓗ, ᐊᖏᑦᖃᐅᔾᔨᒪᔪᓂ ᐊᐅᓭᓯᐊᕐᑲᓇᖅᑐᒃᓴᓕᐅᔾᔪᓂᑎ <ᖁᓇᐅᑎᑦ, ᐅᓄᐊ ᐊᐅᓭᓯᐊᖅᑎᑦᑎᒍᓐᓇᖅᑐᑎ ᑲᑎᕋᖅᑕᐅᔾᔨᒪᔪᓂᑎ ᑐᓴᕐᓯᒪᓂᒃ ᓄᓇᕗᑦ ᐃᓗᐊᓂ ᐊᑐᖅᑕᐅᔾᔨᖏᓪᓕᐊᓂᑎ ᐊᔪᖅᑎᑕᐅᓕᖅᑲᓇᖅᑐᑎ ᐊᐅᓭᓯᐊᕐᑲᓇᖅᑐᒃᓴᓕᐅᔾᔪᓂᓄᑦ ᐊᐅᓭᓯᐊᕐᑲᓇᖅᑐᒃᓴᓕᐅᔾᔪᓂᑎ ᐊᑐᓕᖁᐅᔾᔪᓂᑎ ᐊᓪᒪᓗ ᐊᑐᑎᖅᑭᑎᒃ, ᐱᓕᕆᓂᕐᒪᔾᔪᑎᑦ ᐊᑦᑭᑎᒪᔾᔪᑕᒍ ᐊᐅᓭᓯᐊᕐᑲᓇᖅᑐᒃᓴᓕᐅᔾᔪᓂᑎ <ᖁᓇᐅᑎᑦ ᐅᖁᖅᓂᒪᔨᖅ ᐃᓕᓐᓱ 3.7.

14.4.4 ᑕᐃᑲᓂᓗ ᓄᓇᒥ ᓇᓗᓇᐃᖅᓯᒪᔨᖅᑲᖅᑕᑦ ᒪᓕᕐᓗᒍ 14.4.1, ᐅᓄᐊ ᑲᓇᑕᒥ ᐅᒪᔨᓕᓕᓴᕐᔨᐊᓄᑦ ᑐᐊᖁᑎᓂᐊᖅᑐᖅᑐᑎ ᑎᑎᖁᖅᓯᒪᔾᔪᑎᒃ ᓇᓗᓇᐃᕐᓂᑎᒃᓴᓕᐅᔾᔪᓂᑎ ᐱᔨᔪᑎᑎᕐᒃᑐᑎ ᑕᐃᒪᐢᓇᐅᑦᓴᕆᐊᖅᑐᖅᑐᑎᖅᑭᑐᓂ ᐊᔪᖅᑎᑦᑎᐊᖅᑐᑎᓄᑦ ᐊᐅᓭᓯᐊᕐᑲᓇᖅᑐᒃᓴᓕᐅᔾᔪᓂᑎ >ᓪᓚᕆᔭᖅᒥ, ᑐᔪᖁᒥᔭᖅᒥ ᐅᕐᓂᒍᓐᓂᑎ ᐊᑐᑎᖅᑭᖅᑐᒥ ᐃᓄᖅᑭᖅᑐᒥ ᐊᑕᔪᓄ ᓄᓇᕗᑦ, ᒎᒃᑯᐊᓗ ᐊᑎᖅᑎᑦ ᐊᔪᖅᑎᑦᑎᐊᖅᑐᖅᑐᑎ ᐃᓄᓐᓄᑦ ᐊᐅᓭᓯᐊᕐᑲᓇᖅᑐᒃᓴᓕᐅᔾᔪᓂᑎ ᐊᓪᒪᓗ ᐅᖁᓇᑎᖅᑭᖅᑐᒥ ᐊᐅᓭᓯᐊᕐᑲᓇᖅᑐᒃᓴᓕᐅᔾᔪᓂᑎ <ᖁᓇᐅᑎᑦ.

14.4.5 ᑲᓇᑕᒥ ᐅᒪᔨᓕᓕᓴᕐᔨᐊᓄᑦ ᐃᓕᓕᐅᑎᓗᑎᑦ ᐱᔪᓐᓇᐅᑎᑦ ᓇᓕᓕᐊᖅᑐᑎ ᐊᐅᓭᓯᐊᕐᑲᓇᖅᑐᒃᓴᓕᐅᔾᔪᓂᓪᓚᔪᓄᑦ ᐱᔪᓐᓇᐅᑎᑦ ᑕᐃᒪᐢᓇᐅᑦᓴᕆᐊᖅᑐᖅᑐᑎᖅᑭᑐᓂ ᐊᑎᖅᑎᑦ ᐃᓕᓕᐅᔾᔪᓂᑎᒃ ᓇᓕᓕᐊᖅᑐᑎᐅᔭᓐᓇ ᐊᐅᓭᓯᐊᕐᑲᓇᖅᑐᒃᓴᓕᐅᔾᔪᓂᓪᓚᔪᑦ ᐊᑐᖅᑕᐅᔾᔨᖏᓪᓕᐊᓂᑎ ᐃᓗᐊᓂ ᐊᐅᓭᓯᐊᕐᑲᓇᖅᑐᒃᓴᓕᐅᔾᔪᓂᑎ <ᖁᓇᐅᑎᑦ.

ᐃᓕᓐᖏᑐ 15 — ᐊᑐᓕᖅᑎᑕᐅᓪᓚᕐᓚᐅᕐᓂ, ᖃᑎᖅᑭᐊᖅᓂᖅ ᐊᒻᒪᓗ ᐊᐅᖅᑲᑎᑭᒍᑎᖅᑲᓐᓂᖅ

—

15.1 ᑐᑭᒍᖅᐸᓪᓕᐊᔨᔪᑎᑲᖅᑭᓂ

15.1.1 ᐅᓪᓂ ᐃᓕᓐᒪᓂ ᐊᒋᓂ ᑎᑎᕎᖅᓯᒪᔪᓂ ᑐᑭᒍᖅᐸᓪᓕᐊᔨᔪᑎᑲᖅᑭᓐᓂᓕ:

(a) ᑕᒪᑲᑭᖅᓯᒪᔪᓂ, ᐊᑲᑐᑎᔪᓂ ᐊᒻᒪᓗ ᐃᑲᔪᖅᑲᑎᒋᓴ ᐊᑐᓕᖅᑎᑕᐅᓪᓚᕐᓚᐅᓕᖅᑭᓐᓄᑦᓂ ᐃᓄᐃᓂ ᐊᑲᖅᑕᐅᓂᐊᕐᓂᖅᑭᓂ ᐱᓪᓕᓪᓕᐊᑕᐅᔪᓐᓇᖅᑐᓄᑦᓗ ᐊᖅᑭᑕᐅᑐᒃ;

(b) ᐃᓐᓇᓂᑲᓂ ᖃᑎᖅᑭᐊᖅᑕᐅᖅᑲᖅᑕᕐᓗᑎᒃ ᐃᓄᐃᓂ ᐊᑲᖅᑕᐅᓂᐊᕐᓂᖅᑭᓂᑦ ᐱᓪᓕᓪᓕᐊᑕᐅᔪᓐᓇᖅᑐᓄᑦᓗ ᐊᖅᑭᑕᐅᑐᒃ ᑕᐃᒪᓗ ᐅᐊᓂ ᐃᓄᐃᓂ ᐊᑲᖅᑕᐅᓂᐊᕐᓂᖅᑭᓂᑦ ᐱᓪᓕᓪᓕᐊᑕᐅᔪᓐᓇᖅᑐᓄᑦᓗ ᐊᖅᑭᑕᐅᑐᒃ ᑐᓴᖏᓐᓂᕐᓂ, ᑐᑭᓯᐅᖅᑕᐅᑎᓂᕐᓚᐅᔪᓂ ᐊᒻᒪᓗ ᐱᓕᕆᓚᐊᖅᑲᖅᑕᒋᐊᒃ ᐊᑐᓚᐊᖅᑲᖅᑕᕐᓂ ᑎᑭᐅᑎᓂᑎᐊᖅᓯᒪᓕᖅᑐᓗᒍᓕ;

(c) ᑲᔪᓯᔪᒥᒃ ᓯᓚᕈᐅᓴᐅᔾᐅᓗᑎᒃ ᐃᓄᐃᓂ ᐊᑲᖅᑕᐅᓂᐊᕐᓂᖅᑭᓐᓄᑦᓂ ᐱᓪᓕᓪᓕᐊᑕᐅᔪᓐᓇᖅᑐᓄᑦᓗ ᐊᖅᓯᖅᑕᐅᑐᒃ ᐊᑐᓕᖅᑎᑕᐅᓪᓚᕐᓚᐅᔪᓂ ᑭᓇᐅᔾᔨᓴᒪᓂ ᐅᓪᓗᒥᐅ (ᑎᑎᕎᖅᓯᒪᔪᓂ ᐅᐃᒍᖅ 15-1) ᑕᐃᒪᓗ ᐊᑐᖅᑕᐅᑦᑎᐊᖅᓯᒪᓕᖅᑐᓗᒍᓕ ᑕᒪᑲᑭᖅᓯᒪᔪᓂ, ᑐᑭᓯᐅᖅᑕᐅᑎᓂᕐᓚᐅᔪᓂ, ᑐᑭᓯᐅᖅᑕᐅᑎᓂᕐᓚᐅᓕᖅᑭᓐᓄᑦᓂ ᐊᒻᒪᓗ ᐱᓕᕆᓚᐊᖅᑲᖅᑕᒋᐊᒃ ᐊᑐᓚᐊᖅᑲᖅᑕᕐᓂ ᐃᓄᐃᓂ ᐊᖅᑭᑕᐅᑐᒃ ᐱᓪᓕᓪᓕᐊᑕᐅᔪᓐᓇᖅᑐᓄᑦᓂ ᐊᑐᓕᖅᑎᑕᐅᓪᓚᕐᓚᐅᔪᓂ ᐱᓪᓕᓪᓕᐊᑕᐅᔪᓐᓇᖅᑐᓄᑦᓗ ᐊᑐᓕᖅᑎᑕᐅᓪᓚᕐᓚᐅᔪᓂ ᐊᑲᖅᑕᐅᓂᐊᕐᓂᖅᑭᓐᓄᑦᓂ ᐱᓪᓕᓪᓕᐊᑕᐅᔪᓐᓇᖅᑐᓄᑦᓗ ᐊᑐᓚᐊᖅᑲᖅᑕᕐᓂ ᐊᒻᒪᓗ ᑎᑎᕎᖅᓯᒪ

ᐊᒃᑐᖅᑕᐅᓂᐊᖅᓂᖏᓐᓄᓄᑦ ᐱᓕᕆᓂᖅᑕᐅᔪᓐᓇᖅᑐᓄᓪᓕᐊ ᐊᖏᕈᑎᑦ <ᖅᓇᐃᓂᖅᓗᑦ, ᒥᐊᓂᓚᓂᖅᓗᑦ ᐊᓚᒪᓗ ᐅᓂᒃᑲᓕᐅᖅᓂᖅᓗᑦ, ᑎᑭᓕᐅᖅᑲᕐᔪᐊᑦ ᐅᓂᒃᑲᓕᐅᖅᓂᖅᓗᑦ.

15.2.3 ᑲᓇᑕᒥ ᐅᒪᔪᓕᓚᖅᓴᕐᔪᐊᑯᑦ ᐊᑭᓕᐅᑎᖏᓕᑦ ᑭᓇᐅᖅᑲᓚᓄ ᐃᖏᕐᕋᓯᒪᑕᐅᓂᐊᖅᑐᑦ ᑕᒃᑯᒪᔪᑦ ᓄᓇᕗᑦ ᑐᓴᕆᔨᒃ ᑎᒥᖓ ᑎᑭᐅᑎᓯᒪᓕᖅᐸᑦ ᐊᑐᑎᖃᖅᑐᕐᓂᒃᑯᖓᓕ ᐅᑦᓗᐊ ᐊᓚᒪᓗ ᒪᓕᓚᓗᓂ ᓄᓇᕗᑦ ᐊᑭᓕᖅᑐᑎᑦ.

15.2.4 ᐃᓄᐃᑦ ᐊᒃᑐᖅᑕᐅᓂᐊᖅᓂᖏᓐᓄᓄᑦ ᐱᓕᕆᓂᖅᑕᐅᔪᓐᓇᖅᑐᓄᓪᓕᐊ ᐊᖏᕈᑎᑦ ᐊᑐᓕᖅᑎᑕᐅᓕᕆᓂᐊᓕᓄᑦ ᑭᓇᐅᖅᑲᓚᓄᑦ ᑐᓂᓯᒪᔭᖅᑐᑎᑦ ᐊᑎᓕᖅᐸᑦ ᐊᑐᓕᖅᑎᑕᐅᓕᕆᓂᖅ ᐅᑦᓗᐊ ᐋᖅᑭᒃᑯᐱᑦᑎᐊᓕᖅᑐᑦ ᓄᓇᕗᑦ ᐊᑭᓕᖅᑐᑎᑦ ᐊᑐᓕᖅᑎᑕᐅᓕᕆᓂᐊᖅᑐ

ᐱᓕᕆᓕᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕᐊᑐᖅ ᐊᖏᑎᒃᓴᓂᖅ ᐊᒪᓚᓗ ᑐᓄᓕᐅᑎᑎᓕᕆᓕᔨᓂ, ᐅᑯᐊ ᐱᓇᓱᐊᖅᑎᑎᒋᓂ ᐃᖅᐱᒍᓱᑎᑦᑐᑎᒃ ᐱᑕᖃᓂᐊᖅᑕᐅᓂᖅ, ᐊᒪᓚᓗ ᐊᑐᐃᓐᓇᐅᔾᔨᒃᑐᖅ ᑭᓇᐅᔭᖅ;

(b) ᒪᑕᑭᐊᖅᑲᖅᑐᖅ ᐊᖏᑎᑐᑎ 15.2.1 ᐋᖅᑭᑎᐊᖅᑐᓂᓄ, ᑕᐃᒪᓐᓇᐅᔾᔨᕆᐊᖅᑲᓄ, ᑎᑎᕋᖅᓯᒪᔨᖅ 15-1 ᑕᐃᒪᓗ ᑲᔪᓯᑎᐊᖅᑯᓕᔨᖅ ᐅᑯᐊ ᐃᓇᐃᑦ ᐊᑐᖅᑕᐅᓂᐊᖅᓱᖏᓐᓄᑦ, ᑐᑭᓯᐅᑎᑎᒋᓪᓕᐊᓂᖅᑐᓄᓪ ᐊᖏᑎᓗ ᐱᑕᖃᓂᐊᖅᑕᖏᓐᓄᑦ ᑎᑭᐅᑐᐊᓂᖅᑐᓄᑦ ᐊᑭᑐᑕᐊᓂᖅᑐᓄᑦ ᐱᓯᒪᓕᑎᓄᑦ;

(c) ᑎᑎᕋᖅᓗᑎᑦ ᑐᑭᓯᓇᖅᓯᒪᔪᓕᕆᓂᖅ ᐅᓂᒃᑲᑦ ᐃᖅᑭᐅᑎᔨᓂᖅ ᐅᓇ ᐊᖅᑦᔪᒍᑦ ᖃᑎᖅᑕᐊᖅᑕᐅᓯᒪᔪᕐᒃ ᐊᖏᑎᓗ ᐅᓂᒃᑲᑦᐊᖏᓯᒪᔪᖅ ᐊᑐᐃᓐᓇᐅᔾᔨᓕᓂᖅ ᐃᓄᒃᑎᑦ; ᐊᖏᑎᓗ

(d) ᐃᓯᒪᑎᑦᑎᐊᖅᓗᑎᑦ ᓇᓕᐊᑐᐃᓐᓇᖅ ᐱᓕᕆᔨᓐᓇᖅᑕᑦ ᐅᓇᓗ ᐃᑲᔪᑎᔨᖅ ᐊᑐᑎᖃᓕᖅᓗᓂᓄᑦ ᐊᑐᓕᖅᑎᑕᐅᓂᖏᓐᓄᓕᓂ ᐃᓄᐃᑦ ᐊᖏᑎᑕᐅᓂᐊᖅᓱᖏᓐᓄᑦ ᐱᓕᕆᓕᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕᐊᑐᖅᑐᑦ.

15.4.2 ᐱᓕᕆᓗᑎᑦ ᖃᑎᖅᑕᐊᖅᓗᑎᑦ ᐊᒋᔪᑦ 15.4.1, ᐅᑯᐊ ᐱᓇᓱᐊᖅᑲᑎᑎᒋᓂ ᖃᑎᖅᓗᑎᒃ ᑐᓴᖅᑕᐅᔨᓐᓇᖅᑐᓄᓪ ᐊᖏᑎᓗ ᐅᖅᑲᐅᔾᔨᖅᑕᐅᓗᑎᒃ ᑮᑯᓇᖏᓐᓄᑦ ᐊᐅᓚᕐᑎᑲᑎᑎᒋᓐᓂᕐᔪᑦ ᑲᑎᒪᔨᕐᓂᖅ ᐃᓕᓴᕐᓂᖅ ᐊᑐᓕᖅᑎᑕᐅᓂᖏᓐᓄᑦ ᐃᓄᐃᑦ ᐊᖏᑎᑕᐅᓂᐊᖅᓱᖏᓐᓄᑦ ᐱᓕᕆᓕᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕᐊᑐᖅᑐᑦ.

15.4.3 ᐱᓇᓱᐊᖅᑲᑎᑎᒋᓂ ᐱᓕᕆᖅᑕᕐᓗᑎᑦ ᖃᑎᒌᓕᖅ ᖃᑎᖅᑕᐊᖅᓗᑎᒃ ᑕᐃᒃᑐᓂ ᐅᖃᓖᑦᑐᓂ ᖃᒃᑐᑎᖓᓂ ᐅᑐᐱᓗ ᓴᓂᖅᓯᐃᒃᑯᓂᖅᓗᑎᒃ ᑮᑯᓇᖏᓐᓄᑦ ᐊᒪᓚᓗ ᐊᖅᑭᑎᐊᖅᓗᑎᑦ ᓴᓇᕐᓂᖅᑕᕐᓂᖅ ᐸᖅᓯᐅᖅ.

15.5. ᐊᖏᑎᑐᑎᑦ ᑐᑭᓯᓇᖅᓯᒪᑕᐅᓂᖏᓐᓄᑦ

15.5.1 ᐅᑯᐊ ᐱᓇᓱᐊᖅᑲᑎᑎᒋᓂ ᐊᖏᑎᖅᑲᑎᑎᒋᓪᓗᑎᒃ ᐃᖅᑭᒃᓯᓗᑎᒃ ᓇᓕᓕᐊᖏᓂᒃ ᐊᐃᖏᐅᑕᐅᔨᓂᒃ ᐱᓕᕆᔨᓐᓇᖅᑐᓄᓪᓕᐊᑐᖅᑐᑦ ᑐᑭᓯᓇᖅᓯᒪᔪᓕᕆᓂᖅᑐᓄᓪ ᐱᓕᕆᓕᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕᐊᑐᖅᑐᑦ ᐅᖃᓖᑦᑐᓂ 8.4.4 ᐊᖏᓚᓗ ᑎᑎᕋᖅᓯᒪᔨᓂ 8-3 ᑮᑯᓇᑐᓂ *ᓄᓇᕗᑦ ᓄᓇᕈᑎᑎᓄᑦ ᐊᖏᑎᑐᑎᑦ* ᐅᖃᓖᑦᑐᓂ ᐃᓕᓴᕐᓂᖅ 38 *ᓄᓇᕗᑦ ᓄᓇᕈᑎᑎᓄᑦ ᐊᖏᑎᑐᑎᑦ*, ᐊᑐᑎᕋᖅᑕᑎᒃ.

15.6 ᑕᑦᑕᒪᓄᑦ ᐊᖅᑭᒍ ᐊᓂᒍᕋᖅᐸᑕ ᖃᑎᖅᑕᐊᖅᑕᕐᓗᑎᒃ

15.6.1 ᑭᓯᐊᓂ ᐊᖏᑎᖅᑕᐅᓯᒪᒍᒥ, ᑎᑎᕋᔨᑦ ᐱᑕᖃᓂᐊᖅᑕᖅᑕᑦ ᑲᔪᓯᓗᑎᒃ ᖃᑎᖅᑕᐊᖅᓗᑎᑦ ᐊᑐᑎᖃᓕᖅᓗᑎᓪᓚᕆ ᐃᓄᐃᑦ ᐊᖏᑎᑕᐅᓂᐊᖅᓱᖏᓐᓄᑦ ᐱᓕᕆᓕᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕᐊᑐᖅᑐᑦ ᐊᖏᑎᑐᑎᑦ ᑕᑦᑕᒪᓄᑦ (5) ᐊᖅᑭᒍᓄᑦ ᐊᓂᒍᕋᖅᐸᑕ ᐊᑎᓕᐅᖅᑕᕐᓂᖅᓂᑦ, ᑕᐃᒪᓗ ᑎᑭᐅᑐᐊᓂᖅᑐᓄᑦᓯᒪᔪᑦ ᐅᑯᐊ ᐃᓇᐃᑦ ᑐᑭᓯᓇᖅᓯᒪᔪᓕᕆᓕᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕᐊᑐᖅᑐᑦ 8.4.4 ᑮᑯᓇᑐᓂ *ᓄᓇᕗᑦ ᓄᓇᕈᑎᑎᓄᑦ ᐊᖏᑎᑐᑎᑦ* ᐊᒪᓚᓗ ᐃᓄᐃᑦ ᐊᖏᑎᑕᐅᓂᐊᖅᓱᖏᓐᓄᑦ ᐱᓕᕆᓕᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕᐊᑐᖅᑐᑦ ᐊᑐᑎᕋᖅᑕᐊᖅᓗᑎᒃ.

15.6.2 ᑭᓯᐊᓂ ᐊᖏᑎᖅᑕᐅᓯᒪᓗᑎᓂ, ᐱᓇᓱᐊᖅᑲᑎᑎᒋᓂ ᑎᑎᕋᖅᓗᑎᒃ ᐃᓄᐃᑦ ᐊᖏᑎᑕᐅᓂᐊᖅᓱᖏᓐᓄᑦ ᐱᓕᕆᓕᑕᐅᔨᓐᓇᖅᑐᓄᓪᓕᐊᑐᖅᑐᑦ ᐊᑐᑎᕋᖅᑎᑕᐅᓂᖏᓐᓄᓕᓂ ᐅᑯᐊᑲᓐᓂᖅᓴᓂᖅ ᐃᓕᓴᕐᓂᖅ ᖃᑎᖅᑕᐊᖅᑕᐅᓂᖏᓐᓄᑦ ᒪᑕᑭᐊᖅᑲᖅᑐᖅ 15.6.1-ᒥᑲ.

15.6.3 ᑲᓇᑕᒥ ᐅᒪᔨᓕᓚᓵᔨᐊᑯᓂ ᐊᖏᓚᓗ ᓄᓇᕗᓂ ᑐᓴᖅᓀᐃᖅ ᑎᑎᕋᖅ ᑲᑎᒪᔨᖅᑕᐅᓯᒪᔪᖅ ᑭᓇᐅᔭᖅᑲᖅᓯᕆᓂᖅᓗᓂᑎᓄ ᐱᓕᕆᓗᑎᒃ ᐊᖏᑎᓗ ᐅᓂᒃᑲᑦᐊᓚᓯᒪᔨᓂ ᑎᑎᕋᖅᑕᐅᓂᐊᖅᓱᖏᓐᓄᑦ ᒪᑕᑭᐊᖅᑲᖅᑐᖅ 15.6.1-ᒥᑲ.

15.7 ᐊᐃᔨᖏᑎᒋᒃᑕᐅᖅᓂᕐᓂᓂ ᐃᓄᐃᑦ ᐊᖏᖅᑕᐅᓂᐊᖅᓯᐅᓐᓄᑦ ᐱᓕᕆᓂᑕᒃᓴᓂᖅᓯᓄᑦ ᐊᓯᖏᑐᑎᑦ

15.7.1 ᐅᓇ ᐃᓄᐃᑦ ᐊᖏᖅᑕᐅᓂᐊᖅᓯᐅᓐᓄᑦ ᐱᓕᕆᓂᑕᒃᓴᓂᖅᓯᓄᑦ ᐊᓯᖏᑐᑎᑦ ᑲᔪᓂᐊᖅᑐᖅ ᑭᓱᐊᓂ ᓄᓇᖅᑐᑦ ᐊᓂᖅᑐᖅᑕᐅᔪᓂᓂ ᐊᐅᓚᑦᓂ ᐱᓕᕆᓂᑕᒃᓴᓂᖅᓯᓄᑦ ᐅᕐᓕᔅᓴᓂ ᑭᓱᐊᓂ ᐃᓇᓂᖅᑕᐅᔪᓂᓂ ᓄᑖᒍᑦ ᐃᓄᐃᑦ ᐊᖏᖅᑕᐅᓂᐊᖅᓯᐅᓐᓄᑦ ᐱᓕᕆᓂᑕᒃᓴᓂᖅᓯᓄᑦ ᐊᓯᖏᑐᑎᑦ ᐊᐃᔨᖏᑎᒋᒃᑕᓯᒪᔪᒥᒃ ᒪᓕᒃᑐᖅ 15.7.2-ᒥᒃ.

15.7.2 ᒪᓕᒐᔭᔪ 8.4.7 ᑭᒃᑯᓇᓂ *ᓄᓇᕗᓱᒃ ᓄᓇᑕᑭᑎᓄᑦ ᐊᖏᖅᑭᑎᑦ*, ᐱᓇᓯᐊᖏᖅᑎᑎᑦ ᐃᒪᐃᓇᐅᓂᐊᖅᑐᑦ, ᑭᓱᐊᓂ ᐊᖏᕐᖏᖅᑎᑎᑐᖅ, ᐊᐃᔨᖏᑎᒋᒃᑎᑎᑎᕐᓂᐃᓐᓄᑦ ᐅᓇ ᐃᓄᐃᑦ ᐊᖏᖅᑕᐅᓂᐊᖅᓯᐅᓐᓄᑦ ᐱᓕᕆᓂᑕᒃᓴᓂᖅᓯᓄᑦ ᐊᖏᖅᑭᑎᑦ ᐊᓂᔪᖅᐸᑕ 7 ᐊᕐᕋᔪᑦ; ᐊᐃᔨᖏᑎᒋᒃᑎᑎᑕᐃᓕᓴᕐᓯᑦ ᐱᑕᐊᓪᓕᔭᓂᓂ ᐊᕐᕋᔪᒥ 6-ᖏᓂᓂ.

15.7.3 ᐅᑯᐊ ᐱᓇᓯᐊᖏᖅᑎᑎᑦ ᕿᑭᕐᕕᐊᕐᓗᔪ ᐃᓄᐃᑦ ᐊᖏᖅᑕᐅᓂᐊᖅᓯᐅᓐᓄᑦ ᐱᓕᕆᓂᑕᒃᓴᓂᖅᓯᓄᑦ ᐊᖏᖅᑭᑎᑦ ᐊᑐᓕᖅᑎᑕᐅᔪᑦ ᐅᓂᒃᑲᖏᑦ ᑎᑎᕐᓂᖅᓯᒪᔪᑦ ᒪᓕᒐᔭᔪᑎᒃ 15.6.2, ᑭᒃᑯᓇᖏᑦ ᑐᓴᕐᕕᑎᐅᓇᒪᓂᖅᓯᓂ, ᕿᑐᐅᓴᓇᓱᓐᓂᑎᒃ ᑭᐃᔨᖏᑎᒋᒃᑕᕐᑲᓂᓂᐊᕐᓚᖏᑦ ᐅᓂᒃᑲᖏᑦ ᑭᕕᑭᐊᕐᓂᐊᕐᓚᖏᑦ ᐃᓄᐃᑦ ᐊᖏᖅᑕᐅᓂᐊᖅᓯᐅᓐᓄᑦ ᐱᓕᕆᓂᑕᒃᓴᓂᖅᓯᓄᑦ ᐊᖏᖅᑭᑎᑦ ᑕᐃᒪᓐᓇᐅᔾᔨᖅᑕᐊᕐᕕᖅᑐᖅ.

ᐃᓄᐃᑦ ᐊᑲᑐᖅᑕᐅᓂᐊᖅᓯᖅᑭᓐᓄᑦ ᐱᖓᓕᒫᑕᐅᔪᖅᓯᒪᓕᑦᑐ ᐊᖅᑭᒃᑯᐊᑦ 15-1 (ᒪᓕᒐᖅ 15.2.1)

	ᐊᖅᑭᒍ 1	ᐊᖅᑭᒍ 2	ᐊᖅᑭᒍ 3	ᐊᖅᑭᒍ 4	ᐊᖅᑭᒍ 5	ᐊᖅᑭᒍ 6	ᐊᖅᑭᒍ 7	ᑲᑎᓕᑎᑦ
1. ᐊᐅᓚᑦᑎᖅᑎᖏᑐᓂᒪᑦ ᑲᑎᒪᒃᓴᓕᑦ	$350,000	$350,000	$350,000	$350,000	$200,000	$200,000	$200,000	$2,000,000
2. ᐃᓄᐃᑦ ᑲᑐᔨᖅᑲᑎᖏᑐ ᐃᖅᑭᖅᖃᔪᑎᖅᑭᑦ (ᐊᐅᓚᑦᑎᖅᑎᖏᑐᓂᒪᑦ ᑲᑎᒪᒃᓴᓕᓪᓗᑦ)	$25,000	$25,000	$25,000	$25,000	$20,000	$15,000	$15,000	$150,000
3. ᐱᐅᓯᑐᖅᑲᓚᐱᓂᖅᒪᑦ ᐱᓴᐸᑎᖅᑭᑦ	$300,000	$300,000	$200,000	$150,000	$150,000	$150,000	$150,000	$1,400,000
4. ᑐᑭᓯᓇᖅᓯᐸᑕᐅᔨᓄᓪᓗᑦ ᑐᕐᑐᑦ	$57,142	$57,142	$57,142	$57,142	$57,142	$57,142	$57,148	$400,000
5. ᐃᓄᖏᓄᑦ ᓂᐅᑭᖅᑲᑦᑐᑕᓴᓄᑦ ᑭᓇᐅᔪᖅᑭᖅ [i]	$2,300,000	$300,000	$300,000	$300,000	$300,000	$300,000	$300,000	$4,100,000
6. ᐃᓄᐃᑦ ᐸᐊᑎᑎᐃᖅᑭᓄᑦ ᐃᑲᔪᑎᑦ	$64,285	$64,285	$64,285	$64,285	$64,285	$64,285	$64,290	$450,000
7. ᐃᓄᖏᓄᑦ ᐃᒋᓐᓂᐊᖅᑎᑎᕆᓄᑦ ᒪᓂᒪᔪᖅ [ii]	$50,000	$50,000	$50,000	$50,000	$50,000	$50,000	$50,000	$350,000
8. ᐃᓄᖏᓄᑦ ᐃᑲᔪᖅᑎᑭᐅᔪᓄᑦ ᖃᐅᔨᓴᖅᑎᓄᑦ [iii]	$50,000	$50,000	$50,000	$50,000	$50,000	$50,000	$50,000	$350,000
ᑲᑎᓐᓂᑦ	**$3,196,427**	**$1,196,427**	**$1,096,427**	**$1,046,427**	**$891,427**	**$886,427**	**$886,438**	**$9,200,000**

I. ᐱᓇᓱᐊᖅᑎᑎᑦ ᐊᖏᓂᖅᑐᑦ ᐅᓇ ᑭᓇᐅᔪᑦᑎᖅ ᐊᐃᖏᓐᓄᓂ ᐃᒪᓐᓇ: $2,100,000 ᐄᓇᐅᔪᖅ ᐱᓕᓚᐊᖏᓴᓄᑦ (ᐊᖅᐃᐊᓂ, ᐅᖅᓱᖅᑐᒥ, ᑭᖏᑕᖅᓂ) $2,000,000 ᓄᑦᓴᓄᑦ ᐱᓕᓚᐅᑎᓂᐊᖅᑐᓄᑦ.

II. ᑭᓇᐅᔪᖅᐃᑦ <<ᑕᐅᑎᓂᐊᖅᑐᑦ ᑲᓇᑕᒥ ᐅᒪᔪᓕᓚᐱᖅᓯᐊᑯᓐᓂᑦ ᒪᓕᒃᓴᐃ ᐊᓐᖏᑎᑦ 9.2.2

III. ᑭᓇᐅᔪᖅᐃᑦ <<ᑕᐅᑎᓂᐊᖅᑐᑦ ᑲᓇᑕᒥ ᐅᒪᔪᓕᓚᐱᖅᓯᐊᑯᓐᓂᑦ ᒪᓕᒃᓴᐃ ᐊᓐᖏᑎᑦ 10.3.2

ᐃᓕᓐᓴ 16 — ᐊᐃᖏᐅᑎᓯᓄᑦ ᑐᑭᓯᑎᑦ

16.1 ᑐᑭᓴᖅᓯᑦᓕᐊᓯᑎᕆᓴᑦ

16.1.1 ᐅᖏᓂ ᐃᓕᓐᓄᓂ ᐊᑕᓂ ᑎᑎᕋᖅᓱᒪᓯᑦ ᑐᑭᓴᖅᓯᑦᓕᐊᓯᑎᕆᓴᓐᓂᑦ:

(a) ᓴᕋᑦᑎᓗᑎᒃ ᕋᓕᒫᒥᐅᓯᒥ, ᐊᑭᒃᑭᑦᑐᒥᒃ, ᐊᑐᑎᕋᖅᑐᑐᒥᒃ ᐊᒪᒪᓗ ᐊᕋᑦᑎᐊᖅᓱᓯᓇᖅᑐᓂᒃ ᐊᐃᖏᐅᑕᐅᑎᓯᓄᑦ ᑐᑭᑕᑭᑎᓂᒃ ᐱᖏᓕᓕᐊᓂᖅᓴᓗᑎᒃ; ᐊᒪᒪᓗ

(b) ᓴᐳᒻᒥᓂᖅ ᐊᒪᒪᓗ, ᐊᑐᑎᕋᖅᑎᑦᓗᒍ, ᓴᓐᖏᑎᑎᓴᓂᖅ ᐃᓕᑭᔪᓂᖏᓐᓂᒃ ᐱᓇᓱᕋᖅᑎᑭᑦ.

16.2 ᑐᑭᖏᑦ

16.2.1 ᐅᖏᓂ ᐃᓕᓐᓴᓂ:

(a) “ᐃᓕᓐᓴᕋᖅᑎᑭᖅᑎᑦᑎᓂᖅ” ᑐᑭᓴᕋᖅᑐᖅ ᑐᑭᓱᓇᓱᕋᖅᑎᑭᖅᓂᖅ ᑕᒃᑯᐊ ᐱᓇᓱᐊᕋᖅᑎᑭᑦ ᕋᑎᓗᑐᑎᒃ ᐊᒪᒪᓗ ᐱᑕᕆᓇᓱᒃᑕᒥᖅᓄᑦ, ᐃᒃᑯᑦᑐᐊᓇᓱᒃᑕᒥᖅᓄᑦ ᐊᐃᖏᐅᑕᐅᑎᓯᓄᑦ ᐊᐃᖏᐅᑕᐅᑎᓯᓄᑦ ᐱᓇᓱᐊᕋᖅᑎᑭᑦ ᐃᓕᓴᕐᓂᖅ ᐊᐃᖏᐅᑕᐅᑎᓯᓄ

ᐱᓕᓀᐊᖏᓐᓇᓕᒃᓗᓂ. ᐱᓇᓱᐊᕐᑲᑎᑦ ᓇᓚᒪᓂᖅᑐᑦᑕᐅᕐᕋᓱᐊᕐᓂᐊᓂᖅᑐᑦᑕᑦ ᐊᓂᕐᑲᑎᑦᔪᑕᐅᓱᒪᔪᒃ ᐅᕐᑲᖅᓱᒪᔪᖅ ᐊᐃᓐᐅᑎᓂᖅᑯᑦ ᐊᓂᕐᑲᑎᑦᔪᓂᓇᐃᕐᓱᒪᔪᑦ.

16.3.2 ᐱᑕᕐᑲᖅᑐᑦᖅ ᐅᓐᓂ ᐊᓂᕐᑕᑎᓂ ᐊᔪᕐᑎᑕᑎᓗᑎᒃ ᐋᕐᕿᔪᑕᐅᔪᓇᖅᑐᓂᒃ ᓇᑦᑕᐊᓄᑕᐅᔪᓂᖅᑐᓂᒃ ᐱᓇᓱᐊᕐᑲᑎᑦᔪᓂᖅ.

16.3.3 ᐊᑐᓂ ᐱᓇᓱᐊᕐᑲᑎᑦᔪᑦ ᐊᕿᒋᓂᐊᕐᑲᖅᑐᑦ ᐊᐃᓐᓕᔪᕐᑕᐅᔪᓇᓂᑦ ᑐᕿᑕᕿᑎᓂᕐᓂᓂᑦ ᐱᑕᓚᒃᐅᔪᓇᓂᑦ ᒪᑕᓪᓗᑎᒃ ᐅᓪᒥᖏᓪᓚ ᐊᖏᓪᓗᓐᓇ, ᐊᓚᒪᓐᓇ ᐱᓇᓱᐊᕐᑲᑎᑦᔪᑦ ᑕᒪᕐᒥᒃ ᐊᑐᓂ ᐊᖅᐳᑎᒥᒃ ᐊᕿᒋᓐᓇᑎᒃ ᐃᖅᑲᐅᐃᓇᕐᕕᒋᓐᓇᓂᑦ ᓴᖅᑭᑲᑎᑦᑎᑦᑎᓂᓗ ᒪᑕᓪᓗᑐᑦ 16.3.1 (b) ᐊᓚᒪᓐ ᒪᑕᓪᓗᑐᑦ ᑕᒪᐃᓐᓇᑦ ᐊᓂᕐᑲᑎᑦᔪᓂᓇᐃᕐᓱᒪᔪᖅ ᑕᕐᕈᑦ 16.3.1 (c).

ᐅᐃᒍᖅ 1— ᓂᐦᑭᖏᓂᖅ ᑲᓇᑕᒥ ᐅᒪᔪᖅᑲᐃᑦ

1.0 ᐃᓄᐃᑦ ᓄᓇᕋᐊᑎᖏᑦ

1.1 ᑲᓇᑕᒥ ᐅᒪᔪᑦᑕᓗᔪᕐᓗᑦ ᑐᓴᐅᔪᓂᖅᑐᑦ ᐊᑐᑎᔪᓂᒃ ᐱᔪᓐᓇᐅᑎᓂᒃ ᒪᑦᓴᒍᒍ ᐊᒋᑦ ᑲᓇᑕᒥ ᐅᒪᔪᑦᑕᓂᖓᒍ ᐱᕋᖅᓯᔪᕐᓗᖅ ᐊᔪᖏᑎᑦᓯᐅᑦ ᕿᑭᖅᑕᑦ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑎᒋᔭᖏᑦ ᐅᖓᓚᓂᖓᓂ ᐃᓄᐃᑦ ᓄᓇᓕᒫᑦ ᐊᑐᑎᖅᑎᓂᖅᑐᒃ >ᓕᕋᑐᑦᓕᑕᓂᖅ ᐃᓗᓯᒥᒃ ᐊᒻᒪᓗ ᖃᐅᔨᕋᖅᓯᒪᔭᖅ, ᖃᐅᔨᓕᓴᐃᖅᑎᓂᒃ ᒥᑕᕋᖅ ᐊᒻᒪᓗ ᑐᓕᖅᑕᕋᖅᑐᒃ [Cape Raper].

1.2 ᓇᓂᓐᓂᖓ ᐃᓂᖓ ᐊᒻᒪᓗ ᓴᓇᖅᐅᓂᐊᕐᓂᖏᑦ ᓇᐸᖅᑎᑕᐅᓂᐊᕐᓂᖏᑦ ᐊᒻᒪᓗ ᐊᐅᓕᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐱᑎᐊᕐᕕᖅᓴᖅᑎᑦ ᐱᑦᑐᑎᑦ ᓇᐸᖅᑕᐅᕐᓴᑦ ᓇᑐᐃᐸᖅᓴᕐᓴᑦ ᒪᓕᒐᖅ 1-1 ᐊᖏᕐᖅᑕᐅᓴᓕᑎᑎᒃ ᐊᐢᑕᒋᑦ ᕿᑭᖅᑕᐅᑦ ᐃᓄᐃᑦ ᑲᑐᔾᔨᖅᑎᑎᒋᔭᖏᑦ ᐊᒻᒪᓗ ᑲᓇᑕᒥ ᐅᒪᔪᑦᑕᓗᔪᕐᓗᑦ, ᕿᑎᕐᕉᐊᖅᑐᓂᑎᑦᓯᐅᑦ ᐅᖓᓚᓂᖓᓂ ᐊᐅᓕᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᒪᓕᒐᖅ ᑲᑐᔾᔨᖅᑎᑎᒋᔭᖏᑦ, ᕿᑎᑦᓴᐊᖅᑎᑎᒋᔭᖏᑦ ᓴᖅᑭᑎᓂᖅ ᐊᐅᓕᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐊᒻᒪᓗ ᐊᐅᓕᑕᐅᓂᐊᕐᓂᖏᓐᓄᑦ ᐊᐢᑕᒋ

2.4 ᐱᓕᕆᓂᖅᑐᑦ ᐊᒃᑲᕈᓇᓐᓂᑎᒃ ᐅᖅᑳᖅᓯᒪᔪᓄ 2.1 ᑎᑭᓪᓗᒍ 2.3 ᐊᑐᑎᓂᐊᖏᑦᑐᔪᓄ ᓄᑖᖓᐱᐊᖅᑎᑕᓄᑦ ᐅᖃᓪᓚᑉᓂ ᐊᖅᑭᑎᐊᖅᑕᓄᑦ ᒪᓕᓇᐅᔨᖅ ᐊᑐᖅᑕᐅᔨᓂ ᐱᓕᕆᓂᖅᑎᑦ, ᑕᐃᑲᓂᓗ ᓄᑖᑦᑎᑕᐅᔪᓂ ᐅᖃᓪᓚᑉᓂ ᐊᖅᑭᑎᐊᖅᑕᐅᔪᓂ ᐊᔾᔨᑭᓐᓇᕐᓂᑎᐊᖅᑕᖓᓄᑦ ᐅᖃᓪᓚᑉᓂ ᓇᓂᓐᓂᖓᔪᓂ ᐊᑐᖅᑕᐅᔨᔪᒃ ᐱᓕᕆᓂᖅᑎᑦ ᐅᖃᓪᓚᑉᓂ ᐃᓇᓐᓂᖓᔪᓂ.

2.5 ᐱᖁᐅᔪᓂ ᐱᓕᕆᐅᑎᓄᑦ ᐅᒃᑐᑎᑎ ᐅᖃᓪᓚᑉᓂ ᑎᑎᕐᖅᖅ ᐱᓕᒪᔨᓂᑦ ᐃᓄᐃᓐᓕᕐᓂᑦ ᐅᖃᓪᓚᑉᓂ ᐃᓄᐃᓐᓕᕐᓂᑦ ᓇᓐᒥᓂᖅᑐᖅᓂᑦ, ᑲᓇᑕᒥ ᐅᒪᔨᓕᕆᓂᕐᕕᐊᓯᑦ ᑐᓴᕐᓂᕐᑎᐊᖅᑐᖅ ᐱᒃᑐᑎᓐᓇᓄ ᐅᖃᓪᓚᑉᓂ ᑎᑎᕐᖅᓯᒪᔪᒥᒃ ᐱᒃᑐᑎᓐᓇᓄ ᐅᖃᓪᓚᑉᓂ ᐃᓇᓐᑎᕐᓯᒪᔪᓄ ᑎᑎᕐᖅᓯᒪᔪᒥᒃ ᐱᒃᑐᑎᓐᓇᓄ ᐅᖃᓪᓚᑉᓂ ᑎᑎᕐᖅᓯᒪᔪᓄ. ᖃᐅᖅᑕᒥᓂᖅ ᐃᓄᐃᓐᓕᕐᓂᑦ ᐅᖃᓪᓚᑉᓂ ᐃᓇᓐᑎᕐᓯᒪᔪᓄ ᑐᓴᕐᓂᕐᑎᐊᖅᑐᖅᓂᑦ ᐅᖃᓪᓚᑉᓂ ᐃᓄᐃᓐᓕᕐᓂᑦ ᓇᓐᒥᓂᖅᑐᖅᓂᑦ, ᑲᓇᑕᒥ ᐅᖃᓪᓚᑉᓂ ᐃᓄᐃᓐᓕᕐᓂᑦ ᐃᓇᓐᑎᕐᓯᒪᔪᓄ ᑐᓴᕐᓂᕐᑎᐊᖅᑐᖅᓂᑦ

ᐊᑎᓕᐅᖅᑐᑦ

ᑯᐃᖏ ᐱᓪᓇᒪᖅᓂᖅᑎᖅᓂᖏᓐᓄᑦ ᑲᓇᑕᒥ

ᒥᓴᑕ ᑲᑐᓕᖓ ᒪᑭᓇ
ᒥᓴᑕ ᐊᕐᑎᓕᕆᔨᑯᖓᓄᑦ

ᐅᑉ

ᓄᓇᕗᑦ ᓇᓗᓇᐃᖅᑕᐅᓯᒪᓂᖏᓐᓄᑦ

ᓄᓇᕗᑦ ᑐᖏᓐᓇᕆᕐ ᑎᒥᖏᓐᓄ:

ᑲᑎ ᑕᐅᑐᖏᑎ, ᐊᖏᓐᓇᓪᓚᖅᑲᖅ
ᓄᓇᕗᑦ ᑐᖏᓐᓇᕆᕐ ᑎᒥᖏᓐᓄ:

ᐅᑉ

ᓄᓇᕗᑦ ᑐᖏᓐᓇᕆᕐ ᑎᒥᖏᓐᓄ:

ᓯᑕᖏᓕ ᐊᓇᐸᓕᒃ, ᐊᖏᓐᓇᓪᓚᖅᑲᖅ
ᖃᑎᖅᑎᐅᑦ ᐃᓄᐃᑦ ᑲᑐᔪᖅᑎᑎᕆᖏᑦ:

ᐅᑉ

ᖃᑎᖅᑎᐅᑦ ᐃᓄᐃᑦ ᑲᑐᔪᖅᑎᑎᕆᖏᑦ:

ᑕᐃᕕᑎ ᓂᖏᐅᖏᓐᓇᑦ, ᐊᖏᓐᓇᓪᓚᖅᑲᖅ
ᑭᓴᓕᖅ ᐃᓄᐃᑦ ᑲᑐᔪᖅᑎᑎᕆᖏᑦ:

ᐅᑉ

ᑭᓴᓕᖅ ᐃᓄᐃᑦ ᑲᑐᔪᖅᑎᑎᕆᖏᑦ:

ᐅᑉ

Group	Comment	Contact	Invited
Salliqmiut	Invite via radio announcements and advertisements throughout the community	N/A – radio and advertisements	Radio and advertisements
Elders	Invite via radio announcement	N/A - radio	Radio and advertisements
Youth Center	Coordinator	Sarah Netser Sarahnetser@hotmail.com	Radio and advertisements
CLARC	Inactive CLARC	Ron Ningeongan clocoral@kivalliqinuit.ca	Email 24-Oct-2019
Hamlet	SAO	Leonie Pameolik munch@qiniq.com Phone : (867) 925-8867	Email 24-Oct-2019
Aiviit HTO	Manager	Natasha Hattie Ottokie Aiviit Hunters' and Trappers' Organization Coral Harbour, NU X0C 0C0 (T) 867-925-8622 aiviit@kivalliqhto.ca	Email 24-Oct-2019
Outfitters	Requested contact list from Aiviit HTO Sept 13 and Sept 19, 2019 HTO provided 2 names Sept 20; 1 name Sept 23. *HTO must ask for permission to share contact information.*	Jimmy Ningeocheak jningeocheak@yahoo.ca Aaron Emiktowt aaron_emiktowt@hotmail.com Darcy Nakoolak darcynakoolak@hotmail.ca	Email 24-Oct-2019
Cabin Owners	Requested contact list from Aiviit HTO Sept 13 and Sept 19, 2019	None in MBSs.	N/A
Kivalliq Inuit Association	Director of Implementation and Planning Implementation Officer	Annie Tattuinee atattuinee@kivalliqinuit.ca Phone: (867) 645-5732 Michael Tucktoo mtucktoo@kivalliqinuit.ca Phone: (867) 645-5743	Email 24-Oct-2019
Nunavut Planning Commission	Director of Policy and Planning	Brian Aglukark aglukark@nunavut.ca	Email 24-Oct-2019

		P.O. Box 419, Arviat, Nunavut, X0C 0E0 Phone: (867) 857-2242	
Nunavut Impact Review Board	Executive Director	Ryan Barry rbarry@nirb.ca Phone: (867) 983-4608 29 Mitik Street, PO Box 1360 Cambridge Bay, Nu X0B 0C0	Email 24-Oct-2019
Government of Nunavut – Environment	Regional Biologist, Arviat Conservation Officer in Coral Harbour	Mitch Campbell MCampbell1@gov.nu.ca Phone: (867) 857-3171 Troy Nester TNetser@gov.nu.ca 867-925-8823	Email 24-Oct-2019
Government of Nunavut – Culture and Heritage	Territorial Archaeologist	Sylvie LeBlanc SLeBlanc1@GOV.NU.CA Phone: (867) 934-2040	Email 24-Oct-2019
Kivalliq Wildlife Board	Kivalliq Regional Coordinator	Amy Kaludjak kwb@niws.ca Phone: (867) 645-4860 Nunavut Inuit Wildlife Secretariat Box 219 Rankin Inlet, NU X0C 0G0	Email 24-Oct-2019
Nunavut Water Board	Executive Director In Gjoa Haven	Stephanie Autut Phone: (867) 360-6338 PO Box 119 Gjoa Haven, NU X0B 1J0 info@nwb-oen.ca	Email 24-Oct-2019
Inuit Heritage Trust	Executive Director Traditional Place Names Manager	William Beveridge wbeveridge@ihti.ca Lynn Peplinski lpeplinski@ihti.ca	Email 24-Oct-2019
MLA Aivilik		Patterk Nester pnetser1@gov.nu.ca Phone: (867) 975-5070	Email 24-Oct-2019

CIRNAC	Director of Lands	Erik Allain erik.allain@canada.ca Phone: (867) 975-4295 Box 2200 Iqaluit, NU X0A 0H0	Email 24-Oct-2019
DFO	Oceans Program Division Manager Senior Oceans Biologist	Joel Ingram Joel.Ingram@dfo-mpo.gc.ca 204-983-5006 Charlotte Sharkey Charlotte.sharkey@dfo-mpo.gc.ca 867-979-8027	Email 24-Oct-2019
ECCC researchers		Jim Leafloor jim.leafloor@canada.ca 204-983-5258 Frank Baldwin frank.baldwin@canada.ca 204-984-6670 Paul Smith paulallen.smith@canada.ca 613-998-7362 Grant Gilchrist grant.gilchrist@canada.ca 613-998-7364 Dominique Henry Dominique.henry@canada.ca 514-496-9024	Email 24-Oct-2019 Email 25-Oct-2019
ECCC Wildlife Enforcement		Steve Allan steve.allan@canada.ca 867-669-4788 Quentin Deering quentin.deering@canada.ca 867-669-4761	Email 24-Oct-2019
Nunavut Tunngavik Inc.	Director of Implementation Assistant Director of Implementation	Qilak Kusugak qkusugak@tunngavik.com 867-975-4908 Pacome Lloyd plloyd@tunngavik.com	Email 25-Oct-2019

		867-975-4925	
Academic – University of Ottawa	Researcher – Inuit knowledge	Natalie Carter natalieacarter001@gmail.com	Email 25-Oct-2019

Archived: February 6, 2020 3:26:02 PM
From: Dufour2, Jean-François (EC)
Sent: October 24, 2019 10:58:00 AM
Cc: Noah Kadlak; Dufour2, Jean-François (EC); Ronnie Ningeongan
Bcc: SAO Coral; 'Aiviq HTA'; 'jningeocheak@yahoo.ca'; 'aaron_emiktowt@hotmail.com'; 'darcynakoolak@hotmail.ca'; 'atattuinee@kivalliqinuit.ca'; Michael Tucktoo; 'aglukark@nunavut.ca'; 'rbarry@nirb.ca'; 'MCampbell1@gov.nu.ca'; 'TNetser@gov.nu.ca'; 'SLeBlanc1@GOV.NU.CA'; Qovik Netser; 'info@nwb-oen.ca'; 'wbeveridge@ihti.ca'; 'Lynn Peplinski'; 'pnetser1@gov.nu.ca'; 'erik.allain@canada.ca'; 'Joel.Ingram@dfo-mpo.gc.ca'; 'Charlotte.sharkey@dfo-mpo.gc.ca'; Leafloor, Jim (EC); Baldwin, Frank (EC); Smith, Paul Allen (EC); Gilchrist, Grant (EC); Allan, Steve (EC); Deering, Quentin (EC); 'receptionist@nwmb.com'; Pirie-Dominix, Lisa (EC)
Subject: Irniurviit ACMC Community Meeting Invitation and Review of draft Management Plans
Sensitivity: Normal
Attachments:
IMBS MP_DRAFT2_October2019.pdf;Irniurviit ACMC Community Meeting Invitation and Review of Draft MPs.pdf;QMBS MP_DRAFT2_October2019.pdf;

Good afternoon,

Please find attached an invitation to the Irniurviit Area Co-Management Committee's community meetings 26-28 Nov 2019 in Coral Harbour. Also attached is a copy of the draft management plans for the Qaqsauqtuuq and Ikkattuaq Migratory Bird Sanctuaries for your review and feedback **by 28 November 2019**.

If you have any questions about the meetings or draft management plans, please do not hesitate to contact me.

Qujannamiik!
JF

JF Dufour
Vice-Chair, Irniurviit ACMC
jean-francois.dufour2@canada.ca / Tel: 867-669-4766

ᐃᖃᓗᐃᑦ ᐊᐅᓚᑦᑎᐊᕐᓂᖅᑕᐅᓯᒪᔪᑦ
ᑲᑎᒪᔨᖏᑦᑐᑦ
IRNIURVIIT AREA
CO-MANAGEMENT
COMMITTEE

Archived: February 6, 2020 3:3 :14 PM
From: Dufour2, Jean-François (EC)
Sent: October 25, 2019 8:50:00 AM
Bcc: Henri, Dominique (EC); 'Natalie Carter'; 'qkusugak@tunngavik.com'; 'Pacome Lloyd'
Subject: Irniurviit ACMC Community Meeting Invitation and Review of draft Management Plans
es onse re uested: No
Sensitivity: Normal
Attachments:
IMBS MP_DRAFT2_October2019.pdf ;Irniurviit ACMC Community Meeting Invitation and Review of Draft MPs.pdf ;QMBS MP_DRAFT2_October2019.pdf ;

Good morning,

Please find attached an invitation to the Irniurviit Area Co-Management Committee's community meetings 26-28 Nov 2019 in Coral Harbour. Also attached is a copy of the draft management plans for the Qaqsauqtuuq and Ikkattuaq Migratory Bird Sanctuaries for your review and feedback **by 28 November 2019**.

If you have any questions about the meetings or draft management plans, please do not hesitate to contact me.

Qujannamiik!
JF

JF Dufour
Vice-Chair, Irniurviit ACMC
jean-francois.dufour2@canada.ca / Tel: 867-669-4766

ᐃᖃᓗᐃᑦ ᐃᓕᓴᕐᓂᐊᕐᓂᖅ ᑲᑎᒪᔨᖃᑦᑕᕐᓂᖅ
IRNIURVIIT AREA CO-MANAGEMENT COMMITTEE

ᐊᖏᔪᐊᓗ 24, 2019

ᑎᑎᕋᖅᑕᐅᔾᔨᓚᐊᓂᖓᓂ ᐊᐅᓚᕋᑎᓕᓐᓂᖅ ᕿᖏᓐᓄᕐᕋᐃᑦ ᑕᐅᔾᔭᓂ
ᖃᖅᖢᓴᐅᖅᑐᑎᒥ ᐊᓘᒪᓗ ᐃᑲᔪᑐᐊᖏᑦ ᑎᓐᓕᐊᖅᖃᖅᓴᓐᓄᖅ

ᖃᖅᖢᓴᐅᖅᑐᑎᒥ ᐊᓘᒪᓗ ᐃᑲᔪᑐᐊᖏᑦ ᑎᓐᓕᐊᖅᖃᖅᓴᐃᖅ, ᓴᓪᓕᐅᕐᓂ ᖁᑭᖅᑕᒧᑐᐊᓂᑐᑐᒃ, ᐱᓇᓱᒪᖅᑕᐅᓕᐅᖅᓯᒪᔨᒃᑲ ᑕᒪᓂ 1959-ᖁᑎᓪᓗᐃ. ᒪᓕᓴᕐᑕᐅᔭᕆᐊᖅᑎᖅᓂᖓᓕᓇᓂᓐᓂ ᓇᓇᕐᓕᒻᒥ ᓇᓇᑕᐸᑎᓇᓂ ᐊᖓᕈᕐᑐᕐᓂ ᐊᖏᕐᕈᓐᓇᐅᓕᐅᖅᓯᒪᔨᖏᓪᓗᒃ ᐊᐅᓚᕋᑎᓕᓐᓂᖅᓴᖅᑎᑦᑎᓂᖓᓂᓇᓂᓐᓂ ᐃᓇᓇᐃᓐᓇᖅᑐᓂ ᑲᑎᒪᔨᖏᓪᓗᒃ ᑲᑎᒪᔨᖏᑦ ᐃᓇᓇᐃᓐᓇᖅᑐᓂ

ᐃᖃᓗᖅᓯᐅᑦ ᐊᐅᓚᑦᑎᓕᐅᕆᓂᖅᓯᒪᓂᖅ ᑲᑎᒪᔨᖏᑦᑐᑦ

IRNIURVIIT AREA CO-MANAGEMENT COMMITTEE

24 October 2019

DRAFT MANAGEMENT PLANS FOR QAQSAUQTUUQ (EAST BAY) AND IKKATTUAQ (HARRY GIBBONS) MIGRATORY BIRD SANCTUARIES

The Qaqsauqtuuq (East Bay) and Ikkattuaq (Harry Gibbons) Migratory Bird Sanctuaries, located on Southampton Island, were created in 1959. As required by the *Nunavut Agreement*, an *Inuit Impact and Benefit Agreement for National Wildlife Areas and Migratory Bird Sanctuaries in the Nunavut Settlement Area (IIBA)* was first concluded in 2006 and renegotiated in 2016. Nunavut Tunngavik Inc., the three Regional Inuit Associations, and the federal Minister of the Environment, Environment and Climate Change Canada signed the *IIBA*. The *IIBA* created co-management committees for these protected areas in Nunavut. Inuit, from Coral Harbour, and Environment and Climate Change Canada co-manage the Qaqsauqtuuq and Ikkattuaq Migratory Bird Sanctuaries through the Irniurviit Area Co-Management Committee (ACMC).

Part of the Irniurviit ACMC's mandate is to write management plans for the Migratory Bird Sanctuaries. The Irniurviit ACMC completed draft management plans and is holding community meetings **26-28 November 2019 in Coral Harbour at the Parish Hall** to discuss their content and get feedback. Everyone is welcome! Choose among these options:

Tuesday, 26 November: 7:00PM
Wednesday, 27 November: 2:00PM or 7:00PM
Thursday, 28 November: 2:00PM or 7:00PM

We realize that not everyone can attend these meetings. The Irniurviit ACMC is also welcoming input on the management plans via email. We ask that you provide all written input by **Thursday, 28 November 2019** to JF Dufour, Vice-Chair of the Irniurviit ACMC (jean-francois.dufour2@canada.ca; 867-669-4766).

Sincerely,

Noah Kadlak
Chair, Irniurviit ACMC

Community Meetings

about the draft management plans for Qaqsauqtuuq and Ikkattuaq Migratory Bird Sanctuaries

ᐃᖃᓗᐃᑦ ᐃᓕᓴᕐᕕᐃᑦ ᐊᐅᓚᖑᔪᐊᖅᑕᐅᓯᖏᑦ ᑲᑎᒪᔨᖅᑕᐅᓯᖏᑦ

IRNIURVIIT AREA CO-MANAGEMENT COMMITTEE



- See the content of the draft Management Plans and provide your input
- Meet the Irniurviit Area Co-management Committee members
- Refreshments will be served
- Share your knowledge and stories about Qaqsauqtuuq and Ikkattuaq
- Learn more about the Migratory Bird Sanctuaries
- Door Prizes

Everyone welcome!

Where: **Parish Hall**

When: **Nov 26 - Open House 1:30-4:30**
Nov 27 - Presentation at 2:00pm and 7:00pm
Nov 28 - Presentation at 2:00pm and 7:00pm

~ For more information call Ron Ningeongan (KIA) at 867-925-8136 ~

ᓄᓇᕗᒻᒥ ᑲᑎᒪᓴᒃᑎᑲᓂᖅ

ᑎᑎᕆᖃᐅᔨᓕᕆᓂᖅᐳᑦᓄᑦ ᐊᐅᓚᕆᑎᓴᒃᑎᑲᓂᑦ ᐸᕐᓇᐅᑎᒃᓴᓗᑦ ᖃᖅᖢᓴᐅᖅᑐᕐᒪᑦ ᐊᓪᒪᓗ ᐃᑲᑕᐊᕐᒪᑦ ᑎᒥᕆᐊᖃᖅᓯᒪᓂᑦ

ᐃᓕᓴᕐᓂᖅ | IRNIURVIIT AREA
ᐊᐅᓚᕆᑎᓴᒃᑎᑲᓂᕐᒪᑦ | CO-MANAGEMENT
ᑲᑎᒪᓴᕆᓚᓕᕆ | COMMITTEE



- ᑕᑯᓗᒋᑦ ᐃᓗᓕᖅᓯᒋᑦ ᑎᑎᕆᖃᐅᔨᓴᒃᖢᒪᓕᖅᓯᓗᑦ ᐊᐅᓚᕆᑎᓂᕐᒪᑦ ᐸᕐᓇᐅᑎᓴᑦ, ᐃᓱᒪᒋᔭᑎᑦᓗ ᐅᖃᐅᔨᓱᓚᓄᑦ
- ᑲᑎᖃᑎᒋᓗᒋᑦ ᐃᓂᓕᕆᓂᖅᓴᑦ ᐊᐅᓚᕆᑎᓴᐅᖃᖅᑕᐅᓂᕐᒪᑦ ᑲᑎᒪᓴᕆᓚᓕᖅᓯᖏᓐᓂᑦ
- ᑕᒪᐊᕐᒥᖃᖅᓂᐅᓂᖅᓯᓗᑦ ᐃᒋᑐᓂᓕᓐᓄᓗ ᑭᒋᓗ

- ᖃᐅᐅᓚᒋᔭᓂᒃ ᐅᖃᐅᔨᓱᖃᓂᐊᖅᐳᑎᑦ ᐅᓂᒃᑲᓂᓕᓐᓄᓗ ᑐᓂᒃᑲᓂᓕᓐᓄᓗ ᐅᖃᐅᔨᓱᓚᓄᑦ
- ᐃᓕᓴᐅᒋᔭᖅᓂᒃ ᐊᓪᒪᓗᑭᓇᐅᑎᓴᒃᑎᑲᓂᕐᒪᑦ
- ᖃᒃᐃᓕᒋᔭᖅᓂᒃ ᐊᒧᓱᖃᒋᓂᐊᖅᐳᑎᑦ ᑐᓂᒃᑲᓂᓕᓐᓄᓗ ᐅᓂᒃᑲᓂᓕᓐᓄᓗ

ᑕᒪᐃᓐᓂᑎᐊᖅ ᐃᓄᐃᑦ ᑐᖓᓱᓕᕆᔨᒃᑯᑦ!

ᓴᓪᓕᓐᓂ: ᐊᓱᖏᖅᑐᐃᐅᔭᒃᑯᑦ ᑲᑎᙵᖓᓂ

ᖃᒃᖓᒐ: ᓄᙵᐱᒃ 26 - ᐃᓄᖓᓄᑦ ᒪᑐᐃᖓᑎᓪᓗᑎᓂᖅ 1:30-ᒥᒃ 4:30-ᒧᑦ ᐅᑦᓗᑯᑦ
ᓄᙵᐱᒃ 27 - ᐅᓂᒃᑲᓕᐅᔨᓂᖅ 2:00-ᒥ ᐅᑦᓗᑯᑦ ᐊᓪᒪᓗ 7:00-ᒥ ᐅᖃᓄᑯᑦ
ᓄᙵᐱᒃ 28 - ᐅᓂᒃᑲᓕᐅᔨᓂᖅ 2:00-ᒥ ᐅᑦᓗᑯᑦ ᐊᓪᒪᓗ 7:00-ᒥ ᐅᖃᓄᑯᑦ

ᑐᑭᓯᐊᕆᑎᐊᒃᑲᖃᓂᑕᒪᔭᙵᓕᑦ ᐅᖃᖅᙵᐅᔨᓴᖃᖅᑐᑐᒃᖢᖅ ᖁᓕ ᓂᕆᑭᐅᓚᑦ (ᑭᓲᓕᓕᕐᒥ ᐃᓄᐃᑦ) ᐅᓲᓂ 867-925-8136

EVENT: IRNIURVIIT ACMC COMMUNITY MEETING	LOCATION: PARISH HALL
DATE: NOVEMBER 26, 2019 OPEN HOUSE	

LIST OF ATTENDEES

NAME (PRINT FIRST AND LAST NAME)	NAME (PRINT FIRST AND LAST NAME)
1 Maria Kadlak	26
2 Lucassie Nakoolak	27
3 Elizabeth Nakoolak	28
4 Louisa Kudluk	29
5 Kidlapik Alogut	30
6 Emily Eetuk	31
7 Sandy Ruttat	32
8 Anthony Katalik	33
9 Janet Panigoak	34
10	35
11	36
12	37
13	38
14	39
15	40
16	41
17	42
18	43
19	44
20	45
21	46
22	47
23	48
24	49

EVENT: IRNIURVIIT ACMC COMMUNITY MEETING	**LOCATION:** PARISH HALL
DATE: NOVEMBER 27, 2019 2PM	

LIST OF ATTENDEES

NAME (PRINT FIRST AND LAST NAME)	NAME (PRINT FIRST AND LAST NAME)
1 LOUISA KUDLUK	26
2 Mark Panipak	27
3 Elijah Katsuk.	28
4 Raymond Kudluk	29
5 ELIZABETH NA[illegible]	30
6 Emily Ettuk	31
7 Sandy Pullut	32
8 [illegible]	33
9 NOAH KATALUK	34
10 [illegible]	35
11 [illegible]	36
12	37
13	38
14	39
15	40
16	41
17	42
18	43
19	44
20	45
21	46
22	47
23	48
24	49

EVENT: IRNIURVIIT ACMC COMMUNITY MEETING	LOCATION: ~~[illegible]~~ HAMLET
DATE: NOVEMBER 27, 2019 7PM	

LIST OF ATTENDEES

	NAME (PRINT FIRST AND LAST NAME)		NAME (PRINT FIRST AND LAST NAME)
1	Wayne Kudluk	26	
2	NOAH KATALUK	27	
3	CINDY NINGEONGAN	28	
4	Logan Kadlak	29	
5	UNREAD	30	
6	Lizzie Kadlak	31	
7	LUCASSIE ALAKOMIAK	32	
8		33	
9		34	
10		35	
11		36	
12		37	
13		38	
14		39	
15		40	
16		41	
17		42	
18		43	
19		44	
20		45	
21		46	
22		47	
23		48	
24		49	

EVENT: IRNIURVIIT ACMC COMMUNITY MEETING	**LOCATION:** PARISH HALL
DATE: NOVEMBER 28, 2019 2PM	

LIST OF ATTENDEES

NAME (PRINT FIRST AND LAST NAME)	NAME (PRINT FIRST AND LAST NAME)
1	26
2	27
3	28
4	29
5	30
6	31
7	32
8	33
9	34
10	35
11	36
12	37
13	38
14	39
15	40
16	41
17	42
18	43
19	44
20	45
21	46
22	47
23	48
24	49

EVENT: IRNIURVIIT ACMC COMMUNITY MEETING	**LOCATION:** PARISH HALL
DATE: NOVEMBER 28, 2019 7PM	HAMLET

LIST OF ATTENDEES

NAME (PRINT FIRST AND LAST NAME)	NAME (PRINT FIRST AND LAST NAME)
1 Unnecd	26
2 Lizzie Shapcott	27
3 Lizzie K	28
4	29
5	30
6	31
7	32
8	33
9	34
10	35
11	36
12	37
13	38
14	39
15	40
16	41
17	42
18	43
19	44
20	45
21	46
22	47
23	48
24	49

ᑎᑎᕐᖅᑕᐅᓂᒃᑲᓂᐊᓂᖅᑭᑦ ᓄᓇ

ᐊᐅᓪᓗᑎᓂᕐᒧᑦ ᕿᕐᓇᐅᑎᖅᑲᐃᑦ

ᖃᖅᖁᐅᖅᑐᕐ ᐊᒻᒪᓗ ᐃᑲᑦᑐᐊᕐᒥ ᑎᖅᒥᐊᖃᕐᓂᐊᕐᑎᑦ

Draft Management Plans

Qaqsauqtuuq (East Bay) and Ikkattuaq (Harry Gibbons) Migratory Bird Sanctuaries

ᓴᓪᓗᖅ, ᓄᕆᐱᓗ 26-28, 2019

Coral Harbour, 26-28 November 2019

ᐃᓕᓴᕐᕕᐃᑦ ᐃᓄᐃᑦ ᐊᐅᓚᓂᖅᑐᑦᑕᐅᓂᒃᑲᓂᕐᒧᑦ ᑲᑎᒪᔨᖏᑦ

IRNIURVIIT AREA CO-MANAGEMENT COMMITTEE

ᐅᓂᒃᑲᓕᐊᓂᒃ ᑕᒪᐃᓐᓂᒃ ᑐᑭᓯᑎᕆᑎᑭᐊᕐᓂᖅ
Presentation Overview

- ᖃᐅᔨᒪᑎᕆᑎᐠᓯᑎᒃᓴᐃᓕᕆ ᓴᐳᓐᓂᐊᖅᑕᐅᓯᒪᓂᖏᓐᓂᒃ ᓄᓇᐃᕆ ᓄᓇᕗᓴᒥ
- ᑕᑯᒃᓴᐅᑎᕆᑎᔪᕆ ᐃᓗᓕᖏᓐᓂᒃ ᑎᑎᕋᖅᑕᐅᕙᓪᓕᐊᓂᖏᓐᓂᒃ ᓯᓕ ᐊᐅᓪᓚᕆᑎᓂᕐᒧᑦ ᐸᓇᐅᑎᒃᓴᓂᒃ
- ᐃᓯᒪᒋᔭᐠᓯᓐᓂᒃ ᑕᒪᒃᑯᓂᖓᓐᓄ ᑐᓴᖅᑕᐅᔨᒪᖓᓐᓄᑦ ᑎᑎᕋᑕᐅᕙᓪᓕᐊᓂᖏᓐᓂᒃ ᓯᓕ ᐊᐅᓪᓚᕆᑎᓂᕐᒧᑦ ᐸᖅᓇᐅᑎᒃᓴᐃᓕᕆ

- Information on protected areas in Nunavut
- Show you the content of the Draft Management Plans
- Get your input and comments on the Draft Management Plans



ᒥᑎᕕᒃ Common Eider



ᑐᖏᕕᑎᑕᖅ Ruddy Turnstone

ᖃᐅᔨᒪᑎᑦᑎᓕᐊᕈᑎᑦ ᑲᓇᑕᒥ
ᐅᒪᔨᖃᖅᖁᐅᑎᑕᐅᔨᓂᒃ ᐊᓚᒪᓗ
ᑎᖏᒥᐊᖃᖅᖁᐅᑎᑕᐅᔨᓂᒃ ᓄᓇᕗᓚᒥ

Information on National Wildlife Areas and Migratory Bird Sanctuaries in Nunavut

ᐊᓂᑎᓪᓗᖅᐸᐃᑦ ᑲᓇᑕᒥ ᓄᓪᓕᐅᕐᓂᓗ
ᐊᓯᔾᔨᖅᐸᓪᓚᐊᓂᖏᓐᓂᑦ ᑲᓇᑕᒥ
Environment and Climate Change Canada

ᐱᔾᔨᑕᐅᓂᖅᑎᑦ ᐱᓚᑦᖃᑎᒋᔭᑕᐅᓂᖅᓂᕐᒧᑦ:


- ᓴᐳᓄᐊᖅᑕᐅᓯᒪᔪᑦᐊᖃᖅᓂᖅᑎᓄᑦ
- ᖃᐅᔨᓴᖅᑕᐅᖃᑕᑦᐊᖅᓂᖅᑎᓄᑦ
- ᑐᑭᓯᓇᖅᓯᑎᑕᐅᔪᑦᐊᖃᖅᓂᖅᑎᓄᑦ

ᓄᓇᑕᒥ ᐅᒪᔨᖃᖅᓴᑦ ᓴᐳᔾᔪᐅᓯᒪᓕᕐᓇᖅᑎᑦ
ᑎᖅᑎᐊᖃᖅᓴᑦ ᓴᐳᔾᔪᐅᓯᒪᓕᕐᓂᖅᑎᑦ

Purpose of Network

- Conservation
- Research
- Interpretation

National Wildlife Areas (NWA)
Migratory Bird Sanctuaries (MBS)



ᐊᖑᓕᓂᖅᑎᑦ 12 ᒥᓕᒃᖑᓪ
ᐄᐃᐊᖅᑕᓂᖅ ᑕᒪᖏᖅ ᓄᑎᓪᓗᓕᑦ
ᐊᖑᓕᖑᓕᑦ
12 million hectares in total


ᐊᐢᑎᓕᓗᓴᕐᓯᐊᖏᑦ ᑲᓇᑕᒥ ᓯᓚᐅᑦᓗ ᐊᓱᓴᖅᐸᑦᓕᐊᓂᖅᓯᒪᓂᑦ ᑲᓇᑕᒥ
Environment and Climate Change Canada

	ᑎᖅᑎᐊᖅᑲᐃᑦ ᓴᐳᔪᐅᓯᒪᕐᓇᖅᑐᑦ	ᑲᓇᑕᒥ ᐅᒪᓯᖅᑲᐃᑦ ᓴᐳᔪᐅᓯᒪᕐᓇᖅᑐᑦ
ᓴᐳᓂᐊᑭᑕᐅᓂᖅᔪᑦ ᑐᕆᓚᖅᖃᖅᑐᑦ	ᑎᖅᑎᐊᑦ	ᑕᒪᖅᑭᑲ ᐅᒪᔨᐃᑦ
ᓴᐳᓂᐊᖅᑕᐅᓯᓂᖅᐳᓇᓂᑦ ᐊᑐᖅᓂᖅᑎᑦ	ᐅᓪᓗᖅᑲᖅᓇᖅᑐᑦ /ᐃᓕᓴᓇᖅᑐᑦ/ ᓇᓕᐊᖅᓇᖅᑐᑦ	ᐊᖅᖁᔭᓕᒫᖅ
ᒪᓕᓚᖅᑎᔪᑦ	*ᑎᖅᑎᐊᖅᑲᓕᓚᓂᓂᖅᑭᖅ ᑲᑎᒪᖅᑲᐃᑦᑕᐅᓯᒪᔨᑦ ᒪᓕᓚᐃᐃᑦ*	*ᑲᓇᑕᒥ ᐅᒪᔨᓕᓚᓂᓂᖅᔪᑦ ᒪᓕᓚᐃᐃᑦ*

	MBS	NWA
Protection Focus	Migratory birds	All wildlife
Protection Period	Nesting/breeding season	Year-round
Legal Basis	*Migratory Birds Convention Act*	*Canada Wildlife Act*

ᐃᓄᐃᑦ ᐊᖏᑐᕐᑕᐅᓯᒪᓂᖅᐸᓂᖅ ᐱᓕᕆᓂᕐᑕᐅᔨᓐᓇᖅᑐᓂᒃᑲᓂᕐᓗ ᐊᖏᕈᓐᓇ (2008; 2016)

Inuit Impact and Benefit Agreement (2008; 2016)





ᐃᓄᐃᑦ ᐊᖏᑐᖅᑕᐅᓯᒪᓂᖅᐸᓄᓄᑦ ᐱᓵᓕᕐᑕᐅᔨᓴᖅᑐᓂᒃᑐᓂᒃ ᐊᖏᑎᑦᑎᐊᓕ (2008; 2016)

Inuit Impact and Benefit Agreement (2008; 2016)

- ᐃᓱᒪᓕᐅᑭᑕᐅᓴᖅᑐᑦ ᐱᒻᓱᑎᖅᑲᖅᑐᑦ ᑎᖅᑭᐊᖅᑲᐃᓂᒃ ᐊᒪᓗᓗ ᐅᒪᔨᖅᑲᐃᓂᒃ ᓴᖅᑭᔨᒥᒃ ᐃᓄᐃᑦ ᖃᐅᔨᒪᔭᖅᑐᖅᑲᖅᖢᑦ ᐃᓄᐃᑦ ᖃᐅᔨᓴᖅᑲᖅᑲᓐᓂᖅᓯᒪᔪᖅ

- ᐃᓕᑕᒫᒃᐅᓱᒪᓴᑦ ᐊᐅᓪᓚᑎᐅᖅᑕᐅᒃᓗᐊᖅᑲᓂᖅᑭᑦ ᐃᓄᐃᑦ ᑎᖅᑭᐊᖅᑲᐃᓂᒃ ᐊᒪᓗᓗ ᐅᒪᔨᖅᑲᐃᐅᔨᓂᒃ ᐱᒻᒪᓗᐅᓂᖅᑭᑦ ᐊᖏᑎᑦᑎᐊᓕ ᐃᓄᐃᑦ ᑎᖅᑭᐊᖅᑲᐃᓂᒃᑐᓂᒃ: ᐊᐅᓪᓚᑎᐅᖅᑕᐅᓂᖅᓴᓪᒧᑦ ᑲᑎᕕᔪᕐᓇᑦ


- Decisions for Migratory Bird Sanctuaries and National Wildlife Areas strongly influenced by **Inuit Qaujimajatuqangit**

- Guarantees co-management of Migratory Bird Sanctuary and National Wildlife Areas
  Important feature: **Area Co-Management Committees**

ᐊᐅᓪᓗᕆᐅᖅᑕᐅᓂᖏᓐ ᑲᑎᒪᔨᕋᓪᓕᐅᑦ
Area Co-Management Committees (ACMCs)

- 9-ᖃᓄᑦ ᐊᐅᓪᓗᕆᐅᖅᑕᐅᓂᖏᓐ ᑲᑎᒪᔨᕋᓱᒃᑐᖅᑎᑦ ᓄᓇᕗᒻᒥ, ᐊᐅᓪᓗᕆᐅᖅᑕᐅᓯᒪᔪᑦ 13-ᖃᓄᓂᒃ ᓴᓇᔨᓂᐊᖅᑕᐅᓕᒫᖅᓂᐊᓂᒃ ᓄᓇᓂᒃ
- ᑲᑎᒪᔨᖅᓯᖃᓪᓚᑎᒃ ᑕᒪᓕᒪᓂᒃ 5 ᐃᓄᖃᓂᒃ ᖁᓂᑭᖏᑐᓂ ᓄᓇᓕᓂᖃᖅᓯᒪᔪᓂᒃ ᐊᓪᒪᓗ ᐃᓕᖁᖃᓗᑎᒃ ᐊᑕᐅᓯᕆᑦ 1 ᑲᓇᑕᒥ ᐅᒪᔨᓕᕆᔨᒋᑦ ᐱᓕᕆᓚᖏᑐᓂ (ᐊᓯᑎᓕᕆᔨᒋᐊᓂᒃ ᓯᓚᐅᒪᓗ ᐊᓯᔨᕐᓴᒪᓕᐊᓂᖏᓐᓂᒃ)

- 9 ACMCs were created in Nunavut to co-manage the 13 protected areas
- Made up of 5 people from the associated community and 1 from Canadian Wildlife Service (Environment and Climate Change Canada)



- ᐃᖃᓗᐃᖅᓯᑦ ᐊᐅᓪᓚᕆᐅᖅᑕᐅᓂᖃᑦ ᑲᑎᒪᔨᖅᓯᒪᔪᑦ ᑲᑎᒪᔨᖏᓪᓗᐅᖅᑐᑦ 2012-ᒥ
- ᑕᕐᓕᒪᓂᒃ 5 ᑲᑎᒪᔨᖅᖅᑐᑦ ᓴᕐᓕᖅᑎᐅᑕᓂᒃ
  - ᓄᐊ ᑲᓕᓪᒃ (ᐃᒃᓯᐊᐅᑕᖅ)
  - ᐊᔪᓂ ᓂᖏᑎᐅᖏᓐᓄᑦ
  - ᓗᐃᓴ ᑲᓕᓪᒃ
  - ᐅᐃᓕ ᐊᑕᒪᓴ
  - ᐊᑕᐅᓯᖅ ᐃᓄᖅᖏᑎᑐᖅ
- ᐊᑕᐅᓯᖅᑎᒃ 1 ᑲᓇᑕᒥ ᐅᒪᔨᓕᕆᔨᒃᑯᑦ ᐱᓕᕆᓚᖓᓂ ᑲᑎᒪᔨᖅᕆᓂᕐᒧᑦ: ᖃᐅᔨ-ᐃᐊᐟ ᑐᐅᐊ (ᐃᒃᓯᐊᐅᑕᐅᐟ ᑐᖏᑎᓚᖓᓂᒃ)
- ᐃᑲᔪᖅᑕᐅᔪᖅ ᐅᒧᖏᒐ ᖃᓄ ᓂᖏᐅᖏᓚᓄ (ᑭᐊᓕᓪᒥ ᐃᓄᐃᑦ ᑲᑐᔨᖅᑎᑎᑦ)

- Irniurviit ACMC was formed in 2012
- 5 members from Coral Harbour
  - Noah Kadlak (Chair)
  - Annie Ningeongan
  - Louisa Kudluk
  - Willie Adams
  - Vacant
- 1 member from Canadian Wildlife Service: JF Dufour (Vice-Chair)
- Supported by Ron Ningeongan (KIA)



2012



2018

ᐃᖏᓂᐅᕐᕕᐊᑦ
ᐊᐅᓪᓗᓯᐅᖅᑕᐅᓂᖓᓪᓚᑦ
ᑲᑎᒪᔨᖏᓪᓕᖅ


IRNIURVIIT AREA
CO-MANAGEMENT
COMMITTEE

ᒪᖏᕈᒃ ᑎᖅᑭᐊᖅᑲᓇᐃᒃ
ᓴᐳᓂᐊᖅᑕᐅᓯᒪᖕᓇᐅᔭᓕᑎᒃ:

- ᖃᓄᖅᓴᐅᖅᑐᖅᑎ
- ᐃᓄᓕᑐᐊᖅᑎ

Two Migratory Bird Sanctuaries:

- Qaqsauqtuuq
- Ikkattuaq



ᐊᐅᓪᓗᓂᐅᖅᑕᐅᓂᖏᑦ ᑲᑎᒪᔨᖏᓪᓕᓪᓂ
Area Co-Management Committees (ACMCs)

ᐊᐅᓪᓗᓂᐅᖅᑕᐅᔾᔨᓂ ᑲᑎᒪᔨᖏᓪᓕᓯᓪᓂ
ᐃᒪᓐᓇᐃᓕᐅᖅᑕᕐᓂᓕᕆᖅᑐᑦ:

- ᐅᖅᐅᓴᔨᕆᕕᖅᑎᐅᖅᑕᕐᓂᔭᓐᓄᑦ ᒥᓂᔾᑕᒧᑦ ᑕᒪᐃᓐᓇᓂᕆᕕᖅ ᐱᔾᔪᑎᖅᖅᑐᓂᑦ ᑎᓪᒥᐊᖅᕿᓐᓂᓂᑦ/ᐅᒪᔪᖅᕿᓐᓂᓂᑦ ᐊᐅᓪᓗᓂᑎᓂᕐᒥᑦ
- ᖃᐅᒥᖅᑕᐊᖅᑕᕐᓂᔭᓐᓄᑦ ᐱᔪᓐᓇᐅᑎᑦᓴᑕᕐᓂᔪᓐᓇᕐᓂᖏᑦ ᐅᑦᑐᖅᑕᐊᖅᑎᓂᑦ ᑐᑦᔪᖅᑐᐅᑎᓂᑦ
- **ᓴᓇᖅᑕᕐᓂᔭᓐᓄᑦ ᐊᐅᓪᓗᓂᑎᑦᔪᑎᑦᓴᓂᑦ ᐸᖅᓇᐅᑎᓂᑦ**



ACMC will:

- advise the Minister on all aspects of MBS/NWA management
- review permit applications
- **develop Management Plans**



ᓯᓇᐅᖏᒪᑕ ᐊᐅᓪᓕᑎᐱᔨᑎᖅᓴᓕᓂᑦ ᐸᖕᓇᐅᑎᑦ?

What is a management plan?

- ᐱᔨᓴᓇᐅᑕᐅᐳᖅᑐᑦ ᐊᐅᓪᓕᑎᐱᔭᖑᓇᐅᑕᐅᓂᖕᒧᑦ ᑲᑎᒪᔨᕆᓪᓕᑦ ᐅᖃᐅᔨᖑᕐᓇᖕᓂᖏᓐᓂᖅ ᑕᑯᑎᑕᑎᐱᓴᓇᖕᓂᒃ ᑕᐅᑐᒪᓗᓇᖕᓂᖕᒧᑐ ᖃᓄᐃᓕᖏᓐᓄᓚᐊᖃᖕᓂᖏᓐᓂᖅ ᓴᐳᓴᐊᖃᑕᐅᔨᒪᔨᑦ
- ᓇᓗᓇᐃᔭᐃᔨᐅᐸᓗᓗᑎᒃ ᒪᓕᒃᑕᐅᓂᐊᖃᖕᒥᓂᒃ ᐃᔨᒪᓕᐅᖃᑎᒌᖏᓐᓂᖃᖅ ᐱᔭᐸᑎᖃᖕᒥᓂᒃ ᓴᐳᓴᐊᖃᑕᐅᔨᒪᔨᓂᖅ ᓄᓇᓂᒃ
- ᐅᓂᒃᑲᖅᐸᒃᑐᒥᑦ ᐱᓕᒪᓗᐅᓂᖏᓐᓂᖅ ᐃᓕᓐᓄᔨᑕᖅᑎᑦ ᐊᐸᑎᖏᓐᓂᓕᓪᓗ ᖃᓄᐃᓕᖏᓐᓄᓂᖏᓐᓂᖅ
- ᐅᓂᒃᑲᖅᔨᒪᑕᑎᐊᖅᑐᑦ ᖃᓄᖅ ᐱᓕᓗᐊᐅᔨᓇᖕᓂᖏᓐᓂᖅ ᓄᓇᖏᑦ ᐊᓪᒪᓗ ᖃᓄᖅ ᐱᓕᓗᐊᐅᔨᓇᓐᖏᓐᓂᖏᓐᓂᖅ ᓄᓇᖏᑦ

- Allows the Area Co-Management Committees to share their vision of the protected area
- Guides decision making for the protected area
- Describes important cultural and environmental aspects
- States which activities are permitted and not permitted





ᐊᐅᓪᓗᐊᑎᒃᑲᐃᓕᑦ ᓱᖏᓐᓄᐅᑎᑦ: ᐱᓕᓀᐊᖅᓯᓂᓕᐊᖏᑕᐃᓇᖏᓐᓂᑦ
Management Plans: *Steps*

ᐱᓯᓂᓕᐊᓂᖅᓂᓐᓂ ᓯᓚᓐᓂᖅ 1

ᖃᐅᔨᒪᔨᕐᑎᑲᓇᓂᐊᖏᑕᖅᓂᓂᑦ ᑲᑎᖏᓯᐃᓂᖏᖅ

ᐱᓯᓂᓕᐊᓂᖅᓂᓐᓂ ᑐᑭᓂᖅ 2

ᑐᑭᓂᐅᖏᑕᐅᓗᑎᑦ ᑕᐅᑐᑐᓀᒃᐅᔨᓂ ᓯᓚᓂᓂᒃᓕᒍᑦ, ᑐᖏᓀᓀᒃᐅᔨᓂ, ᐃᓂᖏᑕᐅᓯᒪᓂᐊᖏᑐᑐᓂᓗ

ᐱᓯᓂᓕᐊᓂᖅᓂᓐᓂ ᐱᖓᓱᖓᑦ 3

ᑎᑎᕋᖏᑕᐅᓯᓂᓕᓗᑎᑦ ᐊᐅᓪᓚᑎᓂᓀᓂᐊᖏᑕᖅᓂᕐᓂᓄᑦ ᓴᖃᐅᑎᑦᑕᖅᓂᑦ

ᐱᓯᓂᓕᐊᓂᖅᓂᓐᓂᑦ ᓯᑕᒪᖓᑦ 4

ᓄᓇᓕᖅᓂ ᑲᑎᒪᓂᑦᑎᓂᖅᓂᑦ ᐱᓵᓯᑎᑭᓗᑦ ᑎᑎᕋᖏᑕᐅᔭᖏᖅᓯᒪᓯᑦ ᐊᐅᓪᓚᑦᑎᓯᑲᓇᖅᓂᑦ ᓴᖃᐅᑎᖏᓐᓂᓂᑦ

ᐱᓯᓂᓕᐊᓂᖅᓂᓐᓂ ᑕᖅᓚᒪᖓᑦ 5

ᓄᓇᓄᓐᒥ ᐅᒪᔨᓚᓗᔨᔨᐊᐳᑦ ᑲᑎᒪᔨᖏᓐᓄᓪᓕᑦ ᐊᓐᒪᓗ ᑲᓇᑕᐅᐸ ᒥᓂᑕᖓᓄᓪᓕᑦ ᐊᕿᑎᓚᓗᓂᔪᑦ ᐊᖏᕈᑕᐅᓗᑎᑦ

ᐊᑐᓚᖅᑎᑕᐅᓗᑎᑦ ᐊᐅᓪᓚᑦᑎᓯᑲᓇᖅᓂᑦ ᓴᖃᐅᑎᖏᓐᓂᑦ

Step 1

Collect Information

Step 2

Articulate Vision, Goals, and Objectives

Step 3

Draft the Management Plan

Step 4

Community Meetings
about the draft Management Plan

Step 5

Nunavut Wildlife Management Board and
federal Minister of the Environment Approvals

Implement the Management Plan

ᑕᒪᔨᓂᖅ ᐅᖃᐅᓯᖅ
Where we are

ᐃᓗᓕᓐᖏᑦ ᑎᑎᕐᖅᑕᐅᓂᓕᕆᔨᑦ ᓄᓕ ᐊᐅᓚᑦᑎᐊᕐᓂᖅᓴᓂᑦ ᓴᖅᑭᐅᖅᑎᑦ ᖃᖅᑳᐅᖅᑐᕐᒪᑦ ᐊᒻᒪᓗ ᐃᑲᑦᑐᐊᕐᒪᑦ ᑎᓪᓗᐊᖅᑲᕐᔪᓂᑦ

Content of the Draft Management Plans for the Qaqsauqtuuq and Ikkattuaq Migratory Bird Sanctuaries

ᐃᓗᓕᓐᖏᑦ ᐊᐅᓪᓚᑎᑦᑎᐸᒋᑎᖅᑲᐃᑦ ᓯᖏᐅᑎᒃ
Content of the Management Plans

- ᐅᓂᖅᑲᑦᑎᐊᖅᑐᑦ ᑎᖏᒥᐊᖅᖏᓐᓇᑦ ᖃᓄᐃᓕᓐᖑᓂᖏᓪᓂᖅ
- ᐊᓭᑎᖑᓂ ᓄᓇᒥᐅᑕᐃᓕᓴᐅ ᐅᒪᔪᐃᓕᓴᐅ ᐱᑕᖅᖏᓂᖏᓪᓂᖅ ᑕᒪᓂ
- ᐱᐅᔪᑐᖅᖅᑎᒍᑦ ᐊᓭᑎᒥᐅᑕᖏᓐᖏᑦ ᔪᓇᑕᖅᖏᓂᖏᓪᓕᓴᐅ
- ᑕᐅᑐᒪᓱᐅᔪᑦ, ᑐᖃᓱᓱᐅᔪᑦ ᐊᓚᒪᓗ ᐱᓕᓱᐊᓂᔪᒪᓂᐊᖅᑐᑦ
- ᐊᐅᓪᓚᑎᓂᕐᒧᑦ ᐃᓕᒪᖏᖅᔪᐅᑭᑎᖏᑦᑲᐃᑦ - ᔪᓇᑐᐃᓵᖏᑲᐃᑦ ᐃᓕᒪᖏᖅᔪᐅᑭᑕᐅᓂᖏᓪᑎᑦ ᐊᐅᓪᓚᑎᓂᕐᒧᑦ ᑎᖏᒥᐊᖅᖏᓐᓄᓂᓂ

- Description of the Migratory Bird Sanctuaries
- Ecological resources
- Cultural resources
- Vision, goals and objectives
- Management Considerations – things to consider in managing the Migratory Bird Sanctuaries





ᐱᓪᓕᐅᓴᖅ **Black Guillemot**



ᐃᓗᓕᓐᖏᑦ ᐊᐅᓪᓚᑎᐱᑎᖅᓴᐃᑦ ᐸᖕᓇᐅᑎᓂᑦ
Content of the Management Plans

- ᐊᐅᓪᓚᑎᓂᖕᒪᑦ ᖃᓄᐃᓕᓐᖓᓂᐊᖕᓂᖏᑦ - ᖃᓄᐃᓕᓐᐅᖕᓂᐊᖕᒪᖓᑕ ᑎᑭᐅᑎᕐᒪᓂᖕᒪᑦ ᑐᖓᒪᓗᐅᔪᓄᑦ ᐊᓱᒪᓗ ᐱᓕᕆᐊᖅᐊᓂᔪᒪᓂᐊᖅᑐᑐᓄᑦ
- ᐊᖏᕈᑕᐅᓯᒪᔪᑦ ᐱᓕᕆᐊᖅᔨᓇᖕᓂᖅ ᑎᑭᓯᒪᖕᓇᐅᔪᖏᓐᓄᑦ ᐱᔨᓇᐅᑎᖃᕆᐊᖅᑐᑦ ᐊᓱᒪᓗ ᑭᓇᐅᔭᖓᑦ ᐱᔨᓇᐅᑎᖃᕆᐊᖅᖢᓂᖅᑐᑦ
- ᓄᓇᐃᑦ ᖃᓄᐃᓕᓐᖓᔪᒥᒃ ᑎᒃᓯᐊᖅᑕᐅᓯᒪᓂᖏᑦ: ᐃᔨᓇᐅᔪᖅ ᑎᓪᒥᐊᖅᖃᕐᖢᐅᑎᑕᐅᔪᖅ
- ᐊᐅᓪᓚᑎᐱᑎᖅᓴᐃᑦ ᐸᖕᓇᐅᑎᓂᑦ ᐊᑐᓪᖅᑎᑦᑕᐅᓂᖏᑦ ᐊᓱᒪᓗ ᐱᓕᕆᐊᖅᑐᑐᓂᖏᑦ

- Management Approaches – what we will do to achieve the goals and objectives
- Authorized Activities and Access – what activities require a permit, who requires a permit
- Site designation: currently a Migratory Bird Sanctuary
- Management Plan implementation and collaboration







ᖃᓄᖅ ᐱᓂᓪᓕᐊᒋᐊᓕᓚᐅᖅᓯᒪᓐᒪᓐᓂᑦ
Historical Background





ᖃᓄᐃᓕᐅᖅ 1959-ᓴᑎᓪᓗᒍ ᓴᓇᓂᐊᖅᑕᐅᓇᓯᑎᒃ ᖃᓴᐃᐃᑦ, ᓂᕆᓪᓇᐃᓪᓗ

“ᖃᓄᐃᓕᐅᖅ” - ᖃᓄᐃᓕᐅᖅᑐᓪᓗᒍᒃ ᑕᓯᕋᐊᖅ ᑕᒪᓂ

ᑎᑎᕋᖅᑕᐅᓭᓪᓗᐊᓂᖅᑎᑦ ᓯᓚ ᐊᐅᓪᓚᑦᑎᓗᒃᓴᐃᑦ ᐸᕋᐅᓕᖅᑎᑦ

- Established in 1959 to protect Snow Geese and Brant from disturbance
- “Kouksauktow” – Bay of red-throated loons
- Draft management plan in 1986



ᖃᓄᐃᓕᐅᖅ Red-throated loon

ᖃᓄᖅ ᖃᐅᔨ
Snow goose

ᐃᒃᑲᑦᑐᐊᖅᑎ ᑎᓪᓗᐊᖃᕐᓇᒃ
Ikkattuaq

- ᐱᓪᓕᖅᑎᑕᐅᓚᐅᖅᓯᒪᔪᖅ 1959-ᓴᑎᓪᓗᒍ
  ᓴᐅᓂᐊᖅᑕᐅᓇᓯᓚᑎᒃ ᒃᓴᐃᐃᑦ, ᖃᐅᔨᑦ, ᓂᕆᓪᓇᐃᓪᓗ
- ᐄᐃᐅᓕ ᑭᐱᓪᓴ ᐅᓇᐃᓪᓄᒃ
- ᑎᑎᕋᖅᑕᐅᓭᓪᓗᐊᓂᖅᑎᑦ ᓯᓚ ᐊᐅᓚᑦᑎᓗᒃᓴᐃᑦ ᐸᕋᐅᓕᖅᑎᑦ

- Established in 1959 to protect “Blue” Goose, Snow Goose and Brant from disturbance
- Harry Gibbons Unainnuk (c. 1900-1954)
- Draft management plan in 1986

ᓄᓇᖅᑐᑦ ᑕᐅᓂᓂᒃ ᖃᓄᐃᓕᓐᖅᓯᓂᖅᑐᑦ
Landscape





ᐃᑲᑦᑐᐊᖅᑎ ᑎᖅᑎᐊᖃᖅᖁᓕᑲ
Ikkattuaq

- ᐊᖏᕐᓂᓕᑲ 1,433-ᓂᑲ ᓴᓂᒧᑦ ᑐᑭᒧᑉᓗ ᑭᐸᑦᓴᓐᖅᑐᑎᑲ ᑕᒪᐃᓐᓂ ᑭᓯᒥᑕᓕᖅᓂᑲ, ᐊᓚᒪᓗ 190 ᓴᓂᒧᑦ, ᑐᑭᒧᑉᓗ ᑕᒪᐃᓐᓂᑲ ᑭᓯᒥᑕᓂᑲ ᑕᓗᐅᖏᒥ ᐃᒪᐃᑦ
- ᐳᐊᓴᐅᐸ-Boas ᑯᖏᓚ ᓄᓇᐅᐸ ᐃᓗᑐᓂᖏᓐᓄᑦᑐᖅ ᐊᓚᒪᓗ ᐊᑭᐊᓄᓴᓭᑎᑲ ᑎᓂᖏᓂᐊᓗᓄᖅᑐᑦ ᓭᓴᒥ ᑕᐅᓂᓐᓂᐸ ᐃᑲᑦᑐᐊᐸ ᑕᓗᐅᖏᓐ
- ᓄᓇᖏᓚ ᖃᖃᓴᖃᖏᓕᑦᑐᖅ (ᑕᒫᓂ <60m ᑭᑕᒥᑲ ᐊᖏᕐᓂᖏᒥ) ᐃᒪᐃᓴᖅᖁᖃᑦᑎᐊᖏᕐᑐᐊᓗᓇᓗᓂᓗ

- 1,433 km^2, includes 190 km^2 of marine waters
- Boas River delta and the adjacent tidal flats in the Bay of God’s Mercy
- Land is flat (<60m) & poorly drained

ᖃᖅᖁᓴᐅᖅᑐᖅᑎ ᑎᖅᑎᐊᖃᖅᖁᓕᐅᓂᖏᓐᓄᑦ
Qaqsauqtuuq

- ᐊᖏᕐᓂᓕᑲ 1,124-ᓂᑲ ᓴᓂᒧᑦ ᑐᑭᒧᑉᓗ ᑭᐸᑦᓴᓐᖅᑐᑎᑲ ᑕᒪᐃᓐᓂ ᑭᓯᒥᑕᓕᖅᑎᑲ, ᐃᓕᖃᖅᓯᓂᓗ 286-ᓂᑲ ᑭᐸᑦᓴᓐᓂ ᑕᒪᐃᓐᓂᑲ ᑕᓗᐅᖏᒥ ᐃᒪᖁᓂᑲ
- ᖃᖅᖁᓴᐅᖅᑐᖅᑎ, ᑕᑭᓂᖃᖅᑎᑎᕐᒥᑲ 50-km ᑭᓯᒥᑕᓕᖅᑎᑲ ᑲᖏᕐᓄᖃᖅᖁᖃᖅᓄᖅ
- ᓄᓇᖏᓚ ᖃᖃᓴᖃᖏᓕᑦᑐᖅ (ᑕᒫᓂ <120m ᑭᑕᒥᑲ ᐊᖏᕐᓂᓕᑲ) ᐃᒪᐃᓴᖅᖁᖃᑦᑎᐊᖏᕐᓄᐊᓗᓇᓗᓂᓗ

- 1,124 km^2, includes 286 km^2 of marine waters
- East Bay, a 50-km-long inlet
- Land is flat (<120m) & poorly drained




ᓄᓇᖅᑐᑦ ᑕᐅᔭᓂᒃᑯᓂᖅ
Landscape






ᐃᖃᓪᓗᐊᖅᑎ ᑎᖅᑎᐊᖅᑐᖃᖅ
Ikkattuaq

- ᑕᒪᕆᒃᓴᒃ ᑕᒪᓐᓇ ᓄᓇ ᑲᓇᑕᐅᔨᑦ ᓴᓇᒪᓐᓂᒃᑯᐊᓂᖅ ᓄᓇᕐᔪᑎᑎᕆᔨᓐᓂ
- ᐊᑕᐅᓯᖅ ᑭᓪᓚᐸᓐᓂᕐᒥᖅ ᐃᓄᖅᓄᑦ ᓄᓇᕐᔪᑎᑎᕆᐅᓯᖅ
    - ᓄᓇᐅᔨᑦ ᖃᖅᓴᓂᑕᓂᕐᓄᑦ ᐱᓯᓇᐅᑎᖃᖅᑐᐊᓯᖅ
- Mostly federal crown land
- 1 parcel of Inuit Owned Land
    - Surface rights





ᓄᓇᖅᑐᑦ ᑕᐅᔭᓂ ᖃᓄᐃᓕᖅᑎᑦ

Landscape






ᖃᖅᓴᐅᔪᒃ ᑎᖅᑕᐊᖃᖅᓯᐅᓂᖅᓯᓂ

Qaqsauqtuuq

- ᑕᒪᖁᑲᓴᐃᑦ ᓄᓇᖅᑦ ᑕᒫᓂ ᐃᓄᖅᓄᑦ ᓄᓇᕐᒪᑎᖅᑲᐅᔪᑦ
  - ᓄᓇᐅᔨ ᖃᖅᓱᓂᑐᑐᓂᒃ ᐱᓵᓇᐅᑎᖃᕐᓯᐅᔪᖅ
- ᑕᕆᐅᖁᒥ ᐃᒪᐃᑦ ᐊᓚᓗ ᖁᑭᖅᑕᑯᓗᒃ - ᑲᓇᑕᐅᔨ ᒪᓪᒪᖅᓵᓄᑦ ᓄᓇᒥ
- Mostly Inuit Owned Land
  - Surface rights
- Marine waters and Qikiqtakuluk - federal



ᐊᐢᑎᖏᓂ ᓄᓇᒥᐅᑕᐃᓪᓗ ᐅᒪᔨᐃᓪᓗ ᐱᑕᖅᖃᑐᑦ ᑕᒪᓂ
Ecological Resources

ᐃᒃᑲᑐᐊᖅᒥ ᑎᓯᒥᐊᖅᖁᓕᒃ
Ikkattuaq

- ᓄᓕᐊᖏᐅᓴᓂᓯᓂᖓ ᑕᒪᓂ ᑲᓇᑕᓕᒫᒥ ᐱᓪᒪᓚᐅᓂᖅᑐᑦ ᐱᑕᖅᐃᓚᓇᑲᓚᖏᓚᓄᑦᑐᓚᓂᖅᑲᐃᐃᑦ (5% ᐳᕆᓚᑎᒋᑐᑦ) ᓂᖏᓕᓚᐃᓪᓗ (1% ᐳᕆᓚᑎᒋᑐᑦ)
- ᐱᓪᒪᓚᐊᓗᔭᖅ ᑎᑭᓯᒪᑲᐃᓇᖏᖅᑕᑦᑕᖏᓂᔪᑦ ᐊᓯᖏᓚᓄᓚᓂᐊᓂᖏᓚᓂ ᓯᓴᖅᐸᓯᐅᓂᖏᒋᓯᓯᕐᓂ ᑎᓯᒥᐊᑦ
- ᐅᖃᓕᑕᓂᒥᔪᑦ >50 ᐊᐠᔨᑦᖏᑦᑐᑎᑦ ᑎᓯᒥᐊᑲᓕᐃᑦ ᑕᒪᓂᓯᓯᕐᒥᔪᑦ
- ᖏᓂᐅᔨᑦ 10 ᐅᒪᔨᐃᑦ ᐊᒥᓯᖏᓪᓗᐊᓕᖏᓂᖏᑦ

- Breeding habitat for nationally significant populations of Snow Geese (5%) and Brant (1%)
- Important stopover site in fall for shorebirds
- >50 species of birds
- 10 species at risk



ᓂᖏᓚᓇᐃᑦ **Brant**



ᖃᓄᐃ **Snow Goose**

ᐊᐢᑎᖏᓂ ᓄᓇᒥᐅᑕᐃᓪᓗ ᐅᒪᔨᐃᓪᓗ ᐱᑕᖅᖃᖅᑐᑦ ᑕᒪᓂ
Ecological Resources

ᐃᒃᑲᑐᐊᖅᒥ ᑎᖏᒥᐊᖅᖃᕐᓂᖅ
Ikkattuaq

- 19-ᖏᔪᑦ ᐊᓴᒋᖏᑐᑐᑎᑦ ᐊᒪᒪᒃᑎᑦᑎᐸᒃᑐᑦ ᐱᐊᕆᖏᑦᓂᒃ ᓄᓇᒥᐅᑕᐃᑦ ᐅᒪᔨᐃᑦ ᑕᒪᓐᓇᑦ
- ᓇᓄᖅᓯᖅ ᐊᐅᓴᒃᑯᑦ ᐊᓚᒪᓗ ᑕᒪᓂ ᐃᒡᓗᖅᓴᐅᓯᖏᓪᓗᑎᒃ ᓇᓄᕐᓄᑦ
- ᓇᓪᓴᖅᑐᖅᖏᑦᑐᕈᐅᑕᓄᑦ ᑐᒃᑐᖅᕆᖅᓴᐅᐸᒃᑐᖅ, ᐃᓕᖏᑦᓗ ᓄᕐᕆᕆᐅᖅᑕᑦᑕᖅᓯᒪᔪᓗᑎᒃ
- ᓇᑦᓯᖅᑕᖅᖅᑐᖅ ᐊᓚᒪᓗ ᐊᕐᕆᒃᑕᖅᖅᑐᖅ

- 19 species of mammals
- Polar bear summer and denning habitat
- Barren-ground caribou summer range, portions used as calving grounds in the past
- Seals and Bowhead whale



ᓇᓄᖅ **Polar Bear**



ᑐᕐᑐ **Caribou**

ᐊᐢᑎᖏᓂ ᓄᓇᒥᐅᑕᐃᓪᓗ ᐅᒪᔨᐃᓪᓗ ᐱᑕᖅᑐᖅᑐᑦ ᑕᒪᓂ
Ecological Resources

ᖃᖅᓴᐅᖅᑐᑎᒥ ᑎᖏᒥᐊᖅᖁᕕᐅᓂᖏᓂ
Qaqsauqtuuq

- ᓄᓕᐊᖁᕕᐅᔪᖅᑐᖅ ᐱᓕᓚᐅᕕᔾᒥᒃ ᑲᓇᑕᒥ ᐊᒥᓱᓄᓕ ᑲᖏᖅᓄᓕ (2% ᐳᓴᓂᑎᖏᓂᑕᓄᓕ), ᒥᑎᖁᓄᓕ (5% ᐳᓴᓂᑎᓄᓕ) ᓂᖁᓕᖁᖁᓄᓪᓗ (1% ᐳᓴᓂᑎᑦᑕᓐᓄᓕ)
- ᑲᓇᑕᐅᐸ ᐅᑭᐅᖅᑕᖅᑐᑐᓂᓗᒃᑮᖏᓂ ᐊᖏᕐᓭᖏᖅᖁᖅ ᒥᑎᖅᑕᖃᖁᕕᐅᔪᓐᓂᖅ
- ᐅᖏᓪᓕᓂᒃᓕᕆᔨᓕᑦ >90 ᐊᐢᔨᑲᖏᑦᑐᑎᑲᑦ ᑎᖏᐊᓚᐱᐅᑦ ᑕᒪᓂᓕᒃᓴᖏᖅᓕᕆᔨᓕᑦ
- Breeding habitat for nationally significant populations of Snow Geese (2%), Common Eider (5%) and Brant (1%)
- Arctic Canada’s largest single colony of Common Eiders
- >90 species of birds



ᒥᑎᐹᖅ Common Eider



ᓴᐅᖁᖅ Red Phalarope

ᐊᐢᑎᖏᓂ ᓄᓇᒥᐅᑕᐃᓪᓗ ᐅᒪᔨᐃᓪᓗ ᐱᑕᖅᖃᖅᑐᑦ ᑕᒪᓂ
Ecological Resources

ᖃᖅᓴᐅᖅᑐᖅᒥ ᑎᖏᒥᐊᖅᖁᐅᓂᖏᓂ
Qaqsauqtuuq

- ᖁᓕᐅᔨᑦ 19 ᐅᒪᔨᐃᑦ ᐊᒥᓱᖏᓪᓗᐊᓕᖁᓂᖏᑦ
- ᓄᓇᖁᓇᑦ ᐃᒡᓗᖅᖁᐊᕐᕿᖅᑐᑦ (ᓱᑎᖅᖁᐊᕐᕿᖅᑐᑦ) ᐊᐅᔪᓚᐅᓪᓗ ᓂᖏᖅᓴᖅᓱᐅᖁᐊᕐᕿᖅᑐᖅ ᓄᓇᖁᓇᑦ
- ᓄᓯᑲᖅᑐᖅᖏᑦᑐᖁᐅᑕᓄᑦ ᑐᒃᑐᖅᖁᐊᕐᕿᖅᑐᖅ, ᓇᓇᖁᓇᖅᑕᐅᕐᕿᖅᑐᖁᖁᓪᓗ ᐊᐅᔪᐅᓂᖏᓂ
- ᖃᓄᓗᒃᐃᑦ ᐅᑎᕋᐊᖅᑕᖅᓱᒪᓱᑦ (ᐃᕆᓪᓕᖁᓂᒃ), ᐊᒥᔪᖅᑕᖅᓱᑎᒃᓗᓪᓗ, ᓂᖏᖅᓴᖅᓱᐅᖁᐊᕐᕿᖅᓱᓂᓗ ᑕᒪᒃᑯᓄᖓ
- ᐊᐃᐹᖅᑕᓕᒃ ᐊᐅᔪᒃᑯᑦ ᑕᒪᐢᓭ

- 19 species at risk
- Polar bear denning and summer foraging habitat
- Barren-ground caribou calving and summer range
- Beluga calving, molting and foraging habitat
- Walrus summer foraging habitat



ᐊᐃᐹᖅ Walrus



ᖃᓕᓴᖅ Beluga

ᐊᐢᕈᑎᖏᓂ ᓄᓇᒥᐅᑕᐃᓪᓗ ᐅᒪᔨᐃᓪᓗ ᐱᑕᖃᖅᑐᑦ ᑕᒫᓂ
Cultural Resources



ᐃᓄᐃᑦ ᓴᓪᓕᐅᕐᓂ ᖃᐅᖅᑕᒪᓗᐊᖏᖅᑕᑦᑕᕐᓂᑯᓕᖅᓴᓂ ᑕᐅᓴᓴᒪᓕᕆᐅᓕᖅᑐᓂ ᐊᕐᕿᒍᓂ


- ᐊᒥᓱᒪᒪᓕᕆᐅᕐᓂᒃ ᖃᐅᔨᒪᓴᐅᔨᕐᑎᑦ ᑕᐃᔾᓱᒪᓂᓂᑕᖃᕐᓂᐊᕐᓂᖅ, ᐃᓄᖃᕐᓂᐊᕐᓚᐅᕐᓂᖏᓐᓂᕐᓂᖅ, ᐃᓕᓴᕐᓯᑐᖃᕐᓂᖅᓴ ᓇᓗᓇᐃᑯᑕᐃᑦ ᐱᑕᖃᖅᑐᑦ ᐊᑐᓗ ᐱᓴᓯᑎᐃᓂᖃᕐᓂᐊᕐᓂᖅᓱᓂ ᑕᐅᕐᓂᓐᓇ ᓄᓇᖓ

- ᑕᐃᔾᓱᒪᓂᓂᑕᐃᑦ ᖃᖓᓂᓂᑕᐃᑦ ᒥᑭᓂᑕᐃᑦ ᑎᑭᑎᐊᖃᕐᓂᐊᕐᓚᐅᕐᖅᑐᑦ ᑕᐅᕐᓂᓄᓐ



Inuit have lived on Southampton Island for thousands of years

- Numerous known archaeological sites, cultural features & artifacts in the area
- Historic trap lines



ᐊᐢᓂᑎᓂᓄ ᓄᓇᒥᐅᑕᐃᓪᓗ ᐅᒪᔨᐃᓪᓗ ᐱᑕᖅᑲᖅᑐᑦ ᑕᒪᓄ Cultural Resources



- ᐊᒥᔨᐊᓗᐃᑦ ᔨᓕ ᐃᓄᖅᕿᐅᓂᒪᐅᖅᑐᖅᑐᒃᓗᐃᑦ ᔨᓕ ᑎᑎᕿᑕᐅᔨᒪᖏᑦᑐᑦ
- ᐊᑕᐅᔨᕿᒃ 1 ᑎᑎᕿᑕᐅᔨᒪᓕᖅᑐᖅ ᒪᓕᒐᖅᑎᒍᑦ ᓄᓇᖏ ᑕᐅᓂᓄ ᖃᐹᖅᑕᑯᓗᒃ
- ᒪᕆᐣ 2 ᑎᑎᕿᑕᐅᔨᒪᓕᖅᑐᖅ ᒪᓕᒐᖅᑎᒍᑦ ᓄᓇᕗᑦ ᖁᓂᒃᑯᖏᓐᓂᑦ ᐃᓕᔨᐊᕆᔨᒃᑯᑦ


- Many archaeological sites remain unregistered
- 1 registered site on Qikiqtakuluk
- 2 registered sites near Itijuarjuk





ᑕᐃᔨᓯᒪᓂᑐᓄᖅᑎᒃ ᐃᓐᓇᒧᑦᓗ ᐃᓄᖅᓄᑦ ᓄᓇᖅᑎᑦ ᐊᑐᖅᑕᐅᖅᑕᕐᑐᖅᑐᑦ ᐃᓅᔨᐸᑎᖅᓴᓄᐅᓕᖅᓂᕐᒧᑦ

Past and current subsistence land uses



- Goose hunting and egg collection (early spring)
- Seal and walrus hunting
- Polar bear hunting
- Char fishing from lakes and rivers
- Fox trapping

- Camping
- Char fishing continued (using nets)
- Summer fishing camps were set-up at mouth of the rivers
- Seal, caribou and geese were traditionally hunted from canoes along the shoreline
- Sea mammal hunting from boats
- Collecting berries

- Fox trapping
- Caribou until the 1950s
- Fishing nets set under ice to catch char
- Hunting at the flow edge (seal, walrus)

- Securing meat for winter by hunting walrus, caribou, polar bear, and beluga whales
- Fishing with nets under new ice
- Meat caching
- Fox trapping

ᑕᐃᓴᓯᓂᑐᖅᓯᒪᖅ ᐃᓐᓇᒧᑦᓗ ᐃᓄᖅᓄᑦ ᓄᓇᖅᑎᑦ ᐊᑐᖅᑕᐅᖅᑕᕐᑐᖅᑐᑦ ᐃᓅᔨᐸᑎᖅᓴᓂᐅᕙᖅᓂᕆᒧᑦ

Past and current subsistence land uses

ᐅᓪᓗᒥᐅᓕᖅᑐᖅ, ᐊᑐᖅᑕᐅᕙᒃᑐᖅ ᓯᓚ ᐊᐅᓪᓚᐸᖅᓯᒪᓂᖅᓴᓂᐅᕙᖅᓂᕆᒧᑦ ᐊᖅᑐᓯᒪᔪᓴᖅᐱᐅᕙᖅᓂᕆᒧᑦᓗ ᐃᓚᓂᑯᓂᖅ ᖃᐅᑎᑯᓂᖅ

- ᐊᖅᓯᓇᓯᖅᑏᑦ ᖃᑐᐸᖅᑎᑎᒋᖅᓂᑦ ᐃᒡᓗᓕᕆᓚᖅᑐᑦ ᑕᐅᕙᓂ ᐃᖅᑦᑐᐊᒥᒃ ᐊᑐᖅᑕᐅᖅᑕᕐᑐᖅᑐᑦ ᐊᖅᓯᓇᓯᖅᑐᓄᓕᓂ, ᐱᓗᐊᖅᑐᒥᖅ ᓇᓂᐊᖅᓯᒪᔪᓄᓕᓂ



Today, used for camping & harvesting activities periodically


- HTO cabin located in Ikkattuaq used to support harvesting activities, in particular polar bear hunting




ᐊᐅᓪᓚᓂᖅ ᐃᓱᒪᖏᖅᓯᐅᑭᑎᖏᓚᐃᓪ
Management Considerations

ᐊᐅᓪᓚᓂᖅ ᐃᓱᒪᖏᖅᓯᐅᑭᑎᖏᓚᐃᓪ
ᐱᔨᔾᑎᖅᖅᓴᓪ ᑕᐃᒃᑯᓂᖓ
ᖃᐅᔨᒪᔨᐅᔨᓗᐊᖅᖅᑐᓂᒃ ᐃᓗᐊᓂ
ᐊᓚᓗ ᐊᕐᓗᓐᓯᓂᑐᓂᒃ,
ᓴᓴᓂᐊᖅᑕᐅᓱᒪᓂᓯᓂᓂ ᓄᓇᐃᓪ
ᐋᖏᐅᒪᓪᑎᐊᖅᑐᒥᒃ
ᐊᐅᓪᑕᐅᔨᓂᓕᖅᓯᐊᖅᒪᑕ

Management considerations are things we need to be aware of within, and surrounding, the protected area in order to effectively manage it





ᐊᐅᓪᓗᓂᔭᕐᓗᑎ ᐃᓕᓴᖅᓯᐅᑕᑎᓴᐃᑦ Management Considerations

ᐱᑕᖅᖃᑐᖅᓯᓪᓗᒍ ᐱᒻᒪᓚᐅᔨᓂᖅ ᐄᓐᓇᕆᐅᑎᑭᓕᓐᓂᒃ ᐅᓗᓚᐊᓇᑕᐅᔨᓂᖅ, ᑭᓯᐊᓂᓕᑕᐅᖅ:

1. ᑎᓪᓗᐊᖅᖃᕐᐅᓂᓪᓗᑎᑦ ᐃᓕᑕᑦᓴᐅᓂᓪᓚᑦ ᑎᒃᓯᐊᖅᑕᐅᓯᒪᓂᖅ
    - ᓴᐳᓐᓂᐊᖅᑕᐅᓯᒪᑦᑎᐊᒃᑲᓂᓚᐊᖅᖃᖅᑐᖅ
    - ᑎᓪᓗᐊᕐᓂᓐᓇᐅᔨᓐᓂᖅᑐᖅ
2. ᑲᔪᐃᑦ ᐊᒥᓯᓂᑦ ᐊᒻᒪᓗ ᓄᓇᒋᖅᑕᑦᓂᓂᓪᓚᑦ
    - ᐊᒥᓯᓗᐊᓕᖅᑐᑦ
    - ᐅᒪᔨᕐᓂᒃ ᐊᓯᖏᓐᓂᒃ ᐊᒃᑐᖅᓯᓀᓕᓕᐊᖏᒪᑕ ᐱᔭᕇᑕᐅᓗᑎᒃ ᓇᒻᒪᖏᑦᑐᒥᒃ

No significant or immediate threats, but:

1. MBS designation
    - need better protection
    - not just birds
2. Light geese population and habitat
    - too many
    - impact on other species



ᑐᑦᑐ Caribou





ᖃᓕᓗᒃ Beluga





ᑲᖏᒍ ᖃᐅᔨᖅ Snow goose

ᐊᐅᓪᓕᓂᐅᓐᓴᑦ ᐃᓱᒪᖏᖅᓯᐅᑕᑎᖏᓚᐃᓪᑦ
Management Considerations

3. ᓯᓚᐅᐟ ᐊᓯᔨᖅᐸᑦᓪᓕᐊᓂᖅ
   - ᐊᖏᑐᖅᑕᐅᓴᑦᓪᓕᐊᓴᑦ ᐅᐠᔨᕐᓇᖅᓯᖅᓯᒪᓕᑕ
   - ᐊᐅᓪᓕᓂᐅᓂᖅᖃᕐᑎᑦᑕᓄᑦ
4. ᓄᓇᓗᑦ ᐃᓄᐃᑦ ᐃᓪᓕᐅᕆᓪᓚᐊᖅᖅᑐᑦ ᐊᓖᒪᓗ ᖃᐅᔨᒪᕆᓪᓚᐊᖅᖅᓯᑎᖏ
   - ᖃᐅᔨᔨᖅᑕᐅᔪᓂᖅ
   - ᐃᓪᓕᑎᕐᐅᕆᓪᓚᐊᖅᖃᕐᓂᕐᓂᖏ ᐊᑐᖅᑕᐅᓗᑎᖏ ᐃᓄᐃᑦ ᖃᐅᔨᒪᔨᑐᖅᖃᕐᑐᓂ ᓄᓇᓕᕐᓂ ᐃᑲᔪᕐᓂᖅᓂᕐᓂᐊᖏᓪᓕᑕ

3. Climate Change
   - effects are noticed
   - out of our control
4. Community participation and awareness
   - research
   - inclusion of Inuit Qaujimajatuqangit
   - local benefits







ᐊᐅᓪᓗᓂᔭᕐᒪᑦ ᐃᓕᒪᖏᖅᓯᐅᑕᑎᖏᖅᓴᐃᓪᓗᑦ
Management Considerations







5. ᑭᓇᐅᔭᓄᑐᐃᓇᖅ ᐱᓕᕆᐊᖓᔪᓐᓇᐊᕐᓂᑦ
   - ᐅᒥᐊᖅᓯᐊᖅᑲᓐᓂᐊᓕᖅᓂᖅ
   - ᐊᒥᓱᑭᖅᐸᓐᓂᐊᓂᖓᓂᑦ ᑎᑭᐸᐸᓂᖅᑐᑦ
   - ᐊᑭᓚᑎᑦᑎᔪᓂᖅ ᐊᖓᔪᓇᓱᖓᐊᑭᐅᔨᔪᓂᓂᖅᓴᑦ
6. ᖃᐅᔨᒪᔭᐅᕆᓇᐊᓕᑦ ᐱᑕᖃᖓᓂᓂᖓᑦ
   - ᐱᐅᓱᑐᖃᖅᑎᒍᑦ ᐊᔪᑎᖓᓄᓂ ᓄᓇᒥᐅᑕᐃᓕᓐᓂ ᐅᒪᔨᐃᓕᓐᓂ ᐱᑕᖃᖅᓂᖓᑦ, ᐱᓂᐊᖅᑐᒥᐅᒃ ᐃᓄᖃᖅᓴᐅᔪᓇᒪᐅᖅᓂᖓᓂᓂᒃ ᖃᐅᒪᔪᑐᖃᐃᓕᑦ
   - ᑕᓚᐅᒥᐅᑕᓂᒃ ᐅᒪᔨᖓᑦ, ᐃᖃᓂᓕᕆᐃᓕᓐᓂ ᐊᖅᔨᑎᖓᑦᑐᑦ

5. Human Activities
   - shipping
   - increased access
   - sport hunting
6. Information missing
   - Cultural resources, especially archaeology
   - marine habitat, fish

ᐊᐅᓪᓗᓂᔭᕐᓗᑦ ᐃᓱᒪᖏᖅᓯᐅᑕᑎᖏᓚᐃᓪᓗᑦ (ᖃᖅᖏᖏᖅᑐᕐᕆᓂᖅ) Management Considerations (*Qaqsauqtuuq only*)



ᐅᓪᓚᐊᓇᑕᐅᓂᖅᐳᑦ ᒥᑎᕕᖅᖏᓚᐊᓄᑦ ᖃᐅᖅᑕᓯᔪᓐᓇᖅ

- ᓄᓂᔪᖏᕐᓂᕐᒥᒃ ᑎᖅᒥᐊᕐᓄᑦ ᐊᓂᓂᐊᖅᑕᓚᖏᐅᓂᒃᑐᑦ
- ᓇᓄᐃᑦ


Threats to eider colony at Qikiqtakuluk

- Avian cholera
- Polar bear




ᓇᓄᖅ **Polar Bear**

ᑕᐅᑐᒪᓗᓴᐅᔪᑦ, ᑐᕐᒪᓗᓴᐅᔪᑦ, ᐃᓂᖏᑕᐅᔪᒪᔪᓪᓚ
Vision, Goals, and Objectives



ᑕᐅᑐᓚᓴᐅᔨᖅ ᑎᓐᒥᐊᖅᑲᕆᓐᓄᑦ ᓴᓗᓂᐊᖅᑕᐅᓯᒪᕆᓐᓄᑦ
Vision for the MBSs



“ᑎᓐᒥᐊᖅᑲᓱᑦ ᓴᓗᓂᐊᖅᑕᐅᓯᒪᕆᓐᓄᑦ ᐊᐅᓚᑕᐅᓂᐊᖅᑐᑦ ᐅᓴᖅᑐᑦᑎᐊᕐᒪᓂᕐᒧᑦ ᐊძᓂᐅᓂᐊᖅᑐᖅ ᓴᓗᓂᐊᖅᑕᐅᓂᖏᓐᓄᑦ ᐅᓇᖏᑦ ᐊᓪᒪᓗ ᐃᒪᐃᑦ ᑕᒪᐃᓐᓄᑦ ᐱᑕᖅᑲᓗᑎᒃ ᐅᒪᔨᕐᓄᑦ, ᐃᓄᐃᑦ ᐱᔨᓐᓇᐅᑎᖅᑲᕐᓂᖏᓐᓄᑦ, ᐱᑕᖅᐃᓐᓇᕐᓂᖏᓐᓄᓗ ᑕᐃᒪᓐᓇ ᐃᓄᓂᑦ ᐃᓅᓂᕐᒥᓂᖅ, ᐃᓕᓴᕐᓂᖏᓐᓄᓗ ᑕᐃᒪᓐᓇ ᐊᑐᖅᑕᐅᓂᖅ ᐃᓄᐃᑦ ᖃᐅᔨᒪᔨᑐᖃᖅᑐᓂᖏᓐᓄᑦ ᐃᓕᓴᕐᓂᐊᕐᓗᒍ ᑕᒪᐃᓐᓇ ᐃᓄᐃᑦ ᖃᐅᔨᓴᖃᖅᑐᓂᖏᓐᓄᑦ ᐃᓕᓴᕐᓂᐊᖅᑕᐅᓯᒪᔪᓐᓇᖅᑐᑦ.”


“The MBSs will be managed to ensure the long-term conservation and protection of the land and water for all wildlife, Inuit rights and their heritage while considering Inuit Qaujimajatuqangit in all decisions.”


© Jason Caswell

ᑐᓴᓐᓂᐊᖅ Arctic Fox




ᐊᐅᓪᓂᐹᓂᒃ ᑐᖏᓐᓂᒃ
Management Goals



*“ᑎᓕᒋᐊᖅᑲᓇᑦ ᓴᓇᓂᐊᖅᑕᐅᓂᒡᓕᐊᖅᓯᒪᔪᓂᑦ ᐊᐅᓚᑕᐅᓂᐊᖅᑐᑦ ᐅᖃᖅᑐᑎᐊᕈᒪᓂᖅᓯᑦ ᐊdᓂᐅᓂᐊᖅᑐᖅ ᓴᓇᓂᐊᖅᑕᐅᓂᖅᓂᓇᓂ ᓄᓇᖅᑦ ᐊᓚᒐ ᐃᒪᐃᑦ ᑕᒪᐃᓇᓂᑦ ᐱᑕᖅᑐᑎᓂᑦ ᐅᓪᓗᓂᑦ, ᐃᓄᐃᑦ ᐱᔨᓇᐅᑎᖅᑲᖅᖢᓂᓇᓂᑦ, ᐱᑕᖅᐃᓇᖅᓂᖅᖢᓂᓇᓂᑦ ᑕᐃᒪᖅᑎᑦ ᐃᓕᓴᕐᕕᒃ ᐃᓄᐃᑦ ᖃᐅᔨᓚᕐᖅᑕᐅᓂᖅᑐᑦ ᑕᒪᐃᓇᓂᑦ ᐊᑐᖅᑕᕐᕕᐊᖅ ᐃᓕᓴᕐᓂᐊᖅᑐᓂᑦ ᑕᒪᐃᓇᓂᑦ ᐃᓄᐃᑦ ᖃᐅᔨᓚᐅᖅᑐᓂᑦ ᑕᒪᐃᓇᓂᑦ ᐊᑐᖅᑕᕐᕕᐊᖅ ᐃᓕᓴᕐᓂᐊᖅᑐᓂᑦ ᑕᒪᐃᓇᓂᑦ ᐱᑕᖅᑐᑎᐊᕈᒪᓂᖅᓯᑦ ᐊᐅᓚᑕᐅᓂᐊᖅᑐᑦ ᐃᓄᐃᑦ ᖃᐅᔨᓚᐅᖅᑐᓂᑦ ᑕᒪᐃᓇᓂ


ᐊᐅᓪᓗᓂᔪᓐᓇᖅ ᐃᓂᖅᑕᐅᔪᒪᓂᖅᑎᑦ
Management Objectives



ᓴᓕᓂᖅᑕᐅᓂᖅᑎᑦ ᐃᓕᓴᕐᓂᐊᓂᖅ ᑲᓇᙵᓂᖅ ᑲᑎᒪᔨᓂᖅ ᐃᒪᐃᑦ ᐅᑭᐅᖅᑕᐅᔪᒪᓂᖅᑎᑦ ᑕᐃᔭᑯᓐᓚ ᑲᑎᒪᔨᓂᖅᑎᑦ ᐃᒪᐃᑦ ᐅᑭᐅᖅᑕᐅᔪᒪᓂᖅᑎᑦ ᑕᐃᔭᑯᓐᓚ ᑲᑎᒪᔨᓂᖅᑎᑦ ᐃᒪᐃᑦ ᐅᑭᐅᖅᑕᐅᔪᒪᓂᖅᑎᑦ ᑕᐃᔭᑯᓐᓚ ᑲᑎᒪᔨᓂᖅᑎᑦ ᐃᒪᐃᑦ ᐅᑭᐅᖅᑕᐅᔪᒪᓂᖅᑎᑦ ᑕᐃᔭᑯᓐᓚ

ᓴᓕᓂᖅᑕᐅᓂᖅᑎᑦ ᓄᓇᓕᖕᒥ ᐃᓕᓴᕐᓂᐊᓂᖅ ᑲᑎᒪᔨᓂᖅᑎᑦ ᐃᒪᐃᑦ ᐅᑭᐅᖅᑕᐅᔪᒪᓂᖅᑎᑦ ᑕᐃᔭᑯᓐᓚ

ᐊᐅᓪᓕᓂᐊᓪᓗᑦ ᐃᓂᖅᑕᐅᔪᓂᖏᑦ
Management Objectives



ᐅᓴᖅᑐᑦᑎᐊᓂᖅ ᓄᓇᓕᑦ ᖃᐅᔨᒪᔨᐅᑎᑦᑎᐊᖅᑕᕐᓂᖅᓗᑦ ᐸᕐᓇᐅᑎᓴᓂᖅ ᐊᑐᑎᖅᑕᕐᓂᐊᖅᑐᓂᖅ ᖃᐅᔨᒪᑎᑦᑎᐊᖅᑕᕐᓂᖅᓗᑦ ᐊᒻᒪᓗ ᖃᐅᔨᒪᔨᐅᑎᑦᑎᐊᖅᑕᕐᓂᖅᑎᑦ ᖃᐅᔨᓴᐅᓯᓕᐊᔪᓂᖅ, ᐅᖃᓕᒫᖅᑐᓂᖅᓴᓇᓱᓗ ᐃᔪᒪᓕᑕᐅᓂᖓᓂ ᐃᓄᖅᓄᑦ ᐊᒻᒪᓗ ᐃᓄᐃᑦ ᐃᔪᒪᓕᕆᖅᓂᖓᓂ, ᖃᐅᔨᓴᖅᑕᐅᓂᐊᖅᑐᓂᖅ ᔪᓚᑦᓂᐅᑎᑕᐅᓂᐊᖅᑐᓂᖅ ᐱᓕᕆᓂᖅᓂᓂ

ᐱᔨᒥᓴᐃᖅᑕᕐᓴᓗᑎᖅ ᓴᓪᓂᕐᒥᐅᑕᓂᖅ ᐃᖅᓚᐃᓴᖅᑕᐅᓯᕐᓂᖅᓂᓂᖅ ᐊᒻᒪᓗ ᐃᔪᒥᖅᓴᐃᓴᐅᖅᑕᐅᓯᕐᓂᖅᓂᓂᖅ ᐱᓴᐊᖅᑕᖅᑐᓂᖅ ᖃᐅᔨᓴᖅᑕᐅᓂᐊᖅᑐᓂᖅ ᐊᒻᒪᓗ ᓇᐅᑦᑎᖅᑕᐅᖅᑕᕐᓂᐊᖅᑐᓂᖅ

ᖃᐅᔨᒪᔨᐅᓂᖅᓴᐅᑦᓴᓗᑎᖅ ᓄᓇᕐᓂ ᐱᓕᕆᐃᐅᔨᓇᖅᑐᓯᕐᓂᖅᓂᓂᖅ ᑎᓐᒥᐊᖅᑲᖅᓴᐃᐅᔨᑦ ᓴᓐᓂᐊᖅᑕᐅᔪᒪᖅᓯᕐᓂᖅᓂᓂ


Ensure local awareness, participation and benefits

Develop a communication plan to facilitate information and knowledge sharing, discuss concerns and input into research priorities

Encourage Salliqmiut employment and mentorship in research and monitoring

Increase local awareness of permitted activities within the MBSs

ᐊᐅᓪᓕᓂᐊᓕᒃ ᐃᓂᖅᑕᐅᔪᒪᓂᖅᑎᑦ
Management Objectives



ᐊᖏᑐᕈᓂᕐᔪᖏᑎᑦᑎᑦ ᐃᓕᒪᐃᓐᓇᕐᓂᖅ ᑭᒪᔭᖏᑐᖅ ᐅᖃᐅᔾᔨᒌᖕᖏᑦ

ᐅᔾᔨᖅᑐᑦᑎᐊᕐᓇᕐᔪᓂᖅᑎᑦ ᖃᐅᔨᓴᖅᑎᑦᑎᓂᖅ ᐊᑐᖅᑕᐅᔨᒪᔨᖅᑎᑦᑎᓂᖅ, ᐅᑭᐅᖅᑕᐅᔪᒪᓂᖅᑎᑦᑎᓂᖅ, ᒪᑐᐊᖅᑎᑦᑎᓂᖅ

ᐊᖏᑐᕈᓂᕐᔪᖏᑎᑦᑎᓂᖅ ᐃᓕᒪᐃᓐᓇᕐᓂᖅ ᑭᒪᔭᖏᑐᖅ ᐅᖃᐅᔾᔨᒌᖕᖏᑦ ᐅᖃᐅᓯᒪᔭᕐᓂᖅ ᐃᓕᓴᕐᓂᖅ ᐊᑐᖅᑕᐅᔨᒪᔨᖅᑎᑦᑎᓂᖅ

ᐊᐅᓪᓕᓂᖅᓯᓐᓂ ᖃᓄᐃᓕᓴᕐᓂᓱᐊᕐᓯᓐᓂ
Management Approaches

ᐊᐅᓪᓕᓂᖅᓯᓐᓂᒃ ᖃᓄᐃᓕᓴᕐᓂᓱᐊᖅᑐᑦ ᐅᖃᐅᓯᖃᓕᓯᐅᕙᒃ
ᖃᓄᐃᓕᓴᐅᑭᑕᐅᓂᐊᓕᓴᖅᑐᓂᒃ ᐊᒃᓯᑭᑕᐅᔨᓐᓇᖅᑐᓂᒃ ᐊᐅᓪᓕᓂᕆᓯᓐᓂᕐᒧᑦ,
ᐊᖏᓚᕆᐅᓗᑎᒍᓗ ᐊᐅᓪᓕᓂᖅᓂᕐᒧᑦ ᐱᓕᕆᐊᖏᐊᓂᔪᓚᓯᔨᑦ

- ᐱᐅᓯᑐᖅᓴᒍᑦ ᐊᐸᑎᓐᓴᓂ ᓄᓇᒥᐅᑕᓐᓂᑦ ᐊᓚᒪᓗ ᐅᓚᔨᐃᑦ ᐱᑕᖅᑎᓂᓐᓂᑦ
- ᐅᓚᔨᐃᓪᓗ ᐅᓚᔨᑎᓂᓗ ᓇᔨᓴᕆᐅᕙᐸᓐᓂᓂᓐᓂᒃ ᐊᐅᓪᓕᓂᖅᓂᖅ
- ᓇᐅᑦᑎᖅᑐᐊᖅᑕᐅᖅᑕᑕᐃᓇᖅᓂᑦ ᐊᓚᒪᓗ ᖃᐅᔨᓴᖅᑕᐅᖅᑕᓴᖅᓂᑦ
- ᐃᓄᓕᒋᓂᒃ ᖃᐅᔨᓴᒪᑦᑎᐸᓪᓕᐊᓂᖅ ᐊᓚᒪᓗ ᖃᐅᔨᓴᓕᐅᔨᕐᓴᐅᓂᖅᑐᓂᒃ ᐊᐅᓪᓕᓂᐸᓐᓂᖅ

Management Approaches outline ways to address the Management Challenges, while meeting the Management Goals

- Cultural Resources Management
- Wildlife and wildlife habitat Management
- Monitoring and Research
- Public Awareness and Information Management

ᐊᐅᓪᓂᐊᓂᖅᓯᓂᖅ ᖃᓄᐃᓕᓐᓄᑦᓯᐊᖏᓂᖅ Management Approaches

ᐱᐅᓯᑐᖃᖅᑎᒍᑦ ᐊᕐᕌᑎᓄᑦ ᓄᓇᒥᐅᑕᓂᖅᑐᑦ ᐊᓚᒐᓗ ᐅᒪᔨᐃᑦ ᐱᑕᖃᖏᓂᖅᑐᑦ

ᐅᖃᓕᖅᑐᑦᑎᐊᖃᑕᖏᓂᖅ ᓴᐳᓂᓂᐊᖅᑕᐅᓂᖅᑐᓂᓂᒃ ᐃᓄᖅᖃᖏᐅᕐᓂᐅᔪᓐᓇᖅᑐᖅᑐᑦ ᖃᑭᒋᐃᓂᖅᖃᖏᓂᖅᑐᑦ, ᐃᓄᐃᓪᓗ ᐅᖃᐅᔾᔨᑐᖅᑐᑦ, ᐃᓄᐃᓪᓗ ᓄᓇᒥᓂᒃ ᐃᒪᖏᑎᖅᑐᓂᓗ ᑕᐃᔭᐳᑐᖅᑐᑦ, ᐃᓕᓴᐅᕐᓕᓪᓕᐊᓗᑎᒃ ᖃᐅᔨᒪᔭᐅᕐᓕᓪᓕᐊᔨᓄᑦ, ᑎᑎᕋᖅᑕᐅᕐᓕᓪᓕᐊᖃᑦᑕᐃᓐᓇᓪᓗᑎᒃ ᐊᒪᓚᓗ ᓄᓇᖓᐊᓕᐅᖃᑦᑕᕋᓗᑎᒃ

Cultural Resources Management

Ensure preservation of archeological sites, Inuit language, and Inuit place names and add to knowledge through inventory and mapping projects

ᐊᐅᓪᓕᓂᐅᖅᓯᒪᓂᖅ ᖃᓄᐃᓕᓴᖅᓯᐊᕐᓯᖅᓯᒪᓂᑦ Management Approaches

ᐅᒪᔨᐃᑦᓗ ᐅᒪᔨᕐᓄᑦᓗ ᓇᔨᕕᓴᐅᕙᖅᓯᖅᓯᒪᓂᖅ ᐊᐅᓪᓕᓂᐅᖅᓯᖅ

- ᐊᓯᔾᔨᖅᑕᐅᓗᓂᓂ ᐃᓕᑕᕕᓴᐅᓂᖅᓴᓄᑦ ᑎᒃᑯᐊᖅᑕᐅᓯᒪᓂᖅᓴ ᖃᓇᑕᒥ ᐅᒪᔨᖅᓴᕐᐃᐅᓂᕐᒧᑦ, ᐱᔭᔨᑕᐅᔨᓐᓇᕐᒪᑦ ᐊᓯᔾᔨᖅᑕᐅᔨᓐᓇᕐᓂᖅᓂᓂᒃ ᑭᓪᓕᖅᓴᕐᐃᐅᓂᖅᓯᒪᓂᑦ
- ᐊᒥᓱᓴᖅᓂᒃ ᖃᓄᐃᓕᓕᐅᑕᐅᓂᐊᖅᑐᓂᒃ ᐃᑲᔨᖅᓯᐃᓗᓐᓂᑎᒃ ᐊᐅᓪᓕᑎᒃᔨᐅᑎᑲᓄᑦ ᖃᖅᓴᐃᑦ ᐊᒥᓱᓂᖅᓯᖅᓂᒃ
  - ᐱᔨᒥᓴᐃᓂᖅ ᓄᓇᓕᖅᓂ ᖃᖅᓴᓯᓯᐅᖅᑕᑕᕕᔨᓐᓂᕐᒥᒃ, ᑐᓂᐅᖅᐃᖅᑕᑕᐅᑎᐊᓐᓂᖅ ᐊᐃᑦᑐᓐᓴᒃᓗᑎᒃ ᐃᓄᖅᑎᑎᓄᓗᓄᑎᒃ ᐊᖅᓯᓓᑕᕕᔨᓐᓇᖅᑐᑕᕕᓴᐅᓂᖅ ᐊᓗᒪᓗ ᑭᓇᐅᔭᓕᐅᑕᐅᔨᓐᓇᖅᑐᑕᕕᓴᐅᓂᖅ ᐊᑐᑭᓐᓇᖅᑐᑕᕕᓴᐅᓂᖅ ᐊᑐᔨᒪᓗ ᖃᐅᔨᓴᓂᒃ ᖃᐅᔨᓴᖅᑕᐅᓐᓇᓕᓕᐅᑕᐅᓂᖅ
- ᐊᒃᑐᖅᑕᐅᓂᕐᓗᒃᑎᑦᑎᑕᐃᓕᒪᕐᐊᕐᓂᖅ ᑕᓇᐅᒃᓯᒃ ᐅᒥᐊᕐᔪᐊᕐᓂᒃ ᖃᖅᓱᑕᓯᐅᑦᓯᓕᕐᓂᖅᓴᓄᑦ ᐅᒪᔨᐃᑦ ᐱᓕᕐᓂᑦ

Wildlife and wildlife habitat Management

- Change designation to National Wildlife Area, which may involve a boundary change
- Support several strategies to manage goose population
  - Promote local harvest, sharing, sport hunting, and commercial opportunities, ongoing research
- Mitigate marine and air traffic impacts on wildlife

ᐊᐅᓪᓕᓂᖅᓯᓐᓂᖅ ᖃᓄᐃᓕᓴᕐᓯᐊᕐᓯᓐᓂᒃ
Management Approaches

ᓇᐅᑦᑎᖅᑐᑕᐊᖅᑕᐅᖃᑦᑕᐃᓐᓇᕐᓯᓐᓂᒃ ᐊᓪᓚᓗ ᖃᐅᔨᓴᖅᑕᐅᖃᑦᑕᕐᓯᓐᓂᒃ

- ᐃᑲᔪᖅᓭᐃᓂᖅ ᑲᔪᓭᓐᓂᖅᓂᑉ ᖃᐅᔨᓴᖅᑕᐅᓕᕐᓂᐊᔪᑦ ᐊᓪᓚᓗ ᓇᐅᑦᑎᖅᑐᖅᑕᐅᓴᖅᑐᖃᑦᑕᕐᓯᓐᓂᒃ, ᑎᑎᕐᖅᑕᐅᓕᕐᓂᐊᓗᑎᑦᓗ ᐃᓗᖃᖏᕐᕕᐅᓕᐅᖅᑐᑦ ᖃᑭᒪძᕕᓂᖃᖏᕐᕕᐅᔪᑦ
- ᐃᑲᔪᖅᓭᖅᑕᐅᓗᑎᑉ ᓴᓂᓕᕆᐅᑕᓗᑦ ᖃᐅᔨᓴᖅᑕᐅᕐᔪᐊᕐᓇᔪᑦ ᓭᓄᓂᓕᐅᕐᔪᐊᕐᓂᓗᑎᑉ
- ᐱᐅᓱᕈᐊᖅᑕᐅᓗᑎᑉ ᖃᐅᔨᒪᓴᓱᑦ ᐊᓕᑎᒥᑉ ᐅᒪᔪᓚᕆᐅᓕᖅᑐᑦ ᐊᒪᓗ ᐱᐅᓱᑐᖃᖅᑎᒍᑦ ᐊᓕᑎᖏᓐᓂᒃ ᓄᓇᒥᐅᑕᕐᓂᑦ ᐊᒪᓗ ᐅᒪᔪᐃᑦ ᐱᑕᖃᕐᓯᓐᓂᒃ

Monitoring and Research

- Support ongoing research and monitoring, document archaeological sites
- Support Salliqmiut research priorities
- Improve knowledge of ecological and cultural resources in the MBSs

ᐊᐅᓪᓂᓂᖅᓯᓐᓄᑦ ᖃᓄᐃᓕᖅᓯᐊᕐᓯᓐᓄᑦ
Management Approaches

ᐃᓄᓕᒻᓂᖅ ᖃᐅᔨᒪᑎᑦᑎᐊᓪᓕᐊᓂᖅ ᐊᓪᒪᓗ ᖃᐅᔨᒪᔭᐅᔪᕐᓇᐅᓂᖅᓯᓐᓂᖅ ᐊᐅᓪᓂᑎᐊᖅᓂᖅ

- ᓴᖅᑭᑎᑦᑎᓗᑎᖅ ᑐᓴᐅᒪᔭᐅᑎᑦᑎᖃᑦᑕᕐᓂᕐᒧᑦ <ᕐᓇᐅᑎᖅᓴᓂᖅ ᐊᓪᒪᓗ ᐱᐃᖅᓴᖃᖅᑎᑕᐅᐊᖅᓯᓐᓄᑦᓯᓄᑦ ᐃᓄᐃᑦ ᑐᓴᐅᒪᑎᑦᑎᖃᑕᑭᓇᕐᓂᕐᒥᖅ ᖃᐅᔨᒪᔭᖅᑭᑦᑐᓂᐊᔪᓂᖅ, ᖃᐅᔨᒪᔭᖅᓯᓐᓄᓪᓗ ᓄᓇᒧᖅᑐᐊᐅᓂᑦ, ᐅᖃᐅᔨᔪᓇᖃᖅᑕᕐᕕᑎᓪᓗ ᐃᔪᒪᓗᑕᐅᔪᑦ ᐃᓄᖅᓄᑦ ᐊᓪᒪᓗ ᖃᐅᔨᓴᖃᑕᐅᓂᐊᖅᑐᐊᓂᖅ ᔪᐊᓂᐅᓕᖅᑎᐊᖅᑐᑦ
- ᐱᔪᒥᓴᐃᓂᖅ ᐃᑲᔪᖅᔪᐃᐊᖅᓂᕐᓗ ᐃᓄᖅᓂᖅ ᐊᓪᒪᓗ ᐃᓄᖅᓂᖅ ᐃᓕᓴᓂᐊᖅᑎᐅᔪᓂᖅ ᐱᖃᑕᐅᔪᒪᓂᖅᑐᓂᖅ ᓄᓇᓕᖅᓂ ᐊᔪᖅᑎᓂᖅᑕᑭᓇᕐᓂᕐᒧᑦ ᐃᓄᖅᓂᖅ
- ᐃᑲᔪᖅᔪᐃᓂᖅ ᑲᔪᔭᓇᑭᖅᑐᓂᖅ ᐊᑐᖅᑕᑭᓇᑭᖅᑐᑦ ᐊᓪᒪᓗ ᐃᓄᖅᓂᖅ ᐃᓕᓴᓂᐊᖅᑎᐅᔪᓂᖅ ᐱᖃᑕᐅᔪᒪᓂᖅᑐᓂᖅ ᓄᓇᓕᖅᓂ ᐊᔪᖅᑎᓂᖅᑕᑭᓇᕐᓂᕐᒧᑦ ᐊᐅᓪᓂᑎᐊᖅᓂᖅ ᐃᓗᓐᓂᐊᓂᖅ ᑎᖅᑎᐊᖅᑎᓇᖅᑐᓂᖅ

Public Awareness and Information Management

- Develop a communication plan and provide opportunities to share information and knowledge among users, discuss concerns and research priorities
- Encourage and support Inuit and Inuit student initiatives to build local capacity
- Support sustainable tourism operations within the MBSs

ᐊᐅᓪᓕᓂᐅᖅᓯᖅ ᖃᓄᐃᓕᖅᓱᓕᐊᕐᓯᖅ
Management Approaches

ᐃᓄᓕᒻᒥᐅᖅ ᖃᐅᔨᒪᑎᑦᑎᐊᕐᓕᐊᓂᖅ ᐊᖏᒪᓗ ᖃᐅᔨᒪᓴᐅᔪᖅᓴᐅᓂᖅᓯᖅᓄᑦ ᐊᐅᓪᓕᓂᑎᐊᕐᓂᖅ

- ᑎᑎᕆᖅᑕᐅᔪᒪᓗᑎᑦ ᓄᓇᒋᑦᑕᖅᑐᕆᐊᐅᓪᐊᕐᑐᑦ, ᖃᐅᐸᐊᖅᑕᐅᖃᖅᑎᑦᑕᕆᓗᑎᒃᓗ ᐊᖏᔪᓇᔪᓰᑦ ᑲᑐᔨᖃᑎᒋᖅᓯᖅᓄᑦ ᑎᑦᔨᕆᐊᑎᒋᓄᑦ ᑕᐃᔪᑯᐊ ᓄᑕᓂᓂᑦ ᑐᔪᔪᕐᐅᑎᑦ ᑎᖅᑎᐊᖃᖅᕆᒋᑦᑐᒪᓂᖅᓯᖅᓄᑦ, ᖃᐅᔨᕆᖅᑕᐅᖃᑕᕆᓕᑎᓐᓗ ᐃᓐᓇᓂᓪᓕᐅᔪᖏᖅ
- ᒪᓪᓕᐊᖃᑕᕆᓂᖅᑎᑦ ᐱᔨᒥᕆᐃᐃᓂᖅ - ᑐᕆᐅᒪᑎᑦᑎᖃᑕᕆᓗᑎᒃ ᖃᐅᔨᒪᓴᐅᓪᓕᐊᖅᑐᓂᒃ ᒪᓪᓚᓪᓀᔨᓂᒃ ᐊᒻᒪᓗ ᒪᓪᓚᓵᓗᑐᓴᐅᐊᖅᑐᓂᒃ, ᐊᐅᓪᓕᓂᑎᓂᕐᒧᓵᓗ ᑐᖏᓚᓀᓴᐅᔨᔨᓂᒃ ᐊᒻᒪᓗ ᐱᓪᓀᐊᖏᓂᔪᒪᔨᓂᒃ

Public Awareness and Information Management

- Maintain list of camps, assess new requests inside MBSs with HTO and inspect periodically
- Compliance promotion – share information about laws and regulations, management goals and objectives

ᐊᖏᑭᑕᐅᓯᒪᔪᑦ ᐱᓕᕆᐊᖓᔪᓂᓐᓄᑦ ᑎᑭᓯᒪᖓᐊᖅᓂᓐᓄᑦᑐᓕᓗ
Authorized Activities and Access



- ᐃᓄᐃᑦ **ᐱᔨᖅᓇᐅᑎᖏᕆᐊᖏᖅᑎᑕᐅᖅᖁᓕᑦ** ᑎᑭᓯᒪᔨᖅᓇᓐᓄᑦᓂᓄ ᑎᖅᑕᐊᖏᖅᓴᑦ ᓄᓇᖅᑐᓄᑦ, ᐱᐅᓯᑐᖏᑎᒃ ᐊᑐᓂᐊᖏᑎᓪᓗᑎᑦ, ᓯᑦᓗ ᐊᖓᔪᓇᖅᓂᖅᖁᖅᑐᓂᓄ, ᓴᓇᖅᖁᐊᖓᕆᓴᒃ ᐅᓴᕋᐃᓴᐃᓂᖅᖁᖅᑐᓂᓄ, ᓄᓇᒥᑕᖏᑐᓴᖅᑐᓂᓄ (ᒪᓕᒐᖅᑎᑦ ᒪᓕᒐᓗᑎᑦ ᓄᓇᕗᓴᒥ ᓄᓇᑕᑭᑎᓄᑦ ᐊᖅᑭᑐᑎᑦ ᐊᒡᒪᓗ ᐃᓄᐃᑦ ᐊᒃᑐᖏᑕᐅᓯᒪᓂᖅᑐᓄᑦ ᐃᑲᔪᓯᐊᕐᓂᖅᑐᓄᑦ ᐊᖅᑭᑐᑎᑦ)
- ᐃᓄᐃᑦ **ᐱᔨᖅᓇᐅᑎᖏᕆᐊᖏᖅᑎᑕᐅᖅᖁᓕᑦ** ᑎᑭᓯᒪᔨᖅᓇᓐᓄᑦ ᑎᖅᑕᐊᖏᖅᓴᑦ ᓄᓇᖅᑐᓄᑦ ᐊᖅᑭᑐᑎᖅᑕᐃᔨᓂᖅᑐᓄᑦ ᐅᕝᕙᓚᐅᔪᓂᑦ ᐃᖏᓪᓗᓯᐅᑦᑎᐊᓂᖅᑐᓄᑦ

- Inuit **do not** require a permit to enter the MBSs for traditional activities, including harvesting, removal of carving stone, setting up camps *(Nunavut Agreement* and IIBA)
- Inuit **do not** require a permit to enter the MBSs when working as a hunting or fishing guide



ᐊᖏᑭᑕᐅᓕᕆᓂ ᐱᓕᕆᐊᖏᔪᓐᓇᖅᓂᖅ ᑎᑭᓕᒫᐅᔪᓐᓇᖅᓂᖏᑦᓗ
Authorized Activities and Access

- ᑕᒪᖅᓂᒃ ᐃᓄᒡᒪᓗᐅᖏᑦᑐᑦ **ᐱᔪᓐᓇᐅᑎᖅᑲᖅᑎᑕᐅᔭᑦ** ᑎᑭᓕᔪᓐᓇᖅᓂᖅ ᑎᖏᒥᖅᑲᖅᕕᖓᓄᑦ (ᓄᓇᖏᑦᓂᓄᓐᓂᑦ ᐃᒪᖅᔪᑎᖏᑦᓂᓄᓐᓂᑦ) ᑕᐃᒃᑯᐊᓗ ᐱᔪᓐᓇᐅᑎᖅᑲᖅᑲᖅᑐᑦ ᐃᓄᒡᒪᓗᐅᖏᑦᑐᑦ ᐊᑭᓐᓇᒃᑐᓂᒃᑐᒥᒃ ᐊᖏᔪᓇᓕᐊᑭᓴᓇᒃᑐᑦ, ᖃᐅᔨᓴᖅᑎᓗᑦ, ᐳᓪᓕᑎᓪᓗ ᓂᐅᖅᑕᐊᑦ

- ᑕᒪᖅᑭᒃ ᑭᓇᐅᔭᓪᓕᐅᑎᖃᖅᓂᖅᓗᑦ ᐱᓪᓚᐊᖃᖅᓂᖅᓗᓴᓪᓕᓂᑦ ᐱᓪᓚᓂᖅᓗᑦ (ᐃᓄᐃᑦ ᐊᒻᒪᓗ ᐃᓄᒡᒪᓗᐅᖏᑦᑐᑦᖃᖅ ᓇᖏᑭᓂᖅ ᐱᓪᓚᐃᖅᑯᑎᖃᖅᑐᑦ **ᐱᔪᓐᓇᐅᑎᖅᑲᖅᑎᑕᐅᔭᑦ**. ᑕᐃᒪᓇᔨᖏᑦᑐᑐᓄᑦ ᐃᓄᐃᑦ ᐊᖏᔪᓇᓕᐊᑭᓴᑦ ᐊᒻᒪᓗ ᐃᖃᓪᐊᖅᑭᒪᓕᕆᑦ

- All non-Inuit **do** require a permit to enter the MBSs (land or water) including non-Inuit sport hunters, researchers and tourists

- All commercial or business activities (Inuit and non-Inuit owned businesses) **do** require a permit. Except for Inuit hunting and fishing guides



ᐊᓐᑭᑐᑕᐅᓯᒪᔪᑦ ᐱᓕᕆᐊᓐᔪᐊᓂᓐᓄᖅ ᑎᑭᓯᒪᓴᐃᐅᔪᓂᓐᓂᓯᑐᓕᓪ
Authorized Activities and Access

- ᑕᒪᖅᑭᒃ ᐱᓕᕆᕆᔨᐅᔪᑦ ᐃᓄᒃᒪᒪᕆᐅᓐᕿᑦᑕᔭᑦ ᐱᔪᓂᐊᐅᑎᖅᕆᓚᐅᖅᓴᓪᓕᖅ ᑎᓐᒥᐊᖅᓂᓐᒪᓂ (ᐃᐳᒥᒃ-ᐊᒃᑐᐱᕆᒧᑦ ᑕᖅᑭᓂ)
- ᖃᓄᐃᑦᑐᑐᐃᓐᓇᓂᒃ ᐱᓕᕆᔪᑦ ᓯᕐᖅᑎᕆᑕᐅᔪᓐᓇᖅᑐᓂᒃ ᑎᓐᒥᐊᖅᓂᐹᐃᑦ ᐊᓂᓗᓐᕿᓂᒃ ᐱᔪᓂᐊᐅᑎᖅᕆᓚᐅᖅᓴᓪᓕᖅ (ᐊᖅᕐᓗᒋᓕᒫᖅ)
- ᐃᓄᒃᒪᒪᕆᐅᓐᕿᑦᑕᔭᑦ ᐱᔪᓂᐊᐅᑎᖅᕆᓚᐅᖅᓴᓪᓕᖅ ᖁᔨᐅᓯᕐᓂᔪᓐᓇᓂᓗᑦ ᑎᓐᒥᐊᖅᓂᕐᐹᓐᓂ

- All activities by non-Inuit require a permit when migratory birds are present (APR-OCT)
- Any activities that may result in the destruction of migratory bird habitat requires a permit (YEAR-ROUND)
- Non-Inuit must have a permit to carry a firearm in the MBSs





ᐱᓯᓂᓇᐅᑎᖅᑭᑕᐅᓯᓂᓇᖅᓂᓕᒫᑦ
Permit Authorizations

- ᐃᓄᐃᑦ ᐱᓯᓂᓇᐅᑎᖅᑌᐊᖅᖃᑐᑦ ᑎᑭᓕᒪᓯᓂᓇᖅᓂᖅ ᑎᓪᓗᐊᖅᖃᕐᓂᓄᑦ ᐅᖅᑐᑭᓂᓇᖅᐳᑦ ᐱᓇᓕᔪᓐᓇᑎᒃ ᑲᓇᑕᒥ ᐅᒪᓯᓕᓚᓱᖅᑯᑦ ᐱᔨᑦᑎᕋᐃᓐᓇᕐᓂᖅ:

ec.nupermisscf-cwspermitnu.ec@canada.ca

ᐃᖅᓚᓐᓂ ᐃᖅᓇᐃᒪᖃᖅᓱᒃ: 867-975-4642



- ᐃᖃᓂᐅᖃᖅᓱᒃᑯᑦ ᐊᐅᓪᓚᑦᑎᐸᐅᖅᑕᐅᓂᖃᑦ ᑲᑎᒪᔨᕋᓪᓕᕐᓂᑦ ᖁᒥᖃᕈᐊᖅᑎᐅᐳᓕᑦ ᑕᒪᐃᓂᓂᑎᐊᖅ ᐅᖅᑐᑭᑎᓂᒃ ᐱᓯᓂᓇᐅᑎᑕᕋᓱᐃᑐᓂᒃ ᐊᒻᒪᓗ ᐊᑐᓕᖃᕐᑯᑐᓂᒃ ᑲᑎᒪᔨᕋᓪᓕᕐᓂᑦ ᐃᖅᓇᐃᒪᖃᖅᓱᖅᓱᓪᓗᓂᒃ ᑲᑎᒪᔨᕋᓪᓕᕐᓂᑦ ᐃᖅᓇᐃᒪᖃᖅᓱᖅᓱᑎᑦᓯᓚᖅᓂᒃ ᓇᓗᓇᐃᒪᖃᐃᓕᒃᑯᑦ

- Persons who require a permit to enter the MBSs can apply for a permit from the Canadian Wildlife Service:

ec.nupermisscf-cwspermitnu.ec@canada.ca

Iqaluit office: 867-975-4642

- The Irniurviit ACMC reviews all permit applications and provides recommendations to the Canadian Wildlife Service on whether the permit should be issued or not

ᐱᔨᓇᐅᑎᖅᑕᐅᔨᓇᓂᓐᓕᒃ
Permit Authorizations

- ᐱᔨᓇᐅᑎᓂᒃ ᑐᓂᔭᐅᔨᓇᖅᑐᑦ ᐃᒪᓇᐃᒃᑯᑎᒃ:
  - ᐱᓕᕆᐊᕆᔪᒪᔭᒃᓕᑦ ᓯᕐᖅᑎᕆᓗᑕᐅᓂᐊᖏᓕᑐᑐᒋᐊᕈᑎᒃ ᑎᖏᒥᐊᖅᓄᑦ, ᒪᓇᓂᖏᓇᓄᑦ, ᐅᓪᓗᖏᓇᓄᑦ, ᓇᔨᖅᐸᒃᑕᖏᓇᓄᓪᓗ ᐊᒻᒪᓗ/ᐅᕐᓂᓗᓇᓚᑦ
  - ᐱᓕᕆᐊᕆᓂᐊᖅᑕᖏᑦ ᒪᓕᑎᐊᕈᑎᒃ ᐱᑕᖅᕈᑕᐅᓂᖏᓄᓄᑦ ᑎᖏᒥᐊᖅᑲᐃᑦ ᐊᒻᒪᓗ ᐊᐅᓪᓚᑎᐅᑎᒃᓴᐃᑦ ᑕᐃᒃᑯᐊ ᕿᖏᓇᐅᑎᒃᓴᖏᑦ
- Permits may be issued if:
  - The activity is not harmful to migratory birds, eggs, nests, or habitats and/or
  - The activity is consistent with the purpose of the MBSs and their management plans



ᓱᓴᓕᐊᖅᓯᐊᖅ White-rumped Sandpiper





ᓴᐅᖅᑐᖅ Red Phalarope


ᓄᓇᖅᑐᑦ ᑎᒃᓯᐊᖅᑕᐅᓯᒪᓂᖅᑐᑦ / Site Designation

- ᖃᖅᖁᐅᖅᑐᖅ ᐊᓪᒪᓗ ᐃᑲᑦᑐᐊᖅ ᐃᓐᓇᐅᔪᖅ ᐃᓕᓴᕆᔨᐅᓪᒪᓂᒃ ᑎᒃᓯᐊᖅᑕᐅᓯᒪᔪᓕᒃᑐᓗᑎᒃ ᑎᓂᒥᐊᖅᑎᑦᓴᐃᐅᓕᖅᑎᓪᓗᑦ ᓴᓪᓕᖅᑐᖅ; ᐅᓯᐊᓕᓕ, ᐅᒪᔪᐃᑦ ᐊᓪᒪᓗ ᐱᐅᓯᑐᖅᑎᑦ ᐊᓂᒍᑎᖓᓕ ᓄᓇᒥᐅᑕᓂᑦᑕ ᐊᓪᒪᓗ ᐅᒪᔪᐃᑦ ᐱᑕᖅᑦᓕᓂᑦ ᐊᓪᓕᖅᑐᖅᑎᑦᑎᐊᖅᓕᓂᖅᑐᑦ ᐃᓕᓴᕆᔨᐅᓕᖕᓂᖅ ᐊᓕᖅᑐᖅᓄᓇᓕᒃ ᓄᓇᖓ ᑕᐅᓂᓐᓇ ᖃᓄᑕᒥ ᐅᒪᔪᖅᑦᓴᐃᐅᔪᕐᓂᖅ
- ᐊᐅᓪᓕᑎᔪᕐᓂᓕᒃ <ᑦᓇᐅᑎᖓᑦ ᐅᖅᐅᓯᖅᓴᓂᑦ ᖅᓄᖅ ᒪᓕᖅᑕᓪᓕᐅᔪᑦ ᒪᓕᓵᑕᐅᓯᒪᓂᖕᓂᓵᓂᒃ ᖃᓄᕐᓗᒍ ᐅᒪᔪᖅᑦᓴᐃᐅᓗᑎᒃᓴᖅᓕᑦ, ᓴᓂᐊᓂᒃ ᑎᓂᒥᐊᖅᑦᓴᐃᐅᑎᑕᐅᑐᐃᓐᓇᑦᓂᓕᓂᒃ
- ᐅᖅᐅᓯᐅᓂᖅᑐᑦ ᑎᒃᓯᐊᖅᑕᐅᓯᒪᓂᖅᑎᓂᒃ ᐃᓕᓴᕆᔨᐅᒪᓂᖅᑎᓂᓂ ᑕᐃᒃᓂ ᐊᐅᓪᓕᑎᔪᕐᓂᓂᓂ <ᑦᓇᐅᑎᖓᑎᓂ ᐱᑎᐊᑭᑎᖅᓴᖅᑦᓕᑦ ᐅᖅᐅᓯᐅᓂᖅᑎᓂᓂᒃ ᑎᓂᒥᐊᖅᑦᓴᐃᐅᔨ ᓯᔭᓂᒃᓕᖓᓂᒃ; ᐊᓯᔪᖅᑕᐅᓯᓇᖅᓯᒪᑕ ᐅᖅᐅᓯᐅᓪᓕᐅᖅᓯᒪᓂᖅᑎᓂᒃ, ᑐᑭᓯᔪᓂᑦᓴᐃᐅᓪᓕᖅᓯᒪᓂᖅᑎᓂᓂ ᐃᓄᐃᓂᓂ ᓄᓇᓕᒃ

- Qaqsauqtuuq and Ikkattuaq are currently designated as a Migratory Bird Sanctuary; however the wildlife and cultural resources within them are better suited by protection as a National Wildlife Area
- Management plans discuss how the criteria for a National Wildlife Areas are met by these MBSs
- Talking about the site designation in the management plan begins the discussion for the future; no changes would be made without further consultation with community

	ᑎᒃᓯᐊᖅᑐᑦ ᓴᓪᓕᖅᑐᖅᓄᓇᓕᒃ	ᖃᓄᕐᓗᒍ ᐅᓚᔨᖅᑐᖅᓄᓇᓕᒃ
ᓴᓪᓕᐊᑭᑕᐅᓯᒪᓂᖅ ᑐᖃᓚᖅᑐᖅ	ᑎᓂᒥᐊᖅ	ᑕᒪᖅᑎᖅ ᐅᓚᔨᐃᑦ
ᓴᓪᓕᐊᖅᑕᐅᔪᓂᖅᑎᓂᓄᓇᓕᒃ ᐊᑐᓂᖅᑎᖅ	ᐅᓪᓗᓂᖅᑐᖅᓄᓇᓕᒃ /ᐃᓕᓴᕆᔨᖅᑎᖅᓄᓇᓕᒃ	ᐊᖅᖁᔭᓕᒫᖅ
ᒪᓕᒐᖅᑎᔪᓂᖅ	ᑎᓂᒥᐊᖅᓯᓂᓂᓯᖅᑐᖅ ᑲᑎᒪᖅᓴᐊᑕᐅᔪᓚᔨᓂᖅᑐᑦ ᒪᓕᓵᑕᐃᑦ	ᑲᓇᖅᑐᖅ ᐅᓚᔨᓂᓂᓯᔪᓂᖅᑐᑦ ᒪᓕᓵᑕᐃᑦ

	MBS	NWA
Protection Focus	Migratory birds	All wildlife
Protection Period	Nesting/breeding season	Year-round
Legal Basis	*Migratory Birds Convention Act*	*Canada Wildlife Act*



ᐊᐅᓪᓕᐊᒃᑲᐃᓕᓪ ᐸᖏᓐᓂᐅᓂᒃ ᐊᑐᓪᖅᑎᑎᓂᖅ
Management Plan Implementation

- ᐊᑐᓪᖅᑎᑕᐅᓐᓪᓚᐊᓂᖅᑐᑦ ᐊᑐᒐᒃᑲᐃᓕᓪ (ᐱᕆᒃᑲᖏᕆᐊᖅᖏᓂᔪᓂ ᑭᓇᐅᔭᐃᓪᓗᓂ)
  ᐱᔨᓴᓇᐅᑕᐅᓂᐊᖏᒪᑕ
- ᐸᖏᓇᐅᑎᓯᓂᑦ ᖃᑎᖏᑭᐊᖅᑕᐅᔨᓴᓇᖏᓂᖅ ᑕᓪᓕᒪᐅᓂᐊᖅᑐᑦ 5 ᐊᖏᕿᔪᓂᐊᖅᑐᑦ ᓇᒃᓱᖅᐸᑕ, ᑭᖑᓂᐊᒍᓪᓗ ᑕᐃᑲᖑᓐᓂ ᖁᓕᓪᓂ 10 ᐊᖏᕿᒍᐃᓪ ᓇᒃᕃᖑᓐᓇᑎᒃ



- Implemented as resources (time and money) allow
- Plan will be reviewed after 5 years, and then every 10 years after that

2025 / 2035

ᖃᓗᓐᓇᒋᑦ ᐃᓕᐅᖅᑐᑦᑕᕐᓂᖅ ᐅᓂᒃᑎᓂᓪᓚᑦ!

Thank you for joining us!



ᐅᖅᖃᓕᕆᔨᒃᑐᑦ ᖃᐅᔨᓴᓕᕐᓗᓂ:

ᐃᖃᓗᐅᖃᕐᓕᕆ ᐊᐅᓚᑦᑎᐊᐅᖅᑕᐅᓴᖃᑦᔪᑦ ᑲᑎᒪᔨᖏᓚᑦ

ᑲᒪᔨᐅᓯᖅ: ᐊᖏᓚᔨᖅᑉ ᑐᖏᕐᓂᖏᓐ ᓴᓗᐃ-ᐃᐊᓕ ᑐᓄᐊᕐ

ᖃᓄᑕᐅᓴᕆᔨᒃᑯᑦ ᑎᑎᖅᖃᕐᓕᖏᓂᓗ

jean-francois.dufour2@canada.ca

ᐅᖅᓱᓚᐅᕐᒥ (867) 669-4766

Contact Information:

Irniurviit Area Co-Management Committee

c/o Vice-chairperson JF Dufour

jean-francois.dufour2@canada.ca

(867) 669-4766

Laura McKinnon

ᑐᓴᑭᒪᔨᔪᑦ ᐃᓕᓴᐸᓂᖅ!
We want to hear from you!



- ᐅᖏᐅᓯᖏᓐᓄᓴᓯ ᑐᓴᖏᑕᐅᖃᐅᔨᓴᓯᓗ ᐊᑐᑎᓕᒪᓐᓄᐊᓗᖓᒪᑕ ᐅᐸᕐᑎᓕᓂᒃ
- ᖁᓂᖃᓕ ᐃᓱᒪᒋᐊᕐᑦ ᐃᓗᓕᓐᖑᕐᑦ ᐊᐅᓕᑦᑎᐹᑎᖏᑦᑐᐃᓐ <ᖃᓪᓗᐅᑎᖑᓕᓐᑐᑦ?
  - ᓇᓴᒪᒋᐊᓱᑦ ᐃᓗᓕᓐᖑᕐᑦ ᐊᐅᓕᑦᑎᐹᑎᖏᑦᑐᐃᓐ <ᖃᓪᓗᐅᑎᖑᓕᓐᑐᑦ?
  - ᓱᐻᑎᖑᕐᑦ ᖁᓂᐃᓕᓐᓄᓂᖑᕐᓯᓗᓂᑦ ᓇᓴᒪᒋᖑᐱᒋᑦ ᐃᓗᓕᓐᖑᕐᓂᓂᓗ ᐊᐅᓕᑦᑎᐹᑎᖏᑦᐃᓐ <ᖃᓪᓗᐅᑎᓂ?
- ᖁᐅᓵᒪᔪᖁᖅᐲᑦ ᑎᖑᒥᐊᖁᖃᐊᓕᓚᓱᓂᖏᒻᒪᖅ ᖁᐅᓵᒪᔪᐅᖃᐻᓂᓂᖅ ᐃᓕᑭᔪᐅᓕᑐᓗᓂᖅ ᐊᐅᓕᑦᑎᐹᑎᓄᐃᓐ <ᖃᓪᓗᐅᑎᖑᓂᓗᓕᓐᑐᑦ?

- Your comments and input are very valuable to us
- What do you think of the content of the management plans?
  - Are you comfortable with the content of the management plans?
  - What are you not comfortable with in the management plans?
- Do you have any knowledge about the MBSs that you want to share so it can be included in the management plans?

Irniurviit ACMC Management Plan Presentations:

Day 2: Wednesday, November 27, 2019 2:00PM Session

Start time: 2:05 – Welcome and introduction (Ron Ningeongan)

Start time: 2:06 End Time: 3:16 – Presentation

Questions and Answers:

1. Q: About the avian cholera, will the disease be going on permanently now?

   A: It was first detected in 2005 where there was a huge die-off in the colony. Since then, the disease has only gone down and less and less fatalities are being recorded. It's hard to tell how long it will go on, so it's important to continue monitoring it.

2. Q: Where is the disease coming from? The air or the land?

   A: Avian Cholera is a bacterial disease and it comes from contaminated food or water.

Day 2: Wednesday, November 27, 2019 7:00 PM Session

Start time: 7:33 – Welcome and introduction (Noah Kadlak)

Start time: 7:36 End Time: 8:35 – Presentation

Questions and Answers:

1. Q: Are the number of geese increasing in Qaqsauqtuuq or are they leaving the area?

   A: Survey results show an increase in the population over the last 20-30 years, and a change in distribution as well. They believe the population is stabilizing now, as fewer young are surviving or becoming adults.

2. Q: Are the Migratory Bird Sanctuaries protected in the winter? For example, could a mining company come in the winter and start mining in the area in the winter?

   A: The habitat is protected; you don't need a permit to enter the sanctuaries in the winter because there are no birds present. But the bird habitat is still protected, so you can't go in and destroy any bird habitat in the winter; you would require a permit for that.

3. Q: Regarding the harvesting program; you mentioned in the presentation that there was an amount of 12 million dollars being distributed for the program; Is this amount of money a one time distribution or is it being distributed annually?

   A: The money was related to the Inuit Impact Benefit Agreement signed by ECCC, NTI and Regional Inuit Associations. There's a funding schedule and most of the funds go to NTI and the RIAs.

4. Q: For the spring goose harvesting program, will there be an increase of funding?

   A: GN funded the program. The ACMC coordinated the planning and supported the HTO in getting the funding from GN. The HTO will have to re-apply to GN for funding each year. Should there be more funding allocated?

   Comment: Yes, the rate [$/goose or egg harvested] was too low. It didn't cover enough of the costs to go out and hunt. The program came together late in the spring and the conditions were not normal last spring.

5. Q: They will always have to ask for funding again?

   A: Yes, that's our understanding. Last year was a pilot year and we'd need to discuss with GN about options to renew the program.

6. Q: I am concerned about the numbers and closeness of the geese. I feel like they are becoming more dangerous [to eat??]. I am wondering what's being done or will be done?

   A: The ACMC supports a number of strategies to help with the goose issue, e.g. promote local harvest, sharing, sport hunting, commercial harvest and more monitoring. But our role only extends to the boundaries of the MBSs. The ACMC also actively participated and supported the IQ Snow Goose study, the local Goose Spring Harvest program and new community-based monitoring studies. The federal government opened the hunting season, increased bag and possession limits and changed other hunting strategies for non-Inuit/Indigenous Peoples down south.

   Comment: They told us to hunt them, but we know when they are fat enough to hunt. If we hunted just to kill (while they are skinny), that would not look good for Inuit. Inuit do not hunt to kill; we hunt for food and if we hunted the skinny ones, it would just be wasting food. We do not hunt the skinny geese; we go after the fat ones.

Day 3: Thursday, November 28, 2019 2:00 PM Session

No attendees.


Day 3: Thursday, November 28, 2019 7:00 PM Session



Start time: 7:30 – Welcome and introduction (Noah Kadlak)

Start time: 7:30 End Time: 8:30 – Presentation

Questions and Answers: None.

Management Plan Review and Consultations

Organization	Comment	ACMC Response
Kivalliq Inuit Association Via email 22-Aug-2019	It was reported that barren-ground caribou have used some portions of the MBS as calving grounds in the past, but no details were provided.	Both management plans refer to the Keewatin Regional Land Use Plan (NPC, 2000) for more details. The map on p.51 of the Keewatin Regional Land Use Plan shows caribou calving grounds on Southampton Island. The overlap was determined using the Kivalliq Inuit Association Mapping Tool – see clips below. Ikkattuaq MBS:  Qaqsauqtuuq MBS:

		 More recent information from Draft Nunavut Land Use Plan process suggests that only Qaqsauqtuuq MBS overlaps the main calving grounds on Southampton Island. See the blue line in clip below. 

		The ACMC will add more details to Sections 2.3.1 of the management plans (see in red below). Ikkattuaq: "At least historically, the northwestern portion of the Ikkattuaq MBS overlapped ~~portions of~~ the Southampton Island caribou calving grounds (NPC, 2000). Recent information suggests that important caribou calving grounds now surround the community of Coral Harbour further to the east (NPC, 2012)." Qaqsauqtuuq: "Qaqsauqtuuq MBS also overlies portions of the traditional Southampton Island caribou calving grounds to the northeast (NPC, 2000).~~;~~ Recent information suggests that important caribou calving grounds now surround the community of Coral Harbour and overlap the western portion of the MBS (NPC, 201~~6~~2)."
Kivalliq Inuit Association Via email 22-Aug-2019	Their first objective is to work towards converting the MBS to a National Wildlife Area (under the Canada Wildlife Act) – the major change is year-round protection for more than just migratory birds. This change is supported by the Irniuviit Area Co-Management Committee, and could involve boundary changes. I am sure Government of Canada It is aware that any changes in the actual designation made have an impact in Inuit and required open of the IIBA to address the changes and a full fusibility of that change. (See Article 9 NLCA). I will advised to removed this section because have legal repercussions to the IIBA	Under the IIBA, the Irniurviit ACMC advises the Minister on all aspects of MBS management, including the protection of wildlife and wildlife habitat. The ACMC determined during the management planning process that better protection for all wildlife and cultural resources in the protected areas is required and would be of benefit to Salliqmiut – see Section 8 for more details. The purpose of the management plans is to outline the vision and goals for the protected areas and year-round protection for all wildlife and important cultural resources is the primary goal identified. The Irniurviit ACMC has written letters to CWS, NTI and KivIA (06 July 2015) formally requesting that CWS pursue the option of a status change. Article 13 of the IIBA outlines a process for changes to MBSs and NWAs. This process includes (but is not limited to) notification and consultation with NTI, KivIA, Salliqmiut and the ACMC, and will involve the

		establishment of an assessment group to consider the status change. The IIBA already anticipated potential changes to protected areas in Article 13, and as such all signatories, including the Government of Canada, have agreed to an established process and hence are aware of the implications. For these reasons, the Irniurviit ACMC will keep this section in the management plans.
Kivalliq Inuit Association Via email 22-Aug-2019	Section 5.0, pg 24: There was recognition of inclusion of IQ in research and management decisions. KIA develop a methodology for water and fish to collect IQ and Scientific information at the same time please (see attach document)	Thank you for providing this information. The Irniurviit ACMC will refer to this methodology when implementing the management plans.
Kivalliq Inuit Association Via email 22-Aug-2019	Section 6.2.6, pg 29: Minimum air traffic flight avoidance (650 m altitude and 1.5 km horizontal) is consistent with other Nunavut management plans.	The Irniurviit ACMC's intent throughout the management plans is to be consistent with existing guidance and regulation. Thank you for confirming this.
Kivalliq Inuit Association Via email 22-Aug-2019	The 2016 to 2023 IIBA for NWAs and MBSs in the Nunavut Settlement Area (Section 12.3.1) states that $20,000 compensation should be paid to the local HTO if a polar bear is killed by a CWS employee or person authorized to enter a NWA/MBS. It does not appear to be linked to IOL. There is no mention of compensation in either of the two draft management plans.	The ACMC will add a reference to the IIBA in Section 9.1. See change in red below. Compensation to the Aiviit HTO for a bear kill within the MBS will follow Section 12.3 of the *IIBA*.
Nunavut Impact Review Board Via letter 20-Nov-2019	Sec. 6.3 Monitoring and Research It states that "CWS and the Irniurviit ACMC will screen project proposals to ensure compatibility with management goals and objectives". While the ACMC as a committee make a decision on a project proposal being permitted in the MBS, this statement is a bit misleading as it may lead the	The ACMC will replace "will screen" with "will examine" to avoid confusion with screenings under *NuPPAA*. The intent of Section 6.3 (Management Approaches) is outline strategies that the ACMC and CWS will implement to manage habitat, migratory birds within the protected area to attain the goals and objectives of the management plan. As such, Nunavut's regulatory system will be described in section 7.1.7 of the

	reader to conclude that the screening by the ACMC is the only process. To make this section clearer to the reader the NIRB suggests that an additional statement about the Nunavut Planning Commission (Commission) and the NIRB's role in screening for the ecosystemic and socio-economic impacts as associated with a monitoring or research proposals within the MBS. Additionally, while further in the paragraph it directs readers to Section 7 for information on permitting, in this section there is no mention that there are other regulatory agencies besides CWS that may issue licences/permits. For example; research may also require a scientific research licence from the Nunavut Research Institute (NRI) or potentially a wildlife research permit from the Government of Nunavut, Department of Environment (GN-DOE).	management plans and other examples of licences and permits will be added to section 7.2 – see other responses below.
Nunavut Impact Review Board Via letter 20-Nov-2019	7.1.1 Authorizations by Permit This section describes authorizations of "prohibited activities under the MBSR", however, the title would indicate that this section should describe activities where authorizations are required by permit for activities that are not prohibited (e.g., non-Inuit access to the MBS). Subsection 7.1.4 Non-Inuit Activities Authorized Only by Permit also does not include the access only example.	Section 7.1.1 outlines the conditions under which any permit is possible. The ACMC will add entry and access as as example in section 7.1.4. "All activities, including entry and access, may require a permit for non-Inuit when migratory birds are present (April through October)."
Nunavut Impact Review Board Via letter 20-Nov-2019	7.1.7 CWS Permit Application Process While this section presents a figure that describes the regulatory process that includes NPC and NIRB there is no verbal text to describe regulatory process under the Nunavut Planning and Project Assessment Act (NuPPAA); additional clarity is recommended in two areas for this section. First,	The ACMC will add a paragraph in section 7.1.7 to describe Nunavut's regulatory system. "For project proposals located in the Qaqsauqtuuq MBS, the CWS is a regulatory authority under the *Nunavut Planning and Project Assessment Act*, which defines Nunavut's integrated regulatory system. All project proposals in Nunavut must be first submitted to the Nunavut Planning

	additional text should be provided to ensure the reader has complete information on the regulatory processes applying to a MBS within the territorial boundaries of Nunavut. Second, in the figure which indicates that the ACMC will review the "project comments" received from the NIRB, for clarity, the language should be updated to reflect that the NIRB in its screening decision does not provide comments but rather recommended terms and conditions to the Minister of Environment to be incorporated into the permits or licences subsequently issued.	Commission (NPC). The NPC assesses project proposals for conformity with the Keewatin Land Use Plan. The NPC sends a positive conformity determination and any recommendations to the CWS. The NPC also sends the project proposal, with its determination and any recommendations, to the Nunavut Impact Review Board (NIRB) for screening, unless the project proposal is exempt from screening pursuant to Schedule 12-1 of the *NA*. The NIRB determines whether the project has the potential to result in significant ecosystemic or socio-economic impacts and, accordingly, whether it requires further review. The NIRB submits a screening decision report to the CWS. The CWS cannot issue a permit without a positive conformity determination from the NPC and a screening decision report from the NIRB confirming that a review of the project is not required. In addition, the Irniurviit ACMC reviews CWS permit applications and provides recommendations prior to CWS issuing a permit (*IIBA s.3.3*). Figure 8 illustrates the permitting process." The ACMC will also revise Figure 8 to address NIRB's comments.
Nunavut Impact Review Board Via letter 20-Nov-2019	7.2 Other Federal and Territorial Authorizations and Permits The section notes that there are other federal or territorial authorizations and permits that may be required to undertake activities in the MBS however there are no examples of what types of activities may require permits. It may be helpful to provide these common examples: research may require a Scientific Research Licence from the Nunavut Research Institute or a Wildlife Permit from the Government of Nunavut – Department of Environment.	The ACMC will add the examples provided to section 7.2. See revised text below. "Depending on the type of activity, other federal or territorial authorizations or permits may be required to undertake an activity in the Qaqsauqtuuq MBS. For example, research may require a Scientific Research Licence from the Nunavut Research Institute or a Wildlife Permit from the Government of Nunavut – Department of Environment. Contact your regional federal and territorial permitting offices for more information."

Fisheries and Ocens Canada Via email 5-Dec-2019	p. 2 – Should Settlement Region be “Nunavut Settlement Area” as opposed to “Kivalliq”.	The ACMC will revise the heading to “Region”
Fisheries and Ocens Canada Via email 5-Dec-2019	p. 4 – for the list of species assessed by COSEWIC, Bowhead Whale is mentioned on p. 19 as occurring in waters of East Bay during summer. The Eastern Canada-West Greenland Bowhead Whale population is COSEWIC assessed Special Concern. Also, very recently (earlier this week!), Ringed Seal was COSEWIC assessed Special Concern: https://www.theglobeandmail.com/canada/article-ringed-seals-a-bellwether-species-of-the-arctic-now-deemed-at-risk/ (this doesn’t appear to be updated yet on the Species at risk public registry).	The ACMC added both species to SAR lists and counts.
Fisheries and Ocens Canada Via email 5-Dec-2019	p. 7 – Table 2. Would you want to add the Hudson Bay Complex marine bioregion? https://www.dfo-mpo.gc.ca/oceans/maps-cartes/bioregions-eng.html	The ACMC added the Marine Bioregion to Table 2.
Fisheries and Ocens Canada Via email 5-Dec-2019	p. 8 – mentions the Area of Interest. “The Southampton Island AOI currently includes the marine waters of the Qaqsauqtuuq MBS”, but note that the wording that ECCC preferred for the webpage was Marine conservation values within the Area of Interest extend into two Environment and Climate Change Canada (ECCC) Migratory Bird Sanctuaries	The ACMC notes the comment. The proposed wording for the overlap is less clear. As the management plan serves a different purpose, there is no need for the wording to be identical to the AOI webpage. The ACMC included all the minor revisions to the text. The proposed additional sentences were not kept as the purpose of the paragraph is to describe land ownership and interests within the MBSs, and not describe the MPA establishment process.

	The Department of Fisheries and Oceans (DFO) selected the nearshore waters around Southampton Island and Chesterfield Inlet as an area of interest (AOI). This marks the beginning of a Marine Protected Area (MPA) establishment process. The Kivalliq Inuit Association has endorsed the AOI and will be working closely with DFO to complete the required AOI assessment and MPA establishment steps. Marine Conservation values within the Southampton Island AOI extend into ~~currently includes~~ the marine waters of the Qaqsauqtuuq MBS, however ~~DFO will determine~~ the final boundary of a potential MPA would be decided following further assessments and extensive consultation. The Southampton Island Area of Interest Co-development Committee, which advises on steps in the process to establish a Marine Protected Area, has representation from the Irniurviit ACMC. To learn more about the Southampton Island AOI, please visit the webpage at https://www.dfo-mpo.gc.ca/oceans/aoi-si/southampton-eng.html.	
Fisheries and Ocens Canada Via email 5-Dec-2019	p. 15 – Figure 5. Can the map display the nesting areas so that we can better see the overlap with the MBS.	The overlap is currently visible in Figure 5. The ACMC cannot revise the figure without further information to understand the issue. All figures follow a map template - changes to one figure will likely result in changes to all figures.
Fisheries and Ocens Canada Via email 5-Dec-2019	p. 33 For example, progressive earlier ice break-up in Hudson Bay has forced polar bears ashore much earlier (Lunn, et al., 2016) leading to documented increases in predation of eider nests ~~predation~~ in the MBS (Iverson, et al., 2014) and other breeding	The ACMC revised this sentence.

	birds on Southampton and Coats Islands (Smith, et al., 2010).	
Fisheries and Ocens Canada Via email 5-Dec-2019	p. 38 ~~Management~~ Managers of the MBS and Wildlife Areas of Importance to Inuit will ~~include~~: (1) Minimize disturbance to wildlife and wildlife habitat, and promote the maintenance of vital, healthy wildlife populations; (2) Make all reasonable efforts, consistent with the Minister's jurisdiction under the *MBCA* to respect the cultural significance of Wildlife Areas of Importance to Inuit, taking into account any Inuit Qaujimajatuqangit documented and presented to them ~~it~~ by Inuit, the Irniurviit ACMC and other knowledgeable parties; (3) Investigate public concerns regarding the protection or management of wildlife and document the response, or refer the matter to the wildlife management agency having jurisdiction; and, (4) As appropriate, consult Inuit organizations about issues pertaining to the effective management of wildlife and wildlife habitat within the MBS.	The section was revised for more consistency with *IIBA s.12.2.3*.
Fisheries and Ocens Canada Via email 5-Dec-2019	p. 45 – replace "conversation" with "conservation". "KIA" is used versus "KivIA".	The ACMC revised these words.
Fisheries and Ocens Canada Via email 5-Dec-2019	pp. 48-49 – 7.1.2-7.1.5. It was a little unclear what the rules are for non-Inuit. It's stated that Inuit have a "right of access without permit to guide sport hunters in or across the sanctuary", so I understand that sport hunting with an Inuk guide,	The ACMC revised sections 7.1.2-7.1.5 of the text to address these concerns.

	either by laws of general application or by assignment of rights, can occur within the MBS – but the non-Inuk sport hunter requires a permit. 7.1.2 Nunavut Inuit Activities Authorized Without a Permit Nunavut Inuit have a right of access to MBSs, without fee or permit, for the following activities: □ Harvesting and activities incidental (*NA s.5.7.18* and *IIBA s.5.2*), □ Removal of up to 50 cubic yards of carving stone within the MBS and any amount from IOL within the MBS (*NA s.19.9.4* and *IIBA s.5.4*), and □ Establishment of new outpost camps (so long as consistent with Section 6.4.5, *IIBA s.5.5*, as well as the *NA*). The *IIBA* (*s.5.3*) gives Inuit a further right of access without permit to guide sport hunters in or across the Sanctuary, and to carry a firearm for self-protection or the protection of clients. It also extends the right of access without permit for harvesting and covers activities that are reasonably incidental to harvesting. Non-Inuit cannot be assigned this right of access and exemption to carry a firearm, even if ~~hunting~~ harvesting rights are assigned to that person pursuant to *NA s.5.7.35*. These activities by non-Inuit require a permit. more information on permits. 7.1.3 Nunavut Inuit Activities Authorized Only By Permit Commercial ventures or businesses require a permit to conduct commercial activities within the	

	MBS even if the commercial venture or business is Nunavut Inuit-owned. The one exception to commercial activities is guiding sports hunters or sport fishermen. A Nunavut Inuit guide does not require a permit, but the non-Inuit hunters or fishermen he or she is guiding do require a permit. Refer to Section 7.1.5 for non-Inuit. Other types of guiding (eco-tourism, canoeing, etc.) by Nunavut Inuit may require a permit. Refer to Section 7.1.7 for more information on permits. 7.1.4 Non-Inuit Activities Authorized Without a Permit There are no activities authorized without a permit for non-Inuit when migratory birds are present (April through October). In addition, any activities at any time of year that may result in the destruction of migratory bird habitat may require a permit. All activities by non-Inuit require a permit even if ~~hunting~~ harvesting rights are assigned to that person under *s.5.7.35* of the *NA*. Refer to Section 7.1.7 for more information on permits. The last sentence states that all activities by non-Inuit require a permit. Why is it necessary to also state that all activities by non-Inuit when migratory birds are present require a permit (1st sentence) and all activities by non-Inuit at any time of year that may result in the destruction of migratory bird habitat require a permit (2nd sentence). The 2nd sentence actually says these activities may require a permit, which is in contradiction to the last sentence that says all activities require a permit. The 1st	

	sentence in 7.1.5 below also says non-Inuit may require a permit. 7.1.5 Non-Inuit Activities Authorized Only By Permit Non-Inuit may require a permit to undertake any activity in MBSs in Nunavut. Non-Inuit must have a permit to carry a firearm in a MBS. Non-Inuit also require a permit to shoot and have dead migratory birds in their possession. This includes non-Inuit hunters on guided hunts for any wildlife species (even when the guide is a Nunavut ~~Inuit~~ Inuk and he or she does not require a permit; *IIBA s.5.3.1*; Section 7.1.2). Refer to Section 7.1.7 for more information on permits.	
Fisheries and Ocens Canada Via email 5-Dec-2019	p.9 – change to Minister of Environment and Climate Change	Use of Minister of Environment is intended for consistency with language in the IIBA.
Fisheries and Ocens Canada Via email 5-Dec-2019	p.13 – DFO – change to Fisheries and Oceans Canada	The ACMC changed this.
Fisheries and Ocens Canada Via email 5-Dec-2019	p.15 - How does criteria #2 apply to this MBS? Earlier in the document is stated POTENTIAL exploration/development. Is subsistence hunting considered an area-specific threat?	No, subsistence is not considered a current threat. The MBS supports a large portion of a national population of more than one migratory bird species. Birds are concentrated in the MBS during spring and summer period and therefore may be vulnerable to present or future area-specific threats.
Fisheries and Ocens Canada	p.17 – add -Eastern Canada-West Greenland Bowhead whale COSEWIC special concern Western Hudson Bay Beluga	The ACMC added Bowhead whale to the list of SAR species. Specific populations are described in Table 4.

Via email 5-Dec-2019	Atlantic Walrus - Central-Low Arctic population	
Fisheries and Ocens Canada Via email 5-Dec-2019	p.21 – Change to "Fisheries and Oceans Canada" and revise to: "however the final boundary will be determined by DFO and their partners following further assessments and extensive consultations.	The ACMC revised the text.
Fisheries and Ocens Canada Via email 5-Dec-2019	p.31 – add Ringed seal, Narwhal, walrus The waters off Southampton Island have substantial kelp forests as well. Also a variety of shellfish/marine invertebrates (shrimp, mussels, urchin, sea cucumber, etc.)	The ACMC added Ringed Seal to the SAR lists. However, the ACMC was not provided with any references and unable to locate detailed information to describe how these species use the MBS. Similarly, without a specific reference or information on which areas relative to the MBS these other marine resources are found, the ACMC is unable to determine the importance of including this information.
Fisheries and Ocens Canada Via email 5-Dec-2019	p.43 - Not a threat, but an additional management consideration should be the potential establishment of a Marine Protected Area. This could influence some of the management concerns here (i.e. shipping and other human activities)	The potential establishment of the MPA is described in section 1.3 and also considered in section 6.4.8 (e.g. establishment of conservation areas in the Kivalliq region) as a mechanism by which the ACMC can further regional conservation of migratory birds and other wildlife through its participation in the process. However, until further assessments are conducted to inform the MPA boundary discussions and its associated regulations are published, it is difficult to describe how the MPA might impact the management of the MBS or influence other management concerns.


ᑐᓂᔨᐅᔪᑦ ᐅᒃᖢᓴ ᓄᓇᒃᓗᒥ ᐅᒪᔪᓕᓗᓚᔨᖏᔪᐊᓘᓐᓃᑦ ᑲᑎᒪᔨᖏᑦ

ᐅᒧᓴᖅ

ᑐᐱᓯᕐᓂᕆᑦ: X ᐃᓱᒪᓕᐅᔨᓂᖅ:

ᐱᔨᔪᑎᖓ: ᑐᐱᓯᕐᓂᕆᑦ ᑲᑎᒪᑎᑦᑎᓂᕐᒥᑦ ᓄᓇᒃᓗᒥ ᑐᓴᓯᖏᓐᓇᒧᑦ ᑎᑎᕋᖅᑕᐅᓂᖏᓐᓄᓇᓕᒃ ᓇᓪᓕᖅᑐᖓᖏᑦᑐᒍᑦ ᑐᑦᑐᐃᓐ ᐊᑦᑕᓇᖅᑐᒍᓐᓇᖅᑐᑦ ᐅᒪᓕᔪᐃᓕᑦ ᐊᑦᑕᓇᖅᑐᒍᓐᓇᖅᑐᑦ ᐱᓕᕆᔨᒃᑯᑦ (SARA)





ᓇᓗᓇᐃᖅᓯᒪᓂᖏᑦ:

ᑐᓂᐅᖅᑲᖅᑕᐅᔨᑦ

- ᐅᖓᓯᓂᖏᓂᔪᓂᓕᕆᓚᕐᒥᖏᑦ ᓇᓴᖅᑐᖓᖏᑦᑐᒍᑦ ᑐᑦᑐᐃᓐ ᑲᑎᒪᔨᐃᑦ ᑲᓇᑕᒥ ᐱᑎᐊᖅᑐᓂ ᖁᑭᑦᑐᒍᑦ ᖃᐅᔨᓴᖅᑎᑦ, ᐊᒻᒪ ᓂᑎᖓᓂ ᓴᓯᑲᖏᓐᓂᐊᓕᐅᑉ ᐊᒻᒪ ᒪᓂᓗᑦ.
- ᓄᓇᒃᓗᒥ, ᐱᖅᓯᐅᑦᔨᔨᑦ ᐳᓚᔪᓴᓗ–ᑲᓇᓂᓇ, ᐃᖏᓕᐅᑦ, ᐱᐊᒃᓕᓪ, ᐊᕼᐃᐃᐊᖅ, ᖃᒪᓂᖅᔨᐊᖅ, ᓗᐊᖅᓱᓪᓕᑦ, ᐊᓪᓚᓪᓴᖅᖁᑦ, ᐅᖅᔪᓯᓴᓪᓗ, ᓴᓪᓕᓪᑦ ᖁᑭᖅᑕᖏᓐ, ᐊᑦᐸᑐᑎᔨᐊᖅ, ᐊᒻᒪ ᖁᑭᖅᑕᓪᓗ ᑐᑦᑐᖅᑦᑦ.

ᖃᐅᔨᓴᓂᓂᑦ ᐊᒻᒪ ᐊᑦᑕᓇᒋᑎᐅᔨᑦ

- ᑲᑎᒪᔨᕿᓪᑦ ᖃᓄᐃᓪᓚᖏᓂᓂᔪᓂᓄᑦ ᐊᑦᑕᓇᖅᑐᒍᑎᑦᑐᒍᑦ ᐅᒪᔪᐃᑦ ᑲᓇᑕᒥ (COSEWIC) ᖃᐅᔨᓴᓕᐅᖅᑐᒍᑦ ᓇᐸᖅᑐᖓᖏᑦᑐᒍᓂᐅ ᑐᑦᑐᐃᓂᑦ ᐊᑦᑕᓇᖅᑐᒍᑎᑦᑐᒍᑎᑎᑦ ᐃᓚᐃᓚᓇ 2016. ᐊᑦᑕᓇᖅᑐᒍᑎᑦᑐᒍᑦ ᐅᒪᔪᐃᑦ ᐱᑕᖃᓕᓂᔪᓐᓇᑦ ᐊᑦᑕᓇᖅᑐᒍᑎᑦᑐᓕᖅᑐᒍᑦ ᑭᓯᐊᓂ ᐊᑦᑕᓇᕐᑐᒍᑦ ᐱᓕᓚᐊᖓᔪᖅ.
- ᐊᒥᓱᓂᖅᓴᐃᑦ ᓇᓴᖅᑐᖓᖏᑦᑐᒍᓂ ᑐᑦᑐᐃᑦ ᑕᑯᔭᐅᑎᑎᓯᒪᔨᑦ ᐊᖏᔪᓂᑦ ᖃᑦᑎᐊᓕᕐᓂᐊᕐᓇᖅᑐᒍᑎᑦᑐᒍᑦ 1990-ᒥᓂᑦ. ᑲᓇᑕᒥ, ᓇᓴᖅᑐᖓᖏᑦᑐᒍᓂ ᑐᑦᑐᐃᑦ ᐊᑦᑎᓵᓚᓕᒪᓂᓯᒪᔨᑦ ᐱᑎᐊᖅᑐᒍᓂ ᖃᓄᐃᓪᓚᖏᓐᓂᐊᓕᐅᑉ 2 ᒥᓚᐊᓐ ᑐᑦᑐᐃᑦ ᐱᑎᑦᓴᓂᖏᓐᓂᓂ 1990-ᖏᓐᓂ ᑎᑭᑦᑐᒍᑦ ᖃᓄᐃᓪᓕᔭᓂᖅᑐᑦ 800,000 2016–ᖁᑎᓵᓇᒍ. ᐊᑦᑎᓵᓚᓕᒪᓂᓯᒪᔨᑦ ᒥᓴᐅᓴᑦᑕᐅᔨᑦ 56% ᐱᖓᓱᓂᓂ ᑭᖓᔪᓰᖅᓴᓂᑦ (ᐊᑯᓂᖏᓐᓂᓂ1989 ᐊᒻᒪ 2016). ᐱᑕᖃᑦᓂᖏᓐᓂᓄᑦ ᖃᐅᔨᓴᑦᓂᐅᔨᑦ ᖃᐅᔨᓴᑦᓂᐅᖃᖅᓯᒪᔨᑦ ᑲᑎᒪᔨᕿᓪᑦ

ᖃᓄᐃᓕᓐᓄᑦᓂᖅᓴᓂᐅᓄᑦ ᐊᑦᑕᓇᖅᑐᑦᑕᑦ ᐅᒪᓲᐃᑦ ᑲᓇᑕᒥ ᖃᐅᔨᓴᐃᓚᐅᕆᑎᓪᓗᓂᑦ ᑕᑯᓴᐅᑎᑦᑎᐹᒪᓲᑦ ᐃᒪᓕᕆᓂ ᑐᑦᑕᓂ, ᐱᖃᓯᐅᑎᓪᓗᒍ >ᓗᒃᑲᓴᓪ–ᑲᓇᓱᓇᖅ, ᖃᕆᓚᐅᑦ, ᐊᓴᒪ ᐱᐊᕐᓂ ᑐᑦᑐᕆᑦ.

- ᑐᑦᑐᕆᑦ ᐊᑦᑕᓇᖅᑐᒃᓯᖅᓴᐃᓲᑦ ᐊᓴᒪ ᐊᑦᑐᖅᑕᐅᓴᕿᐃᑦᑐᑎ ᐊᑦᑕᓇᑭᑎᓂᑦ ᐊᒥᓯᓂᕆᑦ ᕿᑎᑦᑖᑦᑐᓪᓗᓂᑦ.
- ᐃᓂᐃᑦ ᖃᐅᔨᓚᕐᓂᑦ ᐅᖃᖅᓯᒪᓲᑦ ᑐᑦᑐᐃᑦ ᐅᑎᖅᑕᖃᖅᑐᑦᑕᓂᖅᓯᓂᑦ ᐊᒥᓯᓂᕆᑦ ᕿᑎᑦᑖᑦᑐᓪᓗᓂᑦ ᐊᒥᓯᓂᕆᑦ ᖃᓄᐃᓕᓐᓄᑦᓂᖅᓴᓂᐅᓄᑦ ᐊᒥᓯᓂᕆᑦ ᕿᑎᑦᑖᑦᑐᓪᓗᓂᑦ ᐊᒥᓯᓂᕆᑦ ᓯᓚᐅᑦᑐ

- ᐱᓕᓀᐊᓂᓂᐊᕐᓂᔪᒍ ᐃᓱᒪᓕᑲᖏᑎᒋᒍᑦᓴᓕᒃ ᐊᒥᓱᓚᓪᓂᓄᓂ ᓄᓇᓕᓪᓂᓄᓂ ᐊᓗᒪ ᐊᐱᖅᑯᑎᖏᓕᒃ ᑭᐅᔾᔨᐅᑎᐊᕐᓂᔪᒍᑎ, ᐃᓱᒪᓕᓂᐅᒍᖅᑐᑐᔪᒍᑦ ᐅᖃᖅᑐᒥᓂᕐᒪᑦᓂᓕᒃ ᓄᓇᓕᖅᓴᒥᑦ.
- ᑲᓇᑕᒥ ᐅᒪᔨᓕᓀᓵᕗᑦ ᐃᖅᑲᓇᐃᔾᖏᑎᖏᑎᐊᓕᐅᖅᑐᑦᑐᑦ ᐅᑯᓂᖅᓯ ᓄᓇᖅᑐᑦ ᓴᓇᖅᑐᑦ ᑐᓵᖏᓚᐱᒃ ᑎᒥᖏᓐᓂ ᔪᖅᑭᑎᑎᓂᔪᓂ <ᕐᓇᐅᓴᒥᑦ ᑐᓴᖅᑎᑎᓂᔪᓂ ᐊᐃᑦᑐᖅᓭᓕᔨᓂᑦ ᐅᒪᔨᓕᓀᓵᕗᑦ ᑲᑎᑦᓴᓕᖅᓂᑦ ᐅᐊᓲᓴᓂ 2019 ᐊᖏᔪᒍᑦᓯ ᑲᑎᑦᓴᓕᖅᓂᑦ (AGM). ᐅᖏᓪᓕᐅᑎᖏᑦᑕᐅᑎᓂᖏᑦ ᖃᑎᖏᒥᐅᑦ ᐅᒪᔨᓕᓀᓵᕗᑦ ᑲᑎᑦᓴᓕᖅᓂᑦ (KRWB) ᐊᖏᔪᒍᑦᓯ ᑲᑎᑦᓴᓕᖅᓂᑦ 2019 ᓴᙵᒪᑐᖅᑲᑦ ᓂᕈᐊᖏᓂᖅᑎᑎᓂᑦ ᑐᑭᓯᐅᑦ ᐅᒪᔨᓕᓀᓵᕗᑦ ᑲᑎᑦᓴᓕᖅᓂᑦ <ᕐᓇᓱᒪᓴᓂᐅ ᓄᖅᑲᖏᑎᑐᑎᐅᑎᖏᑦᑕᐅᓂᖅ ᐊᒥᓱᓕᑲᖏᑎᐅᖅᑐᑦ ᑲᑎᑦᓴᓕᖅᓂᑦ.
- ᑕᐃᒪᓐᖏᓂ, ᑲᓇᑕᒥ ᐅᒪᔨᓕᓀᓵᕗᑦ ᐃᖅᑲᓇᐃᔾᖏᑎᖏᑎᓂᓱᒪᔪᑦ ᓄᓇᓕᖅᑐᑦ ᓴᙵᒪᑦᓂ ᐅᒪᔨᓕᓀᓵᕗᑦ ᖃᐅᔨᔪᖅᑎᓂᓂ ᐊᑐᓂ ᐊᐃᑦᑐᖅᓭᓕᔨᓂᑦ ᓇᓗᓇᐃᔭᓗᒍ ᐱᐅᓪᓗᒥᑦ ᐊᑐᖅᑕᐅᓂᐊᕐᓂᔪᒍᑦᓯ ᐱᐅᓂᖏᑦᑐᒃᑯᓂᖅᑐᐅᑦ ᐊᒥᓱᓕᑲᖏᑎᐅᖅᑐᑦ ᐅᒪᔨᓕᓀᓵᕗᑦ ᐃᓕᓴᐅᔪᖅᑐᑦ.
- ᐅᑯᓇᓂᐅᑎᓪᓗᒍ ᐅᑦᑐᐃᓚᓇ ᐊᓲᒪ ᓄᓇᐃᓚᓇ 2019, ᑕᐃᒪᐃᓕᓴᖅᓪᓗᓂ,ᑐᓂᓯᔪᓕᐅᖅᑐᒍᑦ ᑭᓴᓪᓕᖏᑦ ᐅᒪᔨᓕᓀᓵᕗᑦ ᐅᒪᔨᓕᓂᖅᑐᑦ (KWB) ᐊᓲᒪ ᖁᑭᖅᑕᓪᓕᑦ ᐅᒪᔨᓕᓀᓵᕗᑦ ᐅᒪᔨᓕᓂᖅᑐᑦ (QWB) ᐊᖏᔪᒍᑦᓯ ᐅᒪᔨᓕᓀᓵᕗᑦ ᐅᒪᔨᓕᓂᖅᑐᑦ ᓄᓇᓕᖅᑐᑦ ᓄᓇ

ᑲᑎᒪᑎᑐᓂᐅᓂᐅᓂᐊᖅᑐᑦ

- ᐃᖏᕆᐊᓗ 2020, ᒪᓕᓂᕐᑐᓄ ᓄᓇᕆᑦ ᓱᖏᒪᑲᑦ ᐅᒪᔨᓂᑦ ᖃᐅᔨᔪᖅᑎᖓ ᓄᓇᓕᒫᒧᑦ >ᓕᕆᖅᑎᓕᓄᑕ, <ᖏᓇᓯᒪᔨᔪᑦ ᐊᐅᓕᖏᓄᑕ ᐊᑐᓄ ᑎᕆᒪᐅᔨᓐᖑᓕᖅᑎᑐᑦ ᑭᖏᓕᕐᕆᑦ ᓄᓇᓕᑦ ᐃᓄᖑᑎᑦ ᑲᑎᒪᖃᖅᑐᓂᖏᓕᓄᑕ ᑐᔪᓯᕐᖏᔪᒧᑦ ᑎᑎᕋᖅᑕᐅᓂᖏᓐᓄᑦ. ᑭᓚᖅᑐᐅᔭᑦ ᓄᓇᓕᑦ ᓱᖏᒪᑲᓂᓂᑦ ᐊᓱᒪ ᓄᓇᓕᑦ ᑐᓂᖏᒪᕕᖃ ᑎᒥᖏᑦ ᑲᑎᒪᖃᑕᐅᓂᐊᖅᑐᑦ ᑦᑐᔨᓂᖏᓐ ᐊᑐᓄ ᑲᑎᒪᓂᐅᔨᓂᑦ.
- ᐃᓄᖑᑎᒧᑦ ᑲᑎᒪᖃᑎᒌᖅᑕᐅᓕᑦ ᑭᖏᓕᕐᕆᒥ ᐃᓄᐃᑦ ᑲᑐᔨᖃᑎᑲᓂᑦ ᑲᖏᑦᓯᓂᓂᒥ ᕕᖓᓇᐊᓗ 2020.
- ᐃᐳᑭ 2020, ᑲᑎᖃᑎᒋᓂᐊᖅᑕᐅᓕᑦ ᐱᐊᐅᓕᓂ ᐊᓱᒪ ᖃᒪᓂᖅᔨᐊᖅ ᑐᑦᑐᖓᑦ ᐊᐅᓪᑕᐅᓂᓂᖏᓐᓄᓕᑦ ᑲᑎᒪᔨᖓᑦ (ᑲᑎᒪᔨᐅᔨᑦ ᐃᓪᓕᑭᔨᐅᓪᓗᑎ ᑭᓕᓕᕐᕆᒥ, ᓄᓇᑦᓱᐊᖅᑯᑦ, ᖁᕕᐃᖁᔨᐅᐊᓐ ᐊᓱᒪ ᐃᓂᑐᐳ, ᐊᓱᒪᓗᑦᑕᐅᖅᑯ ᓄᓇᓕᑦᒥ ᐅᒪᔨᓪᓚᕕᖅᑯᓂᑦ ᑲᑎᒪᔨᖓᑦᑕ) ᑲᑎᒪᑎᑎᑲᓂᓂᖅᓗᑦ.
- ᐅᑭᐅᖅ/ᐅᐱᖅᓯᖅ 2020 (ᓇᓗᓇᐃᖅᑕᐅᑭᐊᓪ ᐅᓪᓗᖓ) ᑲᑎᒪᑎᑐᓂᐅᔨᑦ ᖃᑎᖅᑎᐅᓂᑦ, ᐱᑕᖃᓚᐊᖃᑦᑕᓂ.

ᑭᖓᔪᓪᓕᖅᑎ ᐱᓕᓚᐊᖓᔪᓂᐊᖅᑐᑦ:

- ᐱᓴᕆᖅᓗᔪ ᑲᑎᒪᑎᑐᑲᓂᓂᐅᔨᖅ ᑭᓴᓪᓕᖅᑎ (ᐃᖏᕆᐊᓗ 2020) ᐊᓱᒪ ᖃᑎᖅᑎᐅᓂᑦ (ᐅᑭᐅᖅ/ᐅᐱᖅᓯᖅ 2020, ᓇᓗᓇᐃᖅᑕᐅᑭᐊᓪ ᐅᓪᓗᖓ), ᐱᑕᖃᓚᐊᖃᑦᑕᓂ.
- ᑐᓴᔪᕐᕋᐅᑦ ᐃᔪᒪᓪᓕᐅᓂᖅᓗᒍᑦ ᓄᓴᐅᓗᑎ ᐅᒪᔨᓪᓚᕕᖅᑯᓂᑦ ᑲᑎᒪᔨᖓᑦᓂᓂ ᑐᓴᔪᕐᕋᖅᑕᐅᓕᔪᒧᑦ ᑎᑎᕋᖅᑕᐅᓂᐊᖅᑐᑦ (ᓇᓗᓇᐃᖅᑕᐅᑭᐊᓪ ᐅᓪᓗᖓ, ᐃᓚᒪᖃᖅ ᔫᓂ ᐊᐅᓚᑦ).

Appendix A

Summary of feedback received from consultations in Nunavut on the proposed listing of Barren-ground Caribou as Threatened under the federal *Species at Risk Act* (SARA)

February 2018 – January 2020

Qikiqtaaluk Wildlife Board AGM
November 17, 2019

- We presented an overview of what SARA is, discussed the health of the Baffin herd and threats to its recovery. We discussed the implications of SARA-listing, the anticipated timeline for a decision on the listing proposal and what the steps would be if caribou is listed. Following the presentation we had a discussion and asked what the Board suggested is needed to support the Baffin herd to ensure it is healthy for future generations; and whether the Board agrees, disagrees, or has concerns with the SARA listing proposal.
- We presented the slides in both English and Inuktitut. In addition, we provided each member with printed copies of the presentation in both English and Inuktitut, an Inuktitut version of the COSEWIC report, an English summary of the COSEWIC report, a glossary of terms in the COSEWIC report in both languages, and questionnaire sheets in both languages.
- One member noted that the caribou on Baffin Island cross back and forth from the mainland so when it looks like there is a dip in population numbers it's only because they have moved to the mainland.
- There was a question about what makes herds distinct and whether the assessment considered that individuals move around between herds.
- A question from the Kitikmeot Wildlife Board representative asking when we would come to consult with the Kitikmeot.
- A few members noted concern that the knowledge of Inuit elders is being lost as elders pass away and that that knowledge is the best resource for information on caribou cycles, habits and general biology.
- The QWB wildlife advisor noted that the Baffin caribou herd has historically been this low during each of their cycles and that it's not unusual. He noted that it is wrong to blame harvesters for declines.
- One member asked if we personally thought the caribou herds are declining.
- One member noted that there is Inuit knowledge on caribou population cycles and that they are related to the population cycles of lemmings.

- One member asked if there is a listing, if a management plan would address development. He also asked if things got better, at which point could caribou be removed from SARA?
- One member asked about how we would address the threat of climate change to the recovery of BGCA.
- One member noted that the Baffin herd is already managed by the GN and there is already a draft management plan.
- A few members noted that a new survey of Baffin Island is desperately needed. The surveys that were used to establish the current TAH were flawed and they want to see a new one completed with more Inuit involvement.
- One member asked if we follow western science or Inuit knowledge in our processes and noted that IQ indicates that caribou will recover.
- We did not receive any requests for further consultation on the proposed listing from any of the QWB representatives or audience members.

Kivalliq Wildlife Board AGM
October 23, 2019

- We presented an overview on the health of the Kivalliq herds, the main threats to caribou in that region, the implications of SARA-listing, and our proposed plan to revisit communities in the Kivalliq in early 2020.
- Representatives from the NWMB, BQCMB and GN were supportive of our approach to the topic and the material that we presented.
- One Board member asked how HTOs can receive more support if caribou is listed
- One asked how a 'plan' is implemented and if it's the GN management plan that gets adopted
- Another member asked how SARA provisions would be enforceable.
- No members objected to our plan to do further consultations in the region.

Cambridge Bay
HTO and public meetings, February 26, 2018

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.
- HTO members expressed strong concerns about the lack of Inuit participation and traditional knowledge in the assessment of Barren-ground caribou and the decision-making process. They also want to be involved in the drafting of the recovery strategy.

- They would appreciate a herd-by-herd assessment as they are in the opinion that their herd is doing fine.
- Increased predation: would like to see an incentive for hunters to harvest wolves.
- They are worried about their income (subsistence and income hunting).
- Kugluktuk [Bluenose-East and Bathurst herds]:
- HTO members expressed strong concerns about the lack of Inuit participation and traditional knowledge in the assessment of Barren-ground caribou and the decision-making process.
- They would appreciate a herd-by-herd assessment, and worry about the flexibility in prohibitions and how it will be applied to local management.
- They noticed an increase in wolf and wolverine populations.

Gjoa Haven
HTO meeting, March 1, 2018

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.
- HTO members expressed strong concerns about the lack of Inuit participation and traditional knowledge in the assessment of Barren-ground caribou and the decision-making process.
- They are worried about the Caribou declining, the species is critical for their way of life and food security. They are interested in collaborating to help the species recover.
- They mentioned various reasons why they see less Caribou than before :
    - Increased predation (wolves, wolverines and grizzlies)
    - They are seeing a lot more Muskox than before
    - Climate Change: Caribou are vulnerable to migration on thin ice.
- They would like to improve youth education regarding hunting practices.

Kugaaruk
HTO meeting, February 27, 2018

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.
- HTO members expressed strong concerns about the lack of Inuit participation and traditional knowledge in the assessment of Barren-ground caribou and the decision-making process.

- They would appreciate a herd-by-herd assessment, they never heard of any studies done in their region.
- They are worried about harvest restrictions and food security.
- They mentioned various reasons why they see less Caribou than before :
    - Increased predation (wolves)
    - They are seeing a lot more Muskox than before
    - Mining (chemicals, plane/helicopters flying low)
    - Climate Change : They noticed a lot of rain on snow event making the foraging harder for the species.
    - Natural cycle of caribou population
    - Diseases

Rankin Inlet
HTO meeting, March 5, 2018

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.
- HTO members expressed strong concerns about the lack of Inuit participation and traditional knowledge in the assessment of Barren-ground caribou and the decision-making process. They would like to see an Inuit representative on COSEWIC committee.
- They would appreciate a herd-by-herd assessment, they never heard of any studies done in their region.
- Increased predation: They are seeing more wolves and grizzlies than before.
- They want to be involved in drafting the recovery strategy. Management plans already in place in some regions should be recognized.
- Some agreed to the listing of the barren-ground caribou.

Whale Cove
HTO meeting, March 6, 2018

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.
- HTO members expressed some worries about their harvest rights.
- They mentioned various reasons why they see less Caribou than before :
    - Increased predation (eagles, wolverines)
    - They are seeing a lot more Muskox than before
    - Natural cycle of caribou population

- Diseases: They noticed swollen hooves (brucellosis)
- Mining

Arviat
HTO meeting, March 7, 2018

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.
- HTO members would like to see a herd-by-herd assessment as they think their herd is stable.
- They mentioned various reasons why they see less Caribou than before :
  - Sport hunting happening south of Aviat
  - Migration routes have changed
  - Increased predation (wolves, grizzlies, wolverines)
  - Moose habitat range has recently extended to their region
  - Hunting ways have changed (bullet type, transportation)

Baker Lake
HTO meeting, March 8, 2018

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.
- HTO members expressed strong concerns about the lack of Inuit participation and traditional knowledge in the assessment of Barren-ground caribou and the decision-making process. They would like to see an Inuit representative on COSEWIC committee.
- They are worried about their harvest rights and food security.
- They haven't seen the Qamanirjuaq herd for a long time, and mentioned potential causes of the caribou decline, i.e. changing migration routes, natural cycle of the caribou population, forest fires, and increased predation.
- They want to see the recovery strategy being elaborated cooperatively with all territories and provinces sharing the herds. Attention to the migration routes and its protection should be emphasized.

Chesterfield Inlet
HTO meeting, March 9, 2018

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.
- HTO members expressed strong concerns about the lack of Inuit participation and traditional knowledge in the assessment of Barren-ground caribou and the decision-making process.
- The caribou have other predators, like wolves, grizzlies and wolverines that are also responsible for their decrease.
- They are worried about their harvest rights.
- Some noticed the caribou have recently started to increase in the area.

Igloolik
Public meeting, September 25, 2018

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.
- Community and HTO members expressed strong concerns about the lack of Inuit participation and traditional knowledge in the assessment of Barren-ground caribou and the decision-making process.
- They are worried about their harvest rights and food security; they see this proposed listing as an additional hardship done to their community and traditional way of life.
- Most of them believed the caribou population is going through a natural cycle and will eventually come back on its own. Some people mentioned the population numbers were not trustworthy.
- They want more responsibility in regards to managing their own herd, instead of having outsiders getting involved. They also didn't appreciate all the herds being merged together for the listing assessment.

Hall Beach
Public and HTO meetings, September 26, 2018

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.

- Community and HTO members expressed strong concerns about the lack of Inuit participation and traditional knowledge in the assessment of Barren-ground caribou and the decision-making process.
- Most of them believed the caribou population is going through a natural cycle and will eventually come back on its own.
- The caribou have other predators, like wolves that are also responsible for their decrease.
- They are worried about their harvest rights and food security. Some of them thought animals should not be surveyed and didn't like outsiders coming to interfere with their wildlife management.

Naujaat
Public meeting, September 27, 2018

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.
- Community members wanted to see local management of the herd. They didn't like having their caribou lumped in with other herds across Canada as part of the assessment of Barren-ground caribou or in future recovery plans. Some of them mentioned the survey methodology was not trustworthy.
- Most of them believed the caribou population is going through a natural cycle and will eventually come back on its own.
- They are worried about their harvest rights being affected after the listing.

Arctic Bay
Public meeting, October 16, 2018

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.
- There were strong concerns about the lack of Inuit participation in the assessment of the caribou and decision-making regarding the wildlife management. They would like to see more traditional knowledge involved throughout the process.
- Community members mentioned the cause of decline is most likely due to wolf predation and natural cycle of the caribou population. They strongly believe the caribou will come back on their own.
- Community members were also worried about their harvest right and food security.

Pond Inlet
Public meeting, October 17, 2018

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.
- Community and HTO members expressed strong concerns about the lack of Inuit participation in the assessment of Barren-ground caribou and decision-making. They would like to see more local management and are concerned about their harvest rights.
- There were some doubts regarding the numbers of caribou and the survey methodology.
- Most of them believed the caribou population is going through a natural cycle and will eventually come back on its own.
- One person asked for a further investigation on the actual causes of decline of the Baffin herd

Clyde River
Public and HTO meetings, October 18, 2018

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.
- Community and HTO members expressed strong concerns about the lack of Inuit participation in the assessment of Barren-ground caribou and decision-making. They would like to see more local management and are concerned about their harvest rights.
- One member mentioned there was a generational conflict where the younger hunters want to try new wildlife management methods, but the elders disagree. The Inuit's profound respect for elders makes the younger generation hesitant to talk about it.
- Community members also mentioned potential causes of the caribou decline, i.e. the natural cycle of the caribou population, the predation by wolves, female-male ratio allowed for hunting, and the new technology (snowmobiles scare caribou away).

Iqaluit
HTO meeting, October 22, 2018

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.
- Community and HTO members expressed strong concerns about the lack of Inuit participation and traditional knowledge in the assessment of Barren-ground caribou and the decision-making process.

- They would also like to be involved in the scientific research. The survey methodology is not clear to them, and they believe a herd-by-herd assessment would be much more relevant.
- They believe the caribou populations are going through natural cycle and will eventually come back up on their own.
- Many were worried about their harvest rights and would like to see investigation on other threats like predation, industry and impact of research.

Qikiqtarjuaq
Public and HTO meetings, October 23-24, 2018

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.
- HTO board members expressed their concerns about the harvest restrictions and mentioned they want to collaborate to see the caribou population increase again.
- HTO and community members mentioned the helicopters from mining companies were an important threat to the caribou (scaring them away).
- The wolf population is also an important threat to consider.
- Some of them strongly believed the caribou will come back on their own. They migrate long distance and undergo natural cycles of population density.

Coral Harbour
Public and HTO meetings, January 22, 2019

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.
- Community members were concerned about the mining activity and identified industry as one of the main threats to caribou.
- Community members identified climate change as an on-going threat to caribou.
- Several community members believe caribou undergo natural cycles of population density. When populations are too abundant the numbers drop, but increase again when vegetation grows back.
- Community members expressed interest in knowing current local caribou numbers, particularly on Coats Island.
- The lag time between caribou surveys and results is too long. The community members would like to be informed of the health of the herds more quickly, so they can better manage their harvest.
- Community members expressed interest in knowing how caribou herds across Canada were doing, and how they were being managed.

Cape Dorset
Public and HTO meetings, January 23, 2019

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.
- Community members questioned the accuracy of the range of Barren-ground caribou herds shown on the maps
- Community members weren't sure that caribou populations will cycle up and down as they have in the past because of all the things that have changes
- Community members were concerned about the affect of the mines on caribou and want to find ways to protect the caribou from mining. Critical habitat is one way to protect habitat for caribou.
- Community members were concerned about predation from wolves, and suggested wolf control
- Community members were concerned about harassment of caribou by helicopters and airplanes
- Some community members though that caribou are not threatened, and have just moved to another area
- Community members were concerned that caribou are not surveyed often enough
- Community members spoke about the importance of using Inuit Qaujimajungit
- Community members were concerned about the possible impact that listing caribou might have on harvesting

Kimmirut
Public and HTO meetings, January 24, 2019

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.
- HTO members expressed strong concern regarding the listing of all Barren-ground caribou herds as one. They believe South Baffin and North Baffin populations should be considered separately.
- Several HTO and community members do not believe the South Baffin population is in decline.
- Several HTO and community members believe caribou undergo natural cycles of population density. When populations are too abundant the numbers drop, but increase again when vegetation grows back.
- Some community members do believe caribou populations on South Baffin Island are in decline, support the listing, and believe more survey efforts are required.

- Community members expressed concern related to methods used to survey caribou (e.g. helicopter use), and suggest using less intrusive methods.
- Community members identified parasites and wolves as threats, and expressed an interest in better understanding how parasites (e.g. ticks) have arrived and how they impact the caribou.
- HTO and community members expressed concern about their harvest rights and the lack of Inuit participation in the listing process.

Taloyoak
Public and HTO meetings, February 26, 2019

- ECCC presented a PowerPoint describing the federal species at risk process, why barren-ground caribou have been assessed as threatened, and what it would mean if barren-ground caribou are listed as threatened under the federal species at risk act.
- HTO members did not agree with the delineation of the herds. They advised that caribou from various herds (Peary and Ahiak) interbreed and mix with Boothia Peninsula caribou, and that the range of the Boothia Peninsula herd actually extends further south than Taloyoak. Interbreeding may have caused the original Boothia Peninsula caribou to move south, but all caribou in the area are considered as one herd.
- HTO and community members identified wolves and muskox as threats to the Boothia Peninsula herd, and expressed an interest in establishing an incentivised culling program for wolves.
- HTO members were interested in knowing more about the various threats and their impact on caribou herds.
- HTO and community members expressed concern about their harvest rights.
- HTO and community members found the information presented was very broad, and would like to see more concrete survey data showing caribou population estimates in their area before deciding to support the listing. They do not want to rush any decisions.
- Some community members do not believe their local herds are at risk or are declining, while others do believe they are a threatened species.
- Community members strongly support conservation and believe maintaining clean hunting grounds (free of carcasses) and minimizing waste is very important. They believe their people need to take more accountability and think a carcass cleaning program could be beneficial.
- HTO and community members believe growing populations are increasing the harvest pressure on caribou unlike before. For this reason, it is important to work collaboratively, and plan for the future to ensure caribou are conserved.

Kitikmeot Regional Wildlife Board
ᖁᑎᖅᑕᐅᑦ ᓄᓇᓕᖕᓂᓄᑦ ᐅᒪᔨᓕᓗᐊᖅᑯᑦ ᑲᑎᒪᔨᖏᑦ
Kitikmeot Nunaliit Avikhimaniani Angutikhaligiyit Katimayit

Kitikmeot
ᖁᑎᖅᑕᐅᑦ
Kitikmeot
Cambridge Bay
ᐃᖅᑲᓗᑦᑎᐊᖅ
Ikaluktutiak
Gjoa Haven
ᐅᖅᓯᖅᑐᖅ
Uqhuqtuuq
Kugaaruk
ᑰᒐᖅᓴᒃ
Kugaaruk
Kugluktuk
ᖁᑭᑲᖅᑐᖅ
Kugluktuk
Taloyoak
ᑕᓗᔨᐊᖅ
Taloyoak

SUBMISSION TO THE NUNAVUT WILDLIFE MANAGEMENT BOARD
REGULAR MEETING 001-2020
March 11, 2020

for

INFORMATION: **DECISION:** X

ISSUE: Kitikmeot Regional Wildlife Board request for 5 additional Grizzly Bear Sport Hunt Tags

BACKGROUND:

The last study on population survey by the Government of Nunavut was in 2007.

Inuit in Kugluktuk have observed an increase in the grizzly bear population. There are many observations of sows with twins and triplets and increasingly more each year.

Grizzly bears are always present in caribou calving grounds and close to the community.

In 2017, the Kugluktuk HTO had originally requested an additional 5 grizzly bear tags.

An increase in grizzly bear tags would bring more income into the community; it would assist in bringing the population back down and it would assist the caribou herds, as there would be less predation on the calving grounds.

Historically, the Kitikmeot Regional Wildlife Board had 10 grizzly bear tags and they were allocated as: Kugluktuk – 5; Omingmaktok – 2; Burnside – 3.

In June 2019, the Nunavut Wildlife Management Board approved 5 additional tags, for a total of 15 for the Kitikmeot Regional Wildlife Board. It is respected by the Kugluktuk Angoniatit Association that the Ekaluktutiak Hunters and Trappers Organization have put in the request for 5 additional tags. The Kugluktuk HTO does not want to take any tags away from other HTOs. These tags were allocated as follows: Kugluktuk – 5; Cambridge Bay – 5; Omingmaktok – 2; Burnside – 3.

RECOMMENDATION:

Kitikmeot Regional Wildlife Board
ᖁᑎᖅᑕᐅᑦ ᓄᓇᓕᖕᓂᓄᑦ ᐅᒪᔨᓕᓗᕆᔨᐊᒃᑯᑦ ᑲᑎᒪᔨᖕᓂᑦ
Kitikmeot Nunaliit Avikhimaniani Angutikhaligiyit Katimayit

Kitikmeot
ᖁᑎᖅᑕᐅᑦ
Kitikmeot

Cambridge Bay
ᐃᖅᑲᓚᖅᑐᐊᖅ
Ikaluktutiak

Gjoa Haven
ᐅᖅᓱᖅᑐᖅ
Uqhuqtuuq

Kugaaruk
ᑰᒍᖅ
Kugaaruk

Kugluktuk
ᖁᑦᓴᖅᑐᖅ
Kugluktuk

Taloyoak
ᑕᓗᕕᐊᖅ
Taloyoak

During our conference call on October 18th, 2019, the Kitikmeot Regional Wildlife Board heard concerns from the Kugluktuk Angoniatit Association's observation on the increase in grizzly bears in the Kugluktuk area.

We have also heard from KAA on their request for an increase in grizzly bear sport hunt tags that did not go through. We would like to show our support and approval for their request for five (5) additional tags to be used for sport hunts.

Please do not hesitate to contact our office for questions to this request.

Yours Truly,

Bobby Klengenberg

KRWB Chair

Kitikmeot Regional Wildlife Board
ᖃᑎᖏᑦᑕᐅᑦ ᓄᓇᓕᖕᓂᖅᓄᑦ ᐅᒪᔨᓕᓇᕐᔨᕐᕕᒃᑯᑦ ᑲᑎᒪᔨᖏᑦ
Kitikmeot Nunaliit Avikhimaniani Angutikhaligiyit Katimayit

Kitikmeot
ᖃᑎᖏᑦᑕᐅᑦ
Kitikmeot

Cambridge Bay
ᐃᖃᓗᐊᑎᒃ
Ikaluktutiak

Gjoa Haven
ᐅᖅᔪᓂᑐᖅ
Uqhuqtuuq

Kugaaruk
ᑯᒡᖃᕐᒃ
Kugaaruk

Kugluktuk
ᖁᑦᓴᖅᑐᖅ
Kugluktuk

Taloyoak
ᑕᓗᕐᕕᖅ
Taloyoak

ᑐᓂᔪᓕᕐᓂᖅ ᓄᓇᕗᑦ ᐅᒪᔨᓪᓚᕆᔨᕐᕕᐊᑯᓄᑦ ᑲᑎᒪᔨᖏᓐᓂᓄᑦ ᑲᑎᒪᓂᐅᕐᓂᖅ ᓴᖅᑭᑎᓂᖅ **001-2020**

ᐃᑦ 11, 2020

ᑐᖓᔾᔭᐃᓂᖅ: ᐊᙵ

ᐊᑦᑏᖅᓯᒪᔨᖅ: ᖃᑎᖏᑦᑕᐅᑦ ᐊᕐᕌᒍᑎᔪᓄᑦ ᐅᒪᔨᓪᓚᕆᔨᕐᕕᐊᑯᓄᑦ ᑲᑎᒪᔨᖏᓐᓂᒃ ᑐᓴᕆᖏᖏᓐᓂᒃ ᑕᓪᓕᓂᓪ ᐃᓕᓴᕐᕕᐊᓂᖅ ᐊᓯᖏᓐᓂᒃ ᐊᑐᐊᓕᖏᓐᓂᒃ ᐊᑐᖏᓐᓂᒃ ᐊᑐᐊᓕᖏᓐᓂᒃ ᐊᑐᐊᓕᖏᓐᓂᒃ ᐊᑐᐊᓕ

Kitikmeot Regional Wildlife Board
ᖃᑎᖅᑕᐅᓂ ᓄᓇᓕᖕᓂᖅᓄᓂ ᐅᒪᔨᓕᕆᔨᖁᔭᐅᔪᓂ ᑲᑎᒪᔨᖏᑦ
Kitikmeot Nunaliit Avikhimaniani Angutikhaligiyit Katimayit

Kitikmeot
ᖃᑎᖅᑕᐅᓂ
Cambridge Bay
ᐃᖅᑲᓇᐃᑎᒍᖅ
Ikaluktutiak
Gjoa Haven
ᐅᖅᓱᖅᑐᖅ
Uqhuqtuuq
Kugaaruk
ᑰᒐᖏᒃ
Kugluktuk
ᖁᔭᓕᖅᑐᖅ
Taloyoak
ᑕᓗᖏᐊᖅ

15 ᑐᖏᖕᒪᖏᓂᓄᓂ ᖃᑎᖏᑕᐅᓂ ᐊᕐᕋᑐᖏᓱᒪᖏᓂᓄᓂ ᐅᒪᔨᓪᓚᖏᓂᖅ ᑲᑎᒪᔨᖅᑎᓂᓄᓂ. ᖁᔭᓐᖏᑐᖏᑦ ᐊᖓᔪᓂᓂᐊᑎᒍᒃ ᑲᑎᒪᔨᖅᑎᓂᑕ ᐅᒪᔨᓪᓚᖏᓱᐊᓪᓚᒍᖅᑎᓂᓄᓂ ᑐᓴᕐᖏᕐᕋᕐᓂᐅᒪᔨᖅᑎᓂᓄᓂ ᐊᑐᖏᑕᐅᓂᐊᖏᓂᖅᑎᓂᓄᓂ ᐊᒪᓂ ᑐᓴᕐᖏᑕᐅᓕᐅᖏᑐᑐᓂᓄᓂ ᐃᖅᑲᓇᐃᑎᑎᐊᖏᑦ ᐅᒪᔨᓪᓚᖏᓱᐊᓪᓚᒍᖅᑎᓂᓄᓂ ᑕᓪᓕᓂᓄᓂ ᑐᓴᓂᕐᕕᐅᓕᖏᑐᑦ ᐊᑐᖏᑕᐅᓂᐊᖏᓂᖅᑎᓂᓄᓂ. ᑕᒪᔨᐊᑦ ᓴᕕᖅᑕᖅᓂᓄᓂ ᐊᕕᖏᑐᖅᑕᐅᓕᐅᖏᑐᑐᓂ ᐅᔨᐊᑦ ᐅᒪᔨᓪᓚᖏᓱᐊᓪᓚᒍᖅ ᒪᓕᓇᓂᓄ: ᖁᔭᓐᖏᑐᖅ – 5; ᐃᖅᑲᓇᐃᑎᑎᐊᖅ – 5; ᐅᒥᓚᓇᐃᖅ – 2; ᖃᐅᓚᐅᖅ – 3.

ᐊᑐᖃᒥᒃᓴᓂᓄᓂ:

ᐅᖅᑲᐃᑎᔨᑦ ᑲᑎᒪᑐᓄᑕ ᐅᓇᐃᓄᑦ 18th, 2019, ᖃᑎᖏᑕᐅᓂ ᐊᕐᕋᑐᖏᓱᒪᖏᓂᓄᓂ ᐅᒪᔨᓪᓚᖏᓱᐊᓪᓚᒍᖅᑎᓂᓄᓂ ᑲᑎᒪᔨᖅᑎᓂᓄᓂ ᑐᓴᕐᖏᕐᕋᕐᓂᐅᒪᔨᖅᑎᓂᓄᓂ ᖁᔭᓐᖏᑐᖏᑦ ᐅᓂᒪᓕᑎᒃᓴᐅᔪᓂᓄᓂ ᑲᑎᒪᔨᖅᑎᓂᓄᓂ ᐅᒪᔨᓪᓚᖏᓱᐊᓪᓚᒍᖅᑎᓂᓄᓂ ᖃᑎᖏᑕᐅᓂᐊᖏᓂᖅᑎᓂᓄᓂ ᐊᒥᓱᓕᕐᕕᐊᓂᖅᑎᓂᓄᓂ ᐊᑐᖏᑕᐅᓂᐊᖏᓂᖅᑎᓂᓄᓂ ᑕᑎᖏᑕᐅᓱᒪᖏᓂᖅᑎᓂᓄᓂ ᐊᑐᖏᑕᐅᓂᐊᖏᓂᖅᑎᓂᓄᓂ ᐊᕐᕌᒍᓂᐊᓕᖅᑎᓂᓄᓂ ᐊᑐᖏᑕᐅᓂᐊᖏᓂ